

In vitro anti-diabetic activity of stingless bee honey from different botanical origins

Ali, H., *Abu Bakar, M.F., Majid, M., Muhammad, N. and Lim, S.Y.

Faculty of Applied Sciences and Technology, Universiti Tun Hussein Onn Malaysia (UTHM), Pagoh Educational Hub, 84600 Pagoh, Johor, Malaysia

Article history:

Received: 11 December 2019

Received in revised form: 10 March 2020

Accepted: 12 March 2020

Available Online: 7 April 2020

Keywords:

Stingless bee honey,

Phenolic,

Flavonoid,

α -amylase inhibition,

α -glucosidase inhibition

DOI:

[https://doi.org/10.26656/fr.2017.4\(5\).411](https://doi.org/10.26656/fr.2017.4(5).411)

Abstract

The number of patients with diabetes mellitus is increasing at an alarming rate and this situation has triggered the interest of searching natural product as an alternative treatment. Stingless bee honey containing a diverse array of bioactive compounds is commonly utilized as a functional food and also found to possess various therapeutic effects including anti-diabetic through α -amylase and α -glucosidase inhibition. However, the composition level of bioactive compounds varying of geographical origins and botanical sources of honey leads to different enzyme inhibition abilities. Therefore, this study explored the total phenolic, total flavonoid, α -amylase and α -glucosidase inhibition activities of stingless bee honey from various botanical origins. In this study, stingless bee honey was collected from 6 different botanical origins namely, acacia, coconut, mangrove, starfruit, multifruit and multiflower plant. Honey from tualang tree was used as a positive control. Phenolic and flavonoid contents as well as α -amylase and α -glucosidase inhibition activities of honey were studied spectrophotometrically. Stingless bee honey from mangrove was found to have the highest phenolic content (141.74 \pm 0.03 mg GAE/100 g). The honey collected from coconut origin showed the highest flavonoid content with the value of 51.33 \pm 0.02 mg RE/100 g and also achieved the highest percentage inhibition against α -glucosidase (68.33% at 100 μ g/mL). Furthermore, tualang honey and honey samples from mangrove, coconut and Acacia tree were found to have strong α -amylase inhibition abilities as their inhibition percentages were more than 70.00% at 100 μ g/mL. This study showed that the presence of flavonoid and phenolic compounds in honey from different botanical origins yielded different degree of α -amylase and α -glucosidase inhibition and also recommended the uses of stingless bee honey in diabetes treatment.

1. Introduction

Diabetes mellitus is one of metabolic diseases characterized by chronic hyperglycemia that triggers the carbohydrate, fat and protein metabolism disorders in the human body, leading to long-term damage and dysfunction of various organs (Alberti and Zimmet, 1998; Rahmawati *et al.*, 2019). In 2017, approximately 451 million of the population were affected by this disease and it had brought up to 5 million deaths in the worldwide population in the same year (Rutebemberwa *et al.*, 2013). In addition, the number of patients is predicted to increase by years up to 693 million in 2045 (Cho *et al.*, 2018). Failure in recognizing diabetes treatment as priority became the major factor that leads to morbidity and even increases the rate of mortality globally (Shaw *et al.*, 2010). However, the high cost of disease treatment has led to a financial burden to patients

with diabetes, especially those in developing countries. Therefore, there is an urge to encourage people to search for an alternative way such as lifestyle modification including a change of dietary intake to enhance the treatment of patients suffering from diabetes.

Honey, a sugary secretion extracted from honeycombs comprises more than 200 constituents including the main substances such as fructose and glucose (Eteraf-Oskouei and Najafi, 2013). It is known as one of the natural products that have potential health benefits to human and among a variety of honey, stingless bee honey is considered as one of the food substances that used for natural remedies. Basically, it is produced by the stingless bees from the family of *Meliponini*, which made up of various types of genera such as *Melipona*, *Trigona*, and *Heterotrigona* whereas among the stingless bee species, *Geniotrigona thoracica*,

*Corresponding author.

Email: fadzelly@uthm.edu.my

Heterotrigona itama, *Lepidotrigona terminata* and also *Tetragonula laeviceps* are reared commercially in Malaysia (Mohd Rafie *et al.*, 2018). Stingless bee honey has been reported to contain rich phytochemical contents and possess anti-inflammatory, antimicrobial, antioxidant, anticancer and antiseptic properties as well as to boost the immune system (Eteraf-Oskouei and Najafi, 2013; Abd Jalil *et al.*, 2017; Akhir *et al.*, 2017). Furthermore, it displays specific anti-diabetic role in the inhibition of α -amylase and α -glucosidase enzyme activity by controlling the glucose content in the human body (Campone *et al.*, 2014).

However, the nutritional content, physicochemical properties and variety of bioactive compounds of honey might be affected by several factors such as geographical origins, their botanical sources and environmental factor (Majid *et al.*, 2019; Lim *et al.*, 2019). Thus, this study was aimed to determine phenolic and flavonoid contents as well as to evaluate the α -amylase and α -glucosidase inhibition activities of honey collected from the stingless bee, *Heterotrigona itama* in different botanical origins.

2. Materials and methods

2.1 Sample preparation

The sample chosen in this study was stingless bee honey from *H. itama* sp. The stingless bee honey samples were collected from six botanical origins including Acacia tree, mangrove tree, coconut tree, multiflower plant, multifruit plant and starfruit tree while honey from tualang tree was used as a positive control. Then, the honey samples were stored at 4°C for further analysis.

2.2 Determination of total phenolic content

Folin-Ciocalteu method was conducted to determine the total phenolic content of honey samples (Yap *et al.*, 2015; Istasse *et al.*, 2016). The absorbance was measured spectrophotometrically at 725 nm. A calibration curve ($y = 0.009x - 0.0217$; $R^2 = 0.9914$) was used to calculate the total phenolic content. The experiment was carried out in triplicate and the results were expressed as mg of Gallic acid equivalent (GAE) per 100 g of honey.

2.3 Determination of total flavonoid content

Total flavonoid content was determined based on the method as stated in the study of Islam *et al.* (2017) with some modification. The absorbance was measured spectrophotometrically at 510 nm. A calibration curve ($y = 0.0021x + 0.004$; $R^2 = 0.9869$) was used to calculate the total flavonoid content and the result was expressed as mg of Rutin equivalents (RE) per 100 g of honey.

2.4 Determination of α -amylase inhibition assay

α -amylase inhibition activity was carried out based on the method of Inoue *et al.* (2007) with some modification. The reaction mixture containing 500 μ L of appropriate dilutions (20-100 μ g/mL) of the honey sample, 500 μ L of 0.02 M sodium phosphate buffer (pH 6.9) and porcine pancreatic α -amylase (0.5 mg/mL) was incubated at 25°C for 10 mins. Then, 500 μ L of 1% starch solution in 0.02 M sodium phosphate buffer was added to the mixture and the mixture was incubated at 25°C for another 10 mins. After incubation, dinitro salicylic acid (DNSA) solution was added to the mixture. The reaction was stopped by incubating the mixture in boiling water bath for 5 mins and then left to cool at room temperature. A total of 10 mL of distilled water was added to dilute the reaction mixture. The absorbance was measured spectrophotometrically at 540 nm. The reference sample was tested in the same way with the exception of the test sample and the result was expressed as percentage inhibition.

2.5 Determination of α -glucosidase inhibition assay

α -glucosidase inhibitory activity was conducted according to the study of Apostolidis *et al.* (2007) with some modification. 100 μ L of α -glucosidase (0.5 mg/mL) in 0.1 M phosphate buffer (pH 6.0) solution was added to 150 μ L of appropriate dilutions (20-100 μ g/mL) of extracts and incubated at 25°C for 10 mins. Then, the mixture was incubated at 25°C for another 5 mins after the addition of 100 μ L of 5 mM p-nitrophenyl- α -D-glucopyranoside in 0.1 M phosphate buffer (pH 6.9) solution. After incubation, the absorbance was measured spectrophotometrically at 405 nm. The reference sample was tested in the same way with the exception of the test sample and the result was expressed as percentage inhibition.

2.6 Statistical analysis of the data

50% inhibition (IC₅₀) values of α -amylase and α -glucosidase activities were determined from the regression analysis. Correlation analysis was conducted to determine the relationship between the flavonoids, phenolic compounds and inhibition activities of α -amylase and α -glucosidase.

3. Results and discussion

3.1 Total phenolic content

Total phenolic content in this study was ranged from 77.52 \pm 0.02 to 141.74 \pm 0.03 (mg GAE/100 g) at the concentration of 100 μ g/mL (Table 1). Stingless bee honey from mangrove origin showed the highest phenolic content with the value of 141.74 \pm 0.03 (mg

GAE/100 g); followed by the honey from coconut tree with the second highest phenolic content of 101.41 ± 0.06 (mg GAE/100 g). The descending trend of total phenolic content in honey samples from other botanical origins were shown as follow: Acacia (99.74 ± 0.08 mg GAE/100 g) > tualang (89.19 ± 0.01 mg GAE/100 g) > starfruit (85.74 ± 0.02 mg GAE/100 g) > multiflower (83.08 ± 0.04 mg GAE/100 g) > multifruit (77.52 ± 0.02 mg GAE/100 g). The composition of honey varies depending on the botanical origins and the result of this study was in agreement with the study of Shamsudin *et al.* (2019). In general, polyphenols in honey play an important role to maximize the efficiency of some biological activities that take place in the human body (Hoffman and Gerber, 2015). Therefore in this study, the presence of polyphenol compounds in honey might responsible for the α -amylase and α -glycosidase inhibition activity.

Table 1. Total phenolic content for honey samples

Samples	Phenolic content (mg GAE/100 g)
Acacia (K1)	99.74 ± 0.08
Coconut (K2)	101.41 ± 0.06
Mangrove (K3)	141.74 ± 0.03
Starfruit (K4)	85.74 ± 0.02
Multifruit (K5)	77.52 ± 0.02
Multiflower (K6)	83.08 ± 0.04
Tualang (K7)	89.19 ± 0.01

Honey samples were labelled as: K1 represents acacia tree; K2 represents coconut tree; K3 represents mangrove tree; K4 represents starfruit tree; K5 represents multifruit plant; K6 represents multiflower plant; K7 represents tualang tree. Each value was shown in mean \pm standard deviation (n=3).

3.2 Total flavonoid content

Total flavonoid content values were ranged from 13.71 ± 0.02 to 51.33 ± 0.02 (mg RE/100 g) (Table 2). Stingless bee honey from coconut origin displayed the highest flavonoid content with the value of 51.33 ± 0.02 (mg RE/100 g) and the second highest flavonoid content value of 44.66 ± 0.02 (mg RE/100 g) was observed in the honey sample from the mangrove tree. Total flavonoid content of honey samples from other origins was arranged in the descending order as shown follow: multifruit (41.33 ± 0.01 mg RE/100 g) > starfruit (32.28 ± 0.04 mg RE/100 g) > Acacia (29.42 ± 0.01 mg RE/100 g) > multiflower (19.42 ± 0.05 mg RE/100 g) > tualang (13.71 ± 0.02 mg RE/100 g). The overall result showed that the stingless bee honey samples were found to have more total flavonoid content than the tualang honey and this was analogue to the study of Abu Bakar *et al.* (2017) as the total flavonoid content of *H. itama* honey collected from different locations was much higher than tualang honey. The variation on the composition of flavonoid compounds in honey that

observed in this study was likely due to the floral sources but geographical locations, seasonal and environmental factors could also be one of the reasons (Kaskoniene and Venskutonis, 2010). According to Erejuwa *et al.* (2011), elimination of excess free radicals from body might be one of the preventions among diabetic patients. Hence, the result in this study was in compliance with the statement as the presence of flavonoid in all the honey samples could act as a free radical scavenger.

Table 2. Total flavonoid content for honey samples

Samples	Flavonoid content (mg RE/100 g)
Acacia (K1)	29.42 ± 0.01
Coconut (K2)	51.33 ± 0.02
Mangrove (K3)	44.66 ± 0.02
Starfruit (K4)	32.28 ± 0.04
Multifruit (K5)	41.33 ± 0.01
Multiflower (K6)	19.42 ± 0.05
Tualang (K7)	13.71 ± 0.02

Honey samples were labelled as: K1 represents acacia tree; K2 represents coconut tree; K3 represents mangrove tree; K4 represents starfruit tree; K5 represents multifruit plant; K6 represents multiflower plant; K7 represents tualang tree. Each value was shown in mean \pm standard deviation (n=3).

3.3 α -amylase inhibition assay

All honey samples exhibited α -amylase inhibition ability at the concentrations ranged from 20 to 100 μ g/mL. Based on the result obtained in Table 3, the percentage of α -amylase inhibition increased as the concentrations of honey samples increased. At a concentration of 100 μ g/mL, the highest activity was observed in the tualang honey (74.83%), followed by the mangrove (72.15%), coconut (71.65%), Acacia (70.30%), multifruit (55.76%), multiflower (46.76%) and starfruit (36.78%). The percentages inhibition of α -amylase at 80 μ g/mL was recorded with the values from 27.03 to 59.57% while the percentage inhibition beyond 60 μ g/mL was less than 50%. Starfruit honey showed the least α -amylase percentage inhibition (0.17-36.77%) along the concentrations. IC₅₀ values of the honey samples were ranged from 66.86 to 128.75 μ g/mL. Tualang honey achieved the highest inhibition activity with its low IC₅₀ value and this could be related to the presence of more phenolic acid in the honey. Phenolic compounds that present in tualang honey such as kaempferol, caffeic acid and p-coumaric acid were found to have anti-diabetic properties and they might contribute to the α -amylase inhibition activity (Ahmed and Othman, 2013; Bharti *et al.*, 2017). However, stingless bee honey samples from mangrove, coconut and Acacia trees could also be a good α -amylase inhibitor to reduce postprandial blood glucose levels in the body because their inhibition abilities were more than 70% at 100 μ g/mL.

Table 3. α -amylase inhibition of honey samples

Samples	Inhibition activity (%)				
	20 $\mu\text{g/mL}$	40 $\mu\text{g/mL}$	60 $\mu\text{g/mL}$	80 $\mu\text{g/mL}$	100 $\mu\text{g/mL}$
Acacia (K1)	1.23	9.29	39.48	46.38	70.29
Coconut (K2)	0.41	13.6	28.39	45.64	71.65
Mangrove (K3)	21.24	33.2	45.15	56.74	72.14
Starfruit (K4)	0.17	7.44	19.02	27.03	36.77
Multifruit (K5)	6.82	20.13	26.54	50.45	55.75
Multiflower (K6)	3.74	14.71	22.23	28.27	46.75
Tualang (K7)	2.39	22.64	43.02	59.57	74.82

Honey samples were labelled as: K1 represents acacia tree; K2 represents coconut tree; K3 represents mangrove tree; K4 represents starfruit tree; K5 represents multifruit plant; K6 represents multiflower plant; K7 represents tualang tree. Each value was shown in mean \pm standard deviation (n=3).

3.4 α -glucosidase inhibition assay

α -glucosidase inhibition ability of all honey samples was studied at the concentrations ranged from 20 to 100 $\mu\text{g/mL}$ (Table 4). In this assay, stingless bee honey sourced from coconut displayed the highest α -glucosidase inhibition with 68.32% at a concentration of 100 $\mu\text{g/mL}$. This probably related to its high phenolic and flavonoid contents as recorded in this study. 13.82 to 60.60% of α -glucosidase inhibition was recorded among the honey samples at the concentration of 80 $\mu\text{g/mL}$ while at the concentrations from 20 to 60 $\mu\text{g/mL}$, all honey samples achieved less than 50% of inhibition. The least percentage inhibition for α -glucosidase inhibition activity was observed in starfruit honey with the values from 1.08% to 20.52% along with the concentrations from 20 to 100 $\mu\text{g/mL}$. The lowest IC_{50} (77.69 $\mu\text{g/mL}$) was found in the honey from coconut origin whereas the honey from starfruit had the IC_{50} value greater than 100 $\mu\text{g/mL}$. Krishnasree and Mary Ukkuru (2017) showed that the stingless bee honey has a significant effect in α -amylase and α -glucosidase inhibition activities, however different types of bioactive compounds present in the honey from different botanical origins might influence the inhibition ability of honey (Shamsudin *et al.*, 2019).

3.5 Correlation between bioactive compounds, α -amylase inhibition and α -glucosidase inhibition

Total phenolic content was moderately correlated

with α -amylase inhibition activity and α -glucosidase inhibition activity with the r values of 0.5376 and 0.5173, respectively. Meanwhile, low positive correlations were observed between total flavonoid content, α -amylase and α -glucosidase inhibition activities with the r values of 0.1580 and 0.2327, respectively. Both flavonoid and phenolic compounds possess a positive effect on enzyme inhibition activities, but the presence of phenolic compounds contributed more on the inhibition ability of honey. The study of Devarajan and Venugopal (2012) also stated that major phenolic compounds and flavonoid in the honey extract may increase its potential anti-diabetic effects such as in α -amylase and α -glucosidase inhibition activities.

4. Conclusion

In conclusion, phenolic and flavonoid contents of stingless bee honey samples were varied in their botanical sources and the composition of honey resulted in different α -amylase and α -glucosidase inhibitory abilities as there was a positive correlation between enzyme inhibition activities (i.e. α -amylase and α -glucosidase) and bioactive compounds. Tualang honey displayed the highest α -amylase inhibition activity (74.83% at 100 $\mu\text{g/mL}$) but stingless bee honey samples from mangrove, coconut and acacia tree could potentially reduce blood glucose levels in the body, while honey from coconut origin exhibited the highest α -

Table 4. α -glucosidase inhibition of honey samples

Samples	Inhibition activity (%)				
	20 $\mu\text{g/mL}$	40 $\mu\text{g/mL}$	60 $\mu\text{g/mL}$	80 $\mu\text{g/mL}$	100 $\mu\text{g/mL}$
Acacia (K1)	1.89	4.87	12.46	34.46	38.75
Coconut (K2)	3.25	24.93	26.02	56.10	68.32
Mangrove (K3)	8.13	20.33	33.33	45.96	67.51
Starfruit (K4)	1.08	2.16	5.96	13.82	20.52
Multifruit (K5)	5.69	15.71	28.72	35.77	45.52
Multiflower (K6)	4.60	10.29	20.59	42.00	49.32
Tualang (K7)	8.13	33.06	38.75	60.60	62.60

Honey samples were labelled as: K1 represents acacia tree; K2 represents coconut tree; K3 represents mangrove tree; K4 represents starfruit tree; K5 represents multifruit plant; K6 represents multiflower plant; K7 represents tualang tree. Each value was shown in mean \pm standard deviation (n=3).

glucosidase inhibition activity due to its high content of phenolic and flavonoids. Therefore, stingless bee honey is suggested as a good source containing rich bioactive constituents and highly recommended to be utilized for nutraceutical and pharmaceutical applications. Further studies concerning the identification of bioactive compounds of the stingless bee honey and their verification of enzyme inhibition activity through clinical research are needed for effective management of diabetes.

Acknowledgments

The authors would like to acknowledge Universiti Tun Hussein Onn Malaysia (UTHM) for providing the facilities and financial support for this research (H 277).

References

- Abd Jalil, M.A., Kasmuri, A.R. and Hadi, H. (2017). Stingless bee honey, the natural wound healer: A review. *Skin Pharmacology and Physiology*, 30, 66–75. <https://doi.org/10.1159/000458416>
- Abu Bakar, M.F., Sanusi, S.B., Abu Bakar, F.I., Cong, O.J. and Mian, Z. (2017). Physicochemical and antioxidant potential of raw unprocessed honey from Malaysian stingless bees. *Pakistan Journal of Nutrition*, 16(11), 888–894. <https://doi.org/10.3923/pjn.2017.888.894>
- Ahmed, S. and Othman, N.H. (2013). Review of the medicinal effects of Tualang honey and a comparison with Manuka honey. *Malays Journal of Medical Science*, 20(3), 6–13.
- Akhir, R.A.M., Abu Bakar, M.F. and Sanusi, A.B. (2017). Antioxidant and antimicrobial activity of stingless bee bread and propolis extracts. *AIP Conference Proceedings*, 1891(1), 020090. <https://doi.org/10.1063/1.5005423>
- Alberti, K.G. and Zimmet, P.Z. (1998). Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1 Diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabetic medicine: a journal of the British Diabetic Association*, 15(7), 539–553.
- Apostolidis, E., Kwon, Y.I. and Shetty, K. (2007). Inhibitory potential of herb against key enzymes linked to diabetes and hypertension. *Innovative Food Science and Emerging Technology*, 8(1), 46–54. <https://doi.org/10.1016/j.ifset.2006.06.001>
- Bharti, S.K., Krishnan, S., Kumar, A. and Kumar, A. (2018). Antidiabetic phytoconstituents and their mode of action on metabolic pathways. *Therapeutic Advances in Endocrinology and Metabolism*, 9(3), 81–100. <https://doi.org/10.1177/2042018818755019>
- Campone, L., Piccinelli, A.L., Pagano, L., Carabetta, S., Sanzo, R., Russo, M. and Rastrelli, L. (2014). Determination of phenolic compounds in honey using dispersive liquid-liquid microextraction. *Chromatography Analysis*, 1334, 9–15. <https://doi.org/10.1016/j.chroma.2014.01.081>
- Cho, N.H., Karuranga, S., Huang, Y., Rocha, D.F.J.D., Ohlrogge, A.W. and Malanda, B. (2018). IDF diabetes atlas: Global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Research and Clinical Practise*, 138, 271–281. <https://doi.org/10.1016/j.diabres.2018.02.023>
- Devarajan, S. and Venugopal, S. (2012). Antioxidant and α -amylase inhibition activities of phenolic compounds in the extracts of Indian honey. *Chinese Journal of Natural Medicines*, 10(4), 255–259. [https://doi.org/10.1016/S1875-5364\(12\)60051-X](https://doi.org/10.1016/S1875-5364(12)60051-X)
- Erejuwa, O.O., Sulaiman, S.A. and Wahab, M.S. (2011). Oligosaccharides might contribute to the antidiabetic effect of honey: A review of the literature. *Molecules*, 17(1), 248–266. <https://doi.org/10.3390/molecules17010248>
- Eteraf-Oskouei, T. and Najafi, M. (2013). Traditional and modern uses of natural honey in human diseases: A review. *Iranian Journal of Basic Medical Sciences*, 16(6), 731–742.
- Hoffman, R. and Gerber, M. (2015). Food processing and the Mediterranean diet. *Nutrients*, 7(9), 7925–7964. <https://doi.org/10.3390/nu7095371>
- Inoue, R., Suzuki, N. and Nagashima, T. (2007). α -amylase from persimmon honey: Purification and characterization. *Food Properties*, 19(7), 512–521.
- Islam, M.R., Pervin, T., Hossain, H., Saha, B. and Hossain, S.J. (2017). Physicochemical and antioxidant properties of honeys. *Journal of Preventive Nutrition and Food Science*, 22(4), 335–344. <https://doi.org/10.3746/pnf.2017.22.4.335>
- Istasse, T., Jacquet, N., Berchem, T., Haubruge, T., Nguyen, B.K. and Richel, A. (2016). Extraction of honey polyphenols: Method development and evidence of cis isomerization. *Analytical Chemistry Insights*, 11, 49–57. <https://doi.org/10.4137/ACI.S39739>
- Kaskoniene, V. and Venskutonis, P.R. (2010). Floral markers in honey of various botanical and geographic origins: A review. *Comprehensive Reviews in Food Science and Food Safety*, 9(6), 620–634. <https://doi.org/10.1111/j.1541-4337.2010.00130.x>
- Krishnasree, V. and Mary Ukkuru, P. (2017). *In vitro* anti-diabetic activity and glycemic index of bee honeys. *Indian Journal of Traditional Knowledge*,

16(1), 134–140.

- Lim, D.C.C., Abu Bakar, M.F. and Majid, M. (2019). Nutritional composition of stingless bee honey from different botanical origins. *IOP Conference Series: Earth and Environmental Science*, 269, 012025. <https://doi.org/10.1088/1755-1315/269/1/012025>
- Majid, M., Abu Bakar, M.F., Mian, Z., Esa, F. and Yeow, Y.K. (2019). Variations of physicochemical properties of stingless bee honey from different botanical origin in state of Johor, Malaysia. *IOP Conference Series: Earth and Environmental Science*, 269, 012028. <https://doi.org/10.1088/1755-1315/269/1/012028>
- Mohd Rafie, A.Z., Syahir, A., Wan Ahmad, W.A.N., Mustafa, M.Z. and Mariatulqabtiah, A.R. (2018). Supplementation of stingless bee honey from *Heterotrigona itama* improves antiobesity parameters in high-fat diet induced obese rat model. *Evidence-Based Complementary and Alternative Medicine*, 2018, 6371582. <https://doi.org/10.1155/2018/6371582>
- Rahmawati, O., Pratami, D.K. and Raffiudin, R. (2019). Alpha-glucosidase inhibitory activity of stingless bee honey from *Tetragonula biroi* and *Tetragonula laeviceps*. *AIP Conference Proceedings*, 2092(1), 030001. <https://doi.org/10.1063/1.5096705>
- Rutebemberwa, E., Lubega, M., Katureebe, S.K., Oundo, A., Kiweewa, F. and Mukanga, D. (2013). Use of traditional medicine for the treatment of diabetes in Eastern Uganda: a qualitative exploration of reasons for choice. *BioMed Central*, 13, 92-103. <https://doi.org/10.1186/1472-698X-13-1>
- Shamsudin, S., Selamat, J., Sanny, M., Shamsul Bahari A.R., Jambari, N.N. and Khatib, A. (2019). A comparative characterization of physicochemical and antioxidants properties of processed *Heterotrigona itama* honey from different origins. *Molecules*, 24 (21), 1–20. <https://doi.org/10.3390/molecules24213898>
- Shaw, J.E., Sicree, R.A. and Zimmet, P.Z. (2010). Global estimates of the prevalence of diabetes for 2010 and 2030. *Diabetes Research and Clinical Practice*, 87(1), 4-14. <https://doi.org/10.1016/j.diabres.2009.10.007>
- Yap, P., Bakar, M.F.A., Lim, H. and Carrier, D. (2015). Antibacterial activity of polyphenol-rich extract of selected wild honey collected in Sabah, Malaysia. *Journal of Apicultural Research*, 54(3), 163-172. <https://doi.org/10.1080/00218839.2016.1151633>