

## Original Article

# In Vitro Antiamoebic Activity of Iranian *Allium sativum* in Comparison With Metronidazole against *Entamoeba histolytica*

M Behnia<sup>1</sup>, \* A Haghghi<sup>2</sup>, H Komeilizadeh<sup>3</sup>, SJ Seyyed Tabaei<sup>2</sup>, A Abadi<sup>4</sup>

<sup>1</sup> Laboratory Complex, Science and Research Branch, Islamic Azad University, Tehran, Iran

<sup>2</sup> Dept. of Parasitology and Mycology, School of Medicine, Shahid Beheshti University (Medical Campus), Tehran, Iran

<sup>3</sup> Dept. of Medicinal Chemistry, School of Pharmacy, Shahid Beheshti University (Medical Campus), Tehran, Iran

<sup>4</sup> Dept. of Health and Social Medicine, School of Medicine, Shahid Beheshti University (Medical Campus), Tehran, Iran.

(Received 1 Apr 2008; accepted 14 Jul 2008)

## Abstract

**Background:** Amoebiasis is due to infection with the protozoan parasite *Entamoeba histolytica*. The patients infected with *E. histolytica* must be treated right after definite diagnosis and no need to treat infected individuals with *E. dispar* isolates. Metronidazole is used as a drug of choice against amoebiasis. However, like a lot of other chemical agents, this drug has its own side effects. This prompted us to carry out, an in vitro research into antiamoebic effect of Iranian *Allium sativum* (garlic), which has been used for centuries, as an herbal medicine, without harmful side effects.

**Methods:** Hydro-alcoholic, hexanic extracts and essential oil of 100 gram of crushed *A. sativum* was isolated and the minimal inhibitory concentration (MIC) of the extracts and essential oil in comparison with metronidazole were obtained on trophozoite of *E. histolytica*, HM-1: IMSS strain in TYI-S-33 medium.

**Results:** The MIC for *A. sativum* hydroalcoholic, hexanic extracts and essential oil after 24 hours was 60mg mL<sup>-1</sup>, 4mg mL<sup>-1</sup> and 0.4mg mL<sup>-1</sup>, respectively. After 48 hours the MIC for *A. sativum* hexanic extract and essential oil was 3mg mL<sup>-1</sup> and 0.3mg mL<sup>-1</sup>, respectively. MIC for metronidazole was obtained 2µg mL<sup>-1</sup> and 1.5µg mL<sup>-1</sup> after 24 hours and 48 hours, in that order.

**Conclusion:** Iranian *A. sativum* is effective on the trophozoites of *E. histolytica* species and the essential oil exhibited the greatest antiamoebic activity, at the lowest MIC.

**Keywords:** *Entamoeba histolytica*, *Allium sativum*, Antiamoebic

## Introduction

A mong parasitic infections amoebiasis caused by *Entamoeba histolytica* ranks third worldwide in lethal infection, after malaria and schistosomiasis (1, 2). Although it is asymptomatic in 90% of cases, about 50 million people are estimated to suffer from the symptoms of amoebiasis such as hemorrhagic colitis and

amoebic liver abscess. These infections result in 50000–100000 deaths annually (3). Data from some parts of Iran showed that 7.9% of the *E. histolytica*/*E. dispar* isolates were *E. histolytica* or mix infection and 92.15% were *E. dispar* (4). The current treatments of choice are either one of a family of nitroimidazoles (usually metronidazole), nitrofurans, quinacrine or paromomycin (5). However, metronidazole has been reported to cause mutagenicity in bacteria (6) and is

carcinogenic in rodents (7, 8). It has been reported that the human pathogenic bacterium, *Helicobacter pylori*, becomes resistant to metronidazole in vitro (9). Moreover, it seems to act as an immunosuppressive agent in experimental rats, both in cell-mediated and humoral immune responses (10). However, like other chemical agents, metronidazole has some side effects including metallic taste, nausea, transient neutropenia, interaction with warfarin, and peripheral neuropathy (11).

These are the main reasons why it is essential to develop a safe and effective alternative anti-amoebic agent. Garlic is one of the edible plants which have generated a lot of interest throughout human history as a medicinal panacea. A wide range of microorganisms have been shown to be sensitive to crushed garlic preparations. Garlic has been shown to be antibacterial (12), antiviral (13) and antifungal (14), as well as possessing both antitumor (15) and antithrombotic (16) properties. Moreover, garlic has been reported to reduce blood lipids (17). Chemical analyses of garlic have revealed an unusual concentration of sulfur-containing compounds (1–3%) (17, 18). Garlic is a very complex compound consisting not only of allyl components but many other components, some of which, kaempferol and quercetin, have been shown to have anti-giardial activity in vitro (19). The use of garlic and some of its components as antiprotozoals has already been investigated by some authors (20–22).

To our knowledge anti-amoebic of Iranian *A. sativum* have not been studied so far. This prompted us to study the possible anti-amoebic properties of hydroalcoholic, hexanic extract and essential oil of Iranian *A. sativum* (garlic).

## Materials and Methods

### *Plant extract preparation*

*A. sativum* was collected in June 2006 from Hamedan, Iran, identified by the Tehran University Institute of Medicinal Plants (IMP)

and the experimental study was carried out in Shahid Beheshti University of Medical Sciences. One hundred gram of crushed *A. sativum* was subjected to maceration at 25 °C for 14 days using n-hexane and a mixture of ethanol and water (60:40 ratio). The extracts obtained were then filtered using Whatman No. 1 filter paper. The filtrates were concentrated with a rotary evaporator at 40° C to facilitate their further freeze-drying process. The concentrated extracts were finally freeze-dried at -50 °C for 24h and stored at -20 °C. Extracts were redissolved in their solvents before each individual experiment (23). The yield of *A. sativum* using ethanol/water (60:40) as solvent was 13.82% dry extract and with n-hexane was 0.24 % dry extract.

### *Isolation of essential oil*

One hundred gram of crushed garlic were subjected to hydrodistillation using a clevenger-type apparatus for 3 h. The essential oil obtained was dried with anhydrous sodium sulfate and kept at 4 °C (24).

The hydrodistillation of the crushed plant yielded amber-like oil, in a 0.22% yield with a characteristic odor.

### *Parasite culture*

*E. histolytica*, HM-1: IMSS strain was used in all experiments. The trophozoites were cultured axenically in screw-capped tubes at 35.5 °C on Diamond's TYI-S-33 medium, supplemented with 10% (v/v) heat-inactivated bovine serum (25). Subculturing was performed routinely at 48 h intervals by replacing the medium without detaching the monolayer. Cells were harvested by replacing the medium with a fresh one, chilling on ice for 20 min, and inverting gently to detach the monolayer.

### *Preparation of stock solutions*

One hundred gram mL<sup>-1</sup> of each extract was dissolved in its solvent, sterilized through a 0.22 µm Millipore filter and kept as stock solution. The essential oil was sterilized through a 0.22 µm Millipore filter and was then emulsified by 5% Tween 80.

### *Exposure and evaluation*

A concentration range of 0-100mg mL<sup>-1</sup> of plant extract and the essential oil was prepared in test tubes by serial dilution and air-dried under sterile conditions. To the test tubes was added Diamond's TYI-S-33 medium. The contents of the test tubes were then turned into a suspension by using the ultrasound bath for 30 min and were kept at rest for 24 h at 4° C. To the test tubes were subsequently added 1.0×10<sup>3</sup> trophozoites of *E. histolytica* followed by incubation at 35.5° C for 24 h and 48 h.

The tests were also repeated using the standard amoebicidal drug metronidazole (26). The experimental controls were assigned in the following manner: I) Diamond's TYI-S-33 medium+trophozoites II) Diamond's TYI-S-33 medium+trophozoites+5% Tween 80 and III) Diamond's TYI-S-33 medium+trophozoites+metronidazole (0.5, 1, 1.5 and 2µg mL<sup>-1</sup>).

After the 24 h and 48 h periods, the tubes were chilled for 20 min and the attached trophozoites were detached by gentle inversion. The number of viable cells was determined using eosin 0.01% (27) and the microscope (the Neubauer chamber), the criteria for viability being motility and dye exclusion.

The lowest concentration of each plant extract and essential oil which completely inhibited the growth of trophozoites of *E. histolytica* was considered the MIC (26). After determining the MIC, concentrations of ¼, ½ and ¾ of MIC were used. Growth rate (GR) is defined as the difference between the number of viable protozoa counted at 0h and after 24 h and 48 h. The percentage of growth inhibition (% GI) was calculated using the following formula (28):

$$\%GI = \left(1 - \frac{GR_{\text{extract}}}{GR_{\text{control}}}\right) \times 100$$

The experiments were performed in duplicate and repeated three times.

#### **Statistical analysis**

Data were analyzed using two-way analysis of variance (ANOVA) (29).

## **Results**

The inhibitory capacity of the extracts and essential oil of *A. sativum* were assayed using eosin, as a result of which viable trophozoites of *E. histolytica* remained unstained whereas dead cells were light red in color (27). The effect of various concentrations of *A. sativum* extracts and essential oil against trophozoites of *E. histolytica* and the calculated %GI (%growth inhibition) at 24 h and 48 h are shown in the table of results (Table 1).

Both the extracts and the essential oil of *A. sativum* exhibited antiamoebic activity. However, the essential oil proved to be more effective. The MIC of the essential oil after 24 h was 0.4mg mL<sup>-1</sup> and after 48h was 0.3mg mL<sup>-1</sup>. As shown in Table 1, 60mg mL<sup>-1</sup> of hydroalcoholic extract after 24 h exhibited inhibition and therefore had little effect on the trophozoites of *E. histolytica*, whereas 4mg mL<sup>-1</sup> after 24 h (Fig. 1) and 3mg mL<sup>-1</sup> after 48 h (Fig. 2) of the hexanic extract caused complete growth inhibition. On the other hand, metronidazole, the current drug of choice, had a MIC of 2µg mL<sup>-1</sup> after 24 h and 1.5µg mL<sup>-1</sup> after 48 h.


Experiments showed that as the concentration increased, the number of viable trophozoites decreased. Moreover, concentration being kept constant, the results in Fig. 1 and Fig. 2 indicate that longer periods of exposure to the extracts and the essential oil, decreased the number of viable trophozoites.

**Table 1:** Effect of metronidazole, the hydroalcoholic, hexanic extracts and essential oil of *A. sativum* and on trophozoites of *E. histolytica* after 24 h and 48 h incubation


Antiamoebic agent	Concentration mg mL <sup>-1</sup>	Growth Inhibition (%) <sup>*</sup>	
		24 h	48 h
Hydroalcoholic extract of <i>A. sativum</i>	15	65.68 ± 2.67	77.37 ± 1.68
	30	80.25 ± 1.93	97.93 ± 0.76
	45	96.54 ± 1.15	99.48 ± 0.76
	60 <sup>**</sup>	100	100
Hexanic extract of <i>A. sativum</i>	1	35.1 ± 3.08	87.69 ± 1.31
	2	88.2 ± 2.51	99.74 ± 0.6
	3	99.26 ± 1.33	100
	4 <sup>**</sup>	100	100
Essential oil of <i>A. sativum</i>	0.1	54.19 ± 4.88	79.17 ± 1.38
	0.2	85.38 ± 2.16	99.62 ± 0.69
	0.3	95.13 ± 1.44	100
	0.4 <sup>**</sup>	100	100
Metronidazole	0.5 µg mL <sup>-1</sup>	79.09 ± 3.2	97.67 ± 0.8
	1 µg mL <sup>-1</sup>	90.64 ± 2.47	99.74 ± 0.6
	1.5 µg mL <sup>-1</sup>	95.32 ± 1.99	100
	2 µg mL <sup>-1**</sup>	100	100

\* The values are mean values obtained in three assays done in duplicate.

\*\* The lowest concentration of each plant extracts, essential oil and metronidazole which completely inhibited the growth of trophozoites of *E. histolytica* (MIC)



**Fig. 1:** Effects of metronidazole, the hydroalcoholic and hexanic extracts and the essential oil of *A. sativum* on the growth inhibition of *E. histolytica* HM-1: IMSS trophozoites after 24 h



**Fig. 2:** Effects of metronidazole, the hydroalcoholic and hexanic extracts and the essential oil of *A. sativum* on the growth inhibition of *E. histolytica* HM-1: IMSS trophozoites after 48 h

## Discussion

For many years, garlic has been used as an effective therapeutic agent having antibacterial, antifungal, antiviral, antiprotozoal and antihelminthic properties (30).

This study investigated the antiamoebic activity of *A. sativum* extracts and essential oil and compared the MICs of these agents with metronidazole.

Parasite growth depends on the primary number of active parasites, temperature and culture medium. In our study, in similar experimental conditions, the results of the experiments in study group were significantly different from those of the control group ( $P < 0.01$ ).

Mirelman *et al.* investigated the use of allicin, a sulfur compound on *E. histolytica* (20); Soffar & Mohktar assessed its use as an anti-giardial alongside its use as an antihelminthic in a selection of patients (21), whilst Lun *et al.* looked at the effect of diallyl trisulphide on *E. histolytica*, *G. intestinalis* and *Trypanosomes* (22). Harris *et al.* showed that the efficacy of garlic extract as an anti-giardial (23).

In this study, the essential oil, which is composed of allicin, allylpropyl disulphide (31),

exhibited the greatest antiamoebic activity, at the lowest MIC, which the hexanic extract showed 100% lysis of trophozoites of *E. histolytica* at a concentration of 4 mg mL<sup>-1</sup> in comparison to the hydroalcoholic extract which exhibited almost no antiamoebic activity.

From our findings as well as those of other researchers concerning the antiprotozoal property of garlic, it can be concluded that the essential oil and the hexanic extract of garlic seems to be a good antiamoebic candidate for amoebiasis treatment and inhibition of growth of *E. histolytica* by garlic is dose-dependent. However some *in vivo* studies regarding to its effects on animals and human should be investigated.

## Acknowledgements

This work was supported by a grant from Research Affairs, School of Medicine, Shahid Beheshti University (Medical campus) Tehran, Iran. The authors would like to thank Mr. Kamalinejad M. from the School of Pharmacy of Shahid Beheshti University (M.C), Dr. Aberumand Azar P. and Mr. Larijani K. from the science and research branch of Islamic

Azad University for their help with providing the garlic agents and also Ms. Sh. Farnia from the School of Public Health, Tehran University of Medical Sciences for her help with providing the parasite culture. The authors declare that they have no conflict of interests.

## References

1. Walsh JA. Prevalence of *Entamoeba histolytica* infection. In: Ravdin JI, editor. Amoebiasis. New York: John Wiley & Sons; 1988. p. 93-105.
2. Petri Jr WA, Mann BJ. Molecular mechanisms of invasion by *Entamoeba histolytica*. Semin Cell Bio. 1993;4(5): 305-13.
3. Ravdin JI. Amebiasis. Clin Infect Dis. 1995;20(6):1453-64.
4. Hooshyar H, Rezaian M, Kazemi B, Jeddi-Tehrani M, Soleymani-Mohammadi S. The distribution of *Entamoeba histolytica* and *Entamoeba dispar* in northern, central and southern Iran. Parasitol Res. 2004;94(2):96-100.
5. Tracy JW, Webster JLT. Drugs used in the chemotherapy of protozoal infection. In: Hardman JG, Limbird LE, editors. The Pharmacological Basis of Therapeutics. New York: McGraw-Hill; 1996. p. 987-1008.
6. Legator MS, Connor TH, Stoeckel M. Detection of mutagenic activity of metronidazole and niridazole in body fluids of humans and mice Science. 1975;188 (4193):1118-9.
7. Rustia M, Shubik P. Induction of lung tumors and malignant lymphomas in mice by metronidazole. J Natl Cancer Inst. 1972;48(3):721-9.
8. Shubik P. Current status of chemical carcinogens. Proc Natl Acad Sci USA. 1972; 69(4):1052-5.
9. Zwet AA, Thijs JC, Schievink-de Vries W, Schiphuis J, Snijder JAM. In vitro studies on stability and development of metronidazole resistance in *Helicobacter pylori*. Antimicrob Agents Chemother. 1994;38(2):360-2.
10. Saxena A, Chugh S, Vinayak VK. Modulation of host immune responses by metronidazole. Indian J Med Res. 1985; 81:387-90.
11. Kurohara ML, Kwong FK, Leberherz TB, Klaustermeyer WB. Metronidazole hypersensitivity and oral desensitization. J Allergy Clin Immunol. 1991;88(2):279-80.
12. Sharma VD, Sethi MS, Kumar A, Rarotra JR. Antibacterial property of *Allium sativum* Linn: in vivo and in vitro studies. Indian J Exp Biol. 1977;15(6):466-8.
13. Fenwick G R, Hanley AB. The genus *Allium*-part 3. Crit Rev Food Sci Nutr. 1985;23(1):1-73.
14. Yamada Y, Azuma K. Evaluation of the in vitro antifungal activity of allicin. Antimicrob Agents Chemother. 1997;11(4):743-9.
15. Milner JA. Garlic: its anticarcinogenic and anti tumorigenic properties. Nutr Rev. 1996;54(Suppl):82-86.
16. Kiesewetter H, Jung F, Jung EM, Mroweitz C, Koscrelly J, Wenzel E. Effect of garlic on platelet aggregation in patients with increased risk of juvenile ischemic attacks. Eur J Clin Pharmacol. 1993; 45(4):333-6.
17. Darbyshire B, Henry RJ. Differences in fructan content and synthesis in some *Allium* species. New Phytol. 1981;87(2): 249-56.
18. Koch HP, Lawson LD. Garlic, the science and therapeutic application of *Allium sativum* L. and related species. In: Retford DC, editor. Baltimore: Williams and Wilkins; 1996. p. 213-233.
19. Calzada F, Alanis AD, Meckis M, Contreras A, Cedillo-Rivera R. In vitro susceptibility of *Entamoeba histolytica*

- and *Giardia lamblia* to some medicinal plants used by the people of Southern Mexico. *Phyto Res.* 1998;12:70-2.
20. Mirelman D, Monheit D, Varon S. Inhibition of growth of *Entamoeba histolytica* by allicin, the active principle of garlic extract (*Allium sativum*). *J Infect Dis.* 1987;156(1):243-4.
  21. Soffar SA, Mokhtar GM. Evaluation of the antiparasitic effect of aqueous garlic (*Allium sativum*) extract in hymenolepiasis nana and giardiasis. *J Egypt Soc Parasitol.* 1991;21(2):497-502.
  22. Lun ZR, Burri C, Menzinger M, Kaminsky R. Antiparasitic activity of diallyl trisulphide (dasuansu) on human and animal pathogenic protozoa (*Trypanosoma* sp., *Entamoeba histolytica* and *Giardia lamblia*) in vitro. *Ann Soc Belg Med Trop.* 1994;74(1):51-9.
  23. Schaffer S. Understanding local Mediterranean diets: A multidisciplinary pharmacological and ethnobotanical approach. *Pharmacol Res.* 2005;52(4):353-6.
  24. Goodner KL, Mahattanatawee K, Plotto A, Sotomayor JA, Jord'an MJ. Aromatic profiles of *Thymus hyemalis* and Spanish *T. vulgaris* essential oils by GC-MS/GC-O. *Indust Crops Prod.* 2006;24(3):264-268.
  25. Diamond LS, Harlow DR, Cunnick CC. A new medium for the axenic cultivation of *Entamoeba histolytica* and other *Entamoeba*. *Trans Roy Soc Trop Med Hyg.* 1978;72(4):431-2.
  26. Padayachee T, Odhav B. Anti-amoebic activity of plant compounds from *Virgilia oroboides* and *Chlorophora excelsa*. *J Ethnopharmacol.* 2001;78(1):59-66.
  27. Bingham AK, Jarroll EL Jr, Meyer EA, Radulescu S. *Giardia* sp; Physical factors of excystation invitro and excystation VS eosin exclusion as determinants of viability. *Exp Parasitol.* 1979;47(2):284-91.
  28. Muelas-Serrano S, Nogal JJ, Martinez-Diaz RA, Escario JA, Martinez-Fernandez AR, Gomez-Barrío A. In vitro screening of American plant extracts on *Trypanosoma cruzi* and *Trichomonas vaginalis*. *J Ethnopharmacol.* 2000;71(1-2):101-7.
  29. Taran M, Rezaeian M, Izaddoost M. In vitro Antitrichomonas Activity of *Allium hirtifolium* (Persian Shallot) in Comparison with metronidazole. *Iranian J Pub Health.* 2006;35(1):92-94.
  30. Harris JC, Cottrell SL, Plunmer S, Lloyd D. Antimicrobial properties of *Allium sativum* (garlic). *Appl Microbiol Biotechnol.* 2001;57(3):282-6.
  31. Newall CA, Anderson LA, Phillipson JD. In *Herbal Medicines: A Guide for Healthcare Professionals*. London: The Pharmaceutical Press; 1996; p. 129-133.