

www.waavp2011-argentina.com.ar



2011 WAAVP BUENOS AIRES ARGENTINA

TOWARDS GOOD MANAGEMENT
PRACTICES IN PARASITE CONTROL

August 21-25, 2011

Panamericano Buenos Aire Hotel & Resort

aapavet

Asociación Argentina de Parasitología Veterinaria



***In vitro* efficacy of plant extracts and synthesized substances on *Rhipicephalus (Boophilus) microplus* (Acari: Ixodidae)**

L. D. de Barros¹, A. C. de Souza Chagas², F. Cotinguiba³, M. Furlan³, R. Giglioti¹, M. C. de Sena Oliveira², H. Ribeiro Bizzo⁴
¹Faculdade de Ciências Agrárias e Veterinárias/UNESP, Jaboticabal, SP; ²Centro de Pesquisa Pecuária Sudeste/EMBRAPA, São Carlos, SP; ³Instituto de Química/UNESP, Araraquara, SP; ⁴Centro de Tecnologia Agroindustrial de Alimentos/EMBRAPA, Rio de Janeiro, RJ.

Herbal drugs have been widely evaluated as an alternative method of parasite control, aiming to slow development of resistance and obtain low-cost biodegradable parasiticides. The aim of this study was to evaluate the *in vitro* efficacy on *Rhipicephalus (Boophilus) microplus* of *Carapa guianensis*, *Cymbopogon martinii*, *Cymbopogon schoenanthus* and *Piper tuberculatum* extracts and similar synthesized substances. In the immersion test, engorged females were collected from cattle and evaluated in five dilutions ranging from 10% to 0.030625%. In the larval test, around 100 larvae were placed on filter paper impregnated with the dilutions, ranging from 10% to 0.02%. The treatments and controls in both tests were done in three replicates. Phytochemical analysis of oils was performed by gas chromatography. The main compounds were oleic acid (46.8%) for *C. guianensis* and geraniol for *C. martinii* (81.4%) and *C. schoenanthus* (62.5%). The isolated and synthesized substances showed no significant effect on larvae and engorged females. *C. martinii* at a concentration of 5% and *P. tuberculatum* at 10%, showed the best efficacy on the engorged females, causing 75.60% and 91.97% mortality, respectively. The fact that geraniol is present in greater quantities in *C. martinii* explains its higher activity in relation to *C. schoenanthus*. There is still a need to validate the *in vivo* use of safe and effective phytoparasitocidal substances. Efforts should be focused on developing formulations that enhance the efficacy of these extracts *in vivo* and lengthen the residual period.
