

In Vitro Evaluation of Schistosomicidal Activity of Essential Oil of *Mentha x villosa* and Some of Its Chemical Constituents in Adult Worms of *Schistosoma mansoni*

Authors

Thiago José Matos-Rocha^{1,2}, Marília Gabriela dos Santos Cavalcanti^{1,2}, José Maria Barbosa-Filho³, Ana Sílvia Suassuna Carneiro Lúcio³, Dyana Leal Veras^{1,2}, Ana Paula Sampaio Feitosa^{1,2}, José Pinto de Siqueira Júnior³, Reinaldo Nóbrega de Almeida³, Márcia Ortiz Mayo Marques⁴, Luiz Carlos Alves^{1,2}, Fábio André Brayner^{1,2}

Affiliations

The affiliations are listed at the end of the article

Key words

- Lamiaceae
- *Mentha x villosa*
- *Schistosoma mansoni*
- schistosomicidal activity
- natural products

received June 26, 2013
revised July 3, 2013
accepted July 8, 2013

Bibliography

DOI <http://dx.doi.org/10.1055/s-0033-1350732>
Published online August 14, 2013
Planta Med 2013; 79: 1307–1312 © Georg Thieme Verlag KG Stuttgart · New York · ISSN 0032-0943

Correspondence

Fábio André Brayner dos Santos
Molecular and Cell Biology Laboratory
Department of Parasitology Aggeu Magalhães Research Center (FIOCRUZ) and Laboratory of Immunopathology Keizo Asami (LIKA)
Av. Moraes Rego s/n, Campus da Universidade Federal de Pernambuco, 50670-420 Recife, PE Brazil
Phone: + 55 81 21 01 25 00
Fax: + 55 81 21 01 26 71
brayner.santos@gmail.com

Abstract

This study aimed to determine the composition of the essential oil of *Mentha x villosa* and to evaluate its biological effects *in vitro* on adult worms of *S. mansoni*. Rotundifolone (70.96%), limonene (8.75%), trans-caryophyllene (1.46%), and β -pinene (0.81%) were shown to be the major constituents of this oil. Adult worms of *S. mansoni* were incubated with different concentrations of the essential oil (1, 10, 100, 250, 500, and 1000 μ g/mL) and of its constituents rotundifolone (0.7, 3.54, 7.09, 70.96, 177.4, 354.8, and 700.96 μ g/mL), limonene (43.75 μ g/mL), trans-caryophyllene (7.3 μ g/mL), and β -pinene

(4.03 μ g/mL). No schistosomicidal activity was identified at the trans-caryophyllene and β -pinene concentrations studied. However, use of the essential oil (10 μ g/mL), rotundifolone (7.09 μ g/mL), and limonene (43.75 μ g/mL) resulted in decreased worm motility continuing until 96 hours of observation. At higher concentrations (100 and 70.96 μ g/mL, respectively), both the essential oil and rotundifolone caused mortality among adult worms of *S. mansoni*. The positive control praziquantel caused the death of all parasites after 24 h of evaluation. The results from this study suggest that the essential oil of *Mentha x villosa* presents schistosomicidal efficacy.

Introduction

For treating schistosomiasis, WHO recommends praziquantel (PZQ) because it has low toxicity and can be administered in a single dose orally. However, indiscriminate use has resulted in appearance of strains of *Schistosoma mansoni* (Sambon, 1907) refractory to this drug [1]. Besides praziquantel, oxamniquine is also recommended by WHO as an alternative available for treating schistosomiasis, but because of adverse effects, its use has been restricted in Brazil [2].

The search for new effective drugs for treating schistosomiasis has directed studies towards extracts and compounds isolated from plants that can provide schistosomicidal activity [3–6].

M. x villosa Hudson, which is usually reported in the literature as *M. crispata*, belongs to the family Lamiaceae and is known popularly as the mint-leaf-girl, creeping mint, and mint-in-pot. After identification by Dr. R. Harley, of the Royal Botanic Gardens, Kew (UK), it was shown that *M. x villosa* is a hybrid originating from spontaneous crossing of *M. spicata* and *M. suaveolens* [7]. As for bioavail-

ability, there are no studies in the literature that show bioavailability in humans or animals.

Some biological studies have been conducted to evaluate the therapeutic activity of the essential oil of *M. x villosa* (EOMv), and these have also added information about this plant and its main compounds.

Rotundifolone (C₁₀H₁₄O₂; molecular weight 166) is a naturally occurring monoterpenic ketone of plant origin and an important chemical constituent of the essential oils of many *Mentha* species (*M. rotundifolia*, *M. suaveolens*, *M. spicata*, *M. longifolia*, and *M. x villosa*) [8]. This most abundant constituent has been evaluated in relation to the following different biological activities: cardiovascular [8], hypotensive, bradycardic [9,10], antimicrobial, antifungal [11], and antinociceptive [12] properties.

This study is given greater impact to rotundifolone because it represents more than 50% (can reach 65%) of the total oil of *M. x villosa*. It is the substance (monoterpene) present in greater amounts in the leaves of *M. x villosa* [13–17]. The following forms of activity have been reported in the literature with regard to rotundifo-

Substance	(%)	*KI from experiment	KI in literature*
Sabinene	0.23	972	976
β -Pinene	0.81	976	980
Myrcene	3.10	990	991
Limonene	8.75	1029	1031
Cis- β -ocimene	0.85	1035	1040
Rotundifolone	70.96	1363	1363
β -Bourbonene	0.09	1386	1384
β -Elemene	0.24	1393	1391
Trans-caryophyllene	1.46	1420	1418
α -Humulene	0.21	1454	1454
Trans- β -farnesene	0.39	1457	1458
Germacrene D	3.81	1481	1480
Cis-calamenene	0.22	1521	1521
ζ -Elemene	0.31	1523	1524
α -Muurolol	0.49	1653	1645

* Retention Index = Kovats Index (KI). Components are identified based on GC-MS and listed according to elution order in the column

Table 1 Chemical constituents of EOMv, as shown using GC-MS.

lone: as analgesic [13], relaxative [14], hypotensive, bradycardic [15], antinociceptive [16], antimicrobial, antifungal [11], and spasmolytic [17]. Studies demonstrated a possible mechanism of action involved in the relaxative effect exhibited by rotundifolone, which they correlated with the influx of calcium channels [18].

M. x villosa has been used in traditional medicine due to its antiparasitic property. It is known popularly as hortelã-rasteira (creeping mentha), hortelã comum (regular mentha), or hortelã-da-folha-miúda (small-leaved mentha) [19]. Giamebil® is a commercial formulation that has as its active compound the dry extract from the leaves and stem of *M. x villosa*, with amebicidal (*Entamoeba histolytica*) and giardicidal (*Giardia lamblia*) activities [20].

Recent studies have also demonstrated efficacy of *M. x villosa* in *Trichomonas vaginalis* [21]. However, there are no published studies that show the activity of the essential oil from the leaves of *M. x villosa* against adult worms of *S. mansoni*.

New molecules have been harvested from natural sources to fulfill a need for new chemical designs with distinctive pharmacological activities. The present study aimed to evaluate the *in vitro* effect of the essential oil of schistosomicidal *M. x villosa* and each of its main compounds (rotundifolone, limonene, trans-caryophyllene, and β -pinene) in adult worms of *S. mansoni*.

Results and Discussion

Several studies have shown that some natural products or compounds isolated from them present schistosomicidal activity [22–28]. Studies have also reported that some medicinal plants have been used clinically against schistosomiasis, such as myrrh (Mirazid®), a resinous oil obtained from *Commiphora molmol* (*Commiphora myrrha*), which is used in Egypt [29].

However, few studies *in vitro* have demonstrated any schistosomicidal activity of essential oils of medicinal plants against *S. mansoni* [30–33]. No work has been reported demonstrating the antiparasitic activity of EOMv and compounds isolated from it until now.

The EOMv yield was 0.1%, based on the dry weight of the plant, and its composition, analyzed using GC-MS, showed the presence of nineteen substances (Table 1). These were identified

through comparison of their spectral masses, using the GC-MS database system (Nist 62 lib.) and Kovats retention index [34]. The main compounds were: rotundifolone (70.96%), limonene (8.75%), trans-caryophyllene (1.46%), and β -pinene (0.81 μ g/mL). Antiparasitic activity of EOMv and its separate compounds against the adult worms of *S. mansoni* was observed during the biological assays *in vitro*. At the concentration of 100 μ g/mL, worm mortality was observed from 72 h onwards; at the concentration of 250 μ g/mL from 48 h; at the concentration of 500 μ g/mL from 24 h, and at the concentration of 1000 μ g/mL within 24 h of exposure (Table 2).

Using EOMv at concentrations of 1, 5, and 10 μ g/mL, worm mortality was not observed at any of the times examined. However, in terms of relative motility at a concentration of 10 μ g/mL, minor loss of the movements of the tail, suckers, and gynaecophoric canal membrane was observed after 96 h of exposure.

Biological assays evaluating the essential oil of *Ageratum conyzoides* L. (EOAc) demonstrated that it had schistosomicidal activity on adult worms of *S. mansoni* at concentrations of 10 to 100 μ g/mL. At the concentration of 100 μ g/mL, worm mortality was observed when evaluated for up to 120 h [31]. Similarly to the results from the present study, the concentration of 10 μ g/mL of EOMv caused a reduction in the motor activity of adult worms and at the concentration of 100 μ g/mL, worm mortality was observed at times up to 72 h. These results were also reported from a study on the schistosomicidal activity of the essential oil of *Plectranthus neochilus* (EOPn) [32].

Differently from the findings of another study in which the *in vitro* schistosomicidal activity of the essential oil of *Piper cubeba* (EOPc) was evaluated at concentrations of 12.5 and 25 μ g/mL at 120 h of exposure with regard to motility, a viability similar to that of the negative control group of adult worms of *S. mansoni* was observed [33]. At a concentration of 100 μ g/mL, reduction in the movements of adult worms of *S. mansoni* was observed up to 72 h. These results were superior to those observed in other studies evaluating EOAc, in which 75% mortality of the worms after this same time was reported [31].

Differently from the above results, biological assays testing the essential oil of *Baccharis dracunculifolia* found 100% mortality of the worms after 24 h at the concentration of 10 μ g/mL [30]. Also in this same study, 100% mortality of adult worms was demonstrated for concentrations of 250 μ g/mL at 48 h, and of 100 μ g/

Groups	Incubation period (h)	Number of dead worms (%)	Changes in motor activity
Control	120	–	+++
DMSO 1%	120	–	+++
PZQ	24	100	–
1	120	–	+++
5	120	–	+++
10	96	–	++
100	72	100	–
250	48	100	–
500	24	100	–
1000	24	100	–

Negative control (RPMI 1640; RPMI 1640 + DMSO 0.1%); positive control with 0.5 µg/mL. EOMv: +++ normal activity; ++ slight loss of movement, with activity of the tail, suckers, and gynaecophoric canal membrane; + movement of the tail and suckers alone; – no movement

Table 2 *In vitro* effect of different concentrations of EOMv on mortality of adult worms of *S. mansoni*.

Groups	Incubation period (h)	Number of dead worms (%)	Changes in motor activity
Control	120	–	+++
DMSO 1%	120	–	+++
PZQ	24	100	–
0.7	120	–	+++
3.54	120	–	+++
7.09	96	–	++
70.96	72	100	–
177.4	48	100	–
354.8	24	100	–
700.96	24	100	–

Negative control (RPMI 1640; DMSO + RPMI 1640); positive control with 0.5 µg/mL. Rotundifolone: +++ normal activity; ++ slight loss of movement, with activity of the tail, suckers, and gynaecophoric canal membrane; + movement of the tail and suckers alone; – no movement

Table 3 *In vitro* effect of different concentrations of rotundifolone on mortality of adult worms of *S. mansoni*.

Groups	Incubation period (h)	Number of dead worms (%)	Changes in motor activity
Control	120	–	+++
DMSO 1%	120	–	+++
PZQ	24	100	–
(1) 43.75 µg/mL	120	–	++
(2) 7.3 µg/mL	120	–	+++
(3) 4.03 µg/mL	120	–	+++

Negative control (RPMI 1640; RPMI 1640 + DMSO 0.1%); positive control with 0.5 µg/mL. +++ Normal activity; ++ slight loss of movement, with activity of the tail, suckers, and gynaecophoric canal membrane; + movement of the tail and suckers alone; – no movement

Table 4 *In vitro* effect of different concentrations of limonene (1), trans-caryophyllene (2), and β -pinene (3) on mortality of adult worms of *S. mansoni*.

mL at 72 h of observation, and similarly to our results, a dose-dependent effect was observed at the concentrations evaluated.

At the concentration of 70.96 µg/mL, mortality and absence of movement of the worms were observed from 72 h onwards, at the concentration of 177.4 µg/mL from 48 h, at the concentration of 354.8 µg/mL from 24 h, and at the concentration of 700.96 µg/mL within 24 h of exposure (Table 3).

Using rotundifolone at concentrations of 0.7, 3.54, and 7.09 µg/mL, no worm mortality was observed at any of the times examined. Regarding motility at a concentration of 7.09 µg/mL, minor loss of movement of the tail, suckers, and gynaecophoric canal membrane was observed after 96 h of exposure.

Using the limonene concentration of 43.75 µg/mL, no worm mortality was observed at any of the times examined. In relation to motility at this concentration, decreased motor activity was ob-

served after 96 h of exposure. The compounds β -pinene at 4.03 µg/mL and trans-caryophyllene at 7.3 µg/mL did not cause mortality of adult worms of *S. mansoni* at any time analyzed (Table 4). Furthermore, the worms in the negative control groups (RPMI 1640 and RPMI 1640 + DMSO 1.6%) remained viable throughout the analysis period. The positive control PZQ caused the death of all parasites after 24 h of evaluation, as seen in Tables 2, 3, and 4.

After studying the schistosomicidal effect of EOAc in that same study, the authors also evaluated the two major compounds of the essential oils precocene (1) and (E)-caryophyllene, in which schistosomicidal effects similar to that of EOAc were observed [31].

In that study, the major compound of EOMv (i.e., rotundifolone) demonstrated a schistosomicidal effect similar to EOMv [31].

Research and Training in Tropical Diseases, in which: (+ + +) indicates normal activity; (+ +) slight loss of movement, with active tail, suckers, and gynaecophoric canal membrane; (+) movement of tails and suckers alone; and (–) no movement. The worms were considered to be dead when no movement was identified after 3 min of observation using an inverted microscope [38].

Ethical standards

All experiments involving the use of animals were performed in accordance to the ethical standards of Fundação Oswaldo Cruz and were approved by the ethics committee (CEUA-FIOCRUZ, No. 06/2010).

Acknowledgements

We acknowledge LTF/UFPPB, LIKA/UFPE, and FIOCRUZ-PE for support for the experiments and CAPES for a scholarship.

Conflict of Interest

The authors declare that there is no conflict of interest.

Affiliations

- ¹ Molecular and Cell Biology Laboratory, Department of Parasitology, Aggeu Magalhães Research Center, Oswaldo Cruz Foundation, Recife, Pernambuco, Brazil
- ² Keizo Asami Immunopathology Laboratory, Federal University of Pernambuco, Recife, Pernambuco, Brazil
- ³ Federal University of Paraíba, João Pessoa, Paraíba, Brazil
- ⁴ Natural Products Laboratory, Research and Development Center for Plant and Phytochemical Genetic Resources, Campinas Agronomical Institute, Campinas, São Paulo, Brazil

References

- 1 Abdel-Hameed ES, El-Nahas HA, Abo-Sedera SA. Antischistosomal and antimicrobial activities of some Egyptian plant species. *Pharm Biol* 2008; 46: 626–633
- 2 Liu J, Hu W, Xu B, Wang JP, Wang SQ, Wang XN. Schistosomicidal mechanism of auranofin on *Schistosoma japonicum* and its cytotoxicity. *Zhongguo Ji Sheng Chong Xue Yu Ji Sheng Chong Bing Za Zhi* 2012; 30: 450–454
- 3 Allam G, Abuelsaad AS. *In vitro* and *in vivo* effects of hesperidin treatment on adult worms of *Schistosoma mansoni*. *J Helminthol* 2013; 9: 1–9
- 4 Carrara VS, Vieira SC, de Paula RG, Rodrigues V, Magalhães LG, Cortez DA, da Silva Filho AA. *In vitro* schistosomicidal effects of aqueous and dichloromethane fractions from leaves and stems of Piper species and the isolation of an active amide from *P. amalago* L. (Piperaceae). *J Helminthol* 2013; 8: 1–6
- 5 Ramalhetete C, Magalhães LG, Rodrigues V, Mulhovo S, Silva Filho AA, Ferreira MJU. *In Vitro* schistosomicidal activity of Balsaminol F and Karavilagenin C. *Planta Med* 2012; 78: 1912–1917
- 6 Moraes J, Almeida AAC, Brito MRM, Marques THC, Lima TC, Sousa DP, Nakano E, Mendonça RZ, Freitas RM. Anthelmintic activity of the natural compound (+)-limonene epoxide against *Schistosoma mansoni*. *Planta Med* 2013; 79: 253–258
- 7 Lorenzi H, Matos FJA. Plantas medicinais no Brasil: nativas e exóticas. Nova Odessa: Instituto Plantarum; 2002
- 8 Lahlou S, Carneiro-Leão RFL, Leal-Cardoso JH, Toscano CF. Cardiovascular effects of the essential oil of *Mentha x villosa* and its main constituent, piperitenone oxide, in normotensive anaesthetized rats: role of the autonomic nervous system. *Planta Med* 2001; 61: 638–643
- 9 Guedes DN, Silva DF, Barbosa-Filho JM, Medeiros IA. Calcium antagonism and the vasorelaxation of the rat aorta induced by rotundifolone. *Braz J Med Biol Res* 2004; 37: 1881–1887
- 10 Guedes DN, Silva DF, Barbosa-Filho JM, Medeiros IA. Endothelium-dependent hypotensive and vasorelaxant effects of the essential oil from aerial parts of *Mentha x villosa* in rats. *Phytotherapy* 2004; 11: 490–497
- 11 Arruda TA, Antunes RMP, Catão RMR, Lima EO, Sousa DP, Nunes XP, Pereira MSV, Barbosa-Filho JM, Cunha EMV. Estudo preliminar da atividade antimicrobiana do óleo essencial de *Mentha x villosa* Hudson, rotundifolona e seus análogos. *Rev Bras Farmacogn* 2006; 16: 307–311
- 12 Sousa PJC, Linard CFBM, Azevedo-Batista D, Oliveira AC, Coelho-de-Souza NA. Antinociceptive effects of the essential oil of *M. x villosa* leaf and its major constituent piperitone oxide in mice. *Braz J Med Biol Res* 2009; 42: 655–659
- 13 Almeida RN, Hiruma CA, Barbosa-Filho JM. Analgesic effect of rotundifolone in rodents. *Fitoterapia* 1996; 57: 334–338
- 14 Sousa PJC, Magalhães PJC, Lima CC, Oliveira VS, Leal-Cardoso JH. Effects of piperitone oxide on the intestinal smooth muscle of the guinea pig. *Braz J Med Biol Res* 1997; 30: 787–791
- 15 Guedes DN, Silva DF, Barbosa-Filho JM, Medeiros IA. Muscarinic agonist properties involved in the hypotensive and vasorelaxant responses of rotundifolone in rats. *Planta Med* 2002; 68: 700–704
- 16 Sousa DP, Júnior EVM, Oliveira FS, Almeida RN, Nunes XP, Barbosa-Filho JM. Antinociceptive activity of structure analogues of rotundifolone: structure-activity relationship. *Z Naturforsch* 2007; 62: 39–42
- 17 Sousa DP, Júnior GAS, Andrade LN, Calasans FR, Nunes XP, Barbosa-Filho JM, Batista JS. Structure and spasmolytic activity relationships of monoterpenes analogues found in many aromatic plants. *Z Naturforsch* 2009; 63: 808–812
- 18 Silva DF, Araújo IGA, Albuquerque JGF, Porto DL, Dias KLG, Cavalcante KVM, Veras RC, Nunes XP, Barbosa-Filho JM, Araújo DAM, Cruz JS, Correia NA, De Medeiros IA. Rotundifolone-induced relaxation is mediated by BK_{Ca} channel activation and Ca_v channel inactivation. *Basic Clin Pharmacol Toxicol* 2011; 109: 465–475
- 19 Matos FJA, Machado MI, Craveiro AA, Barbosa-Filho JM, Alencar JW, Cunha EVL, Hiruma CA. Essential oil of *Mentha x villosa* Huds. from Northeastern Brazil. *J Essent Oil Res* 1999; 11: 41–44
- 20 Teles NSB, Fachine FV, Viana FAC, Viana IOL, Nascimento DF, Leite ALAS, Frota Bezerra FA, Moraes MO, Moraes MEA. Evaluation of the therapeutic efficacy of *Mentha crispa* in the treatment of giardiasis. *Contemp Clin Trials* 2011; 32: 809–813
- 21 Moraes ME, Cunha GH, Bezerra MM, Fachine FV, Pontes AV, Andrade WS, Frota Bezerra FA, Moraes MO, Cavalcanti PP. Efficacy of the *Mentha crispa* in the treatment of women with *Trichomonas vaginalis* infection. *Arch Gynecol Obstet* 2012; 286: 125–130
- 22 Magalhães LG, Machado CB, Moraes ER, Moreira EBC, Soares CS, Da Silva SH, Da Silva Filho AA, Rodrigues V. *In vitro* schistosomicidal activity of curcumin against *Schistosoma mansoni* adult worms. *Parasitol Res* 2009; 104: 1197–1201
- 23 Fahmy ZH, El-Shennawy AM, El-Komy W, Ali E, Hamid SA. Potential antiparasitic activity of Pomegranate Extracts against schistosomules and mature worms of *Schistosoma mansoni*: *in vitro* and *in vivo* study. *Aust J Basic Appl Sci* 2009; 3: 4634–4643
- 24 Neves JK, Sarinho S, de Melo CM, Pereira VR, De Lima MC, Pitta IR, Albuquerque MC, Galdino SL. Immunological studies and *in vitro* schistosomicidal action of new imidazolidine derivatives. *J Venom Anim Toxins Incl Trop Dis* 2011; 17: 277–286
- 25 Moraes JD, Nascimento C, Lopes PO, Nakano E, Yamaguchi LF, Kato MJ, Kawano T. *Schistosoma mansoni*: *in vitro* schistosomicidal activity of piperitone. *Exp Parasitol* 2011; 127: 357–364
- 26 Magalhães LG, Kapadia GJ, Tonuci LRS, Caixeta SC, Parreira NA, Rodrigues V, Da Silva Filho AA. *In vitro* schistosomicidal effects of some phloroglucinol derivatives from *Dryopteris* species against *Schistosoma mansoni* adult worms. *Parasitol Res* 2010; 106: 395–401
- 27 Lima CMBL, Freitas FIS, Morais LCSL, Cavalcanti MGS, Silva LF, Padilha RJR, Barbosa CGS, Santos FAB, Alves LC, Diniz MFFM. Ultrastructural study on the morphological changes to male worms of *Schistosoma mansoni* after *in vitro* exposure to allicin. *Rev Soc Bras Med Trop* 2011; 29: 327–330
- 28 Miranda MA, Magalhães LG, Tiossi RFJ, Kuehn CC, Oliveira LGR, Rodrigues V, McChesney JD, Bastos JK. Evaluation of the schistosomicidal activity of the steroidal alkaloids from *Solanum lycocarpum* fruits. *Parasitol Res* 2012; 111: 257–262
- 29 Abdul-Ghani R, Loutfy N, El-Sahn A, Hassan A. Current chemotherapy arsenal for schistosomiasis mansoni: alternatives and challenges. *Parasitol Res* 2009; 104: 955–965
- 30 Parreira NA, Magalhães LG, Morais DR, Caixeta SC, De Sousa, JPB, Bastos JK, Cunha WR, Silva MLA, Nanayakkara NPD, Rodrigues V, Da Silva Filho AA. Antiprotozoal, schistosomicidal, and antimicrobial activities of the

- essential oil from the leaves of *Baccharis dracunculifolia*. Chem Biodivers 2010; 7: 993–1001
- 31 Melo NI, Magalhães LG, Carvalho CE, Wakabayashi KAL, Aguiar GP, Ramos RC, Mantovani ALL, Turatti ICC, Rodrigues V, Groppo M, Cunha WR, Veneziani RCS, Crotti AEM. Schistosomicidal activity of the essential oil of *Ageratum conyzoides* L. (Asteraceae) against adult *Schistosoma mansoni* worms. Molecules 2011; 16: 762–773
- 32 Caixeta SC, Magalhães LG, Melo NI, Wakabayashi KAL, Aguiar GP, Aguiar DP, Mantovani ALL, Morais JA, Oliveira PF, Tavares DC, Groppo M, Rodrigues V, Cunha WR, Veneziani RCS, Silva Filho AA, Crotti AE. Chemical composition and *in vitro* schistosomicidal activity of the essential oil of *Plectranthus neochilus* grown in Southeast Brazil. Chem Biodivers 2011; 8: 2149–2157
- 33 Magalhães LG, Souza JM, Wakabayashi KAL, Caixeta SC, Laurentiz RS, Vinhólis AHC, Rezende KCS, Simaro GV, Bastos JK, Rodrigues V, Esperandim VR, Ferreira DS, Crotti AEM, Cunha WR, Silva MLA. *In vitro* efficacy of the essential oil of *Piper cubeba* L. (Piperaceae) against *Schistosoma mansoni* adult worms. Parasitol Res 2011; 106: 395–401
- 34 Adams RP. Identification of essential oil components by gas chromatography/mass spectroscopy, 4th edition. Carol Stream: Allured Publ. Corp; 2007
- 35 Smithers SR, Terry RJ. Infection of laboratory hosts with cercariae of *Schistosoma mansoni* and the recovery of adult worms. Parasitology 1965; 55: 695–700
- 36 Almeida LMS, Farani LA, Tosta LA, Silvério MS, Sousa OV, Matos ACA, Coelho PMZ, Vasconcelos EG, Faria-Pinto P. *In vitro* evaluation of the schistosomicidal potential of *Eremanthus erythropappus* (DC) McLeisch (Asteraceae) extracts. Nat Prod Res 2011; 26: 1–7
- 37 Neves JK, Botelho SP, De Melo CM, Pereira VR, DE Lima MC, Pitta IR, Albuquerque MC, Galdino SL. Biological and immunological activity of new imidazolidines against adult worms of *Schistosoma mansoni*. Parasitol Res 2010; 107: 531–538
- 38 Ramirez B, Bickle Q, Yousif F, Fakorede F, Mouries MA, Nwaka S. Schistosomes: challenges in compound screening. Expert Opin Drug Discov 2007; 2: 53–61