

Inactivation of *Escherichia coli* O157:H7 on Radish Seeds by Sequential Treatments with Chlorine Dioxide, Drying, and Dry Heat without Loss of Seed Viability[∇]

Jihyun Bang,¹ Haeyoung Kim,¹ Hoikyung Kim,² Larry R. Beuchat,³ and Jee-Hoon Ryu^{1*}

Graduate School of Life Sciences and Biotechnology, Korea University, Anam-dong, Sungbuk-ku, Seoul 136-701, Republic of Korea¹; Division of Human Environmental Sciences, Wonkwang University, Shinyong-dong, Iksan, Jeonbuk 570-749, Republic of Korea²; and Center for Food Safety and Department of Food Science and Technology, University of Georgia, 1109 Experiment Street, Griffin, Georgia 30223-2797³

Received 3 June 2011/Accepted 19 July 2011

We developed and validated a treatment to inactivate *Escherichia coli* O157:H7 on radish seeds without decreasing seed viability. Treatments with aqueous ClO₂ followed by drying and dry-heat treatments were evaluated for efficacy to inactivate the pathogen. Conditions to dry radish seeds after treatment with water (control) or ClO₂ were established. When treated seeds with high water activity (a_w) (>0.99) were stored at 45°C and 23% relative humidity (RH), the a_w decreased to <0.30 within 24 h. Drying high-a_w seeds before exposing them to dry-heat treatment (≥60°C) was essential to preserve seed viability. The germination rate of radish seeds which had been immersed in water for 5 min, dried at 45°C and 23% RH for 24 h, and heated at 70°C for 48 h or at 80°C for 24 h was not significantly decreased (*P* ≤ 0.05) compared to that of untreated radish seeds. Sequential treatments with ClO₂ (500 µg/ml, 5 min), drying (45°C, 23% RH, 24 h), and dry heating (70°C, 23% RH, 48 h) eliminated *E. coli* O157:H7 (5.9 log CFU/g) on radish seeds and, consequently, sprouts produced from them without decreasing the germination rate. These sequential treatments are recommended for application to radish seeds intended for sprout production.

In the Republic of Korea and some other Asian countries, consumption of vegetable seed sprouts has increased in recent decades. Worldwide, the number and frequency of sprout-associated outbreaks of disease have increased during this period. There were at least 40 outbreaks implicating vegetable sprouts reported between 1973 and 2006 (30). The majority of these outbreaks were linked to alfalfa, mung bean, clover, radish, mustard, and cress sprouts (20, 23, 29). Pathogens most frequently involved in causing outbreaks were *Salmonella* and *Escherichia coli* O157:H7 (20, 22, 29). The largest outbreak was associated with radish sprouts in Japan (11). This outbreak involved more than 6,000 culture-confirmed cases of *E. coli* O157:H7 infections.

The source of pathogenic bacteria on sprouts is thought to originate largely from seeds rather than the contamination of sprouts during or after production (20). Thus, the National Advisory Committee on Microbiological Criteria for Foods (NACMCF) has recommended applying treatments to achieve a 5-log CFU/g reduction of pathogens on seeds. Even a 5-log CFU/g reduction, however, may not guarantee the absence of pathogens in sprouts. Pathogens that remain on seeds after treatment, even if present in very low numbers, can multiply rapidly to high levels during the sprouting process (1, 12, 19, 20). The goal of decontamination treatments should be to eliminate food-borne pathogens on seeds intended for sprout production.

Many studies have evaluated the effectiveness of various sanitizers, such as hypochlorites, organic acids, ozonated water, ethanol, and hydrogen peroxide, for reducing and eliminating *Salmonella* and *E. coli* O157:H7 on seeds (5, 9, 10, 17, 18, 24, 25, 27, 28). However, most treatments of seeds with a single chemical solution have not consistently reduced populations of pathogens by more than 3 log CFU/g (29). Noted exceptions are mung bean decontamination by treatment with acetic acid vapor at 45°C for 12 h (7) and elimination of *E. coli* O157:H7 and *Salmonella* in mung beans, soybeans, alfalfa seeds, and cress seeds without decreasing germination yields by treating with an oxychloro-based sanitizer (16). To achieve greater reductions in numbers of food-borne pathogens, sequential or simultaneous treatments with chlorine-based sanitizers, organic acids, heat, high pressure, and irradiation have been evaluated (6, 13, 17, 21, 23, 31). Although most studies using multiple treatments have been shown to result in greater reductions in pathogens than those using a single treatment, with some exceptions (3, 4), elimination of pathogens without decreasing the germination rate of seeds has been difficult. Bari et al. (4) reported that *E. coli* O157:H7 was eliminated from alfalfa, mung bean, and radish seeds without decreasing the germination rate and yield by applying dry heat (50°C for 1 h) and irradiation (2.5 kGy).

In a recent study, we observed a synergistic lethal effect of ClO₂ treatment (50 or 200 µg/ml, 5 min) and subsequent air drying (25°C, 40% relative humidity [RH], 24 h) in killing *E. coli* O157:H7 on radish seeds (14). In a follow-up study, a combination of ClO₂ (500 µg/ml, 5 min), air drying (25°C, 40% RH, 2 h), and dry-heat (55°C, 23% RH, 36 h) treatments reduced the total aerobic bacteria (TAB) count by >5 log CFU/g and *E. coli* O157:H7 by >4.8 log CFU/g without sig-

* Corresponding author. Mailing address: Graduate School of Life Sciences and Biotechnology, Korea University, Anam-dong, Sungbuk-ku, Seoul 136-701, Republic of Korea. Phone: 82 2 3290 3409. Fax: 82 2 3290 3918. E-mail: escheri@korea.ac.kr.

[∇] Published ahead of print on 29 July 2011.

nificantly decreasing seed viability (1). In the research reported here, we extended the latter study with the aim of eliminating *E. coli* O157:H7 on radish seeds without substantially decreasing the germination rate. We demonstrated the importance of the drying procedure between ClO₂ and dry-heat treatments in preserving seed viability. Conditions for dry-heat treatment to eliminate *E. coli* O157:H7 without substantially reducing the viability of seeds were established.

MATERIALS AND METHODS

Bacterial strains and preparation of inoculum. Five strains of *E. coli* O157:H7 were used: ATCC 43895 (isolated from hamburger), E0018 (isolated from bovine feces), F4546 (isolated from a patient in an alfalfa sprout-associated outbreak), H1730 (isolated from a lettuce-associated outbreak), and 932 (isolated from a patient with hemorrhagic colitis). We prepared the inoculum as described in earlier studies (1, 14) with some modifications. In brief, *E. coli* O157:H7 strains were adapted to grow in tryptic soy broth (TSB; Difco, BD Diagnostics, Sparks, MD) containing 50 µg/ml nalidixic acid (TSBN) at 37°C. After three consecutive transfers at 24-h intervals, 150 ml of a five-strain cocktail was prepared by combining 30 ml of culture of each strain. The cocktail (150 ml) was centrifuged at 2,000 × g for 15 min at 25°C. The supernatant was decanted, and cells in the pellet were resuspended in sterile distilled water (1,500 ml) to give a population of ca. 8 log CFU/ml.

Inoculation of *E. coli* O157:H7 on radish seeds. Radish seeds (350 g) purchased from Saessakmart (Seoul, Republic of Korea) were immersed in 1,050 ml of *E. coli* O157:H7 suspension (ca. 8 log CFU/ml) with gentle swirling for 5 min at 25 ± 2°C. Seeds were then placed on a sterile sieve (203-mm diameter by 41-mm depth; 600-µm pore size) and held for 2 h at 25 ± 2°C in a laminar flow biosafety hood before using in experiments.

Establishment of a 23% RH environment. To create an atmosphere with 23% RH, 250 ml of saturated, filter-sterilized (bottle-top filter, 0.2-µm pore size; Corning Costar, Lowell, MA) potassium acetate (Sigma-Aldrich Inc., Milwaukee, WI) solution was deposited in a propylene container (2.1 liters; 25 cm long by 18 cm wide by 8 cm high; Lock & Lock, Seoul, Republic of Korea). The container was sealed with polyethylene film (Seven Wrap; Cleanson, Seoul, Republic of Korea) and incubated at 45, 60, 70, or 80°C for at least 48 h before using in experiments.

Preparation of ClO₂ solution. An aqueous solution of ClO₂ was prepared by combining 450 ml of sodium chlorite solution (10,000 µg/ml) with 21 ml of hydrochloric acid (1 N) and incubating the mixture at 25 ± 2°C for 1 h. The solution was diluted in sterile distilled water to give a ClO₂ concentration of 500 µg/ml. The pH of the ClO₂ solution was 8.5 ± 0.4. The concentration of ClO₂ was measured immediately before experiments using a chlorine colorimeter (model Dr/820; Hach, Loveland, CO).

Determination of germination rate. Radish seeds (*n* = 100) were placed on sterile cheesecloth in a commercial sprout cultivator (225 by 325 by 150 mm; Shinhan Innovation & Creative, Suwon, Republic of Korea) containing sterile distilled water. The seeds were incubated at 25°C for 5 days, and the number of seeds that germinated and grew normally was counted. The germination percentage was calculated.

Optimization of temperature and time for drying seeds with high a_w. Radish seeds (40 g) were immersed in sterile distilled water (120 ml) with intermittent swirling for 5 min, spread on the surface of a sterile sieve (88.9-mm diameter by 41-mm depth; 600-µm pore size), placed above the surface of saturated potassium acetate solution (140 ml) in a propylene container (1.2 liters; 16 cm long by 16 cm wide by 9 cm high; Lock & Lock), and incubated at 25°C or 45°C for up to 48 h. After drying for 4, 8, 12, 24, 36, and 48 h, the water activity (a_w) of seeds (3 g) was measured using a water activity meter (AquaLab Series 3TE; Decagon Devices, Inc., Pullman, WA).

Optimization of temperature and time for dry-heat treatments. Radish seeds (40 g) were immersed in sterile distilled water (120 ml) for 5 min with gentle swirling. Seeds were then placed on a sterile sieve (88.9-mm diameter by 41-mm depth; 600-µm pore size), dried at 45°C in air containing 23% RH for 24 h, and heated at 60, 70, or 80°C and 23% RH for up to 48 h. Radish seeds (40 g) were also treated with water (120 ml) for 5 min and, without drying at 45°C, heated at 60, 70, or 80°C and 23% RH for up to 48 h. Germination percentages were determined after dry-heat treatment for 24 and 48 h.

Inactivation of *E. coli* O157:H7 on seeds by sequential treatments with ClO₂, drying, and dry heat. Immersion-inoculated seeds held for 2 h at 25 ± 2°C contained *E. coli* O157:H7 at a population of 5.9 log CFU/g. Seeds (220 g) were

immersed in 660 ml of sterile distilled water or ClO₂ solution (500 µg/ml) in a sterile glass bottle for 5 min, with intermittent swirling, and rinsed twice in sterile distilled water (660 ml) for 1 min. Treated seeds (40 g) were placed on a sterile sieve (88.9-mm diameter by 41-mm depth; 600-µm pore size) and positioned on a rack above 250 ml of saturated potassium acetate in a propylene container. The container was sealed with plastic wrap and held at 45°C at an internal RH of 23% for 24 h. After drying, seeds were incubated at 70°C and 23% RH for 24 and 48 h or at 80°C and 23% RH for 6, 12, 24, and 48 h.

Microbiological analyses of seeds. Populations of total aerobic bacteria (TAB), *E. coli* O157:H7, and molds and yeasts (MY) on radish seeds were determined before treatment with water (control) or ClO₂ (0 h), after treatment with water or ClO₂ for 5 min, after drying (45°C, 23% RH, 24 h), and after dry-heat treatment (70°C, 23% RH, 24 and 48 h; 80°C, 23% RH, 6, 12, 24, and 48 h). At each sampling time, seeds (5 g) were deposited in TSB (45 ml) in a polyolefin stomacher bag (400 ml; Interscience, St. Nom La Breteche, France) and pummeled for 1 min. The TSB in the TSB-seed mixtures was serially diluted in 0.1% peptone water (or not diluted) and surface plated on tryptic soy agar (TSA) for enumerating TAB, MacConkey sorbitol (Difco, BD Diagnostics) agar supplemented with nalidixic acid (50 µg/ml) (MSAN) for enumerating *E. coli* O157:H7 cells, and dichloran rose bengal chloramphenicol (DRBC; Difco, BD Diagnostics) agar for enumerating MY. TSA and MSAN plates were incubated at 37°C for at least 24 h, and DRBC plates were incubated at 25°C for 5 days before colonies were counted. For seeds heated at 70°C for 24 or 48 h or at 80°C for 6, 12, 24, or 48 h, mixtures of seeds and TSB were incubated at 37°C for 48 h to enrich for *E. coli* O157:H7. The enriched suspension was streaked on TSA and MSAN and incubated at 37°C for 24 h. Colonies presumptive for *E. coli* O157:H7 that formed on TSA and MSAN were randomly selected and tested using an *E. coli* O157:H7 latex agglutination test (Oxoid, Basingstoke, United Kingdom). The detection limit by direct plating was 9 CFU/g of seeds (0.95 log CFU/g); the limit by enrichment was 1 CFU/5 g of seeds (-0.70 log CFU/g). Germination percentages were determined after dry-heat treatment, as described above.

Microbiological analysis of sprouts. Radish seeds (20 g) treated with ClO₂ solution (500 µg/ml) for 5 min, dried at 45°C and 23% RH for 24 h, and dry heated at 70°C and 23% RH for 24 h and 48 h or at 80°C and 23% RH for 6, 12, 24, and 48 h were soaked in sterile water at 35°C for 2 h, placed in a commercial sprout cultivator, and incubated at 25°C for 5 days. Sprouts (10 g) were aseptically collected, combined with 90 ml of TSB in a stomacher bag, and pummeled for 1 min. The homogenate was serially diluted in sterile 0.1% peptone, surface plated (0.1 ml in duplicate) on TSA, MSAN, and DRBC agar, and incubated at 37°C for 24 h (TSA, MSAN) or 25°C for 5 days (DRBC agar). To enrich for *E. coli* O157:H7 on sprouts, the remaining sprout and TSB mixture was incubated at 37°C for 48 h, streaked on TSA and MSAN, and incubated at 37°C for 48 h. Several colonies that formed on TSA and MSAN were randomly selected and tested for *E. coli* O157:H7 using an *E. coli* O157:H7 latex agglutination test. The detection limit by direct plating was 10 CFU/g sprout (1.0 log CFU/g); the limit by enrichment was 1 CFU/10 g sprout (-1.0 log CFU/g).

Statistical analysis. All experiments were replicated at least three times. Data were analyzed using the general linear model of the Statistical Analysis Systems procedure (SAS; SAS Institute, Cary, NC). Analysis to determine the effects of drying, heating temperature, and heating time on germination rate and the effect of sequential ClO₂, drying, and dry-heat treatments on TAB, *E. coli* O157:H7, and MY populations recovered from radish seeds and sprouts was done using Fisher's least significant difference (LSD) test. Significant differences are presented at a 95% confidence level (*P* ≤ 0.05).

RESULTS

Optimization of temperature and time for drying seeds with high a_w. To minimize the adverse effect of wet heat on seed viability, radish seeds with high a_w were dried before being exposed to dry-heat treatment. Figure 1 shows the a_w of radish seeds which were immersed in water for 5 min and incubated at 25°C or 45°C and 23% RH for 24 or 48 h. The a_w of seeds was >0.99 after treatment with water for 5 min. When these seeds were stored at 25°C and 23% RH, the a_w decreased to <0.30 within 48 h. When stored at 45°C, the a_w was <0.30 within 24 h and remained constant for an additional 24 h.

Optimization of temperature and time for dry-heat treatments. Tests were done to determine dry-heat conditions that minimize decreases in the germination rate of radish seeds.




FIG. 1. Water activity of radish seeds dried at 25°C (○) or 45°C (□). Radish seeds were immersed in water for 5 min and incubated at 25 or 45°C for up to 48 h at 23% relative humidity (RH).

The influence of drying seeds between immersing in water and dry-heat treatment on the germination rate was also evaluated. Table 1 shows the germination rate of radish seeds as affected by drying, dry-heat temperature, and dry-heat treatment time. Seeds immersed in water for 5 min and, without drying, heated at 60°C and 23% RH for 24 or 48 h had germination rates of 57.3 and 63.7%, respectively. When radish seeds were immersed in water for 5 min, dried at 45°C and 23% RH for 24 h, and heated at 60°C and 23% RH for 24 or 48 h, germination rates were 87.7 or 84.5%, respectively. When water-treated radish seeds were heated without drying at 70°C and 23% RH for 24 or 48 h, germination rates were only 35.0 or 31.7%, respectively. However, when seeds were dried between water treatment and dry-heat treatment, at 70°C for 24 or 48 h, the germination rate was 84.3%. When the dry-heat temperature was increased to 80°C, the germination rate of seeds was 89.0% after dry-heat treatment for 24 h but decreased significantly to 69.5% after treatment for 48 h.

Inactivation of *E. coli* O157:H7 on seeds by sequential treatments with ClO₂, drying, and dry heat. The effects of the sequential treatments (ClO₂, drying, and dry heat at 70°C or 80°C) on populations of microorganisms on radish seeds and germination rates were determined.

Table 2 shows the TAB, *E. coli* O157:H7, and MY populations on seeds treated with water or ClO₂ (500 µg/ml) for 5 min, dried at 45°C and 23% RH for 24 h, and dry heated at 70°C and 23% RH for 24 or 48 h. The initial populations of TAB, *E. coli* O157:H7, and MY on seeds (6.1, 5.9, and 3.5 log CFU/g, respectively) were not significantly reduced by treatment with water for 5 min. Drying seeds at 45°C did not significantly reduce TAB and *E. coli* O157:H7 populations, but the number of MY was significantly lower (1.0 log CFU/g) than the number recovered from seeds treated with water. Subsequent dry-heat treatment at 70°C and 23% RH for 48 h decreased the populations of TAB and *E. coli* O157:H7 to 4.1 and 3.4 log CFU/g, respectively, but the population of MY did not significantly change.

When inoculated radish seeds were treated with ClO₂ (500 µg/ml) for 5 min, populations of TAB, *E. coli* O157:H7, and MY were significantly decreased from 6.1, 5.9, and 3.5 log CFU/g to 5.3, 5.0, and 1.7 log CFU/g, respectively (Table 2). When the ClO₂-treated seeds were dried at 45°C and 23% RH

for 24 h, populations of TAB and *E. coli* O157:H7 significantly decreased by >4.0 log CFU/g. The *E. coli* O157:H7 population on seeds was decreased to an undetectable level (<0.95 log CFU/g) by direct plating but was detected by enrichment (≥1 CFU/5 g). The number of MY was not significantly decreased by drying seeds at 45°C. Dry-heat treatment of seeds at 70°C and 23% RH for 48 h caused TAB and MY to decrease to levels approaching the detection limit by direct plating. *E. coli* O157:H7 was not detected by enrichment in seeds exposed to dry heat for 48 h.

Figure 2A shows the germination rates of radish seeds initially containing *E. coli* O157:H7 (5.9 log CFU/g) after sequential treatments with water or ClO₂ (500 µg/ml, 5 min), drying (45°C, 23% RH, 24 h), and dry heat (70°C, 23% RH, 24 or 48 h). The germination rate of seeds exposed to the harshest conditions (ClO₂ [500 µg/ml, 5 min], drying [45°C, 23% RH, 24 h], and dry heating [70°C, 23% RH, 48 h]) was not significantly different than that of untreated radish seeds.

Table 3 shows the populations of TAB, *E. coli* O157:H7, and MY on radish sprouts cultivated at 25°C for 5 days using seeds which had been subjected to sequential treatments with ClO₂ (500 µg/ml, 5 min) followed by drying (45°C, 23% RH, 24 h) and dry heat (70°C, 23% RH, 24 or 48 h). Populations of TAB on radish sprouts were 7.4 to 8.4 log CFU/g, regardless of treatment with water or ClO₂ or heating time applied to seeds. Sprouts produced using seeds that had been exposed to ClO₂, drying, and dry-heat treatments were negative for *E. coli* O157:H7 by enrichment. However, sprouts cultivated from seeds that had been treated with water rather than ClO₂ and dry heated for 48 h contained 6.1 log CFU/g; sprouts produced from seeds treated with ClO₂ but heated for only 24 h contain *E. coli* O157:H7 at 7.2 log CFU/g. These results showed that *E. coli* O157:H7 can rapidly increase to a high population during cultivation, even after significant reduction on treated seeds. Sprouts produced from radish seeds treated with water had MY counts of 7.8 to 8.2 log CFU/g. Sprouts produced from seeds treated with ClO₂ and subsequently subjected to dry-heat treatments contained only 3.0 to 3.9 log CFU/g.

Shown in Table 4 are populations of TAB, *E. coli* O157:H7, and MY on radish seeds treated with water or ClO₂ (500

TABLE 1. Effects of drying, dry-heat temperature, and heating time on the germination rate of radish seeds^a

Dry-heat temp (°C)	Dry-heat time (h)	Germination rate ± SD (%) ^b	
		Without drying	With drying
60	24	a 57.3 ± 9.0 B	a 87.7 ± 2.3 A
	48	a 63.7 ± 4.7 B	a 84.5 ± 5.1 A
70	24	a 35.0 ± 15.7 B	a 84.3 ± 7.6 A
	48	a 31.7 ± 11.5 B	a 84.3 ± 4.6 A
80	24	a 14.5 ± 12.0 B	a 89.0 ± 2.8 A
	48	a 0.0 ± 0.0 B	b 69.5 ± 4.9 A

^a Seeds were immersed in water for 5 min, dried at 45°C and 23% RH for 24 h (or not dried), and incubated at 60, 70, or 80°C and 23% RH for 24 or 48 h before the germination rates were determined. The germination rate of untreated radish seeds (control) was 89.3 ± 6.7%.

^b Values in the same row that are not followed by the same uppercase letter are significantly different ($P \leq 0.05$). Within the same temperature, values in the same column that are not preceded by the same lowercase letter are significantly different ($P \leq 0.05$).

TABLE 2. Populations of TAB, *E. coli* O157:H7, and MY on radish seeds treated with water or ClO₂ (500 µg/ml) for 5 min, dried at 45°C and 23% RH for 24 h, and dry heated at 70°C and 23% RH for 24 or 48 h

Microorganism	Water or ClO ₂ treatment	Population ± SD (log CFU/g) ^a				
		Treatment time		Dried at 45°C for 24 h	Dry-heat treatment at 70°C	
		0 h	5 min		24 h	48 h
TAB	Water	a 6.1 ± 0.2 A	a 5.7 ± 0.1 A	a 5.8 ± 0.6 A	a 4.8 ± 0.5 B	a 4.1 ± 0.2 C
	ClO ₂	a 6.1 ± 0.2 A	b 5.3 ± 0.1 B	b <1.3 ± 0.3 C	b <1.0 ± 0.0 C	b <1.4 ± 0.4 C
<i>E. coli</i> O157:H7	Water	a 5.9 ± 0.2 A	a 5.5 ± 0.1 AB	a 5.0 ± 0.8 B	a 4.2 ± 0.4 C	a 3.4 ± 0.2 D
	ClO ₂	a 5.9 ± 0.2 A	b 5.0 ± 0.1 B	b <1.0 (3/3) ^b C	b <1.0 (3/3) C	b <1.0 (0/3) C
MY	Water	a 3.5 ± 0.5 A	a 3.3 ± 0.3 A	a 2.3 ± 0.1 B	a 2.0 ± 0.6 B	a 1.7 ± 0.4 B
	ClO ₂	a 3.5 ± 0.5 A	b 1.7 ± 0.3 B	b 1.1 ± 0.2 B	a <1.6 ± 0.7 B	a <1.2 ± 0.2 B

^a Values in the same row that are not followed by the same uppercase letter are significantly different (*P* ≤ 0.05). Within the same microorganism, values in the same column that are not preceded by the same lowercase letter are significantly different (*P* ≤ 0.05).

^b None detected by direct plating. Values in parentheses represent the number of samples out of three analyzed in three replicate trials that were positive for *E. coli* O157:H7 as determined by enrichment. Detection limit by direct plating was 9 CFU/g of seeds; detection limit by enrichment was 1 CFU/5 g of seeds.

µg/ml) for 5 min, dried at 45°C and 23% RH for 24 h, and dry heated at 80°C and 23% RH for 6, 12, 24, or 48 h. The initial populations of TAB, *E. coli* O157:H7, and MY on radish seeds were 6.3, 5.9, and 5.2 log CFU/g, respectively. The populations of TAB, *E. coli* O157:H7, and MY on seeds after treatment with water and drying for 24 h at 45°C decreased significantly

to 4.8, 4.1, and 3.3 log CFU/g, respectively. Dry-heat treatment at 80°C and 23% RH for 48 h reduced the population of TAB to 2.2 log CFU/g; *E. coli* O157:H7 and MY populations were reduced to levels below the detection limit (<0.95 log CFU/g) by direct plating but were detected by enrichment (≥1 CFU/5 g). The number of TAB on seeds treated with ClO₂ for 5 min and dried for 24 h significantly decreased from 6.3 log CFU/g to 1.9 log CFU/g. *E. coli* O157:H7 and MY counts were significantly decreased from 5.9 and 5.2 log CFU/g, respectively, to below the detection limit for direct plating; however, both were detected by enrichment. When seeds were subsequently dry heated at 80°C and 23% RH for 6, 12, 24, or 48 h, TAB and MY populations decreased to numbers approaching the detection limit by direct plating. *E. coli* O157:H7 was detected by enrichment of seeds dry heated at 80°C for 6, 12, or 24 h. When the dry-heating time was extended to 48 h, the pathogen was not detected by enrichment.





FIG. 2. Germination rate of radish seeds after sequential ClO₂, drying, and heat treatments. Radish seeds were immersed in water (white bars) or 500 µg/ml of ClO₂ (gray bars) for 5 min, dried at 45°C and 23% relative humidity (RH) for 24 h, and dry heated at 70°C (A) or 80°C (B) and 23% RH for 24 or 48 h. Bars indicate standard deviations. For the same dry-heat treatment time, bars not noted by the same letter are significantly different (*P* ≤ 0.05).

TABLE 3. Populations of TAB, *E. coli* O157:H7, and MY on radish sprouts^a

Microorganism	Water or ClO ₂ treatment	Population ± SD (log ₁₀ CFU/g) ^b after dry-heat treatment	
		24 h	48 h
TAB	Water	a 8.4 ± 0.4 A	a 7.8 ± 0.5 A
	ClO ₂	a 7.6 ± 0.3 A	a 7.4 ± 1.0 A
<i>E. coli</i> O157:H7	Water	a 6.7 ± 0.2 A	a 6.1 ± 0.5 A
	ClO ₂	a 7.2 ± 0.6 A	b <1.0 (0/3) ^c B
MY	Water	a 8.2 ± 0.5 A	a 7.8 ± 0.3 A
	ClO ₂	b 3.9 ± 0.6 A	b 3.0 ± 2.6 A

^a Radish seeds inoculated with *E. coli* O157:H7 (5.9 log CFU/g) were treated with water or ClO₂ (500 µg/ml) for 5 min, dried at 45°C and 23% RH for 24 h, and dry heated at 70°C and 23% RH for 24 or 48 h. Treated radish seeds were cultivated at 25°C for 5 days.

^b Values in the same row that are not followed by the same uppercase letter are significantly different (*P* ≤ 0.05). Within the same microorganism, values in the same column that are not preceded by the same lowercase letter are significantly different (*P* ≤ 0.05).

^c None detected by direct plating. Values in parentheses represent the number of samples out of three analyzed in three replicate trials that were positive for *E. coli* O157:H7 as determined by enrichment. Detection limit by direct plating was 10 CFU/g of sprouts; detection limit by enrichment was 1 CFU/10 g of sprouts.

TABLE 4. Populations of TAB, *E. coli* O157:H7, and MY on radish seeds after treatment with water or ClO₂ (500 µg/ml) for 5 min, dried at 45°C and 23% RH for 24 h, and dry heated at 80°C and 23% RH for up to 48 h

Microorganism	Water or ClO ₂ treatment	Population ± SD (log ₁₀ CFU/g) ^a						
		Treatment time		Drying at 45°C for 24 h	Dry-heat treatment			
		0 h	5 min		6 h	12 h	24 h	48 h
TAB	Water	a 6.3 ± 0.4 A	a 6.2 ± 0.2 A	a 4.8 ± 0.3 B	a 4.4 ± 0.5 BC	a 4.4 ± 0.3 BC	a 4.0 ± 0.6 C	a 2.2 ± 0.2 D
		a 6.3 ± 0.4 A	a 4.9 ± 0.6 B	b 1.9 ± 0.9 C	b <1.0 (3/3) ^b D	b <1.0 (3/3) D	b <1.1 ± 0.2 D	b <1.0 (1/3) D
<i>E. coli</i> O157:H7	Water	a 5.9 ± 0.3 A	a 5.6 ± 0.3 A	a 4.1 ± 0.4 B	a 3.5 ± 0.4 B	a 3.0 ± 0.7 B	a 3.0 ± 1.4 B	a <1.0 (3/3) C
		a 5.9 ± 0.3 A	a 4.5 ± 0.6 B	b <1.0 (3/3) C	b <1.0 (2/3) C	b <1.0 (0/3) C	a <1.0 (1/3) C	a <1.0 (0/3) C
MY	Water	a 5.2 ± 0.2 A	a 3.6 ± 0.2 B	a 3.3 ± 0.4 B	a 2.9 ± 0.4 BC	a 2.5 ± 0.3 C	a <1.3 ± 0.6 D	a <1.0 (3/3) D
		a 5.2 ± 0.2 A	a 3.0 ± 1.3 B	b <1.0 ± 0.0 C	b <1.1 ± 0.1 C	b <1.1 ± 0.2 C	a <1.1 ± 0.2 C	a <1.1 ± 0.2 C

^a Values in the same row that are not followed by the same uppercase letter are significantly different ($P \leq 0.05$). Within the same microorganism, values in the same column that are not preceded by the same lowercase letter are significantly different ($P \leq 0.05$).

^b None detected by direct plating. Values in parentheses represent the number of samples out of three analyzed in three replicate trials that were positive for *E. coli* O157:H7 as determined by enrichment. Detection limit by direct plating was 9 CFU/g of seeds, and detection limit by enrichment was 1 CFU/5 g of seeds.

Figure 2B shows the germination rates of radish seeds previously inoculated with *E. coli* O157:H7 and subjected to sequential treatments (water or ClO₂ [500 µg/ml, 5 min], drying [45°C, 23% RH, 24 h], and dry heating [80°C, 23% RH, 6, 12, 24 or 48 h]). The germination rates of seeds exposed to water, drying, and dry-heat treatment were 82.7 to 91.3% and were not significantly different from that of untreated seeds (89.3%). However, when radish seeds were treated with ClO₂, dried, and dry heated, the germination rates were significantly lower (63.3 to 77.0%) than those of untreated seeds and seeds exposed to sequential treatments with water, drying, and dry heating.

Populations of TAB, *E. coli* O157:H7, and MY on radish sprouts cultivated at 25°C for 5 days after treatment of seeds with water or ClO₂ (500 µg/ml) for 5 min, dried at 45°C and 23% RH for 24 h, and dry heated at 80°C and 23% RH for 6, 12, 24, or 48 h are shown in Table 5. The number of TAB on radish sprouts was ca. 7.9 to 8.3 log CFU/g, regardless of treatment with water or ClO₂ or dry-heat time. When sprouts were produced using seeds which had been subjected to sequential treatments with ClO₂ (500 µg/ml, 5 min), drying (45°C, 23% RH, 24 h), and dry heat (80°C, 23% RH, 48 h), *E. coli* O157:H7 was not detected by enrichment. When radish seeds were treated with ClO₂, dried, and dry heated for 24 h or less, cultivated radish sprouts contained *E. coli* O157:H7 at

populations of 1.7 to 3.3 log CFU/g. The population of MY was 1.6 log CFU/g of sprouts produced from seeds treated with ClO₂ (500 µg/ml, 5 min), followed by drying (45°C, 23% RH, 24 h) and heating (80°C, 23% RH, 48 h). Populations of MY on sprouts produced from seeds exposed to other dry-heat treatments were 6.1 to 8.2 log CFU/g.

DISCUSSION

This study evolved from earlier observations that treatment of radish seeds with ClO₂ followed by drying and dry-heat treatment has a synergistic effect in killing *E. coli* O157:H7 (1, 2, 14). Kim et al. (14) reported that ClO₂ treatment had a synergistic lethal effect on *E. coli* O157:H7 when combined with air drying. However, the treatment did not achieve the 5-log CFU/g reduction in pathogens recommended by NACMCF (20). Thus, an additional dry-heat treatment was applied in a follow-up study to increase lethality (1). We examined dry-heat treatment for its efficacy in killing *E. coli* O157:H7 because it is known to be less detrimental to seed germination than wet-heat treatment (8, 26). Also, lethality of dry heat may not be affected by crevices or wrinkles on seed testae, which are thought to protect pathogens from contact with chemical treatment solutions (8). Bang et al. (1) reported that treatment of radish seeds with ClO₂ (500 µg/ml) for 5 min, air drying at

TABLE 5. Populations of TAB, *E. coli* O157:H7, and MY on radish sprouts^a

Microorganism	Water or ClO ₂ treatment	Population ± SD (log ₁₀ CFU/g) ^b after dry-heat treatment			
		6 h	12 h	24 h	48 h
TAB	Water	a 8.2 ± 0.2 AB	a 7.9 ± 0.2 B	a 8.1 ± 0.2 AB	a 8.3 ± 0.1 A
		a 8.3 ± 0.2 A	a 8.3 ± 0.2 A	a 8.3 ± 0.2 A	a 8.2 ± 0.0 A
<i>E. coli</i> O157:H7	Water	a 6.9 ± 0.8 A	a 6.6 ± 0.2 A	a 6.4 ± 0.2 AB	a 5.5 ± 0.6 B
		a 3.3 ± 2.5 A	a 3.1 ± 3.6 A	b 1.7 ± 1.1 A	b <1.0 (0/3) ^c A
MY	Water	a 8.0 ± 0.1 A	a 7.9 ± 0.2 A	a 8.1 ± 0.2 A	a 8.2 ± 0.2 A
		a 7.8 ± 0.5 A	a 6.1 ± 3.4 A	a 6.7 ± 1.6 A	b 1.6 ± 0.4 B

^a Radish seeds inoculated with *E. coli* O157:H7 (5.9 log CFU/g) were treated with water or ClO₂ (500 µg/ml) for 5 min, dried at 45°C and 23% RH for 24 h, and dry heated at 80°C and 23% RH for up to 48 h. After dry-heat treatment for 6, 12, 24, or 48 h, radish seeds were cultivated at 25°C for 5 days.

^b Values in the same row that are not followed by the same uppercase letter are significantly different ($P \leq 0.05$). Within the same microorganism, values in the same column that are not preceded by the same lowercase letter are significantly different ($P \leq 0.05$).

^c None detected by direct plating. Values in parentheses represent number of samples out of three analyzed in three replicate trials that were positive for *E. coli* O157:H7 as determined by enrichment. Detection limit by direct plating was 10 CFU/g of sprouts; detection limit by enrichment was 1 CFU/10 g of sprouts.

25°C for 2 h, and dry heating at 55°C and 23% RH for 36 h reduced TAB and *E. coli* O157:H7 counts by 5.1 log CFU/g and >4.8 log CFU/g, respectively. The *E. coli* O157:H7 population on radish seeds was reduced to levels below the detection limit by direct plating (0.8 log CFU/g). However, sprouts grown from treated seeds at 25°C for 5 days contained *E. coli* O157:H7 at ca. 4.3 log CFU/g. In a follow-up study, we attempted to enhance the lethality of sequential treatments by increasing the dry-heat temperature from 55°C to 60°C (2). We treated radish seeds containing *E. coli* O157:H7 at a population of 5.5 log CFU/g with ClO₂ (500 µg/ml) for 5 min, without drying, followed by dry-heat treatment at 60°C and 23% RH for up to 48 h (2). Using these treatments, *E. coli* O157:H7 was eliminated from radish seeds and sprouts produced from them; however, the germination rate of seeds was decreased significantly. We suspected that the high moisture content of ClO₂-treated seeds may have caused loss in germinability during the early stages of dry-heat treatment.

In the study reported here, we increased the lethality of sequential treatments (ClO₂, drying, and dry-heat treatments) without decreasing seed viability by optimizing conditions for drying and dry-heat treatment. We hypothesized that, when ClO₂ and dry-heat treatments are sequentially applied to radish seeds, the drying treatment after ClO₂ treatment (before dry-heat treatment) is critical to minimizing the adverse effect of wet heat on seed viability during the early stage of dry-heat treatment. To test this hypothesis, the influence of drying conditions on seed viability was further investigated. When radish seeds with high a_w were stored at 25 or 45°C with 23% RH, it took 48 h or 24 h, respectively, to decrease the a_w to <0.30. Because an extended drying period may be undesirable by sprout producers, we decided to dry seeds at 45°C for 24 h.

After establishing drying conditions, the effect of the drying procedure in preserving seed viability was confirmed, and the optimum dry-heat temperature and time were established. Dry-heat treatment of radish seeds with high a_w significantly decreased the germination rate. However, when seeds were dried at 45°C preceding dry heating, the germination rate was not compromised, even after treatment at 70°C for 48 h or at 80°C for 24 h. This indicated that drying seeds between ClO₂ and dry-heat treatments is essential to preserve seed viability. Based upon these results, 70°C for 24 and 48 h and 80°C for 6, 12, 24, and 48 h were selected as conditions for dry-heat treatment. Sequential treatments were applied to radish seeds containing *E. coli* O157:H7 (5.9 log CFU/g). When radish seeds were treated with ClO₂ (500 µg/ml, 5 min), dried (45°C, 23% RH, 24 h), and dry heated (70°C, 23% RH, 48 h), the pathogen was not detected in seeds, even after enrichment.

The germination rate (84.7%) of radish seeds that had been exposed to the sequential treatments was not significantly different from that of untreated radish seeds (89.3%). *E. coli* O157:H7 was not detected in sprouts produced from those seeds. However, when the dry-heating time was reduced to 24 h, sprouts contained *E. coli* O157:H7 at 7.2 log CFU/g. These results indicated that, even if the population of *E. coli* O157:H7 on radish seeds was reduced to a very low level, the population after cultivation of those seeds could be high. Similar results have been reported by several researchers (1, 12, 19, 20). This emphasizes the need to eliminate *E. coli* O157:H7 from seeds used to produce sprouts. As the number of MY on

radish seeds was reduced, final populations on sprouts also decreased significantly.

Based on these results, we conclude that sequential treatment of radish seeds with ClO₂ (500 µg/ml, 5 min), drying (45°C, 23% RH, 24 h), and dry heating (70°C, 23% RH, 48 h) inactivates *E. coli* O157:H7 at populations of at least 5.9 log CFU/g without significantly decreasing the germination rate. To determine if an increase in dry-heating temperature could be used without lowering the germination rate, seeds were dry heated at 80°C. Results suggest that that the increased temperature did not substantially decrease the time required to eliminate *E. coli* O157:H7 on radish seeds. In addition, the germination rate of radish seeds treated with ClO₂ decreased significantly by dry-heat treatment at 80°C compared to that of seeds that had been treated with water. We concluded that treatment at 70°C was superior to 80°C as a dry-heating temperature to preserve seed viability.

In summary, sequential treatments to eliminate *E. coli* O157:H7 from radish seeds without decreasing the germination rate have been developed. The optimum conditions for drying radish seeds with high a_w were established. The effects of drying in combination with treatments with aqueous ClO₂ and dry heat in preserving seed viability were determined. Finally, we confirmed that high numbers of *E. coli* O157:H7 (5.9 log CFU/g) were eliminated on radish seeds and sprouts produced from them by applying sequential treatments of ClO₂ (500 µg/ml, 5 min), drying (45°C, 23% RH, 24 h), and dry heating (70°C, 23% RH, 48 h).

In future studies, conditions of ClO₂ treatment, such as concentration and treatment time, should be optimized. Decreasing the concentration of ClO₂ may allow the use of a higher dry-heat temperature to eliminate *E. coli* O157:H7 on radish seeds without substantially decreasing the germination rate. The use of organic acid-based ClO₂ solution at pH 5 to 6 should also be considered, since its lethality may be better than that of HCl-based ClO₂ (15). The efficacy of the decontamination procedure developed in this study should be validated using commercial-scale sprout production practices. The treatment should also be tested for efficacy in eliminating *Salmonella* on radish seeds.

ACKNOWLEDGMENTS

This study was carried out with the support of the Cooperative Research Program for Agricultural Science and Technology Development (project no. PJ007387), Rural Development Administration, Republic of Korea, and a Korea University grant.

REFERENCES

- Bang, J., H. Kim, H. Kim, L. R. Beuchat, and J.-H. Ryu. 2011. Combined effects of chlorine dioxide, drying, and dry heat treatments in inactivating microorganisms on radish seeds. *Food Microbiol.* **28**:114–118.
- Bang, J., et al. 2011. Reduction of *Escherichia coli* O157:H7 on radish seeds by sequential application of aqueous chlorine dioxide and dry heat treatment. *Lett. Appl. Microbiol.* [Epub ahead of print.] doi:10.1111/j.1472-765X.2011.03125.x.
- Bari, M. L., D. Nei, K. Enomoto, S. Todoriki, and S. Kawamoto. 2009. Combination treatments for killing *Escherichia coli* O157:H7 on alfalfa, radish, broccoli, and mung bean seeds. *J. Food Prot.* **72**:631–636.
- Bari, M. L., E. Nazuka, Y. Sabina, S. Todoriki, and K. Ishiki. 2003. Chemical and irradiation treatments for killing *Escherichia coli* O157:H7 on alfalfa, radish, and mung bean seeds. *J. Food Prot.* **66**:767–774.
- Beuchat, L. R. 1997. Comparison of chemical treatments to kill *Salmonella* on alfalfa seeds destined for sprout production. *Int. J. Food Microbiol.* **34**:329–333.
- Beuchat, L. R., and A. J. Scouten. 2002. Combined effects of water activity,

- temperature and chemical treatments on the survival of *Salmonella* and *Escherichia coli* O157:H7 on alfalfa seeds. *J. Appl. Microbiol.* **92**:382–395.
7. **Delaquis, P. J., P. L. Sholberg, and K. Stanich.** 1999. Disinfection of mung bean seed with gaseous acetic acid. *J. Food Prot.* **62**:953–957.
 8. **Feng, G., J. J. Churey, and R. W. Worobo.** 2007. Thermal inactivation of *Salmonella* and *Escherichia coli* O157:H7 on alfalfa seeds. *J. Food Prot.* **70**:1698–1703.
 9. **Gabriel, A. A.** 2005. Microbial quality of chlorine soaked mung bean seeds and sprouts. *Food Sci. Technol. Res.* **11**:95–100.
 10. **Gandhi, M., and K. R. Matthews.** 2003. Efficacy of chlorine and calcinated calcium treatment of alfalfa seeds and sprouts to eliminate *Salmonella*. *Int. J. Food Microbiol.* **87**:301–306.
 11. **Gutierrez, E.** 1997. Japan prepares as O157 strikes again. *Lancet* **349**:1156.
 12. **Holliday, S. L., A. J. Scouten, and L. R. Beuchat.** 2001. Efficacy of chemical treatments in eliminating *Salmonella* and *Escherichia coli* O157:H7 on scarified and polished alfalfa seeds. *J. Food Prot.* **64**:1489–1495.
 13. **Jaquette, C. B., L. R. Beuchat, and B. E. Mahon.** 1996. Efficacy of chlorine and heat treatment in killing *Salmonella stanley* inoculated onto alfalfa seeds and growth and survival of the pathogen during sprouting and storage. *Appl. Environ. Microbiol.* **62**:2212–2215.
 14. **Kim, H., H. Kim, J. Bang, L. R. Beuchat, and J.-H. Ryu.** 2010. Synergistic effect of chlorine dioxide and drying treatments for inactivating *Escherichia coli* O157:H7 on radish seeds. *J. Food Prot.* **73**:1225–1230.
 15. **Kim, H., Y. Kang, L. R. Beuchat, and J.-H. Ryu.** 2008. Production and stability of chlorine dioxide in organic acid solutions as affected by pH, type of acid, and concentration of sodium chlorite, and its effectiveness in inactivating *Bacillus cereus* spores. *Food Microbiol.* **25**:964–969.
 16. **Kumar, M., R. Hora, M. Kostrzynska, W. M. Waites, and K. Warriner.** 2006. Inactivation of *Escherichia coli* O157:H7 and *Salmonella* on mung beans, alfalfa, and other seed types destined for sprout production by using an oxychloro-based sanitizer. *J. Food Prot.* **69**:1571–1578.
 17. **Lang, M. M., B. H. Ingham, and S. C. Ingham.** 2000. Efficacy of novel organic acid and hypochlorite treatments for eliminating *Escherichia coli* O157:H7 from alfalfa seeds prior to sprouting. *Int. J. Food Microbiol.* **58**:73–82.
 18. **Lim, J.-H., J.-W. Jeong, J.-H. Kim, and K.-J. Park.** 2008. Efficacy of aqueous chlorine dioxide and citric acid in reducing *Escherichia coli* on the radish seeds used for sprout production. *Food Sci. Biotechnol.* **17**:878–882.
 19. **Muñoz, M., B. De Ancos, C. Sánchez-Moreno, and M. Pilar Cano.** 2006. Evaluation of chemical and physical (high-pressure and temperature) treatments to improve the safety of minimally processed mung bean sprouts during refrigerated storage. *J. Food Prot.* **69**:2395–2402.
 20. **National Advisory Committee on Microbiological Criteria for Foods.** 1999. Microbiological safety evaluations and recommendations on sprouted seeds. *Int. J. Food Microbiol.* **52**:123–153.
 21. **Peñas, E., R. Gómez, J. Frias, and C. Vidal-Valverde.** 2009. Efficacy of combinations of high pressure treatment, temperature and antimicrobial compounds to improve the microbiological quality of alfalfa seeds for sprout production. *Food Control* **20**:31–39.
 22. **Saroj, S. D., et al.** 2007. Radiation processing for elimination of *Salmonella* Typhimurium from inoculated seeds used for sprout making in India and effect of irradiation on germination of seeds. *J. Food Prot.* **70**:1961–1965.
 23. **Scouten, A. J., and L. R. Beuchat.** 2002. Combined effects of chemical, heat and ultrasound treatments to kill *Salmonella* and *Escherichia coli* O157:H7 on alfalfa seeds. *J. Appl. Microbiol.* **92**:668–674.
 24. **Sharma, R. R., A. Demirci, L. R. Beuchat, and W. F. Fett.** 2003. Application of ozone for inactivation of *Escherichia coli* O157:H7 on inoculated alfalfa sprouts. *J. Food Process. Preserv.* **27**:51–64.
 25. **Sharma, R. R., and A. Demirci.** 2003. Treatment of *Escherichia coli* O157:H7 inoculated alfalfa seeds and sprouts with electrolyzed oxidizing water. *Int. J. Food Microbiol.* **86**:231–237.
 26. **Siddiqui, S. U., A. Ali, and M. F. Chaudhary.** 2008. Germination behavior of wheat (*Triticum aestivum*) varieties to artificial ageing under varying temperature and humidity. *Pak. J. Bot.* **40**:1121–1127.
 27. **Taormina, P. J., and L. R. Beuchat.** 1999. Behavior of enterohemorrhagic *Escherichia coli* O157:H7 on alfalfa sprouts during the sprouting process as influenced by treatments with various chemicals. *J. Food Prot.* **62**:850–856.
 28. **Taormina, P. J., and L. R. Beuchat.** 1999. Comparison of chemical treatments to eliminate enterohemorrhagic *Escherichia coli* O157:H7 on alfalfa seeds. *J. Food Prot.* **62**:318–324.
 29. **Taormina, P. J., L. R. Beuchat, and L. Slutsker.** 1999. Infections associated with eating seed sprouts: an international concern. *Emerg. Infect. Dis.* **5**:626–634.
 30. **Waje, C., and J. H. Kwon.** 2007. Improving the food safety of seed sprouts through irradiation treatment. *Food Sci. Biotechnol.* **16**:171–176.
 31. **Weissinger, W. R., and L. R. Beuchat.** 2000. Comparison of aqueous chemical treatments to eliminate *Salmonella* on alfalfa seeds. *J. Food Prot.* **63**:1475–1482.