

SCIENTIFIC REPORTS

OPEN

Increased expression of circulating miRNA-93 in women with polycystic ovary syndrome may represent a novel, non-invasive biomarker for diagnosis

T. Sathyapalan^{1,2}, R. David², N.J. Gooderham² & S.L. Atkin³

MicroRNAs (miRNA) are a novel class of small noncoding single-stranded RNA molecules that regulate gene expression. There is increasing evidence of their importance in polycystic ovary syndrome (PCOS). The objective was to determine if miRNA-93 and miRNA-223 are differentially expressed in the circulation of women with PCOS compared to age matched women. A case-control study comparing women with PCOS ($n = 25$) to age and weight matched controls ($n = 24$) without PCOS was performed. miRNA-93 and miRNA-223 were determined by total RNA reverse transcription. Both miRNA-93 and miRNA-223 were significantly increased relative to the control group ($p < 0.01$, $p = 0.029$ respectively). In both groups there was no correlation of either miRNA-93 or miRNA-223 with insulin, HOMA-IR, HOMA- β or testosterone levels. The area under the receiver operator characteristic curve for miR-223 and miR-93 was 0.66 and 0.72 respectively, suggesting miR-93 is a more efficient biomarker than miR-223 for diagnosis of PCOS. The combination of the two miRNAs together, tested using multiple logistic regression analysis, did not improve the diagnostic potential. In conclusion, circulating miRNA-93 and miRNA-223 were higher in women with PCOS compared to age and weight matched controls independent of insulin resistance and testosterone levels, and miR-93 may represent a novel diagnostic biomarker for PCOS.

Polycystic ovary syndrome (PCOS) is one of the most common endocrine disorders and affects 6–20% of reproductive-aged women^{1–3}. Seventy five to ninety per cent of PCOS patients demonstrate insulin resistance (IR) above and beyond that predicted by body mass, race, or age^{4,5}, resulting in compensatory hyperinsulinemia⁶ and an increased risk for type 2 diabetes mellitus (T2DM)⁷ and cardiovascular disease⁸. Cellular mechanisms leading to IR in PCOS remain unclear although a post-binding defect in receptor signalling has been suggested⁹. However, tissue changes in the adipocyte function, including the stimulation of glucose transport¹⁰ and GLUT4 production¹¹, have been described in women with PCOS^{12–14}.

MicroRNAs (miRNAs) are a novel class of small noncoding single-stranded RNA molecules 18–24 nucleotides long that regulate gene expression at the posttranscriptional level. Evolutionarily conserved, miRNAs bind to the 3' un-translated regions of messenger RNAs (mRNAs), and induce degradation or inhibition of protein translation. miRNAs possess many critical regulatory functions in a wide range of biological processes such as cell proliferation, differentiation, survival and apoptosis, and the stress

¹Academic Diabetes, Endocrinology and Metabolism, Hull York Medical School, University of Hull, UK. ²Faculty of Medicine, Department of Surgery & Cancer, Imperial College, London, UK. ³Weill Cornell Medical College Qatar, Education City PO Box 24144, Doha, Qatar. Correspondence and requests for materials should be addressed to T.S. (email: Thozhukat.Sathyapalan@hyms.ac.uk)

response¹⁵. Any one particular miRNA has the potential to modulate the expression and functions of hundreds of downstream target genes¹⁶. In addition, the existence of feedback regulation mechanisms between miRNA, their targets, and their products allows for amplification or inhibition of a specific signal. Hence, alteration of even a handful of miRNAs can possibly result in dramatic deregulation of physiologic cellular functions. Emerging evidence suggests an increasing role for miRNA in both type 1 and type 2 diabetes with the potential for their use as novel disease biomarkers¹⁷. Tissue changes, follicular fluid alterations and circulating miRNA have been described in PCOS^{18–21} with the suggestion that the expression of three miRNAs were elevated compared to controls that may act as novel biomarkers¹⁸. However, their expression is likely to be complex with evidence that miRNA-21, miRNA-27b, miRNA-103, and miRNA-155 may be differentially expressed in obesity and in PCOS²¹. Recently a study identified the down regulation of the insulin sensitive-glucose transporter GLUT4 gene expression by miRNA-93 in adipose tissue¹⁹, with the suggestion that miRNA-223 may have an undefined role in insulin resistance in PCOS. A subsequent study showed that miRNA-93 in adipose tissue was overexpressed in PCOS patients with insulin resistance though discordant for expression of the host gene MCM7²². In view of these findings this study was performed to determine if miRNA-93 and miRNA-223 were found in the circulation and to determine their correlation to the metabolic indices found in PCOS compared to weight matched normal controls.

Materials and Methods

Twenty-five medication naïve women with PCOS and biochemical hyperandrogenaemia (age 18–45 years) who presented sequentially to the department of endocrinology and who fulfilled the criteria of the study were recruited from the local PCOS biobank (ISRCTN70196169). Twenty-five normal women (age 20–44 years) were recruited from the PCOS biobank and were age and body mass index (BMI) matched to the PCOS subjects for inclusion into this study. All of the control women had regular periods, no clinical or biochemical hyperandrogenemia, no significant background medical history and none of them were on any medications including oral contraceptive pills or over the counter medications. All women with PCOS and control women were Caucasian. Subject demographics are shown in Table 1.

The diagnosis of PCOS was based on all three diagnostic criteria of the Rotterdam consensus, namely clinical and biochemical evidence of hyperandrogenemia (Ferriman-Gallwey score >8; free androgen index >4 respectively), oligomenorrhea or amenorrhea and polycystic ovaries on transvaginal ultrasound²³. Liver ultrasound was performed at the same time to exclude non alcoholic fatty liver disease. Study participants had no concurrent illness, were not on any medication for the preceding nine months including oral contraceptive pills or over the counter medications and were all non-smokers. None of the patients had successful pregnancy or miscarriage at least five year prior to the study entry. Diabetes was excluded by a 75 g oral glucose tolerance test. Non-classical 21-hydroxylase deficiency, hyperprolactinaemia, Cushing's disease and androgen-secreting tumours were excluded by appropriate tests. All patients gave informed consent. This study was approved by the Newcastle & North Tyneside Ethics committee. Methods were carried out in accordance with the approved guidelines.

Blood samples were taken after an overnight fast and serum was stored frozen at -80°C pending analysis. Serum testosterone and androstenedione were measured by isotope dilution liquid chromatography-tandem mass spectrometry (Waters Corporation, Manchester, UK). Sex hormone binding globulin (SHBG) by an immunometric assay with fluorescence detection on the DPC Immulite 2000 analyzer using the manufacturer's recommended protocol. The free androgen index was obtained as the total testosterone $\times 100/\text{SHBG}$. Serum insulin was assayed using a competitive chemiluminescent immunoassay performed on the manufacturer's DPC Immulite 2000 analyzer (Euro/DPC, Llanberis, UK). The analytical sensitivity of the insulin assay was $2\mu\text{U}/\text{ml}$, the coefficient of variation was 6%, and there was no stated cross-reactivity with proinsulin. Plasma glucose was measured using a Synchron LX 20 analyzer (Beckman-Coulter), using the manufacturer's recommended protocol. The coefficient of variation for the assay was 1.2% at a mean glucose value of 5.3 mmol/liter during the study period. The insulin resistance was calculated using the HOMA method [$\text{HOMA-IR} = (\text{insulin} \times \text{glucose})/22.5$], and pancreatic beta cell sensitivity measured by $\text{HOMA-}\beta$ [$\text{HOMA-}\beta = (20 \times \text{insulin})/\text{glucose} - 3.5$]. Serum CRP was measured by the high-sensitivity method on a Beckman DXC analyzer.

RNA extraction. Total RNA was extracted from plasma from 25 women with PCOS or 25 healthy women matched for age and BMI using the MiRVana PARIS kit (Life Technologies, Paisley, UK) following manufacturer's instructions with the following modifications. Prior to addition of the acid-phenol:chloroform, 150Amoles of *c-elegans* miRNA-39 (Ambion, Life Technologies) spike-in was added to the sample as an RNA carrier. Total RNA was eluted by adding $10\mu\text{l}$ of RNase-free water to the membrane of the spin column and incubating for 2 min before centrifugation at 10000g for 30 s at room temperature, which was repeated with a further $10\mu\text{l}$. The RNA was stored at -80°C .

Reverse transcription and qPCR. Total RNA was reverse transcribed using the TaqMan MicroRNA Reverse Transcription Kit (Life Technologies, Paisley, UK) following manufacturer's instructions. Mature miRNA TaqMan assays were purchased from Life Technologies and the generated cDNA amplified

	Normal n = 25	PCOS n = 25	P value
	Mean (SD)	Mean (SD)	
Age (years)	32.2(7.7)	32.1(9.0)	0.97
Weight (kg)	76.0(18.8)	77.4(16.3)	0.79
BMI (kg/m ²)	27.1(5.8)	28.8(5.4)	0.31
Fasting glucose (mmol/l)	4.7(0.4)	4.8(0.6)	0.56
2 Hour glucose (mmol/l)	4.9(1.2)	5.7(1.3)	0.06
Androstenedione (nmol/l)	8.4(5.1)	10.7(6.7)	0.25
ALT (IU/L)	24.3(14.4)	23.3(12.7)	0.79
Insulin (μ U/ml)	6.8(3.7)	10.2(6.4)	0.03
HOMA-IR	1.5(0.9)	2.3(1.7)	0.07
HOMA- β	38.4(66.1)	158.8(89.7)	0.001
Testosterone (nmol/L)	1.2(0.7)	2.3(1.6)	0.03
SHBG (mmol/L)	81.8(105.8)	52.6(53.1)	0.29
FAI	2.6(1.7)	11.1(18.9)	0.13
hsCRP (mg/l)	2.3(3.8)	2.8(3.9)	0.65
miR-93 expression (relative to control group)	1.0(0.6)	2.0(1.6)	0.009
miR-223 expression (relative to control group)	1.0(0.7)	2.01(2.2)	0.029

Table 1. Demographics, biochemical and clinical markers for the PCOS and control group (unpaired t test) (BMI, body mass index; FAI, free androgen index; SHBG, sex hormone binding globulin; HOMA-IR, homeostatic model assessment-insulin resistance; HOMA- β , homeostatic model assessment-insulin beta cell sensitivity; hsCRP, high sensitivity C-reactive protein.)

using the Taqman 2x Universal PCR master mix, No AmpErase UNG (Life Technologies), with each reaction performed in triplicate. The amplification was performed in a StepOnePlus PCR System (Life Technologies) in 96 well plates. The amplification curves were analysed using the StepOne software (Life Technologies) and the comparative Ct Method (Δ CT Method)²⁴. Calibration was based on the expression of the *c-elegans* miR-39 spike-in.

Statistical analysis. For a significant change in miRNA measurements power was based on a 50% increase or decrease of the relative change to control, with a common standard deviation of 0.5¹⁹. A sample size of 23 participants per group to has 90% power, 1% significance (two-tailed) was calculated (n-query software, USA).

An unpaired t test was used to compare changes from baseline for the biochemical data and clinical observations within groups. The Wilcoxon signed rank test was applied to biochemical data that violated the assumptions of normality when tested using the Kolmogorov-Smirnov test. For all analysis, a two-tailed $P \leq 0.05$ was considered to indicate statistical significance. Statistical analysis was performed using SPSS for Windows NT, version 22.0 (SPSS Inc., Chicago, IL). Data are reported as mean (SEM).

A one-way ANOVA with post-hoc Dunnett's test performed (GraphPad Software, La Jolla California USA) was used to test for statistical significance between miRNA expression in plasma from patients with PCOS compared to the control group. A ROC curve was plotted for each miRNA individually, and in combination, to determine their discriminating effects. The sensitivity and specificity of detecting cases and controls were assessed by the area under the ROC curve (AUC) and 95% confidence interval (CI). Multiple logistic regression and ROC curve analysis was performed in R²⁵.

In silico analysis was carried out using miRWalk, which predicts targets based on the mRNA 3' UTR region produced by the miRWalk program and also presents results from other programs with target gene prediction, of which miRanda, TargetScan and DIANA were used. Only targets predicted by all 4 databases were used for pathway analysis by Ingenuity Pathway Analysis (IPA; Qiagen, Redwood City).

Results

Both groups were well matched for age, weight and BMI (Table 1). Of the 25 PCOS patients 8 had a BMI of 25 kg/m² or less and of those only 1 of 8 had biochemical hyperandrogenemia with an elevated FAI, whilst of those 17 PCOS patients with a BMI greater than 25 kg/m², only 3 of 17 did not have a raised FAI. In the PCOS group insulin was significantly increased ($p = 0.03$), HOMA-IR did not differ ($p = 0.07$), there was a significant increase in HOMA- β ($p = 0.001$) and testosterone was significantly elevated ($p = 0.03$) (Table 1).




Figure 1. Increased expression of microRNA(miR)-223 and miR-93 in plasma from patients with PCOS (n = 25) relative to that expressed in the control age and BMI matched population (n = 24). Data are means \pm SEM, significance compared to control (one-way ANOVA with Dunnett's post-test; *P \leq 0.05, **P \leq 0.01).

The expression of both miRNA-93 and miRNA-223 was significantly increased in PCOS patients relative to the control group (p = 0.009 and p = 0.029, respectively) (Fig. 1). In the control group there was no correlation of either miRNA-93 or miRNA-223 with any of the metabolic and hormonal parameters including insulin (0.4;p = 0.09, 0.2;p = 0.5 respectively), HOMA-IR (0.4;p = 0.07, 0.2;p = 0.3 respectively), HOMA- β (0.1;p = 0.8, 0.2;p = 0.3 respectively) or testosterone (0.04;p = 0.9, 0.1;p = 0.7 respectively). In the PCOS group there was no correlation of either miRNA-93 or miRNA-223 with any of the parameters including insulin (0.2;p = 0.4, 0.1;p = 0.5 respectively), HOMA-IR (0.2;p = 0.4, 0.1;p = 0.7 respectively), HOMA- β (0.07;p = 0.8, 0.2;p = 0.3 respectively) or testosterone (0.3;p = 0.1, 0.1;p = 0.6 respectively).

To evaluate the diagnostic value of miR-223 and miR-93, ROC curves were plotted and the AUC calculated (Fig. 2). The sensitivity and specificity for miR-223 and miR-93 were calculated at the cut-off values of 1.2 and 1.1 respectively, at which the largest sensitivity and specificity was defined as the optimal diagnostic point. The sensitivity, specificity and AUC of miR-223 and miR-93 were 0.56, 0.72 and 0.60 (95% CI: 0.51–0.82), and 0.64, 0.76 and 0.72 (95% CI: 0.58–0.86) respectively. The results show that miR-93 is more efficient than miR-223 as a biomarker for the diagnosis of PCOS. Multiple logistic regression and ROC curve analysis was used to determine whether the combination of miR-93 with miR-223 had improved diagnostic potential over miR-93 alone. The results showed that the combination of the two miRNAs did not improve beyond the diagnostic value of miR-93 alone (sensitivity, specificity and AUC of miR-223 and miR-93 combined were 0.64, 0.76 and 0.73 (95% CI: 0.59–0.87) respectively).

Following target gene prediction for miR-223 and miR-93 using miRWalk, miRanda, TargetScan and DIANA, a list of possible target genes identified by all 4 programs was obtained and analysed using IPA to generate a list of the top canonical pathways likely to be specifically controlled by these miRNAs. As shown in Table 2 and Table 3, genes that are predicted to be targeted by these miRNAs are involved in pathways including cancer, stem cell pluripotency and hepatic growth factor (HGF) signalling.

Discussion

This is the first study to show the elevated plasma levels of both miRNA-93 and miRNA-223 in women with PCOS particularly when weight was accounted for. These increased miRNA levels were independent of insulin, insulin resistance and of hyperandrogenaemia in women with PCOS.

It may have been anticipated that a strong correlation to insulin resistance would be found given the recent report of miRNA-93 and miRNA-223 up-regulation in PCOS adipocytes, and the *in vitro* demonstration of down-regulation of GLUT4 gene expression¹⁹. In a subsequent study miRNA-93 in adipose tissue correlated with PCOS and insulin resistance though its host gene MCM7 was discordantly down-regulated²². However, the PCOS and control women were not matched for BMI or insulin resistance, which may explain the discrepancies between that study and ours. Other reasons for this apparent lack of correlation are likely to be multifactorial as it is unclear how circulating levels of miRNA may relate to tissue function and effect. However, the top canonical pathways identified peroxisome proliferator receptor (PPAR), insulin like growth factor- 1 (IGF-1) and angiopoietin signaling as being potentially regulated by miR-223, all of which have been shown to play a role in PCOS. PPAR signaling has been shown to be important in insulin resistance, hyperandrogenism, endometrial response and in the ovary in PCOS,^{26,27} with PPAR gamma stimulation by pioglitazone being proposed as a potential therapeutic modality²⁶. IGF-1 is a potent mitogenic factor and it appears necessary for ovarian follicle development²⁸, and has been shown to be involved in pre-antral follicle growth in PCOS²⁹. Angiopoietin 1 and 2 are




Figure 2. ROC curve analysis of (A) miR-223 and (B) miR-93 to discriminate women with PCOS from healthy controls.

Name	p-value
Top Canonical Pathways	
PPAR Signalling	6.24E-05
Prostate Cancer Signalling	1.57E-04
IGF-1 Signalling	1.71E-04
Angiotensin Signalling	1.76E-04
Molecular Mechanisms of Cancer	3.69E-04

Table 2. Top canonical pathways predicted by *in silico* analysis using 3' UTR target prediction software to be regulated by microRNA-223.

Name	p-value
Top Canonical Pathways	
Molecular Mechanisms of Cancer	2.38E-06
Nerve Growth Factor Signalling	3.53E-06
Human Embryonic Stem Cell Pluripotency	4.62E-05
Hepatic Growth Factor Signalling	5.76E-05
Pancreatic Adenocarcinoma Signalling	8.21E-05

Table 3. Top canonical pathways predicted by *in silico* analysis using 3' UTR target prediction software to be regulated by microRNA-93

essential for ovarian function and are preferentially increased in ovarian stimulation in PCOS³⁰. The canonical pathways identified both Nerve Growth Factor (NGF) Signalling and Hepatic Growth Factor (HGF) Signalling in the top 5 canonical pathways for potential regulation by miR-93. NGF may have an important role in the ovary with excessive production being associated with cystic ovarian morphology in rodent models akin to that found in women with PCOS^{31,32}. HGF is an adipokine that is also found in the ovary³³ but has been associated with obesity and metabolic syndrome^{34,35,36} with the suggestion that it has importance in insulin resistance associated compensatory mechanisms leading to an increase in insulin secretion through enhanced beta cell mass³⁵. Furthermore, of those identified, molecular mechanisms of cancer was common for both miRNAs. The current study indicates that analysis of miRNAs offers potential insights into pathways that are affected in PCOS. The data, although limited to 2 miRNAs in

the present study, warrant further in depth investigation into the role of miRNA in PCOS. One approach could be to examine the circulating miRNAome in such patients.

The canonical pathways in this study identified different targets for miR-223 and miR-93 to previous studies, but this is not surprising given that miRNAs differ in their tissue specific actions that are likely to be tissue and circumstance specific. Examples include miR-223 overexpression in cardiomyocytes increased GLUT4 expression³⁷, reduced expression of miR-223 in breast cancer cell affects mediators of invasion³⁸, and miRNA-93 and miRNA-223 have been associated with dysregulation in diabetes¹⁷.

Circulating miRNAs are very stable to enzymatic degradation and physical conditions such as freeze/thaw, and can be stored for months without their degradation³⁹. As such they are attractive as biomarkers for disease diagnosis and monitoring. Therefore specific miRNA or combination of miRNAs may become a reliable biomarker for the diagnosis of PCOS, which may help distinguish the differing phenotypes identified by the Rotterdam criteria. Whilst the sensitivity and the specificity shown in the receiver operator curves are too low as a single diagnostic test, miRNA-93 may be a biomarker to support a positive diagnosis of PCOS rather than the diagnosis being one of exclusion that is currently the case, which would be of value in adolescence or in the perimenopausal/menopausal woman where diagnosis is difficult⁴⁰. Our study demonstrates that plasma levels of miR-93 can distinguish patients with PCOS from healthy controls with more sensitivity and specificity than miR-223, and as such, circulating miR-93 may represent a biomarker for PCOS.

In a recent study, miR-222, miR-146a and miR-30c were found to be significantly increased in PCOS patients with the combination of the three miRNAs showing potential promise for the more efficient diagnosis of PCOS¹⁸. In the current study, however, the combination of miR-93 with miR-223 did not improve the diagnostic potential of miR-93 alone. The scope of miRNA involved in the pathophysiology of PCOS is unknown and likely extensive; in a rat PCOS model 24% of 349 miRNAs were differentially expressed compared to control following dihydrotestosterone treatment⁴¹.

The strengths of the current study included age and weight matching of the PCOS with normal subjects that circumvented miRNA expression associated with weight differences in PCOS²¹. Other strengths were the inclusion of all 3 criteria for PCOS diagnosis, the study was powered to detect a difference in miRNA, and liver ultrasound and alanine transferase measurements excluded non-alcoholic fatty liver disease that is associated with miRNA expression⁴². There were no changes in markers of inflammation, hsCRP or white cell count in this study, that may be elevated in PCOS when obesity has not been accounted for, which is important as both miRNA-93 and miRNA-223 have been associated with inflammation^{43,44}.

This study is limited by the small cohort that was employed and a larger study specifically powered for each of the phenotypes that are described in the Rotterdam diagnostic criteria is needed for validation, and comparison with the NIH and Androgen Excess PCOS Society guidelines⁴⁰. This study included women who satisfied all of the three Rotterdam diagnostic criteria for rigorous phenotyping; therefore, the expression of these miRNA may only define the most severe metabolic phenotype and may not be generalizable to those PCOS women diagnosed from only two out of three criteria and further studies are necessary to clarify this. Caution needs to be expressed in extrapolating the results of insulin resistance determined by the HOMA method reported here to measures of insulin resistance by other methods, such as insulin clamps or intravenous glucose tolerance tests. However, there is evidence to show that there is strong correlation between HOMA and other methods^{45–48}.

In summary, PCOS subjects with elevated insulin and testosterone levels, showed higher circulating levels of miR-93 and miR-223 compared to age and weight matched controls, though the elevation in the miRNA levels was not related to either insulin resistance or hyperandrogenaemia. Moreover, miR-93 elevation in the plasma may represent a novel, non-invasive biomarker for the diagnosis of PCOS.

References

1. Yildiz, B. O., Bozdag, G., Yapici, Z., Esinler, I. & Yarali, H. Prevalence, phenotype and cardiometabolic risk of polycystic ovary syndrome under different diagnostic criteria. *Human reproduction* **27**, 3067–3073, doi: 10.1093/humrep/des232 (2012).
2. Teede, H. J. *et al.* Longitudinal weight gain in women identified with polycystic ovary syndrome: results of an observational study in young women. *Obesity (Silver Spring)* **21**, 1526–1532, doi: 10.1002/oby.20213 (2013).
3. March, W. A. *et al.* The prevalence of polycystic ovary syndrome in a community sample assessed under contrasting diagnostic criteria. *Human reproduction* **25**, 544–551, doi: 10.1093/humrep/dep399 (2010).
4. Stepto, N. K. *et al.* Women with polycystic ovary syndrome have intrinsic insulin resistance on euglycaemic-hyperinsulinaemic clamp. *Hum Reprod* **28**, 777–784, doi: 10.1093/humrep/des463 (2013).
5. Azziz, R. *et al.* The Androgen Excess and PCOS Society criteria for the polycystic ovary syndrome: the complete task force report. *Fertil Steril* **91**, 456–488, doi: 10.1016/j.fertnstert.2008.06.035 (2009).
6. Legro, R. S., Finegood, D. & Dunaif, A. A fasting glucose to insulin ratio is a useful measure of insulin sensitivity in women with polycystic ovary syndrome. *The Journal of clinical endocrinology and metabolism* **83**, 2694–2698 (1998).
7. Ovalle, F. & Azziz, R. Insulin resistance, polycystic ovary syndrome, and type 2 diabetes mellitus. *Fertil Steril* **77**, 1095–1105 (2002).
8. Shaw, L. J. *et al.* Postmenopausal women with a history of irregular menses and elevated androgen measurements at high risk for worsening cardiovascular event-free survival: results from the National Institutes of Health–National Heart, Lung, and Blood Institute sponsored Women's Ischemia Syndrome Evaluation. *The Journal of clinical endocrinology and metabolism* **93**, 1276–1284, doi: 10.1210/jc.2007-0425 (2008).
9. Diamanti-Kandarakis, E. & Dunaif, A. Insulin Resistance and the Polycystic Ovary Syndrome Revisited: An Update on Mechanisms and Implications. *Endocrine reviews* **33**, 981–1030, doi: 10.1210/er.2011-1034 (2012).

10. Ciaraldi, T. P. *et al.* Cellular mechanisms of insulin resistance in polycystic ovarian syndrome. *The Journal of clinical endocrinology and metabolism* **75**, 577–583 (1992).
11. Rosenbaum, D., Haber, R. S. & Dunaif, A. Insulin resistance in polycystic ovary syndrome: decreased expression of GLUT-4 glucose transporters in adipocytes. *Am. J. Physiol.* **264**, E197–202 (1993).
12. Marsden, P. J., Murdoch, A. P. & Taylor, R. Tissue insulin sensitivity and body weight in polycystic ovary syndrome. *Clinical endocrinology* **55**, 191–199, doi: cen1303 (2001).
13. Biyasheva, A., Legro, R. S., Dunaif, A. & Urbanek, M. Evidence for association between polycystic ovary syndrome (PCOS) and TCF7L2 and glucose intolerance in women with PCOS and TCF7L2. *The Journal of clinical endocrinology and metabolism* **94**, 2617–2625, doi: jc.2008-1664 10.1210 (2009).
14. Lord, J., Thomas, R., Fox, B., Acharya, U. & Wilkin, T. The effect of metformin on fat distribution and the metabolic syndrome in women with polycystic ovary syndrome—a randomised, double-blind, placebo-controlled trial. *BJOG: an international journal of obstetrics and gynaecology* **113**, 817–824, doi: BJO966 10.1111/j.1471-0528.2006.00966.(2006).
15. Flynt, A. S. & Lai, E. C. Biological principles of microRNA-mediated regulation: shared themes amid diversity. *Nature reviews. Genetics* **9**, 831–842, doi: 10.1038/nrg2455 (2008).
16. Lim, L. P. *et al.* Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature* **433**, 769–773, doi: nature03315 10.1038/nature03315 (2005).
17. Guay, C. & Regazzi, R. Circulating microRNAs as novel biomarkers for diabetes mellitus. *Nature reviews. Endocrinology* **9**, 513–521, doi: 10.1038/nrendo.2013.86 (2013).
18. Long, W. *et al.* Characterization of serum microRNAs profile of PCOS and identification of novel non-invasive biomarkers. *Cellular physiology and biochemistry: international journal of experimental cellular physiology, biochemistry, and pharmacology* **33**, 1304–1315, doi: 10.1159/000358698 (2014).
19. Chen, Y. H. *et al.* miRNA-93 inhibits GLUT4 and is overexpressed in adipose tissue of polycystic ovary syndrome patients and women with insulin resistance. *Diabetes* **62**, 2278–2286, doi: 10.2337/db12-0963 (2013).
20. Roth, L. W. *et al.* Altered microRNA and gene expression in the follicular fluid of women with polycystic ovary syndrome. *Journal of assisted reproduction and genetics* **31**, 355–362, doi: 10.1007/s10815-013-0161-4 (2014).
21. Murri, M., Insenser, M., Fernandez-Duran, E., San-Millan, J. L. & Escobar-Morreale, H. F. Effects of polycystic ovary syndrome (PCOS), sex hormones, and obesity on circulating miRNA-21, miRNA-27b, miRNA-103, and miRNA-155 expression. *The Journal of clinical endocrinology and metabolism* **98**, E1835–1844, doi: 10.1210/jc.2013-2218 (2013).
22. Wu, H. L. *et al.* The expression of the miR-25/93/106b family of micro-RNAs in the adipose tissue of women with Polycystic Ovary Syndrome. *The Journal of clinical endocrinology and metabolism*, jc20134435, doi: 10.1210/jc.2013-4435 (2014).
23. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome (PCOS). *Hum Reprod* **19**, 41–47 (2004).
24. Livak, K. J. & Schmittgen, T. D. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{(-Delta Delta C(T))} Method. *Methods* **25**, 402–408, doi: 10.1006/meth.2001.1262 (2001).
25. R. Core Team. R: A language and environment for statistical computing. *R Foundation for Statistical Computing, Vienna, Austria*, URL <http://www.R-project.org/> (2013).
26. Cho, L. W., Kilpatrick, E. S., Keevil, B. G., Coady, A. M. & Atkin, S. L. Effect of metformin, orlistat and pioglitazone treatment on mean insulin resistance and its biological variability in polycystic ovary syndrome. *Clinical endocrinology* **70**, 233–237, doi: 10.1111/j.1365-2265.2008.03309. (2009).
27. Froment, P. *et al.* Peroxisome proliferator-activated receptors in reproductive tissues: from gametogenesis to parturition. *J. Endocrinol.* **189**, 199–209, doi: 10.1677/joe.1.06667 (2006).
28. Zhou, J., Kumar, T. R., Matzuk, M. M. & Bondy, C. Insulin-like growth factor I regulates gonadotropin responsiveness in the murine ovary. *Molecular endocrinology* **11**, 1924–1933, doi: 10.1210/mend.11.13.0032 (1997).
29. Stubbs, S. A. *et al.* Role of Insulin-like growth factors in initiation of follicle growth in normal and polycystic human ovaries. *The Journal of clinical endocrinology and metabolism* **98**, 3298–3305, doi: 10.1210/jc.2013-1378 (2013).
30. Tal, R., Seifer, D. B., Grazi, R. V. & Malter, H. E. Angiopoietin-1 and angiopoietin-2 are altered in polycystic ovarian syndrome (PCOS) during controlled ovarian stimulation. *Vasc Cell* **5**, 18, doi: 10.1186/2045-824x-5-18 (2013).
31. Wilson, J. L. *et al.* Excess of nerve growth factor in the ovary causes a polycystic ovary-like syndrome in mice, which closely resembles both reproductive and metabolic aspects of the human syndrome. *Endocrinology* **155**, 4494–4506, doi: 10.1210/en.2014-1368 (2014).
32. Dissen, G. A. *et al.* Excessive ovarian production of nerve growth factor facilitates development of cystic ovarian morphology in mice and is a feature of polycystic ovarian syndrome in humans. *Endocrinology* **150**, 2906–2914, doi: 10.1210/en.2008-1575 (2009).
33. Sahin, N. *et al.* The levels of hepatocyte growth factor in serum and follicular fluid and the expression of c-Met in granulosa cells in patients with polycystic ovary syndrome. *Fertility and sterility* **99**, 264–269, doi: 10.1016/j.fertnstert.2012.08.059 (2013).
34. Vistoropsky, Y., Trofimov, S., Malkin, I., Kobylansky, E. & Livshits, G. Genetic and environmental determinants of hepatocyte growth factor levels and their association with obesity and blood pressure. *Ann. Hum. Biol.* **35**, 93–103, doi: 10.1080/03014460701822003 (2008).
35. Araujo, T. G. *et al.* Hepatocyte growth factor plays a key role in insulin resistance-associated compensatory mechanisms. *Endocrinology* **153**, 5760–5769, doi: 10.1210/en.2012-1496 (2012).
36. Hiratsuka, A. *et al.* Strong association between serum hepatocyte growth factor and metabolic syndrome. *The Journal of clinical endocrinology and metabolism* **90**, 2927–2931, doi: 10.1210/jc.2004-1588 (2005).
37. Lu, H., Buchan, R. J. & Cook, S. A. MicroRNA-223 regulates Glut4 expression and cardiomyocyte glucose metabolism. *Cardiovasc. Res.* **86**, 410–420, doi: cvq010 10.1093/cvr/cvq010 (2010).
38. Gong, B. *et al.* Caprin-1 is a novel microRNA-223 target for regulating the proliferation and invasion of human breast cancer cells. *Biomed. Pharmacother.* **67**, 629–636, doi: 10.1016/j.biopha.2013.06.006 (2013).
39. Mraz, M., Malinova, K., Mayer, J. & Pospisilova, S. MicroRNA isolation and stability in stored RNA samples. *Biochemical and biophysical research communications* **390**, 1–4, doi: 10.1016/j.bbrc.2009.09.061 (2009).
40. Legro, R. S. *et al.* Diagnosis and Treatment of Polycystic Ovary Syndrome: An Endocrine Society Clinical Practice Guideline. *Journal of Clinical Endocrinology & Metabolism* **98**, 4565–4592, doi: 10.1210/jc.2013-2350 (2013).
41. Hossain, M. M. *et al.* Altered expression of miRNAs in a dihydrotestosterone-induced rat PCOS model. *Journal of ovarian research* **6**, 36, doi: 10.1186/1757-2215-6-36 (2013).
42. Tan, Y., Ge, G., Pan, T., Wen, D. & Gan, J. A Pilot Study of Serum MicroRNAs Panel as Potential Biomarkers for Diagnosis of Nonalcoholic Fatty Liver Disease. *PLoS one* **9**, e105192, doi: 10.1371/journal.pone.0105192 (2014).
43. Li, F., Liang, X., Chen, Y., Li, S. & Liu, J. Role of microRNA-93 in regulation of angiogenesis. *Tumour Biol.*, doi: 10.1007/s13277-014-2605-6 (2014).
44. Haneklaus, M., Gerlic, M., O'Neill, L. A. & Masters, S. L. miR-223: infection, inflammation and cancer. *Journal of internal medicine* **274**, 215–226, doi: 10.1111/joim.12099 (2013).

45. Malita, F. M. *et al.* Surrogate indexes vs. euglycaemic-hyperinsulinemic clamp as an indicator of insulin resistance and cardiovascular risk factors in overweight and obese postmenopausal women. *Diabetes Metab* **32**, 251–255 (2006).
46. Matthews, D. R. *et al.* Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia* **28**, 412–419 (1985).
47. Hanson, R. L. *et al.* Evaluation of simple indices of insulin sensitivity and insulin secretion for use in epidemiologic studies. *American journal of epidemiology* **151**, 190–198 (2000).
48. Sarafidis, P. A. *et al.* Validity and reproducibility of HOMA-IR, 1/HOMA-IR, QUICKI and McAuley's indices in patients with hypertension and type II diabetes. *J Hum Hypertens* **21**, 709–716, doi: 10.1038/sj.jhh.1002201 (2007).

Acknowledgements

This study was supported by the Diabetes Charitable funds, Hull and East Yorkshire Hospitals Trust, from Weill Cornell Medical College, Qatar and the Qatar Foundation.

Author Contributions

S.L.A. drafted the manuscript, T.S., R.D., N.J.G. and S.L.A. revised the manuscript, All authors reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Sathyapalan, T. *et al.* Increased expression of circulating miRNA-93 in women with polycystic ovary syndrome may represent a novel, non-invasive biomarker for diagnosis. *Sci. Rep.* **5**, 16890; doi: 10.1038/srep16890 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>