COMMENTS AND RESPONSES

Increased Matrix Metalloproteinase-9 Predicts Poor Wound Healing in Diabetic Foot Ulcers

Response to Muller et al.

e thank the authors for their comment (1) on our study (2) and we agree that the pathophysiology of diabetic foot ulcers is complex and not well understood. We recognize that we are not the first to establish a potential role for matrix metalloproteinase-9 (MMP-9) in chronic wound healing. However, we report that its measurement in wound fluids from diabetic foot ulcers has predictive value (2). We acknowledge that the study by Muller et al. (3), published in parallel with ours, also describes a predictive role for MMPs, in particular MMP-1-to-tissue inhibitor of metalloproteinase (TIMP-1) ratio in wound healing. We did not report on MMP-1 levels in our study because they were not reliable

in measurement, being at the lower limit of detection of our assay, while in contrast MMP-9 is found in considerably higher levels (8- to 10-fold). In our work (2), we found a predictive role for pro- and active MMP-9 and its ratio with TIMP-1 in wound healing outcome, whereas Muller et al. (3) found that pro-MMP-9 had no predictive value.

In conclusion, we concur with Muller et al. that it is increasingly clear that MMPs play an important role in wound healing and that their levels are altered in diabetic wound healing. There are many different MMPs, and their expression and activation will depend on the stage of healing, as well as factors such as bacterial load, treatment, local inflammation, etc. A better understanding of how MMPs are affected in diabetes will permit evaluation of their prognostic value, as well as their utility for treatment targets to improve wound healing rates.

> Yu Liu, md¹ Danging Min, Phd² THYRA BOLTON, RN³ Vanessa Nubé, mscpod³ Stephen M. Twigg, phd, $md^{1,2,3}$ Dennis K. Yue, Phd, $md^{1,2,3}$ SUSAN V. McLennan, Phd^{1,2}

From the ¹Discipline of Medicine, University of Sydney, Sydney, Australia; the ²Department of Endocrinology, Royal Prince Alfred Hospital, Sydney, Australia; and the ³Diabetes Centre, Royal Prince Alfred Hospital, Sydney, Australia.

Corresponding author: Susan McLennan, smclennan@ med.usyd.edu.au.

DOI: 10.2337/dc09-1394

© 2009 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. See http:// creativecommons.org/licenses/by-nc-nd/3.0/ for

Acknowledgments—No potential conflicts of interest relevant to this article were

References

- 1. Muller M, Trocme C, Morel F, Halimi S, Benhamou PY. Increased matrix metalloproteinase-9 predicts poor wound healing in diabetic foot ulcers (Letter). Diabetes Care 2009:32:e138. DOI: 10.2337/dc09-
- 2. Liu Y, Min D, Bolton T, Nubé V, Twigg SM, Yue DK, McLennan SV. Increased matrix metalloproteinase-9 predicts poor wound healing in diabetic foot ulcers. Diabetes Care 2009;32:117-119
- 3. Muller M, Trocme C, Lardy B, Morel F, Halimi S, Benhamou PY. Matrix metalloproteinases and diabetic foot ulcers: the ratio of MMP-1 to TIMP-1 is a predictor of wound healing. Diabet Med 2008;25: 419-426

e137