

Increased Risk for Family Violence During the COVID-19 Pandemic

Kathryn L. Humphreys, PhD, EdM,^{a,b} Myo Thwin Myint, MD,^b Charles H. Zeanah, MD^b

Although the public health benefits of social distancing, isolation, and quarantines are well-established and essential for reducing risk of transmitting the coronavirus disease (COVID-19), the disease caused by the novel coronavirus (severe acute respiratory syndrome coronavirus 2), there are also likely consequences for these practices when considering the impact of violence in the home. Reports of increased domestic violence after quarantine orders in China have revealed the interpersonal violence risks of isolation. Indeed, in a recent review of the psychological impact of quarantine published in *The Lancet*, the authors indicated increased anger, confusion, and posttraumatic stress symptoms, as well as evidence of increases in substance use, in those subjected to quarantine.¹

These kinds of dysregulated emotions and substance use can increase violent behavior, especially within the family. Children's exposure to intimate partner violence, whether directly witnessed or overheard, is harmful and may lead to posttraumatic stress disorder and other serious emotional and behavioral problems.² Furthermore, intimate partner violence and child abuse often co-occur,³ and it is likely that children will experience increased risk for maltreatment when isolated at home. In typical (ie, nonpandemic) circumstances, rates of child maltreatment are alarming. In the United States, 1 in 8 children have confirmed maltreatment by child protective services (CPS) in their lifetime.⁴ Among these, recurrence of maltreatment is high.⁵

INCREASED RISK FOR CHILD EXPOSURE TO FAMILY VIOLENCE DURING PERIODS OF CRISIS

The authors are increasingly concerned about the risks for children and vulnerable families during this unprecedented period of isolation as child care centers and schools necessarily close their doors. The risks are compounded by added pressures that many parents continue to work full-time during these periods. If parents must leave their home to work, children face an increased risk for supervisory neglect (ie, not having adequate supervision to keep children from harm). If working from home, parents with young children are forced to try to meet work demands while

^aDepartment of Psychology and Human Development, Vanderbilt University, Nashville, Tennessee; and ^bDepartment of Psychiatric and Behavioral Sciences, School of Medicine, Tulane University, New Orleans, Louisiana

Dr Humphreys conceptualized the initial idea for the manuscript and wrote the first draft of the manuscript; Drs Myint and Zeanah made substantial contributions to the manuscript conception and reviewed and revised the manuscript; and all authors approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

DOI: <https://doi.org/10.1542/peds.2020-0982>

Accepted for publication Apr 13, 2020

Address correspondence to Kathryn L. Humphreys, PhD, EdM, Department of Psychology and Human Development, Vanderbilt University, Nashville, TN 37203.
 E-mail: k.humphreys@vanderbilt.edu

PEDIATRICS (ISSN Numbers: Print, 0031-4005; Online, 1098-4275).

Copyright © 2020 by the American Academy of Pediatrics

FINANCIAL DISCLOSURE: Dr Humphreys reports grants from the National Institute of Mental Health, Jacobs Foundation, Caplan Foundation for Early Childhood, Vanderbilt Kennedy Center, Peabody College (Vanderbilt University), and Vanderbilt Institute for Clinical and Translational Research. Dr Zeanah reports grants from the National Institute of Mental Health, the Lumos Foundation, the Inter-American Development Bank, the Substance Abuse & Mental Health Services Administration, and the Irving Harris Foundation. Dr Myint receives no external funding.

FUNDING: No external funding.

POTENTIAL CONFLICT OF INTEREST: The authors have indicated they have no potential conflicts of interest to disclose.

To cite: Humphreys KL, Myint MT, Zeanah CH. Increased Risk for Family Violence During the COVID-19 Pandemic. *Pediatrics*. 2020;146(1):e20200982

simultaneously caring for young children. Changes in routine are upsetting, confusing, and difficult for young children. Increased oppositional behavior and limit testing are expected, and these behaviors are most likely to elicit harsh responses from parents. Coupled with parental anxiety and stress about financial, logistic, and existential concerns, these interactions are likely a recipe for temper outbursts and verbal and physical abuse. Young children are the most vulnerable to abuse, with the highest abuse related fatalities among those <12 months.⁶ Unfortunately, school closures mean that the largest source of reports to CPS will disappear, resulting in reduced detection of maltreatment. Furthermore, given that well-child visits and other routine medical care are being postponed because of the pandemic, clinicians are losing the opportunities to both detect and prevent maltreatment.

RECOMMENDATIONS FOR PROTECTING CHILDREN FROM HARM

Several avenues for reducing risk to children are achieved indirectly through the advocacy efforts of clinicians. These include stimulus payments from the federal government to provide parents financial relief and legislation mandating guaranteed paid sick leave for all workers. Employers must have clearly communicated and reasonable expectations for employees responsible for caring for others. In particular, caring for young children will reduce productivity, and hours of work will not align with typical “office hours.” Additional funding and support is needed for child welfare agencies and law enforcement, who employ essential workers unable to socially distance during their investigations, placements, and supervision of children needing protection.

TABLE 1 Recommendations for Clinicians in Their Patient Interactions and Recommendations To Make to Parents

<p>Recommendations for clinicians</p> <ul style="list-style-type: none"> Make violence potential part of every assessment. Inquire about family stress levels and how parents manage stress. Inquire about the coparenting relationship. Inquire about social supports available to and being used by the family. Inquire about substance use and any recent increases. Look for signs of stress, irritability, or depression in the parent. Look for harsh responses to child behaviors in parents. Look for signs of fearfulness or dysregulation in children. Look for indicators of controlling behaviors by one partner. Identify families who are more at risk for violence based on previous encounters and conduct check-ins if there are no scheduled appointments for them in the near-term to reduce the likelihood that otherwise high-risk families would be undetected. <p>Recommendations to make to parents</p> <ul style="list-style-type: none"> Recognize that feelings of stress, anger, worry, and irritability are expected given the demands of care of young children, especially if coupled with job demands, income loss, or job uncertainty. Use consistent wake, bed, and mealtimes to provide structure for both children and parents. Consider structuring the day in specific segments, for example, reading, inside or outside playtime, naptime, exercise, screen time, etc. If 2 parents are home, consider a “tag team” approach to child care. Understand that increases in children’s challenging behaviors and limit testing are developmentally typical responses that likely reflect distress and disruptions from typical routines. Identify when feeling activated and use a coping strategy that helps (eg, deep breaths, considering the things that they are grateful for), and if those do not help, institute a household “quiet time” for rest. Develop a plan to call a friend or family member to avoid engaging in violent behaviors. Use reliable resources for parenting during the COVID-19 pandemic, such as www.healthychildren.org and https://www.nctsn.org/resources/parent-caregiver-guide-to-helping-families-cope-with-the-coronavirus-disease-2019.

For pediatricians and other health care professionals who interact directly with children and families, there is a need to maintain continuity of practice during times of disaster.⁷ Although clinicians should discuss best practices to maintain hygiene, respond to school and child care closures, and discuss COVID-19 with children, the authors urge the inclusion of monitoring risk for violence in the home during these visits. Doing so while following social distancing practices means conducting many aspects of well-child visits using telehealth rather than postponing these important points of contact. Flexibility may be required to enable those without Internet to participate in telehealth visits. Telehealth limits the ability to assess children or parents in a space in which they have privacy from an abuser, and thus clinicians should be attuned to nonverbal cues or other

signs that violence is occurring. Clinicians should continue to report concerns about suspected abuse or neglect to local CPS.

It is likely too early to detect large increases in reports of maltreatment, but because the economic stresses of the pandemic and disruptions of families’ usual sources of support will likely extend well beyond the period of “stay at home” orders, the risks of family violence will persist for some time. Recognizing that risk for family violence is high may help increase monitoring of the families being served and provide timely anticipatory guidance (see recommendations in Table 1).

In addition, given the increased risk for trauma exposure, as well as anxiety and grief, during and after this crisis, identifying and managing

anger and stress that affect family interactions, screening for posttraumatic stress symptoms, and providing practical resources such as those published by the American Academy of Pediatrics and National Child Traumatic Stress Network are warranted. This role is consistent with the American Academy of Pediatrics statements regarding the pediatrician's role in supporting the well-being of children and families, particularly during challenging times, and in reducing risk for intimate partner violence and in child maltreatment prevention. Overburdened families will experience the highest risks of violence and will require the most support. Systems ordinarily available to them are likely to be compromised but need to remain sensitive and responsive to these needs. This is important because our collective response to COVID-19 through social distancing and isolation may

require these efforts for a prolonged period and will be unlike any challenge we have yet encountered.

ABBREVIATIONS

COVID-19: coronavirus disease
CPS: child protective services

REFERENCES

1. Brooks SK, Webster RK, Smith LE, et al. The psychological impact of quarantine and how to reduce it: rapid review of the evidence. *Lancet*. 2020;395(10227): 912–920
2. Vu NL, Jouriles EN, McDonald R, Rosenfield D. Children's exposure to intimate partner violence: a meta-analysis of longitudinal associations with child adjustment problems. *Clin Psychol Rev*. 2016;46:25–33
3. Herrenkohl TI, Sousa C, Tajima EA, Herrenkohl RC, Moylan CA. *Intersection of Child Abuse and Children's Exposure*

to Domestic Violence. Trauma, Violence, Wetter, Germany: Abus; 2008

4. Wildeman C, Emanuel N, Leventhal JM, Putnam-Hornstein E, Waldfogel J, Lee H. The prevalence of confirmed maltreatment among US children, 2004 to 2011. *JAMA Pediatr*. 2014;168(8): 706–713
5. Kim H, Drake B. Cumulative prevalence of onset and recurrence of child maltreatment reports. *J Am Acad Child Adolesc Psychiatry*. 2019;58(12): 1175–1183
6. US Department of Health and Human. Child maltreatment 2018. 2020. Available at: <https://www.acf.hhs.gov/cb/research-data-technology/statistics-research/child-maltreatment>. Accessed March 16, 2020
7. Disaster Preparedness Advisory Council; Committee on Pediatric Emergency Medicine. Ensuring the health of children in disasters. *Pediatrics*. 2015;136(5). Available at: www.pediatrics.org/cgi/content/full/136/5/e1407