

 Open access • Journal Article • DOI:10.1038/NATURE10176

Increased soil emissions of potent greenhouse gases under increased atmospheric CO₂ — [Source link](#)

Kees Jan van Groenigen, Kees Jan van Groenigen, Craig W. Osenberg, Bruce A. Hungate

Institutions: Northern Arizona University, Trinity College, Dublin, University of Florida

Published on: 14 Jul 2011 - Nature (Nature)

Topics: Greenhouse gas, Carbon sequestration, Radiative forcing, Carbon dioxide in Earth's atmosphere and Carbon sink

Related papers:

- [The meta-analysis of response ratios in experimental ecology](#)
- [Assessing the effect of elevated carbon dioxide on soil carbon: a comparison of four meta-analyses](#)
- [Effects of Elevated Carbon Dioxide and Increased Temperature on Methane and Nitrous Oxide Fluxes: Evidence from Field Experiments](#)
- [What have we learned from 15 years of free-air CO₂ enrichment \(FACE\)? A meta-analytic review of the responses of photosynthesis, canopy properties and plant production to rising CO₂.](#)
- [Nitrous oxide emissions from soils: how well do we understand the processes and their controls?](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/increased-soil-emissions-of-potent-greenhouse-gases-under-4ulrcpql6>

Increased soil emissions of potent greenhouse gases under increased atmospheric CO₂

Kees Jan van Groenigen, Craig W. Osenberg & Bruce A. Hungate

Nature 475, 214–216 (14 July 2011) doi:10.1038/nature10176

Abstract

Increasing concentrations of atmospheric carbon dioxide (CO₂) can affect biotic and abiotic conditions in soil, such as microbial activity and water content^{1, 2}. In turn, these changes might be expected to alter the production and consumption of the important greenhouse gases nitrous oxide (N₂O) and methane (CH₄) (refs 2, 3). However, studies on fluxes of N₂O and CH₄ from soil under increased atmospheric CO₂ have not been quantitatively synthesized. Here we show, using meta-analysis, that increased CO₂ (ranging from 463 to 780 parts per million by volume) stimulates both N₂O emissions from upland soils and CH₄ emissions from rice paddies and natural wetlands. Because enhanced greenhouse-gas emissions add to the radiative forcing of terrestrial ecosystems, these emissions are expected to negate at least 16.6 per cent of the climate change mitigation potential previously predicted from an increase in the terrestrial carbon sink under increased atmospheric CO₂ concentrations⁴. Our results therefore suggest that the capacity of land ecosystems to slow climate warming has been overestimated.


Introduction

By burning fossil fuels, cutting down forests and changing land use in other ways, humans are rapidly increasing the amount of CO₂ in the atmosphere and warming the planet⁵. Plant growth is known to increase after an abrupt surge in CO₂ levels⁶. Because stimulated assimilation of carbon by plants can increase soil carbon input and soil carbon storage, terrestrial ecosystems could help to reduce the increase in atmospheric CO₂ and thereby slow climate change⁷. However, the radiative forcing of land ecosystems is not determined by their uptake and release of CO₂ alone; increased CO₂ can also alter soil emissions of N₂O and CH₄ (ref. 2). Although both of these gases occur in far lower atmospheric concentrations than does CO₂, their global warming potentials are much higher: 298 times higher for N₂O and 25 times higher for CH₄ (ref. 5). Agricultural soils are the main source of human-induced N₂O emissions⁸. Soils under natural vegetation produce roughly the same amount of N₂O as all anthropogenic sources combined⁸. Wetlands, including rice paddies, contribute 32–53% to the global emissions of CH₄ (ref. 8). Upland soils, on the other hand, act as a sink for atmospheric CH₄ through oxidation by methanotrophic bacteria⁹. Thus, changes in N₂O and CH₄ fluxes could greatly alter how terrestrial ecosystems influence climate¹⁰.

Studies of greenhouse-gas (GHG) emissions span a variety of ecosystem types, and vary in experimental design and results, making it difficult to determine their global response to increased CO₂ from individual experiments. A quantitative synthesis of results across multiple studies can overcome this problem. Therefore, we used meta-analysis¹¹ to summarize the effect of atmospheric CO₂ enrichment on fluxes of CH₄ and N₂O from soil, using 152 observations from 49 published studies (see Supplementary Table 1, Supplementary Data 1 and 2, Supplementary Notes 1). We also summarized the effect of increased CO₂ on possible drivers of altered CH₄ and N₂O fluxes, using standing root biomass and soil water content from the studies in which the observations on N₂O and CH₄ fluxes were collected (Supplementary Data 3 and 4). All observations were analysed using three different weighting functions (see Methods). As CH₄ and N₂O emissions were not correlated with the concentration of CO₂ used for enrichment (Methods), we treat ‘increased CO₂’ as a category.

Overall, increased concentrations of atmospheric CO₂ stimulated emissions of N₂O by 18.8% (Fig. 1a). This positive response was significant for studies receiving little or no fertilizer, for non-pot studies and for studies on natural vegetation—that is, studies that most closely resembled real-world conditions (Supplementary Table 2). Increased CO₂ stimulated CH₄ emissions in wetlands by 13.2% (Fig. 1a, Supplementary Table 3). In rice paddies, increased CO₂ stimulated CH₄ emissions by 43.4% (Fig. 1a, Supplementary Table 4). In upland systems, increased CO₂ caused on average a small and insignificant net uptake of CH₄ (Supplementary Table 5).


Figure 1: Results of a meta-analysis of the response of GHG emissions and their potential drivers to rising levels of atmospheric CO₂.



To compare the relative importance of changed GHG fluxes in uplands, wetlands and rice paddies, we expressed the absolute effect of increased CO₂ on CH₄ and N₂O fluxes from these ecosystem types (Supplementary Tables 5–8) scaled by their respective total land area. For upland soils, we distinguished fertilized agricultural ecosystems and ecosystems receiving little or no fertilizer. Our estimates of total GHG fluxes under ambient (that is, present-day) CO₂ conditions correspond well to independent global syntheses of modern GHG fluxes (Supplementary Table 9), supporting our scaling approach.

The estimated stimulation by increased CO₂ of total soil N₂O emissions corresponds to an additional source of 0.33 Pg CO₂ equivalents (equiv.) yr⁻¹ from agricultural ecosystems (1 Pg = 10¹⁵ g), and of 0.24 Pg CO₂ equiv. yr⁻¹ for all other upland ecosystems (Fig. 2). The CO₂-stimulation of CH₄ emissions corresponds to an additional source of 0.25 Pg CO₂ equiv. yr⁻¹ from rice paddies and of 0.31 Pg CO₂ equiv. yr⁻¹ from natural wetlands. Our data indicate a small and non-significant effect of CO₂ on global CH₄ fluxes from upland soils for agricultural ecosystems (0.003 Pg CO₂ equiv. yr⁻¹) and for all other upland ecosystems (-0.011 Pg CO₂ equiv. yr⁻¹). The combined effect of increased CO₂ on emissions of these GHGs is 1.12 Pg CO₂ equiv. yr⁻¹.

Figure 2: The effect of rising atmospheric CO₂ on GHG emissions, expressed on the global scale.



For N₂O fluxes, the results for natural and agricultural soils were based on 35 and 19 observations, respectively. For CH₄ fluxes, the results for natural wetlands, rice paddies, natural upland soils and agricultural upland soils were based on 16, 21, 10 and 8 observations, respectively. Effect sizes in all meta-analyses were weighted by replication. Error bars, 95% confidence intervals.

Rising atmospheric CO₂ is expected to increase soil C storage in terrestrial ecosystems, which may contribute to the current residual C sink on land⁷. Meta-analysis of CO₂ enrichment experiments indicates that the sink is larger for ecosystems receiving fertilizer¹². Scaled up by the total area of agricultural and non-fertilized ecosystems, these meta-analyses suggest that increased atmospheric CO₂ levels may increase the soil C sink by as much as 4.0 Pg CO₂ yr⁻¹. Results presented here indicate that enhanced GHG emissions under increased CO₂ reduce the C mitigation effect of soil C storage by 28% (1.12 Pg/4.0 Pg). The magnitude and significance of this result is insensitive to the choice of the weighting function used in the meta-analysis (Supplementary Fig. 1, Supplementary Table 10).

Experiments included in our database increased atmospheric CO₂ concentration to 630 p.p.m.v. on average, a level expected for the second half of this century¹³. Biogeochemical models predict that at that time, the terrestrial C sink may be as much as 6.8 Pg CO₂ yr⁻¹ stronger than it is today⁴ (when considering forcing by rising CO₂ alone). On the basis of our analysis, a CO₂-induced rise in GHG fluxes could negate 16.6% (1.12 Pg/6.8 Pg) of the expected increase of the entire terrestrial C sink (Supplementary Table 10).

This estimate (16.6%) is likely to be an underestimate for three reasons. First, most of the studies in our data set measured GHG fluxes during the growing season only, but we assumed these applied to the entire year. Winter emissions

of CH₄ in wetlands and rice paddies are typically small⁹; however, winter emissions of N₂O during freeze-thaw cycles can contribute substantially to annual N₂O fluxes¹⁴, and available data indicate that winter emissions of N₂O are stimulated under increased CO₂ (ref. 15). A recently published data set¹⁶ suggests that N₂O emissions outside the growing season amount to 88% and 64% of the emissions during the growing season in agricultural systems and natural ecosystems, respectively (see Methods). Assuming that increased CO₂ affects N₂O emissions proportionately throughout the year, its effect on N₂O emissions outside the growing season would therefore amount to 0.29 Pg CO₂ equiv. yr⁻¹ from agricultural systems and 0.15 Pg CO₂ equiv. yr⁻¹ from natural ecosystems. Together, these fluxes negate an additional 7% of the expected increase of the terrestrial C sink.

Second, atmospheric N deposition is predicted to increase during this century¹⁷. Because average CO₂ responses of N₂O emissions were higher in studies receiving additional N (Supplementary Tables 2 and 6), the positive effect of CO₂ on N₂O emissions may strengthen as ecosystems become enriched in N.

Last, CO₂ effects on N₂O emissions showed a weak but significant correlation with experiment duration (Supplementary Fig. 2), suggesting that CO₂ effects on N₂O emissions may increase over time.

Why do GHG emissions respond positively to rising levels of atmospheric CO₂? Atmospheric CO₂ enrichment increased soil water contents for the studies contributing to our N₂O database (Fig. 1b, Supplementary Table 11); this result is probably due to improved efficiency of water use by plants, which reduces soil water loss through transpiration¹⁸. Moreover, increased CO₂ has been shown to enhance soil biological activity across a broad range of ecosystems^{1, 2}. Both responses promote soil anoxia, and thus stimulate denitrification¹⁹ (anaerobic microbial respiration of nitrate), one of the major sources of N₂O from soils³. Increased CO₂ also enhanced root biomass in all three habitats (Fig. 1b, Supplementary Table 12). As denitrification is generally stimulated by high availability of labile C as a source of energy²⁰, and because new C enters mineral soil mainly through the root system, this increase in root biomass would stimulate denitrification rates—and N₂O emissions—even further.

Methane is produced only under anaerobic conditions, which are common in soils of rice paddies and natural wetlands but not uplands. Because methanogenic archaea rely on C assimilation by plants as their ultimate source of organic substrates⁹, increased rates of soil C input with increased CO₂ can also stimulate CH₄ emissions. Indeed, the positive correlation between CH₄ emission rates and net ecosystem production in wetlands²¹ suggests that plant productivity is a key process in the regulation of CH₄ emission from these

ecosystems. The response to increased CO₂ of CH₄ emissions from rice paddies and wetlands showed significant correlation with the CO₂ response of root biomass ($r^2 = 0.17$, $P = 0.02$, Supplementary Fig. 6); this further suggests that increased CO₂ stimulates CH₄ production through its positive effect on plant growth and soil C input.

Global changes in climate and atmospheric composition have previously been suggested to affect GHG emissions from natural ecosystems. For instance, a global rise in temperature of 3.4 °C has been predicted to increase CH₄ emissions from wetlands by 78% (ref. 22). In addition to its direct effect on the global climate through radiative forcing, our results identify two indirect mechanisms through which rising atmospheric CO₂ amplifies climate change: by stimulating the release of N₂O from terrestrial ecosystems, and by enhancing CH₄ release from wetlands and rice paddies. The meta-analytic approach used here, synthesizing results across 49 studies, shows that increased N₂O and CH₄ emissions are both general and quantitatively important. Future assessments of terrestrial feedbacks to climate change should therefore consider these indirect effects of increased atmospheric CO₂ on the production by soil of trace gases like N₂O and CH₄.

Methods

Main Methods References Acknowledgements Author information
Supplementary information

Data collection

We extracted results for soil fluxes of CH₄ and N₂O, root biomass and soil water contents from atmospheric CO₂ enrichment studies, conducted in the field, in growth chambers or in glass houses. We used Google Scholar (Google Inc.) for an exhaustive search of journal articles published before January 2011, using as search terms either “elevated CO₂” or “CO₂ enrichment”, and either “N₂O” and “soil”, or “CH₄”. Further papers were added from a comparable search using Web of Science. For a study to be included in our data set, the atmospheric CO₂ concentration for the ambient and elevated treatments had to be in the range 350–450 p.p.m.v. and 450–800 p.p.m.v., respectively. Means and sample sizes had to be reported for both ambient and elevated CO₂ treatments.

For each study, we noted experimental duration, plant species, N fertilization rates and the type of experimental facility. Estimates of standard deviation were tabulated when available, but were not required for inclusion in the analysis. We included studies involving experiments in pots (that is, any container with dimensions <1 m) or in the field, and studies on natural or planted vegetation. We only considered studies in which soil under both CO₂ treatments had the same treatment history. One study was discarded for this reason. Studies on soil water content and root biomass were only included if data on N₂O or CH₄ fluxes

were available from the same site. When root biomass and soil water content were reported for multiple soil depths, we calculated the overall treatment effects across the entire soil profile. We included separate observations of increased CO₂ effects from a single ecosystem under different experimental treatments (that is, in multifactorial studies). Because wetlands are mostly anaerobic and therefore produce CH₄, whereas upland soils are mostly aerobic and oxidize CH₄, these two groups of ecosystems were considered in separate data sets. We also distinguished studies conducted in rice paddies, which like wetlands produce CH₄. Because the low number of studies on N₂O fluxes from rice paddies (1) and wetlands (3) did not warrant the construction of separate data sets, these studies were not included in our analysis.

We divided the studies into two categories of N availability based on N fertilization rates, that is, more or less than 30 kg N ha⁻¹ yr⁻¹. This cut-off point was chosen because it is comparable to maximum atmospheric N depositions in the US and most of the EU23. We also distinguished between studies on natural or planted vegetation. Agricultural ecosystems were defined as grassland and cropland that received between 30 and 300 kg N ha⁻¹ yr⁻¹. The upper cut-off point was based on reported average fertilization rates for croplands in the world's most intensively fertilized region (that is, East Asia, at 150 kg N ha⁻¹ yr⁻¹)¹⁶, and the assumption that average fertilizer N use per hectare will be twofold higher in 2050³⁰.

Response metrics

We evaluated our data sets by using meta-analysis. As a metric for the response of GHG emissions to increased CO₂, we used the natural log of the response ratio²⁴. This metric starts with an estimate of the relative change in GHG emissions between ambient and increased CO₂ treatments, and log-transforms it to improve its statistical behaviour.

$$\ln R = \ln(\text{GHG}_i - \text{GHG}_a)$$

where GHG is the flux of either CH₄ or N₂O under increased (i) or ambient (a) conditions. We also used lnR as a metric for CO₂ responses of root biomass and soil water contents. Fluxes of CH₄ from upland soils could not be analysed using this metric, because our data set included both sites with negative (that is, CH₄ uptake) and positive (CH₄ emissions) fluxes. For this reason, we also used the difference in annual emissions, expressed on an areal basis (U) as a metric:

$$U = (\text{GHG}_i - \text{GHG}_a)$$

with GHG_i and GHG_a as before. All but one study on wetland soils found net CH₄ emissions under both ambient and increased CO₂ conditions (Supplementary

Data 2). This one study, which reported that increased CO₂ turned wetland soils from a net sink of CH₄ into a net source, was therefore excluded when calculating lnR, but included when calculating U.

Several studies only measured N₂O and CH₄ fluxes during the growing season. In these cases, we assumed that the effect of increased CO₂ on annual fluxes occurred entirely during this period. When the length of the growing season was not explicitly indicated, we assumed a growing season of 150 days. When studies measured gas fluxes for multiple years, fluxes were averaged over time.

Weighting functions

We performed analyses using non-parametric weighting functions and generated confidence intervals (CIs) on weighted effects sizes using bootstrapping. Because effect size estimates and subsequent inferences in meta-analysis may depend on how individual studies are weighted¹², we used three different weighting functions. First, weighted by replication: $WR = (n_a \times n_i) / (n_a + n_i)$, where n_a and n_i are the number of replicates under ambient and increased CO₂, respectively²⁵. For pot studies, n equalled the number of replicate experimental facilities (that is, growth chambers, glass houses, and so on), rather than the number of pots per CO₂ treatment. Second, unweighted. Each observation was assigned an equal weight: $WU = 1$. Third, weighted by the inverse of the pooled variance, the weighting function conventionally used in meta-analyses²⁶: $WV = 1 / (var_a / GHG_a^2 + var_i / GHG_i^2)$, with GHG_a and GHG_i as before, and var_a and var_i as their respective variance.

When variance estimates were missing for a study, we calculated the average coefficient of variation (CV) within each data set, and then approximated the missing variance by multiplying the reported mean by the average CV and squaring the result.

When multiple effects were extracted from the same experimental site, we adjusted the weights defined above by the total number of observations from that site. This approach ensured that all experimental comparisons in multifactor studies could be included in the data set without dominating the overall effect size. For three experimental sites, multiple studies were done on the same GHG fluxes at different points in time. We adjusted the weights of observations from these studies by the total number of observations per site. Thus, the final weights used in the analyses were $w_{f,i} = W_{f,i} / n_c$, where n_c was the number of observations from the same site as the i th observation, and f was the index that referred to one of the three weighting functions defined above.

Mean effects sizes (,) for different categories of studies were estimated as:

We used METAWIN 2.127 to generate these mean effect sizes and 95% bootstrapped CIs (4,999 iterations). Treatment effects were considered significant if the 95% CI did not overlap with 0. The results for the analyses on lnR were back-transformed and reported as percentage change under increased CO₂ (that is, $100 \times (R - 1)$) to ease interpretation.

We tested whether lnR for GHG emissions was correlated with lnR for root biomass using the statistical package SPSS 19. Similarly, we tested whether lnR for GHG emissions was correlated with experiment duration or the level of CO₂ enrichment. The effect of increased CO₂ on soil emissions of N₂O, but not CH₄, showed a weak positive correlation with experiment duration (Supplementary Figs 2 and 3). lnR was not significantly correlated with the degree of CO₂ enrichment for either N₂O or CH₄ emissions (Supplementary Figs 4 and 5). This result is probably due to the large variation in treatment effects between studies, masking effects of the degree in CO₂ enrichment. Alternatively, the results may reflect that plant growth is a saturating function of CO₂ concentrations. Since experiments increased atmospheric CO₂ to a similar extent for all data sets (Supplementary Table 13), we did not normalize effect sizes for the level of CO₂ enrichment.

Results using the different weighting functions were qualitatively similar. However, the variance-based weighting function, W_v , yielded weights that varied over 1,000 times in magnitude (Supplementary Data 1 and 2). By assigning extreme importance to individual observations, average effect sizes were largely determined by a small number of studies. Because variance estimates are notoriously unreliable (especially given the small samples common in many of these studies), we favoured the use of the alternative weighting functions (which assigned less extreme weights). In this Letter, we provide results of the analyses on effect sizes that were weighted by replication; results for all weighting functions can be found in Supplementary Tables 2–8, 11 and 12.

Scaling of results

We scaled up the results from the experiments by multiplying them by the total land area covered by the particular type of habitat that was being summarized. In other words, we took the mean effects and confidence intervals for U calculated above and scaled them:

where F is expressed in $\text{Pg CO}_2 \text{ equiv. yr}^{-1}$, and H is the amount of habitat in uplands, wetlands, or rice paddies (103.1, 5.7, and 1.3 million km^2 , respectively^{28, 29}). Because N fertilization increases N_2O emissions^{16, 17} and enhances plant growth, we distinguished between upland agricultural ecosystems (that is, 19.0 million km^2 of fertilized grasslands and croplands¹⁶, minus 1.3 million km^2 of rice paddies²⁸) and ecosystems receiving little or no fertilizer ($103.1 - 19.0 + 1.3 = 85.4$ million km^2).

We estimated the contribution of winter N_2O emissions to total N_2O emissions from a recently published data set¹⁶. For agricultural soils and soils under natural vegetation, studies conducted over the growing season and lasting 100–200 days were compared to studies conducted over the entire year (that is, lasting >300 days). Because tropical and subtropical systems do not experience marked growing seasons, we excluded studies from those regions. For agricultural soils, we only considered studies on grassland and cropland receiving 30–300 $\text{kg N ha}^{-1} \text{ yr}^{-1}$ (that is, the same restrictions that applied to our data sets 1 and 2 for the global extrapolation shown in Fig. 2). The difference in mean N_2O emissions between the two categories of study duration was assumed to be representative of N_2O emissions outside the growing season.

To estimate the CI for the combined effect of increased CO_2 on all six GHG fluxes shown in Fig. 2, we calculated the square root of the sum of the squared CIs. Because the original CIs were asymmetric, we did this separately for the upper and lower CIs. All studies on rice paddies were conducted on planted vegetation, experimental conditions resembling real-world conditions. When we combined our extrapolated data to calculate the overall CO_2 effect on CH_4 emissions, we therefore included all available data from rice paddies (Fig. 2, Supplementary Fig. 1). To compare the emissions of GHG with soil C sequestration under increased CO_2 , we used results from the analyses weighted by replication and from unweighted analyses as reported in ref. 12, applying the same study selection criteria as for studies in our current data set. These results were expressed as a function of total land area, using the same approach that was used to scale up our results on GHG fluxes.

References

Zak, D. R., Pregitzer, K. S., King, J. S. & Holmes, W. E. Elevated atmospheric CO_2 , fine roots and the response of soil microorganisms: a review and hypothesis. *New Phytol.* 147, 201–222 (2000)

Pendall, E. et al. Below-ground process responses to elevated CO_2 and temperature: a discussion of observations, measurement methods, and models. *New Phytol.* 162, 311–322 (2004)

Smith, K. A. et al. Exchange of greenhouse gases between soil and atmosphere: interactions of soil physical factors and biological processes. *Eur. J. Soil Sci.* 54, 779–791 (2003)

Thornton, P. E., Lamarque, J.-F., Rosenbloom, N. A. & Mahowald, N. M. Influence of carbon-nitrogen cycle coupling on land model response to CO₂ fertilization and climate variability. *Glob. Biogeochem. Cycles* 21 GB4018 doi:10.1029/2006GB002868 (2007)

Forster, P. et al. in *Climate Change 2007: The Physical Science Basis* (eds Solomon, S. et al.) 129–234 (Cambridge Univ. Press, 2007)

Long, S. P., Ainsworth, E. A., Rogers, A. & Ort, D. R. Rising atmospheric carbon dioxide: plants FACE the future. *Annu. Rev. Plant Biol.* 55, 591–628 (2004)

Gifford, R. M. The global carbon cycle: a viewpoint on the missing sink. *Aust. J. Plant Physiol.* 21, 1–15 (1994)

Denman, K. L. et al. in *Climate Change 2007: The Physical Science Basis* (eds Solomon, S. et al.) 499–587 (Cambridge Univ. Press, 2007)

Le Mer, J. & Roger, P. Production, oxidation, emission and consumption of methane by soils: a review. *Eur. J. Soil Biol.* 37, 25–50 (2001)

Schulze, E. D. et al. Importance of methane and nitrous oxide for Europe's terrestrial greenhouse-gas balance. *Nature Geosci.* 2, 842–850 (2009)

Osenberg, C. W., Sarnelle, O., Cooper, S. D. & Holt, R. D. Resolving ecological questions through meta-analysis: goals, metrics, and models. *Ecology* 80, 1105–1117 (1999)

Hungate, B. A. et al. Assessing the effect of elevated carbon dioxide on soil carbon: a comparison of four meta-analyses. *Glob. Change Biol.* 15, 2020–2034 (2009)

Meehl, G. A. et al. in *Climate Change 2007: The Physical Science Basis* (eds Solomon, S. et al.) 747–845 (Cambridge Univ. Press, 2007)

Matzner, E. & Borren, W. Do freeze-thaw events enhance C and N losses from soils of different ecosystems? A review. *Eur. J. Soil Sci.* 59, 274–284 (2008)

Kammann, C., Müller, C., Grünhage, L. & Jäger, H.-J. Elevated CO₂ stimulates N₂O emissions in permanent grassland. *Soil Biol. Biochem.* 40, 2194–2205 (2008)

Stehfest, E. & Bouwman, L. N₂O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modelling of global annual emissions. *Nutr. Cycl. Agroecosyst.* 74, 207–228 (2006)

Galloway, J. N. et al. Transformation of the nitrogen cycle: recent trends, questions, and potential solutions. *Science* 320, 889–892 (2008)

Wullschleger, S. D., Tschaplinski, T. J. & Norby, R. J. Plant water relations at elevated CO₂ – implications for water-limited environments. *Plant Cell Environ.* 25, 319–331 (2002)

Smith, M. S. & Tiedje, J. M. Phases of denitrification following oxygen depletion in soil. *Soil Biol. Biochem.* 11, 261–267 (1979)

Weier, K. L., Doran, J. W., Power, J. F. & Walters, D. T. Denitrification and the dinitrogen/nitrous oxide ratio as affected by soil water, available carbon, and nitrate. *Soil Sci. Soc. Am. J.* 57, 66–72 (1993)

Whiting, G. J. & Chanton, J. P. Primary production control of methane emission from wetlands. *Nature* 364, 794–795 (1993)

Shindell, D. T., Walter, B. P. & Faluvegi, G. Impacts of climate change on methane emissions from wetlands. *Geophys. Res. Lett.* 31 L21202
doi:10.1029/2004GL021009 (2004)

Holland, E. A., Braswell, B. H., Sulzman, J. & Lamarque, J.-F. Nitrogen deposition onto the United States and western Europe: synthesis of observations and models. *Ecol. Appl.* 15, 38–57 (2005)

Hedges, L. V., Gurevitch, J. & Curtis, P. S. The meta-analysis of response ratios in experimental ecology. *Ecology* 80, 1150–1156 (1999)

Adams, D. C., Gurevitch, J. & Rosenberg, M. S. Resampling tests for meta-analysis of ecological data. *Ecology* 78, 1277–1283 (1997)

Hedges, L. V. & Olkin, I. *Statistical Methods for Meta-Analysis* (Academic, 1985)

Rosenberg, M. S., Adams, D. C. & Gurevitch, J. *METAWIN, Statistical Software for Meta-Analysis Version 2* (Sinauer, 2000)

World Resources Institute. Land area classification by ecosystem type. left fence <http://earthtrends.wri.org/datatables/index.php?theme=9> right fence (2003)

Aselmann, I. & Crutzen, P. J. Global distribution of natural freshwater wetlands and rice paddies, their net primary productivity, seasonality and possible methane emissions. *J. Atmos. Chem.* 8, 307–358 (1989)

Tilman, D. et al. Forecasting agriculturally driven global environmental change. *Science* 292, 281–284 (2001)

Download references

Acknowledgements

Main Methods References Acknowledgements Author information
Supplementary information

We thank S. A. Prior, G. B. Runion, F. Hagedorn, A. Niboyet, J. C. Blankinship, W. Cheng, T. Kanerva, R. S. Nowak, S. F. Zitzer, F. A. Dijkstra and J. P. Megonigal for sharing their data. Financial support for this study was provided by DOE-NICCR, NSF (DEB-0949460) and the Irish Research Council for Science, Engineering and Technology, co-funded by Marie Curie Actions under FP7.

Author information

Main Methods References Acknowledgements Author information
Supplementary information

Affiliations

Department of Biological Sciences, Northern Arizona University, Flagstaff, Arizona 86011, USA

Kees Jan van Groenigen & Bruce A. Hungate

Merriam-Powell Center for Environmental Research, Northern Arizona University, Flagstaff, Arizona 86011, USA

Kees Jan van Groenigen & Bruce A. Hungate

Department of Botany, School of Natural Sciences, Trinity College Dublin, Dublin 2, Ireland

Kees Jan van Groenigen

Department of Biology, University of Florida, Gainesville, Florida 32611-8525, USA

Craig W. Osenberg

Contributions

K.J.v.G. and B.A.H. designed the investigation. K.J.v.G. extracted the data from the literature and constructed the database. K.J.v.G. and C.W.O. performed the statistical analyses. All authors contributed to writing the paper.

Competing financial interests

The authors declare no competing financial interests.

Corresponding author

Correspondence to: Kees Jan van Groenigen