

# Increasing importance of deposition of reduced nitrogen in the United States

Yi Li<sup>a</sup>, Bret A. Schichtel<sup>b</sup>, John T. Walker<sup>c</sup>, Donna B. Schwede<sup>d</sup>, Xi Chen<sup>c</sup>, Christopher M. B. Lehmann<sup>e</sup>, Melissa A. Puchalski<sup>f</sup>, David A. Gay<sup>e</sup>, and Jeffrey L. Collett Jr.<sup>a,1</sup>

<sup>a</sup>Department of Atmospheric Science, Colorado State University, Fort Collins, CO 80523; <sup>b</sup>National Park Service, Cooperative Institute for Research in the Atmosphere, Colorado State University, Fort Collins, CO 80523; <sup>c</sup>Office of Research and Development, National Risk Management Research Laboratory, US Environmental Protection Agency, Research Triangle Park, NC 27711; <sup>d</sup>Office of Research and Development, National Exposure Research Laboratory, US Environmental Protection Agency, Research Triangle Park, NC 27711; <sup>e</sup>National Atmospheric Deposition Program, University of Illinois Urbana-Champaign, Champaign, IL 61820; and <sup>f</sup>Clean Air Markets Division, US Environmental Protection Agency, Washington, DC 20460

Edited by John H. Seinfeld, California Institute of Technology, Pasadena, CA, and approved March 31, 2016 (received for review December 31, 2015)

**Rapid development of agriculture and fossil fuel combustion greatly increased US reactive nitrogen emissions to the atmosphere in the second half of the 20th century, resulting in excess nitrogen deposition to natural ecosystems. Recent efforts to lower nitrogen oxides emissions have substantially decreased nitrate wet deposition. Levels of wet ammonium deposition, by contrast, have increased in many regions. Together these changes have altered the balance between oxidized and reduced nitrogen deposition. Across most of the United States, wet deposition has transitioned from being nitrate-dominated in the 1980s to ammonium-dominated in recent years. Ammonia has historically not been routinely measured because there are no specific regulatory requirements for its measurement. Recent expansion in ammonia observations, however, along with ongoing measurements of nitric acid and fine particle ammonium and nitrate, permit new insight into the balance of oxidized and reduced nitrogen in the total (wet + dry) US nitrogen deposition budget. Observations from 37 sites reveal that reduced nitrogen contributes, on average, ~65% of the total inorganic nitrogen deposition budget. Dry deposition of ammonia plays an especially key role in nitrogen deposition, contributing from 19% to 65% in different regions. Future progress toward reducing US nitrogen deposition will be increasingly difficult without a reduction in ammonia emissions.**

ammonia | dry deposition | wet deposition | nitrogen oxides | agriculture

Beginning in the mid-20th century, emissions of anthropogenic reactive nitrogen ( $N_r$ ) to the atmosphere accelerated rapidly due to increased fossil fuel combustion and intensive agricultural activities (1–4). Once emitted to the atmosphere,  $N_r$  compounds are deposited to terrestrial and aquatic ecosystems through dry and wet processes. Although nitrogen is an essential and often limiting element for ecosystems, increases in  $N_r$  deposition resulting from increased emissions have raised concerns around the world due to its adverse environmental impacts, including decreased biological diversity, increased soil acidification, and lake eutrophication (5–9). Critical loads (CLs) have been widely used to quantify levels of  $N_r$  deposition that ecosystems can sustain without significant harmful effects (10, 11). Twenty-four of the 45 national parks in the contiguous United States were estimated to receive  $N_r$  deposition in 2013 exceeding the local CL (12). Atmospheric  $N_r$  (an important ingredient of ozone and fine particle formation) has also been linked with climate change and human health degradation (8, 13, 14).

Atmospheric  $N_r$  sources are dominated by emissions of nitrogen oxides ( $NO_x = NO + NO_2$ ) and ammonia ( $NH_3$ ) (8).  $NO_x$  is produced by a wide range of high temperature processes including lightning and the combustion of fossil fuels by vehicles, electric power generating units, and other industrial and natural combustion sources.  $NO_x$  is oxidized in the atmosphere and converted to a variety of forms, including nitric acid, with a short timescale (typically 1 d or less). Reis et al. (15) attributed more than 80% of the  $NH_3$  emissions in the United States to the

agricultural sector, including emissions from livestock waste and volatilization of N-based fertilizer.

During the last two decades, US  $NO_x$  emissions have steadily declined due to effective regulations designed to decrease  $NO_x$  contributions to ozone, fine particles, and acid deposition (16). Data from the National Emissions Inventory (NEI) Air Pollutant Emissions Trends (<https://www.epa.gov/air-emissions-inventories/air-pollutant-emissions-trends-data>) indicate that  $NO_x$  emissions decreased by nearly 41% from 1990 to 2010. Further reductions are expected in coming years due to additional policy actions (e.g., the Cross-State Air Pollution Rule, reductions in mobile source emissions, and 5-y reviews of the National Ambient Air Quality Standards). Meanwhile,  $NH_3$  emissions have been reported to increase by 11% between 1990 and 2010 (17), with contributing factors including regional growth in livestock numbers and increased application of  $NO_x$  controls such as selective catalytic reduction. Unlike  $NO_x$ ,  $NH_3$  emissions in the United States are not regulated. In 2006, US anthropogenic  $NH_3$  emissions were estimated at 2.8 Tg N/y; they are projected to increase to between 3.3 and 4.2 Tg N/y by 2050, mainly due to increases in N fertilizer application and livestock growth (12).

## Significance

Human activities have greatly increased emissions of reactive forms of nitrogen to the atmosphere. This perturbation to the nitrogen cycle has produced large increases of nitrogen deposition to sensitive ecosystems. Over recent decades, attention has focused on wet and dry deposition of nitrate stemming from fossil fuel combustion emissions of nitrogen oxides. Successful decreases in nitrogen oxides emissions in the United States have substantially decreased nitrate deposition. By contrast, emissions of ammonia, an unregulated air pollutant, and resulting deposition of ammonium have grown. Expanded observations demonstrate that deposition of reactive nitrogen in the United States has shifted from a nitrate-dominated to an ammonium-dominated condition. Recognition of this shift is critical to formulating effective future policies to protect ecosystems from excess nitrogen deposition.

Author contributions: Y.L., B.A.S., J.T.W., and J.L.C. designed research; Y.L., J.T.W., D.B.S., and J.L.C. performed research; X.C., C.M.B.L., M.A.P., and D.A.G. contributed data; Y.L., J.T.W., and D.B.S. conducted model simulations; Y.L., B.A.S., J.T.W., D.B.S., C.M.B.L., M.A.P., D.A.G., and J.L.C. analyzed data; and Y.L., J.T.W., and J.L.C. wrote the paper with review and input from all authors.

The authors declare no conflict of interest.


This article is a PNAS Direct Submission.

Freely available online through the PNAS open access option.

Data deposition: All datasets related to this manuscript have been deposited in a repository operated by the Colorado State University, <https://dspace.library.colostate.edu/handle/10217/171382>.

<sup>1</sup>To whom correspondence should be addressed. Email: [collett@atmos.colostate.edu](mailto:collett@atmos.colostate.edu).

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1525736113/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1525736113/-DCSupplemental).



**Fig. 1.** Comparisons of the 3-y average  $\text{NH}_4^+$  percentage of wet inorganic nitrogen deposition across the United States in 1990–1992 (*Left*) and 2010–2012 (*Right*). To help visualize spatial patterns, isopleths were produced by interpolating  $\text{NH}_4^+$  mole percentages at individual monitoring sites using a cubic inverse-distance weighting of sites within 500 km of each observation station. The black dots on the map represent locations of sites with 3-y data available for each time period. The  $\text{NH}_4^+$  percentage on a molar basis [ $(\text{NH}_4^+ \%) = (\text{NH}_4^+) / (\text{NO}_3^- + \text{NH}_4^+) \times 100\%$ ] is noted at each site.

Through dry deposition processes, gaseous  $\text{HNO}_3$  and  $\text{NH}_3$  are rapidly removed from the atmosphere and deposited to surface ecosystems.  $\text{HNO}_3$  and  $\text{NH}_3$  can both be incorporated into atmospheric particles; this includes their reaction with each other to form fine particle ammonium nitrate, reaction of ammonia with sulfuric acid to form fine particle ammonium sulfate, and reactions of nitric acid with soil dust or sea salt to form coarse particle nitrate (18), among other species. These aerosol particles can also deposit nitrate and/or ammonium via dry deposition. Nitric acid, ammonia, and particulate nitrate and ammonium are scavenged by clouds and precipitation, producing wet deposition of ammonium and nitrate.  $\text{HNO}_3$  and  $\text{NO}_3^-$  are generally referred to as oxidized N, whereas  $\text{NH}_3$  and  $\text{NH}_4^+$  are termed reduced N; both oxidized and reduced N represent significant  $\text{N}_r$  inputs in unmanaged ecosystems (19, 20).

Publications from the 1990s and early 2000s recognized that deposition of oxidized N dominated the US atmospheric reactive N deposition budget (21, 22). With the ensuing emissions reductions of  $\text{NO}_x$  and emissions increases of  $\text{NH}_3$ , relative contributions of oxidized and reduced N have likely changed in recent years. Modeling studies (12, 23) have suggested this change and point to its likely continuation. A recent analysis of US National Atmospheric Deposition Program (NADP) observations (24) illustrates decreases in US wet nitrate deposition and increases in wet ammonium deposition in many regions. Focusing on the Midwest and Eastern United States, Sickles and Shadwick (25) reported that reductions in  $\text{NO}_x$  emissions have decreased oxidized nitrogen dry and wet deposition and that particulate ammonium concentrations now exceed concentrations of particulate nitrate plus gaseous nitric acid in the region. Ammonia gas was not considered in their study.

From an observational perspective, the absence of gaseous ammonia observations has prevented a broad, national understanding of the overall contributions of oxidized and reduced forms of inorganic N to the total  $N_r$  deposition budget. Making use of longer-term wet and dry deposition records and newly available  $NH_3$  measurements from regions across the country, we examine the overall contributions of oxidized and reduced forms of inorganic nitrogen to wet, dry, and total deposition budgets to better inform discussions of strategies to decrease N deposition to sensitive ecosystems.


## Results and Discussion

Analyses of wet deposition records provide important insight into the shift from oxidized to reduced nitrogen deposition across the contiguous United States. Recently expanded measurements of gas and particle phase reactive nitrogen species permit an assessment

of current contributions of oxidized and reduced nitrogen to the US  $N_r$  dry deposition budget. By combining these analyses, we gain a better understanding of the importance of both oxidized and reduced nitrogen to the total (wet + dry)  $N_r$  deposition budget across much of the United States.

**Oxidized vs. Reduced N in Wet N Deposition.** Although wet N deposition was dominated by oxidized N ( $\text{NO}_3^-$ ) across much of the country in the early 1990s, most locations now receive a majority of their wet N deposition as reduced N ( $\text{NH}_4^+$ ) (Fig. 1), a trend also recently reported by Du et al. (24). During the period 1990–1992, 69% of the observation sites were subjected to oxidized N contributions in excess of 50%; 20 y later, 69% of the sites instead received wet deposition of reduced N greater than 50%.


Changes in fractional contributions of oxidized and reduced N depend on the combined changes in wet deposition fluxes of  $\text{NH}_4^+$  and  $\text{NO}_3^-$ . Fig. 2 examines these changes for 45 of the 48 contiguous United States with available data. In every state but North Dakota, nitrate wet deposition fluxes decreased, with an average decrease of 29%. The nationwide decrease of oxidized N in wet deposition is consistent with the downward trend of US  $\text{NO}_x$  emissions. With the successful implementation of the Clean Air Act (CAA) and the 1990 Amendments,  $\text{NO}_x$  emissions were estimated to decline by 36% between 1990 and 2008 (13).



**Fig. 2.** Absolute percentage change of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  in wet deposition across the country.  $\text{C}_{10-12}$  is the average  $\text{NH}_4^+$  or  $\text{NO}_3^-$  flux (kg N/ha per year) in each state between 2010 and 2012 and  $\text{C}_{90-92}$  is the average  $\text{NH}_4^+$  or  $\text{NO}_3^-$  flux (kg N/ha per year) between 1990 and 1992. Only sites in Fig. 1 with both 1990–1992 and 2010–2012 data available are used to calculate the average flux for each state.








**Fig. 4.** Ratio of annual  $\text{NH}_3$  dry deposition rates estimated using the MLM vs. bidirectional approaches. Regions are indicated at top of graph. \*Due to a lack of meteorological data, the bidirectional flux model is not parameterized appropriately for site KY98. \*\*Due to vegetation type, the bidirectional flux model is not parameterized appropriately for site IL11.

here suppresses summertime atmospheric concentrations and therefore dry deposition of reduced and oxidized N species. In Florida wet N deposition contributed more than 75% of total (wet + dry) inorganic N deposition during summer when there was more precipitation (Fig. S1); dry deposition of reduced N was the dominant input during the drier winter season.

At the annual scale, ammonia dry deposition rates estimated using the multilayer model (MLM) approach are larger than those derived from the bidirectional model by a factor of 1.90 (median MLM/bidirectional flux ratio of 35 sites listed in Fig. 4). A reduction in  $\text{NH}_3$  dry deposition rates, relative to the unidirectional flux framework, was also observed on implementation of  $\text{NH}_3$  bidirectionality in the Community Multiscale Air Quality Model (33). MLM vs. bidirectional model differences vary across regions but generally result from stomatal and ground compensation points, as well as the effects of surface acidity, represented in the bidirectional framework. The net result of these processes is to reduce the atmosphere-surface  $\text{NH}_3$  concentration gradient,

and therefore the flux, relative to the unidirectional MLM deposition velocity approach, which assumes a zero surface concentration. Model differences are generally greatest in summer (Fig. S2) when temperature driven stomatal and soil compensation points are at a maximum. On average, the bidirectional and MLM approaches yield comparable net fluxes during winter when compensation points are lowest and surfaces are more acidic. Further discussion of the MLM vs. the bidirectional model is included in *SI Methods*.

The relatively large overall differences between the MLM and bidirectional  $\text{NH}_3$  flux estimates warrant brief discussion of the significant uncertainties that persist in modeling dry deposition of reactive nitrogen. MLM  $\text{HNO}_3$  deposition velocities, on which the  $\text{NH}_3$  deposition velocity used here is based, may contain up to  $\pm 25\%$  uncertainty related to error in the measurements that drive the model and underlying process parameterizations (34). Models of  $\text{HNO}_3$  deposition velocity also differ substantially themselves. For example, a multisite evaluation of the MLM and Big Leaf Model used in the Canadian Air and Precipitation Monitoring Network (CAPMoN) showed a median bias in hourly  $\text{HNO}_3$  deposition velocities of  $-35\%$ , with MLM yielding lower values (35). Regarding bidirectional  $\text{NH}_3$  models, comparisons to average measured  $\text{NH}_3$  fluxes over nonagricultural ecosystems generally demonstrate agreement within  $\pm 30\%$  using site-specific process parameterizations (36–39), although differences can be much larger under specific meteorological and surface conditions. Uncertainty may also be significantly larger when applying generalized parameterizations as done here. In that regard, recent versions of bidirectional models (40–42) have not yet been rigorously compared with each other or against flux measurements for natural ecosystems in North America. Although our analyses use commonly used approaches for both  $\text{HNO}_3$  deposition velocity and bidirectional  $\text{NH}_3$  flux, the abovementioned uncertainties are included to emphasize that the dry deposition component of the N deposition budget is significantly more uncertain



**Fig. 5.** Spatial trends in total reactive inorganic N deposition across the United States from July 2011 to June 2013. Fractional reduced N contributions to total N deposition (dry + wet) at the 37 sites are represented by circle color. The total inorganic nitrogen deposition is indicated by circle size. The pie charts show average fractional contributions of individual reduced and oxidized N deposition pathways for the eight regions, with each pie area proportional to the average total inorganic nitrogen deposition (also listed under each pie).

than the wet fraction, a point that should be considered in the interpretation of our results.

**Fractional Reduced N Contributions to the Total Inorganic N Deposition Budget.** With wet and dry deposition estimates available for 37 locations, the total wet plus dry nitrogen deposition budgets can be estimated across the United States (Fig. 5). Fractional deposition contributions by each wet and dry deposition pathway for each of the eight regions are also illustrated in Fig. 5. Reduced N deposition fractions in the eight regions range from 58% (Washington, I) to 78% (Southwest, VIII), with dry  $\text{NH}_3$  deposition alone contributing between 19% (Northwest, II) and 63% (Southwest, VIII). Fractional reduced N contributions at individual sites range from 42% at CO99 (Mesa Verde National Park) to 84% at CA83 in California's Central Valley. Ammonia dry deposition fractions ranged from 11% (PA27) to 74% (CA83). The spatial patterns of reduced N deposition fraction generally reflect spatial variations in agricultural activity including animal husbandry. Assuming that the biases between the MLM deposition velocity and bidirectional flux approaches shown in Fig. 4 are generally representative, a full assessment using the bidirectional approach would, at many sites, reduce overall deposition rates and the relative fraction of  $\text{NH}_3$  dry deposition. However, the general pattern observed in Fig. 5 remains consistent;  $\text{NH}_x$  still contributes the majority of inorganic N deposition at the national scale.

The site-specific circle sizes in Fig. 5 indicate the combined wet plus dry inorganic N deposition fluxes. Some regions exhibit majority dry deposition [e.g., dry deposition contributions of 58% and 79% in the Upper Midwest (IV) and Southwest (VIII), respectively], whereas others are more strongly influenced by wet deposition [e.g., wet deposition contributions of 66% and 72% in the Washington (I) and Southeast (VI) regions, respectively]. The largest deposition fluxes at individual sites tend to be observed at locations where fractional reduced N contributions are large. The maximum regional average inorganic N deposition flux (12.1 kg N/ha per year) was observed in the Upper Midwest region (IV); relatively large deposition fluxes were also observed for California and the eastern United States. These spatial patterns are similar to those identified in recent model simulations (43).

**Implications and Summary.** Increases in agricultural emissions of ammonia and the success of regulatory policies in decreasing  $\text{NO}_x$  emissions over the last two decades are changing the face of US reactive nitrogen deposition. Although US wet inorganic N deposition was once dominated by nitrate, wet inorganic N deposition now comes mostly from ammonium at nearly 70% of US monitoring sites. Although estimates of dry deposition fluxes of inorganic N inherently contain more uncertainty, dry and total (wet plus dry) inorganic N deposition fluxes also appear to be dominated by reduced N in most parts of the country. Decreases in wet and dry deposition fluxes of oxidized inorganic N species are expected to continue into the future as the United States continues to lower  $\text{NO}_x$  emissions. Current projections of ammonia emissions growth, meanwhile, suggest that reduced N deposition levels will grow in the future. In addition to the adverse impacts of reduced N deposition on ecosystem health, ammonia is an important precursor to fine particle formation. Fine particles decrease visibility (44) and negatively impact human health and increase health care costs (45, 46). Reductions in US ammonia emissions from agricultural and nonagricultural sources, whether by regulation or voluntary actions (e.g., agricultural producer adoption of best management practices), would yield a variety of positive

benefits for ecosystems and society. Increased study of atmospheric ammonia concentrations and improved measures of ammonia dry deposition fluxes are needed to design optimal strategies for achieving such benefits.

## Methods

Weekly precipitation concentrations of  $\text{NH}_4^+$  and  $\text{NO}_3^-$  were obtained from the NADP National Trends Network (NTN; [nadp.isws.illinois.edu/ntn/](http://nadp.isws.illinois.edu/ntn/)). Weekly gaseous  $\text{HNO}_3$  concentrations and particulate  $\text{NH}_4^+$  and  $\text{NO}_3^-$  concentrations were obtained from the Clean Air Status and Trends Network (CASTNET; <https://www.epa.gov/castnet>). Biweekly concentrations of gaseous  $\text{NH}_3$  were taken from the NADP AMoN ([nadp.isws.illinois.edu/AMoN/](http://nadp.isws.illinois.edu/AMoN/)). To gain greater spatial coverage of airborne  $\text{NH}_3$  concentrations, especially in the western United States,  $\text{NH}_x$  ( $\text{NH}_3 + \text{NH}_4^+$ ) measurements from a pilot IMPROVE  $\text{NH}_x$  monitoring network (28) were also used. More detailed information about these observation networks can be found in Table S2.

Wet deposition data were obtained from NTN sites for the periods 1990–1992 and 2010–2012. The number of sites analyzed changed due to network development over this period. From 1990 to 1992, there were 195 sites; 238 sites were available for the 2010–2012 period. Sites were not included if data were unavailable for  $\geq 1$  y in either period examined.

Oxidized and reduced N gas and particle concentrations were obtained for 37 sites (Table S1) where NTN and CASTNET monitoring stations were collocated with AMoN and/or IMPROVE  $\text{NH}_x$  sites. At 30 of these locations, 2 y of measurements (July 2011 to June 2013) were available. The remaining sites had data availability of at least 1 y. Concentrations of all species that contribute to the N<sub>i</sub> deposition budget are not measured at these sites. Important missing compounds include inorganic (e.g.,  $\text{NO}_2$ ) and organic N (e.g., alkyl nitrates, peroxyacetyl nitrate, and amines) species. Wet deposition of organic N is also not routinely measured and therefore not considered in this analysis.

Wet N deposition was determined from the amount of total precipitation and the aqueous concentrations of  $\text{NH}_4^+$  and  $\text{NO}_3^-$ . Dry N deposition was calculated for each species as the product of the N species concentration and a deposition velocity. Deposition velocities of gaseous  $\text{HNO}_3$  and particulate  $\text{NH}_4^+$  and  $\text{NO}_3^-$  were provided by CASTNET for each of its measurement sites based on the MLM (47), with input of on-site meteorology and local site characteristics. Gaps in the meteorological data were addressed by using the CASTNET substitution method (48). The deposition velocity of  $\text{NH}_3$  is difficult to determine due to the bidirectional nature of the dry  $\text{NH}_3$  flux that depends strongly on local conditions (40). To estimate dry  $\text{NH}_3$  deposition here, its deposition velocity was calculated as 70% of the  $\text{HNO}_3$  deposition velocity following previous estimates (49–51). A review of field observations suggests that the  $\text{NH}_3$  deposition velocity is at least half and perhaps as high as the  $\text{HNO}_3$  deposition velocity. Our choice of 70% agrees well with the findings of Neirynck et al. (52) and Nemitz et al. (39).

This  $\text{NH}_3$  deposition velocity approach is a simple approximation of unidirectional air-surface exchange, ignoring important bidirectional exchange processes that influence the magnitude and direction of the flux. To assess the potential importance of these bidirectional exchange processes and their impact on annual reactive N deposition budgets,  $\text{NH}_3$  fluxes derived from the unidirectional approach were compared with fluxes estimated using a two-layer bidirectional flux model (53). The bidirectional model uses hourly CASTNET meteorology and 2-wk integrated AMoN  $\text{NH}_3$  concentrations to estimate  $\text{NH}_3$  exchange with soil and vegetation, as well as net fluxes above the vegetation. Ammonia compensation points and leaf surface resistances were parameterized following the recommendations of Massad et al. (40) for natural vegetation. Development of this modeling framework is ongoing. Thus, the comparison is constrained to the dominant natural vegetation type at each site for which the Massad et al. (40) parameterizations are applicable, which excludes fertilized and nitrogen fixing crops and some other surfaces specified by CASTNET, including water, sand, and rock. Details of the bidirectional model and comparison are included in SI Methods.

**ACKNOWLEDGMENTS.** We thank Amy Sullivan, Katie Benedict, Aohan Tang, Qijing Bian, Arsineh Hecobian, and Yixing Shao (Colorado State University); Brian Michael (University of Illinois); and Nate Hyde (WaveMetrics) for helpful suggestions and support. Funding for this work was provided by the National Park Service.

- Galloway JN, et al. (2008) Transformation of the nitrogen cycle: Recent trends, questions, and potential solutions. *Science* 320(5878):889–892.
- Erisman JW, Galloway J, Seitzinger S, Bleeker A, Butterbach-Bahl K (2011) Reactive nitrogen in the environment and its effect on climate change. *Curr Opin Environ Sustain* 3(5):281–290.

- Galloway JN, Cowling EB (2002) Reactive nitrogen and the world: 200 years of change. *AMBIO* 31(2):64–71.
- Liu X, et al. (2013) Enhanced nitrogen deposition over China. *Nature* 494(7438):459–462.
- Clark CM, Tilman D (2008) Loss of plant species after chronic low-level nitrogen deposition to prairie grasslands. *Nature* 451(7179):712–715.

6. Phoenix GK, et al. (2012) Impacts of atmospheric nitrogen deposition: Responses of multiple plant and soil parameters across contrasting ecosystems in long-term field experiments. *Glob Change Biol* 18(4):1197–1215.
7. Holtgrieve GW, et al. (2011) A coherent signature of anthropogenic nitrogen deposition to remote watersheds of the Northern Hemisphere. *Science* 334(6062):1545–1548.
8. Galloway JN, et al. (2004) Nitrogen cycles: Past, present, and future. *Biogeochemistry* 70(2):153–226.
9. Janssens IA, et al. (2010) Reduction of forest soil respiration in response to nitrogen deposition. *Nat Geosci* 3(5):315–322.
10. Nilsson J (1988) *Critical Loads for Sulphur and Nitrogen*. Air Pollution and Ecosystems (Springer, New York), pp 85–91.
11. Porter E, Blett T, Potter DU, Huber C (2005) Protecting resources on federal lands: Implications of critical loads for atmospheric deposition of nitrogen and sulfur. *Bioscience* 55(7):603–612.
12. Ellis RA, et al. (2013) Present and future nitrogen deposition to national parks in the United States: Critical load exceedances. *Atmos Chem Phys* 13(17):9083–9095.
13. Davidson E, et al. (2012) Excess nitrogen in the U.S. environment: Trends, risks, and solutions. *Issues Ecol* 15(Winter 2012):1–16.
14. Gruber N, Galloway JN (2008) An Earth-system perspective of the global nitrogen cycle. *Nature* 451(7176):293–296.
15. Reis S, Pinder R, Zhang M, Lijie G, Sutton M (2009) Reactive nitrogen in atmospheric emission inventories. *Atmos Chem Phys* 9(19):7657–7677.
16. Hand JL, et al. (2014) Widespread reductions in haze across the United States from the early 1990s through 2011. *Atmos Environ* 94:671–679.
17. Xing J, et al. (2013) Historical gaseous and primary aerosol emissions in the United States from 1990 to 2010. *Atmos Chem Phys* 13(15):7531–7549.
18. Lee T, et al. (2008) Observations of fine and coarse particle nitrate at several rural locations in the United States. *Atmos Environ* 42(11):2720–2732.
19. Galloway JN, Cowling EB, Seitzinger SP, Socolow RH (2002) Reactive nitrogen: Too much of a good thing? *AMBIO* 31(2):60–63.
20. Fowler D, et al. (1998) Regional mass budgets of oxidized and reduced nitrogen and their relative contribution to the nitrogen inputs of sensitive ecosystems. *Environ Pollut* 102(1 suppl):337–342.
21. Bytnerowicz A, Fenn ME (1996) Nitrogen deposition in California forests: A review. *Environ Pollut* 92(2):127–146.
22. Holland EA, Braswell BH, Sulzman J, Lamarque J-F (2005) Nitrogen deposition onto the United States and Western Europe: A synthesis of observations and models. *Ecol Appl* 15(1):38–57.
23. Paulot F, Jacob DJ, Henze DK (2013) Sources and processes contributing to nitrogen deposition: An adjoint model analysis applied to biodiversity hotspots worldwide. *Environ Sci Technol* 47(7):3226–3233.
24. Du E, de Vries W, Galloway JN, Hu X, Fang J (2014) Changes in wet nitrogen deposition in the United States between 1985 and 2012. *Environ Res Lett* 9(9):095004.
25. Sickles II, Shadwick D (2015) Air quality and atmospheric deposition in the eastern US: 20 years of change. *Atmos Chem Phys* 15(1):173–197.
26. Lehmann CM, Gay DA (2011) Monitoring long-term trends of acidic wet deposition in US precipitation: Results from the National Atmospheric Deposition Program. *Powerpl Chem* 13(7):386–393.
27. Lehmann CM, Bowersox VC, Larson RS, Larson SM (2007) *Monitoring Long-Term Trends in Sulfate and Ammonium in US Precipitation: Results from the National Atmospheric Deposition Program/National Trends Network*. Acid Rain-Deposition to Recovery (Springer, New York), pp 59–66.
28. Chen X, et al. (2014) Seasonal ambient ammonia and ammonium concentrations in a pilot IMPROVE NH<sub>3</sub> monitoring network in the western United States. *Atmos Environ* 91:118–126.
29. Rodriguez MA, Barna MG, Moore T (2009) Regional impacts of oil and gas development on ozone formation in the western United States. *J Air Waste Manag Assoc* 59(9):1111–1118.
30. Brunke R, Alvo P, Schuepp P, Gordon R (1988) Effect of meteorological parameters on ammonia loss from manure in the field. *J Environ Qual* 17(3):431–436.
31. Sommer SG, Olesen JE, Christensen BT (1991) Effects of temperature, wind speed and air humidity on ammonia volatilization from surface applied cattle slurry. *J Agric Sci* 117(01):91–100.
32. Li Y, et al. (2014) Observations of ammonia, nitric acid, and fine particles in a rural gas production region. *Atmos Environ* 83(0):80–89.
33. Bash JO, Cooter EJ, Dennis RL, Walker JT, Pleim JE (2013) Evaluation of a regional air-quality model with bidirectional NH<sub>3</sub> exchange coupled to an agroecosystem model. *Biogeosciences* 10(3):1635–1645.
34. Cooter EJ, Schwede DB (2000) Sensitivity of the National Oceanic and Atmospheric Administration multilayer model to instrument error and parameterization uncertainty. *J Geophys Res, D, Atmospheres* 105(D5):6695–6707.
35. Schwede D, Zhang L, Vet R, Lear G (2011) An intercomparison of the deposition models used in the CASTNET and CAPMoN networks. *Atmos Environ* 45(6):1337–1346.
36. Flechard CR, Fowler D, Sutton MA, Cape JN (1999) A dynamic chemical model of bi-directional ammonia exchange between semi-natural vegetation and the atmosphere. *Q J R Meteorol Soc* 125(559):2611–2641.
37. Neirynck J, Ceulemans R (2008) Bidirectional ammonia exchange above a mixed coniferous forest. *Environ Pollut* 154(3):424–438.
38. Kruit RJW, et al. (2010) Modeling the surface-atmosphere exchange of ammonia. *Atmos Environ* 44(7):945–957.
39. Nemitz E, Sutton M, Wyers G, Jongejan P (2004) Gas-particle interactions above a Dutch heathland: I. Surface exchange fluxes of NH<sub>3</sub>, SO<sub>2</sub>, HNO<sub>3</sub> and HCl. *Atmos Chem Phys* 4(4):989–1005.
40. Massad R-S, Nemitz E, Sutton M (2010) Review and parameterisation of bi-directional ammonia exchange between vegetation and the atmosphere. *Atmos Chem Phys* 10(21):10359–10386.
41. Zhang L, Wright L, Asman W (2010) Bi-directional air-surface exchange of atmospheric ammonia: A review of measurements and a development of a big-leaf model for applications in regional-scale air-quality models. *J Geophys Res, D, Atmospheres* 115:D20310.
42. Pleim JE, Bash JO, Walker JT, Cooter EJ (2013) Development and evaluation of an ammonia bidirectional flux parameterization for air quality models. *J Geophys Res, D, Atmospheres* 118(9):3794–3806.
43. Schwede DB, Lear GG (2014) A novel hybrid approach for estimating total deposition in the United States. *Atmos Environ* 92(0):207–220.
44. Malm WC (1999) *Introduction to Visibility* (Cooperative Institute for Research in the Atmosphere, NPS Visibility Program, Colorado State University, Fort Collins, CO).
45. Paulot F, Jacob DJ (2014) Hidden cost of U.S. agricultural exports: Particulate matter from ammonia emissions. *Environ Sci Technol* 48(2):903–908.
46. Stokstad E (2014) Air pollution. Ammonia pollution from farming may exact hefty health costs. *Science* 343(6168):238.
47. Meyers TP, Finkelstein P, Clarke J, Ellestad TG, Sims PF (1998) A multilayer model for inferring dry deposition using standard meteorological measurements. *J Geophys Res, D, Atmospheres* 103(D17):22645–22661.
48. Bowker GE, Schwede DB, Lear GG, Warren-Hicks WJ, Finkelstein PL (2011) Quality assurance decisions with air models: A case study of imputation of missing input data using EPA's multi-layer model. *Water Air Soil Pollut* 222(1-4):391–402.
49. Benedict KB, et al. (2013) A seasonal nitrogen deposition budget for Rocky Mountain National Park. *Ecol Appl* 23(5):1156–1169.
50. Beem KB, et al. (2010) Deposition of reactive nitrogen during the Rocky Mountain Airborne Nitrogen and Sulfur (RoMANS) study. *Environ Pollut* 158(3):862–872.
51. Benedict KB, et al. (2013) Observations of atmospheric reactive nitrogen species in Rocky Mountain National Park and across northern Colorado. *Atmos Environ* 64:66–76.
52. Neirynck J, et al. (2007) Fluxes of oxidised and reduced nitrogen above a mixed coniferous forest exposed to various nitrogen emission sources. *Environ Pollut* 149(1):31–43.
53. Nemitz E, Milford C, Sutton MA (2001) A two-layer canopy compensation point model for describing bi-directional biosphere-atmosphere exchange of ammonia. *Q J R Meteorol Soc* 127(573):815–833.
54. Langford A, Fehsenfeld F, Zachariassen J, Schimel D (1992) Gaseous ammonia fluxes and background concentrations in terrestrial ecosystems of the United States. *Global Biogeochem Cycles* 6(4):459–483.
55. Farquhar GD, Firth PM, Wetselaar R, Weir B (1980) On the gaseous exchange of ammonia between leaves and the environment: Determination of the ammonia compensation point. *Plant Physiol* 66(4):710–714.
56. Hicks B, Baldocchi D, Meyers T, Hosker R, Jr, Matt D (1987) A preliminary multiple resistance routine for deriving dry deposition velocities from measured quantities. *Water Air Soil Pollut* 36(3-4):311–330.
57. Schuepp PH (1977) Turbulent transfer at the ground: On verification of a simple predictive model. *Boundary-Layer Meteorol* 12(2):171–186.
58. Wesely M (1989) Parameterization of surface resistances to gaseous dry deposition in regional-scale numerical models. *Atmospheric Environ* (1967) 23(6):1293–1304.
59. Hill PW, Raven JA, Loubet B, Fowler D, Sutton MA (2001) Comparison of gas exchange and bioassay determinations of the ammonia compensation point in *Luzula sylvatica* (Huds.) Gaud. *Plant Physiol* 125(1):476–487.