

Inequalities of Čebyšev Type for Lipschitzian Functions in Banach Algebras

Marius V. Boldea

Abstract. In this paper we give some Čebyšev type norm inequalities for two Lipschitzian functions on Banach algebras. Some examples for power function, exponential and the resolvent functions are also provided.

AMS Subject Classification (2000). 47A63; 47A99.

Keywords. Banach algebras, Power series, Exponential function, Resolvent function, Norm inequalities

1 Introduction

Let \mathcal{B} be an algebra. An *algebra norm* on \mathcal{B} is a map $\|\cdot\| : \mathcal{B} \rightarrow [0, \infty)$ such that $(\mathcal{B}, \|\cdot\|)$ is a normed space, and, further

$$\|ab\| \leq \|a\| \|b\|$$

for any $a, b \in \mathcal{B}$. The normed algebra $(\mathcal{B}, \|\cdot\|)$ is a *Banach algebra* if $\|\cdot\|$ is a *complete norm*. We assume that the Banach algebra is *unital*, this means that \mathcal{B} has an identity 1 and that $\|1\| = 1$.

Let \mathcal{B} be a unital algebra. An element $a \in \mathcal{B}$ is *invertible* if there exists an element $b \in \mathcal{B}$ with $ab = ba = 1$. The element b is unique; it is called the *inverse* of a and written a^{-1} or $\frac{1}{a}$.

The set of invertible elements of \mathcal{B} is denoted by $\text{Inv}\mathcal{B}$. If $a, b \in \text{Inv}\mathcal{B}$ then $ab \in \text{Inv}\mathcal{B}$ and $(ab)^{-1} = b^{-1}a^{-1}$.

For a unital Banach algebra we also have:

- (i) If $a \in \mathcal{B}$ and $\lim_{n \rightarrow \infty} \|a^n\|^{1/n} < 1$, then $1 - a \in \text{Inv}\mathcal{B}$;
- (ii) $\{a \in \mathcal{B}: \|1 - a\| < 1\} \subset \text{Inv}\mathcal{B}$;
- (iii) $\text{Inv}\mathcal{B}$ is an *open subset* of \mathcal{B} ;
- (iv) The map $\text{Inv}\mathcal{B} \ni a \longmapsto a^{-1} \in \text{Inv}\mathcal{B}$ is continuous.

For simplicity, we denote $\lambda 1$, where $\lambda \in \mathbb{C}$ and 1 is the identity of \mathcal{B} , by λ . The *resolvent set* of $a \in \mathcal{B}$ is defined by

$$\rho(a) := \{\lambda \in \mathbb{C}: \lambda - a \in \text{Inv}\mathcal{B}\};$$

the *spectrum* of a is $\sigma(a)$, the complement of $\rho(a)$ in \mathbb{C} , and the *resolvent function* of a is $R_a: \rho(a) \rightarrow \text{Inv}\mathcal{B}$, $R_a(\lambda) := (\lambda - a)^{-1}$. For each $\lambda, \gamma \in \rho(a)$ we have the identity

$$R_a(\gamma) - R_a(\lambda) = (\lambda - \gamma) R_a(\lambda) R_a(\gamma).$$

We also have that $\sigma(a) \subset \{\lambda \in \mathbb{C}: |\lambda| \leq \|a\|\}$. The *spectral radius* of a is defined as $\nu(a) = \sup \{|\lambda| : \lambda \in \sigma(a)\}$. If a, b are *commuting* elements in \mathcal{B} , i.e. $ab = ba$, then

$$\nu(ab) \leq \nu(a)\nu(b) \text{ and } \nu(a+b) \leq \nu(a) + \nu(b).$$

Let f be an analytic functions on the open disk $D(0, R)$ given by the *power series* $f(\lambda) := \sum_{j=0}^{\infty} \alpha_j \lambda^j$ ($|\lambda| < R$). If $\nu(a) < R$, then the series $\sum_{j=0}^{\infty} \alpha_j a^j$ converges in the Banach algebra \mathcal{B} because $\sum_{j=0}^{\infty} |\alpha_j| \|a^j\| < \infty$, and we can define $f(a)$ to be its sum. Clearly $f(a)$ is well defined and there are many examples of important functions on a Banach algebra \mathcal{B} that can be constructed in this way. For instance, the *exponential map* on \mathcal{B} denoted \exp and defined as

$$\exp a := \sum_{j=0}^{\infty} \frac{1}{j!} a^j \text{ for each } a \in \mathcal{B}.$$

If \mathcal{B} is not commutative, then many of the familiar properties of the exponential function from the scalar case do not hold. The following key formula is valid, however with the additional hypothesis of commutativity for a and b from \mathcal{B}

$$\exp(a+b) = \exp(a)\exp(b).$$

In a general Banach algebra \mathcal{B} it is difficult to determine the elements in the range of the exponential map $\exp(\mathcal{B})$, i.e. the elements which have a

"logarithm". However, it is easy to see that if a is an element in B such that $\|1 - a\| < 1$, then a is in $\exp(\mathcal{B})$. That follows from the fact that if we set

$$b = - \sum_{n=1}^{\infty} \frac{1}{n} (1 - a)^n,$$

then the series converges absolutely and, as in the scalar case, substituting this series into the series expansion for $\exp(b)$ yields $\exp(b) = a$.

It is known that if x and y are commuting, i.e. $xy = yx$, then the exponential function satisfies the property

$$\exp(x)\exp(y) = \exp(y)\exp(x) = \exp(x+y).$$

Also, if x is invertible and $a, b \in \mathbb{R}$ with $a < b$ then

$$\int_a^b \exp(tx) dt = x^{-1} [\exp(bx) - \exp(ax)].$$

Moreover, if x and y are commuting and $y - x$ is invertible, then

$$\begin{aligned} \int_0^1 \exp((1-s)x + sy) ds &= \int_0^1 \exp(s(y-x)) \exp(x) ds \\ &= \left(\int_0^1 \exp(s(y-x)) ds \right) \exp(x) \\ &= (y-x)^{-1} [\exp(y-x) - I] \exp(x) \\ &= (y-x)^{-1} [\exp(y) - \exp(x)]. \end{aligned}$$

Inequalities for functions of operators in Hilbert spaces may be found in the papers [?SSD], [?SSDG] and in the recent monographs [?SSDBook1], [?SSDBook2] and the references therein. Let α_n be nonzero complex numbers and let

$$R := \frac{1}{\limsup |\alpha_n|^{\frac{1}{n}}}.$$

Clearly $0 \leq R \leq \infty$, but we consider only the case $0 < R \leq \infty$.

Denote by:

$$D(0, R) = \begin{cases} \{\lambda \in \mathbb{C} : |\lambda| < R\}, & \text{if } R < \infty \\ \mathbb{C}, & \text{if } R = \infty, \end{cases}$$

consider the functions:

$$\lambda \mapsto f(\lambda) : D(0, R) \rightarrow \mathbb{C}, f(\lambda) := \sum_{n=0}^{\infty} \alpha_n \lambda^n$$

and

$$\lambda \mapsto f_A(\lambda) : D(0, R) \rightarrow \mathbb{C}, f_A(\lambda) := \sum_{n=0}^{\infty} |\alpha_n| \lambda^n.$$

Let \mathcal{B} be a unital Banach algebra and 1 its unity. Denote by

$$B(0, R) = \begin{cases} \{x \in \mathcal{B} : \|x\| < R\}, & \text{if } R < \infty \\ \mathcal{B}, & \text{if } R = \infty. \end{cases}$$

We associate to f the map

$$x \mapsto \tilde{f}(x) : B(0, R) \rightarrow \mathcal{B}, \tilde{f}(x) := \sum_{n=0}^{\infty} \alpha_n x^n.$$

Obviously, \tilde{f} is correctly defined because the series $\sum_{n=0}^{\infty} \alpha_n x^n$ is absolutely convergent, since $\sum_{n=0}^{\infty} \|\alpha_n x^n\| \leq \sum_{n=0}^{\infty} |\alpha_n| \|x\|^n$.

The following result has been obtained in [?DBM].

Theorem 1.1. *Let $f(\lambda) = \sum_{n=0}^{\infty} \alpha_n \lambda^n$ be a power series that is convergent on the open disk $D(0, R)$, with $R > 0$. If $x, y \in \mathcal{B}$ with $xy = yx$ and $\|x\|, \|y\| \leq 1$, then we have for $\lambda \in \mathbb{C}$ with $|\lambda| < R$ the inequality:*

$$\begin{aligned} & \left\| \tilde{f}(\lambda \cdot 1) \tilde{f}(\lambda xy) - \tilde{f}(\lambda x) \tilde{f}(\lambda y) \right\| \\ & \leq \|x - 1\| \|y - 1\| \left\{ f_A(|\lambda|) \left[|\lambda| f'_A(|\lambda|) + |\lambda|^2 f''_A(|\lambda|) \right] - [|\lambda| f'_A(|\lambda|)]^2 \right\}. \end{aligned} \quad (1.1)$$

For other similar results see [?DBB] and [?DBM]. Motivated by the above results, in this paper we consider the similar problem to provide upper bounds for the Čebyšev quantity

$$\left\| P_n \sum_{i=1}^n p_i F(x_i) G(y_i) - \sum_{i=1}^n p_i F(x_i) \sum_{i=1}^n p_i G(y_i) \right\|,$$

where F, G are some functions defined on a Banach algebra \mathcal{B} , $p_i \geq 0$ with $P_n := \sum_{i=1}^n p_i > 0$ and $x_i, y_i \in \mathcal{C}$, $i = 1, 2, \dots, n$ where \mathcal{C} is an appropriate subset of \mathcal{B} . For various results on Čebyšev's inequality in different settings, see [?A1]-[?SSD], [?cfo6b]-[?SSDBook2] and [?6b]-[?ZC].

2 Some New Inequalities

We say that the function $F : \mathcal{C} \subset \mathcal{B} \rightarrow \mathcal{B}$ is a Lipschitzian with constant $L > 0$ on the subset \mathcal{C} of the Banach algebra $(\mathcal{B}, \|\cdot\|)$ if

$$\|F(x) - F(y)\| \leq L \|x - y\|$$

for any $x, y \in \mathcal{C}$.

Theorem 2.1. *Let $F, G : \mathcal{C} \subset \mathcal{B} \rightarrow \mathcal{B}$ be Lipschitzian functions with constants $L > 0$ and $K > 0$. If $x_i, y_i \in \mathcal{C}$, $i = 1, 2, \dots, n$ and $p_i \geq 0$, $i = 1, 2, \dots, n$ with $P_n := \sum_{i=1}^n p_i > 0$, then we have:*

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i F(x_i) G(y_i) - \sum_{i=1}^n p_i F(x_i) \sum_{i=1}^n p_i G(y_i) \right\| \\ & \leq LK \sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\| \end{aligned} \quad (2.1)$$

where Δx_l is the forward difference, namely: $\Delta x_l = x_{l+1} - x_l$, $l = 1, \dots, n-1$.

Proof. Since F and G are Lipschitzian with the constants L and K we have:

$$\|F(x_i) - F(x_j)\| \leq L \|x_i - x_j\|$$

and

$$\|G(y_i) - G(y_j)\| \leq K \|y_i - y_j\|$$

for any $i, j \in \{1, \dots, n\}$. If we multiply these inequalities we get

$$\|F(x_i) - F(x_j)\| \|G(y_i) - G(y_j)\| \leq LK \|x_i - x_j\| \|y_i - y_j\|.$$

Using the property of Banach algebra \mathcal{B} , $\|uv\| \leq \|u\| \|v\|$, that for any $u, v \in \mathcal{B}$ we have,

$$\|(F(x_i) - F(x_j))(G(y_i) - G(y_j))\| \leq LK \|x_i - x_j\| \|y_i - y_j\|$$

for any $i, j \in \{1, \dots, n\}$, that is equivalent to:

$$\begin{aligned} & \|F(x_i)G(y_i) - F(x_j)G(y_i) - F(x_i)G(y_j) + F(x_j)G(y_j)\| \\ & \leq LK \|x_i - x_j\| \|y_i - y_j\| \end{aligned} \quad (2.2)$$

for any $i, j \in \{1, \dots, n\}$. If we multiply (??) by $p_i p_j > 0$ sum over i and j from 1 to n and use the generalised triangle inequality, we get:

$$\left\| \sum_{i,j=1}^n p_i p_j (F(x_i)G(y_i) - F(x_j)G(y_i) - F(x_i)G(y_j) + F(x_j)G(y_j)) \right\| \quad (2.3)$$

$$\begin{aligned} &\leq \sum_{i,j=1}^n p_i p_j \|F(x_i)G(y_i) - F(x_j)G(y_i) - F(x_i)G(y_j) + F(x_j)G(y_j)\| \\ &\leq LK \sum_{i,j=1}^n p_i p_j \|x_i - x_j\| \|y_i - y_j\|. \end{aligned}$$

Now, observe that

$$\begin{aligned} &\sum_{i,j=1}^n p_i p_j (F(x_i)G(y_i) - F(x_j)G(y_i) - F(x_i)G(y_j) + F(x_j)G(y_j)) \quad (2.4) \\ &= P_n \sum_{i=1}^n p_i F(x_i)G(y_i) - \sum_{j=1}^n p_j F(x_j) \sum_{i=1}^n p_i G(y_i) \\ &\quad - \sum_{i=1}^n p_i F(x_i) \sum_{j=1}^n p_j G(y_j) + P_n \sum_{j=1}^n p_j F(x_j)G(y_j) \\ &= 2 \left(P_n \sum_{i=1}^n p_i F(x_i)G(y_i) - \sum_{i=1}^n p_i F(x_i) \sum_{i=1}^n p_i G(y_i) \right). \end{aligned}$$

We also have, by the symmetry in the sum:

$$\sum_{i,j=1}^n p_i p_j \|x_i - x_j\| \|y_i - y_j\| = 2 \sum_{1 \leq i < j \leq n} p_i p_j \|x_j - x_i\| \|y_j - y_i\|. \quad (2.5)$$

Moreover, we have $x_j - x_i = \sum_{l=1}^{j-1} \Delta x_l$ and $y_j - y_i = \sum_{s=1}^{j-1} \Delta y_s$ where $\Delta x_l := x_{l+1} - x_l$, $l = i, \dots, j-1$ and by the generalized triangle inequality we have:

$$\|x_j - x_i\| \leq \sum_{l=i}^{j-1} \|\Delta x_l\|$$

and

$$\|y_j - y_i\| \leq \sum_{s=i}^{j-1} \|\Delta y_s\|. \quad (2.6)$$

By the inequality (??) and by (??) we then have:

$$\sum_{i,j=1}^n p_i p_j \|x_i - x_j\| \|y_i - y_j\| \leq 2 \sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\|. \quad (2.7)$$

Now, by making use of (??), (??) and (??) we get the desired result (??). \square

Corollary 2.2. *With the assumptions of Theorem 1 we have:*

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i F(x_i) G(y_i) - \sum_{i=1}^n p_i F(x_i) \sum_{i=1}^n p_i G(y_i) \right\| \\ & \leq \frac{1}{2} LK \left(P_n^2 - \sum_{i=1}^n p_i^2 \right) \sum_{k=1}^{j-1} \|\Delta x_k\| \sum_{k=1}^{n-1} \|\Delta y_k\|. \end{aligned} \quad (2.8)$$

Proof. It is obvious that for all $1 \leq i < j \leq n$ we have that:

$$\sum_{l=i}^{j-1} \|\Delta x_l\| \leq \sum_{k=1}^{n-1} \|\Delta x_k\|$$

and

$$\sum_{s=i}^{j-1} \|\Delta y_s\| \leq \sum_{k=1}^{n-1} \|\Delta y_k\|.$$

Then

$$\sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\| \leq \sum_{1 \leq i < j \leq n} p_i p_j \sum_{k=1}^{n-1} \|\Delta x_k\| \sum_{k=1}^{n-1} \|\Delta y_k\|.$$

Now, let observe that:

$$\begin{aligned} \sum_{1 \leq i < j \leq n} p_i p_j &= \frac{1}{2} \left[\sum_{i,j=1}^n p_i p_j - \sum_{i=j} p_i p_j \right] \\ &= \frac{1}{2} \left[\sum_{i=1}^n p_i \sum_{j=1}^n p_j - \sum_{i=1}^n p_i^2 \right] \\ &= \frac{1}{2} \left(P_n^2 - \sum_{i=1}^n p_i^2 \right). \end{aligned}$$

Using (??) we deduce (??). \square

Corollary 2.3. *With the assumptions of Theorem 1 we have:*

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i F(x_i) G(y_i) - \sum_{i=1}^n p_i F(x_i) \sum_{i=1}^n p_i G(y_i) \right\| \\ & \leq LK \sum_{1 \leq i < j \leq n} (j-i) p_i p_j \left(\sum_{k=1}^{n-1} \|\Delta x_k\|^p \right)^{\frac{1}{p}} \left(\sum_{k=1}^{n-1} \|\Delta y_k\|^q \right)^{\frac{1}{q}} \end{aligned} \quad (2.9)$$

when $p, q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$.

Proof. Using Hölder's discrete inequality, we can state that

$$\sum_{l=i}^{j-1} \|\Delta x_l\| \leq (j-i)^{\frac{1}{q}} \left(\sum_{l=i}^{j-1} \|\Delta x_l\|^p \right)^{\frac{1}{p}}$$

and

$$\sum_{s=i}^{j-1} \|\Delta y_s\| \leq (j-i)^{\frac{1}{p}} \left(\sum_{s=i}^{j-1} \|\Delta y_s\|^q \right)^{\frac{1}{q}}$$

where $j-1 \geq i, p, q > 1$ and $\frac{1}{p} + \frac{1}{q} = 1$, and then, by multiplication we have

$$\begin{aligned} & \sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\| \\ & \leq \sum_{1 \leq i < j \leq n} p_i p_j (j-i)^{\frac{1}{p} + \frac{1}{q}} \left(\sum_{l=i}^{j-1} \|\Delta x_l\|^p \right)^{\frac{1}{p}} \left(\sum_{s=i}^{j-1} \|\Delta y_s\|^q \right)^{\frac{1}{q}} \\ & = \sum_{1 \leq i < j \leq n} p_i p_j (j-i) \left(\sum_{l=i}^{j-1} \|\Delta x_l\|^p \right)^{\frac{1}{p}} \left(\sum_{s=i}^{j-1} \|\Delta y_s\|^q \right)^{\frac{1}{q}} \\ & \leq \sum_{1 \leq i < j \leq n} p_i p_j (j-i) \left(\sum_{k=1}^{n-1} \|\Delta x_k\|^p \right)^{\frac{1}{p}} \left(\sum_{k=1}^{n-1} \|\Delta y_k\|^q \right)^{\frac{1}{q}} \end{aligned}$$

and the corollary is proved. \square

Corollary 2.4. *With the assumptions of Theorem 1 we have:*

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i F(x_i) G(y_i) - \sum_{i=1}^n p_i F(x_i) \sum_{i=1}^n p_i G(y_i) \right\| \\ & \leq LK \left[P_n \sum_{i=1}^n i^2 p_i - \left(\sum_{i=1}^n i p_i \right)^2 \right] \max_{1 \leq k \leq n-1} \|\Delta x_k\| \max_{1 \leq k \leq n-1} \|\Delta y_k\|. \quad (2.10) \end{aligned}$$

Proof. We have

$$\sum_{l=i}^{j-1} \|\Delta x_l\| \leq (j-i) \max_{l=1, n-1} \|\Delta x_l\|$$

and

$$\sum_{s=i}^{j-1} \|\Delta y_s\| \leq (j-i) \max_{s=1, n-1} \|\Delta y_s\|.$$

Then

$$\begin{aligned} & \sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\| \\ & \leq \sum_{1 \leq i < j \leq n} p_i p_j (j-i)^2 \max_{l=1, n-1} \|\Delta x_l\| \max_{s=1, n-1} \|\Delta y_s\| \\ & = \left[\sum_{i=1}^n p_i \sum_{i=1}^n i^2 p_i - \left(\sum_{i=1}^n i p_i \right)^2 \right] \max_{l=1, n-1} \|\Delta x_l\| \max_{s=1, n-1} \|\Delta y_s\| \end{aligned}$$

and the corollary is proved. \square

3 Some Applications

In [SSD11] S. S. Dragomir proved the following result:

Lemma 3.1. *Let $f(z) = \sum_{n=0}^{\infty} \alpha_n z^n$ be a function defined by power series with complex coefficients and convergent on the open disk $D(0, r) \subset \mathbb{C}$, $R > 0$ and let $f_a(\lambda) := \sum_{n=0}^{\infty} |\alpha_n| \lambda^n$. For any $x, y \in \mathcal{B}$ with $\|x\|, \|y\| \leq M < R$ we have:*

$$\|f(y) - f(x)\| \leq f'_a(M) \|y - x\|.$$

If we take $f(z) = z^p$, $p \in \mathbb{N}$, $p \geq 1$ and assume that $x, y \in \mathcal{B}$ with $\|x\|, \|y\| \leq M$ then we have

$$\|y^p - x^p\| \leq pM^{p-1} \|y - x\|.$$

If we take $f(z) = \exp(\mu z)$, with $\mu \in \mathbb{R}$, $\mu \neq 0$ and assume that $x, y \in \mathcal{B}$ with $\|x\|, \|y\| \leq M$ then we have

$$\|\exp(\mu y) - \exp(\mu x)\| \leq |\mu| \exp(\mu M) \|y - x\|.$$

If we write the inequality (??) for $F(z) = z^p, G(z) = z^q$, with $p, q \in \mathbb{N}$, $p, q \geq 1$ then we have for $\|x_i\|, \|y_i\| \leq M$, $i \in \{1, 2, \dots, n\}$ that

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i x_i^p y_i^q - \sum_{i=1}^n p_i x_i^p \sum_{i=1}^n p_i y_i^q \right\| \\ & \leq pq M^{p+q-2} \sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\|. \end{aligned} \quad (3.1)$$

If $p = q$ and x_i commutes with y_i for each $i \in \{1, 2, \dots, n\}$, then we get from (??) that:

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i (x_i y_i)^p - \sum_{i=1}^n p_i x_i^p \sum_{i=1}^n p_i y_i^p \right\| \\ & \leq p^2 M^{2p-2} \sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\|. \end{aligned} \quad (3.2)$$

In particular, for $x_i = y_i$ and $p = q$, $i \in \{1, 2, \dots, n\}$, we get

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i (x_i)^{2p} - \left(\sum_{i=1}^n p_i x_i^p \right)^2 \right\| \\ & \leq p^2 M^{2p-2} \sum_{1 \leq i < j \leq n} p_i p_j \left(\sum_{l=i}^{j-1} \|\Delta x_l\| \right)^2. \end{aligned} \quad (3.3)$$

Now, if we write the inequality (??) for the functions $F(z) = \exp(\mu z)$ and $G(z) = \exp(\nu z)$, with $\mu, \nu \in \mathbb{R}$, $\mu, \nu \neq 0$ we get for $\|x_i\|, \|y_i\| \leq M$, for $i \in \{1, 2, \dots, n\}$ that

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i \exp(\mu x_i) \exp(\nu y_i) - \sum_{i=1}^n p_i \exp(\mu x_i) \sum_{i=1}^n p_i \exp(\nu y_i) \right\| \\ & \leq |\mu\nu| \exp[(\mu + \nu)M] \sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\|. \end{aligned} \quad (3.4)$$

If x_i commutes with y_i for each $i \in \{1, 2, \dots, n\}$, then from (??) we get:

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i \exp(\mu x_i + \nu y_i) - \sum_{i=1}^n p_i \exp(\mu x_i) \sum_{i=1}^n p_i \exp(\nu y_i) \right\| \\ & \leq |\mu\nu| \exp[(\mu + \nu)M] \sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\|. \end{aligned} \quad (3.5)$$

In particular, for $x_i = y_i$ and $\mu = \nu$, $i \in \{1, 2, \dots, n\}$, we get:

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i \exp(2\mu x_i) - \left(\sum_{i=1}^n p_i \exp(\mu x_i) \right)^2 \right\| \\ & \leq |\mu|^2 \exp(2\mu M) \sum_{1 \leq i < j \leq n} p_i p_j \left(\sum_{l=i}^{j-1} \|\Delta x_l\| \right)^2. \end{aligned} \quad (3.6)$$

Consider now the function $F_p(z) = (1 - z^p)^{-1} = \sum_{n=0}^{\infty} z^{np}$ for $p \geq 1$ a natural number. Then $F'_p(z) = -(1 - z^p)^{-2} (-pz^{p-1}) = p(1 - z^p)^{-2} z^{p-1}$. If $M < 1$, then $F'_p(M) = p(1 - M^p)^{-2} M^{p-1}$.

We have then $F_p(x) = (1 - x^p)^{-1}$, $x \in \mathcal{B}$, $\|x\| < 1$, $p \geq 1$ is Lipschitzian with the constant $p(1 - M^p)^{-2} M^{p-1}$ and the function $G_q(x) = (1 - x^q)^{-1}$, $x \in \mathcal{B}$, $\|x\| < 1$, $q \geq 1$ is Lipschitzian with the constant $q(1 - M^q)^{-2} M^{q-1}$. Therefore, by using Theorem 2.1 for $x_i, y_i \in \mathcal{B}$ with $\|x_i\|, \|y_i\| \leq M < 1$, $i \in \{1, 2, \dots, n\}$, we get for $p_i \geq 0$, $i \in \{1, 2, \dots, n\}$ that

$$\left\| P_n \sum_{i=1}^n p_i (1 - x_i^p)^{-1} (1 - y_i^q)^{-1} - \sum_{i=1}^n p_i (1 - x_i^p)^{-1} \sum_{i=1}^n p_i (1 - y_i^q)^{-1} \right\|$$

$$\leq pq (1 - M^p)^{-2} (1 - M^q)^{-2} M^{p+q-2} \sum_{1 \leq i < j \leq n} p_i p_j \sum_{l=i}^{j-1} \|\Delta x_l\| \sum_{s=i}^{j-1} \|\Delta y_s\|. \quad (3.7)$$

In particular for $p = q$ and $x_i = y_i$ for each $i \in \{1, 2, \dots, n\}$, we get from (??) that:

$$\begin{aligned} & \left\| P_n \sum_{i=1}^n p_i (1 - x_i^p)^{-2} - \left(\sum_{i=1}^n p_i (1 - x_i^p)^{-1} \right)^2 \right\| \\ & \leq p^2 (1 - M^p)^{-4} M^{2p-2} \sum_{1 \leq i < j \leq n} p_i p_j \left(\sum_{l=i}^{j-1} \|\Delta x_l\| \right)^2. \end{aligned} \quad (3.8)$$

Finally, we notice that if one uses the upper bounds from Corollaries 2.2 - 2.4 that one can get further upper bounds in the examples outlined above. The details are omitted.

References

- [1] **G. A. Anastassiou**, Grüss type inequalities for the Stieltjes integral, *Nonlinear Funct. Anal. Appl.*, **12** (4), (2007), 583-593
- [2] **G. A. Anastassiou**, Chebyshev-Grüss type and comparison of integral means inequalities for the Stieltjes integral, *Panamer. Math. J.*, **17** (3), (2007), 91-109
- [3] **G. A. Anastassiou**, Chebyshev-Grüss type inequalities via Euler type and Fink identities, *Math. Comput. Modelling*, **45** (9-10), (2007), 1189-1200
- [4] **M. Biernacki**, Sur une inégalité entre les intégrales due à Tchebyscheff, *Ann. Univ. Mariae Curie-Sklodowska*, **A5**, (1951), 23-29
- [5] **K. Boukerrioua and A. Guezane-Lakoud**, On generalization of Čebyšev type inequalities, *J. Inequal. Pure Appl. Math.*, **8** (2, Article 55), (2007), 4 pp.
- [6] **P. L. Čebyšev**, O približennyy vyraženijah odnih integralov čerez drugie, *Soobščenija i protokoly zasedaniĭ Matemmatičeskogo občestva pri Imperatorskom Har'kovskom Universitete*, **2(93-98)**
- [7] **P.L. Čebyšev**, Ob odnom rjade, dostavljajuščem predel'nye veličiny integralov pri razloženii podintegral'noj funkcii na množeteli, *Priloženi k 57 tomu Zapisok Imp. Akad. Nauk*, **4**
- [8] **P. Cerone**, On a Čebyšev-type functional and Grüss-like bounds, *Math. Inequal. Appl.*, **9** (1), (2006), 87-102

- [9] **P. Cerone**, On Chebyshev functional bounds, *Differential & difference equations and applications*, (2006), 267-277
- [10] **P. Cerone**, On some results involving the Čebyšev functional and its generalisations, *J. Inequal. Pure Appl. Math.*, **4** (3, Article 55), (2003), 17 pp.
- [11] **P. Cerone and S. S. Dragomir**, A refinement of the Grüss inequality and applications, *Tamkang J. Math.*, **38** (1), (2007), 37-49
- [12] **P. Cerone and S. S. Dragomir**, Chebychev functional bounds using Ostrowski seminorms, *Southeast Asian Bull. Math.*, **28** (2), (2004), 219-228
- [13] **P. Cerone and S. S. Dragomir**, New bounds for the Čebyšev functional, *Appl. Math. Lett.*, **18** (6), (2005), 603-611
- [14] **S. S. Dragomir**, Inequalities for the Čebyšev functional of two functions of selfadjoint operators in Hilbert spaces, *Aust. J. Math. Anal. & Appl.*, **6** (1, Article 7), (2009), 1-58
- [15] **S. S. Dragomir**, Some inequalities for power series of selfadjoint operators in Hilbert spaces via reverses of the Schwarz inequality, *Integral Transforms Spec. Funct.*, **20** (9-10), (2009), 757-767
- [16] **S. S. Dragomir**, Inequalities for power series in Banach Algebras, *SUT J. of Math.*, **50** (1), (2014), 25-45
- [17] **S. S. Dragomir**, On some improvements of Čebyšev's inequality for sequences and integrals, *Studia Univ. Babeş-Bolyai, Mathematica (Romania)*, **XXXV** (4), (1990), 35-40
- [18] **S. S. Dragomir**, Some improvement of Čebyšev's inequality for isotonic functionals, *Atti. Sem. Mat. Fis. Univ. Modena (Italy)*, **41**, (1993), 473-481
- [19] **S.S. Dragomir**, A generalization of Grüss' inequality in inner product spaces and applications, **237**, (1999), 74-82
- [20] **S. S. Dragomir**, Some discrete inequalities of Grüss type and applications in guessing theory, *Honam Math. J.*, **21** (1), (1999), 145-156
- [21] **S. S. Dragomir**, Grüss inequality in inner product spaces, *The Australian Math Soc. Gazette*, **26** (2), (1999), 66-70
- [22] **S. S. Dragomir**, A Grüss type integral inequality for mappings of r -Hölder's type and applications for trapezoid formula, *Tamkang J. of Math.*, **31** (1), (2000), 43-47
- [23] **S.S. Dragomir**, Some integral inequalities of Grüss type, *Indian J. of Pure and Appl. Math.*, **31** (4), (2000), 397-415
- [24] **S. S. Dragomir**, On the Ostrowski inequality for the Riemann-Stieltjes integral $\int_a^b f(t) du(t)$, where f is of Hölder type and u is of bounded variation and applications, *J. KSIAM*, **5** (1), (2001), 35-45
- [25] **S. S. Dragomir**, Some Grüss type inequalities in inner product spaces, *J. Inequal. Pure & Appl. Math.*, **4** (2), (2003), Article 42, (Online: <http://jipam.vu.edu.au/article.php?sid=280>)

- [26] **S. S. Dragomir**, Reverses of Schwarz, triangle and Bessel inequalities in inner product spaces, *J. Inequal. Pure & Appl. Math.*, **5** (3), (2004), Article 76. (Online: <http://jipam.vu.edu.au/article.php?sid=432>)
- [27] **S.S. Dragomir**, New reverses of Schwarz, triangle and Bessel inequalities in inner product spaces, *Austral. J. Math. Anal. & Applics.*, **1** (1), (2004), Article 1. (Online: <http://ajmaa.org/cgi-bin/paper.pl?string=nrstbiips.tex>)
- [28] **S. S. Dragomir**, On the Čebyšev's inequality for weighted means, *Acta Math. Hungar.*, **104** (4), (2004), 345-355
- [29] **S.S. Dragomir**, On Bessel and Grüss inequalities for orthonormal families in inner product spaces, *Bull. Austral. Math. Soc.*, **69** (2), (2004), 327-340
- [30] **S. S. Dragomir**, *Advances in Inequalities of the Schwarz, Grüss and Bessel Type in Inner Product Spaces*, Nova Science Publishers Inc, New York, 2005
- [31] **S. S. Dragomir**, Reverses of the Schwarz inequality in inner product spaces generalising a Klamkin-McLenaghan result, *Bull. Austral. Math. Soc.*, **73** (1), (2006), 69-78
- [32] **S. S. Dragomir**, Some new Grüss' type inequalities for functions of selfadjoint operators in Hilbert spaces, *RGMIA Res. Rep. Coll.*, **11** (e), (2008), Art. 12
- [33] **S. S. Dragomir**, Inequalities for the Čebyšev functional of two functions of selfadjoint operators in Hilbert spaces, *RGMIA Res. Rep. Coll.*, **11** (e), (2008), Art. 17
- [34] **S. S. Dragomir**, Some inequalities for the Čebyšev functional of two functions of selfadjoint operators in Hilbert spaces, *RGMIA Res. Rep. Coll.*, **11** (e), (2008), Art. 8
- [35] **S. S. Dragomir**, Springer Briefs in Mathematics. Springer, New York, 2012
- [36] **S. S. Dragomir**, *Operator Inequalities of Ostrowski and Trapezoidal Type*, Springer Briefs in Mathematics. Springer, New York, 2012
- [37] **S. S. Dragomir, M. V. Boldea, C. Buşe, and M. Megan**, Norm inequalities of Čebyšev type for power series in Banach algebras, *J. Inequal. Appl.*, **2014**, 2014:294
- [38] **S. S. Dragomir, M. V. Boldea, and M. Megan**, New bounds for Čebyšev functional for power series in Banach algebras via a Grüss'-Lupaş type inequality, *Preprint RGMIA Res. Rep. Coll.*, **17**, (2014)
- [39] **S. S. Dragomir and G.L. Booth**, On a Grüss-Lupaş type inequality and its applications for the estimation of p-moments of guessing mappings, *Mathematical Communications*, **5**, (2000), 117-126
- [40] **S. S. Dragomir and I. Fedotov**, An inequality of Grüss' type for Riemann-Stieltjes integral and applications for special means, *Tamkang J. of Math.*, **29** (4), (1998), 286-292
- [41] **S. S. Dragomir and B. Mond**, Some mappings associated with Čebyšev's inequality for sequences of real numbers, *Bull. Allahabad Math. Soc.*, **8/9**, (1993/94), 37-55
- [42] **S.S. Dragomir and J.E. Pečarić**, Refinements of some inequalities for isotonic linear functionals, *L'Anal. Num. Théor de L'Approx.(Romania)*, **18** (1), (1989), 61-65

- [43] **S. S. Dragomir, J. Pečarić, and J. Sándor**, The Chebyshev inequality in pre-Hilbertian spaces. II., *Proceedings of the Third Symposium of Mathematics and its Applications (Timișoara, 1989)*, (1990), 75-78
- [44] **S. S. Dragomir and J. Sándor**, The Chebyshev inequality in pre-Hilbertian spaces. I., *Proceedings of the Second Symposium of Mathematics and its Applications (Timișoara, 1987)*, (1988), 61-64
- [45] **A. M. Fink**, A treatise on Grüss' inequality, *Analytic and Geometric Inequalities, Math. Appl.*, **478**, (1999), 93-113
- [46] **G. Grüss**, Über das Maximum des absoluten Betrages von $\frac{1}{b-a} \int_a^b f(x)g(x)dx - \frac{1}{(b-a)^2} \int_a^b f(x)dx \int_a^b g(x)dx$, *Math. Z.*, **39**, (1935), 215-226
- [47] **Z. Liu**, Refinement of an inequality of Grüss type for Riemann-Stieltjes integral, *Soochow J. Math.*, **30** (4), (2004), 483-489
- [48] **A. Matković, J. Pečarić, and I. Perić**, A variant of Jensen's inequality of Mercer's type for operators with applications, *Linear Algebra Appl.*, **418** (2-3), (2006), 551-564
- [49] **D. S. Mitrinović and J.E. Pečarić**, History, variations and generalizations of the Čebyšev inequality and question of some priorities. II, *Rad Jugoslav. Akad. Znan. Umjet. No.*, **450**, (1990), 139-156
- [50] **D. S. Mitrinović and J.E. Pečarić**, On an identity of D.Z. Djoković, *Prilozi Mak. Akad.Nauk. Umj. (Skopje)*, **12** (1), (1991), 21-22
- [51] **D. S. Mitrinović and P.M. Vasić**, History, variations and generalisations of the Čebyšev inequality and the question of some priorities, *Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz.*, **461-497**, (1974), 1-30
- [52] **B. G. Pachpatte**, A note on Čebyšev type inequalities, *An. Științ. Univ. Al. I. Cuza Iași. Mat. (N.S.) (Romania)*, **53** (1), (2007), 97-102
- [53] **B. G. Pachpatte**, New Čebyšev type inequalities involving functions of two and three variables, *Soochow J. Math.*, **33** (4), (2007), 569-577
- [54] **B. G. Pachpatte**, A note on Grüss type inequalities via Cauchy's mean value theorem, *Math. Inequal. Appl.*, **11** (1), (2008), 75-80
- [55] **J. Pečarić**, Remarks on Biernacki's generalization of Čebyšev's inequality, *Ann. Univ. Mariae Curie-Sklodowska Sect. A*, **47**, (1993), 116-122
- [56] **J. E. Pečarić and S.S. Dragomir**, Some remarks on Čebyšev's inequality, *L'Anal. Num. Théor de L'Approx.*, **19** (1), (1990), 58-65
- [57] **C.-J. Zhao and W.-S. Cheung**, On multivariate Grüss inequalities, *J. Inequal. Appl.*, **2008**, Art. ID 249438, 8 pp.

Marius V. Boldea

Department Mathematics and Statistics,
Banat University of Agricultural Sciences and Veterinary Medicine Timișoara
119 Calea Aradului, Timișoara, România

E-mail: marius.boldea@usab-tm.ro

Received: 7.08.2015

Accepted: 26.10.2016