

# An Inequality Related to Vizing's Conjecture

W. Edwin Clark and Stephen Suen

Department of Mathematics, University of South Florida,

Tampa, FL 33620-5700, USA

eclark@math.usf.edu    suen@math.usf.edu

Submitted November 29, 1999; Accepted May 24, 2000

## Abstract

Let  $\gamma(G)$  denote the domination number of a graph  $G$  and let  $G \square H$  denote the Cartesian product of graphs  $G$  and  $H$ . We prove that  $\gamma(G)\gamma(H) \leq 2\gamma(G \square H)$  for all simple graphs  $G$  and  $H$ .

*2000 Mathematics Subject Classifications:* Primary 05C69, Secondary 05C35

We use  $V(G)$ ,  $E(G)$ ,  $\gamma(G)$ , respectively, to denote the vertex set, edge set and domination number of the (simple) graph  $G$ . For a pair of graphs  $G$  and  $H$ , the Cartesian product  $G \square H$  of  $G$  and  $H$  is the graph with vertex set  $V(G) \times V(H)$  and where two vertices are adjacent if and only if they are equal in one coordinate and adjacent in the other. In 1963, V. G. Vizing [2] conjectured that for any graphs  $G$  and  $H$ ,

$$\gamma(G)\gamma(H) \leq \gamma(G \square H). \tag{1}$$

The reader is referred to Hartnell and Rall [1] for a summary of recent progress on Vizing's conjecture. We note that there are graphs  $G$  and  $H$  for which equality holds in (1). However, it was previously unknown [1] whether there exists a constant  $c$  such that

$$\gamma(G)\gamma(H) \leq c \gamma(G \square H).$$

We shall show in this note that  $\gamma(G)\gamma(H) \leq 2 \gamma(G \square H)$ .

For  $S \subseteq V(G)$  we let  $N_G[S]$  denote the set of vertices in  $V(G)$  that are in  $S$  or adjacent to a vertex in  $S$ , *i.e.*, the set of vertices in  $V(G)$  dominated by vertices in  $S$ .

**Theorem 1** For any graphs  $G$  and  $H$ ,

$$\gamma(G)\gamma(H) \leq 2\gamma(G \square H).$$

**Proof.** Let  $D$  be a dominating set of  $G \square H$ . It is sufficient to show that

$$\gamma(G)\gamma(H) \leq 2|D|. \quad (2)$$

Let  $\{u_1, u_2, \dots, u_{\gamma(G)}\}$  be a dominating set of  $G$ . Form a partition  $\{\Pi_1, \Pi_2, \dots, \Pi_{\gamma(G)}\}$  of  $V(G)$  so that for all  $i$ : (i)  $u_i \in \Pi_i$ , and (ii)  $u \in \Pi_i$  implies  $u = u_i$  or  $u$  is adjacent to  $u_i$ . This partition of  $V(G)$  induces a partition  $\{D_1, D_2, \dots, D_{\gamma(G)}\}$  of  $D$  where

$$D_i = (\Pi_i \times V(H)) \cap D.$$

Let  $P_i$  be the projection of  $D_i$  onto  $H$ . That is,

$$P_i = \{v \mid (u, v) \in D_i \text{ for some } u \in \Pi_i\}.$$

Observe that for any  $i$ ,  $P_i \cup (V(H) - N_H[P_i])$  is a dominating set of  $H$ , and hence the number of vertices in  $V(H)$  not dominated by  $P_i$  satisfies the inequality

$$|V(H) - N_H[P_i]| \geq \gamma(H) - |P_i|. \quad (3)$$

For  $v \in V(H)$ , let

$$Q_v = D \cap (V(G) \times \{v\}) = \{(u, v) \in D \mid u \in V(G)\}.$$

and  $C$  be the subset of  $\{1, 2, \dots, \gamma(G)\} \times V(H)$  given by

$$C = \{(i, v) \mid \Pi_i \times \{v\} \subseteq N_{G \square H}[Q_v]\}.$$

Let  $N = |C|$ . By counting in two different ways we shall find upper and lower bounds for  $N$ . Let

$$\begin{aligned} L_i &= \{(i, v) \in C \mid v \in V(H)\}, \text{ and} \\ R_v &= \{(i, v) \in C \mid 1 \leq i \leq \gamma(G)\}. \end{aligned}$$

Clearly

$$N = \sum_{i=1}^{\gamma(G)} |L_i| = \sum_{v \in V(H)} |R_v|.$$

Note that if  $v \in V(H) - N_H[P_i]$ , then the vertices in  $\Pi_i \times \{v\}$  must be dominated by vertices in  $Q_v$  and therefore  $(i, v) \in L_i$ . This implies that  $|L_i| \geq |V(H) - N_H[P_i]|$ . Hence

$$N \geq \sum_{i=1}^{\gamma(G)} |V(H) - N_H[P_i]|$$

and it follows from (3) that

$$\begin{aligned} N &\geq \gamma(G)\gamma(H) - \sum_{i=1}^{\gamma(G)} |P_i| \\ &\geq \gamma(G)\gamma(H) - \sum_{i=1}^{\gamma(G)} |D_i|. \end{aligned}$$

So we obtain the following lower bound for  $N$ .

$$N \geq \gamma(G)\gamma(H) - |D|. \quad (4)$$

For each  $v \in V(H)$ ,  $|R_v| \leq |Q_v|$ . If not,

$$\{u \mid (u, v) \in Q_v\} \cup \{u_j \mid (j, v) \notin R_v\}$$

is a dominating set of  $G$  with cardinality

$$|Q_v| + (\gamma(G) - |R_v|) = \gamma(G) - (|R_v| - |Q_v|) < \gamma(G),$$

and we have a contradiction. This observation shows that

$$N = \sum_{v \in V(H)} |R_v| \leq \sum_{v \in V(H)} |Q_v| = |D|. \quad (5)$$

It follows from (4) and (5) that

$$\gamma(G)\gamma(H) - |D| \leq N \leq |D|,$$

and the desired inequality (2) follows. ■

## References

- [1] Bert Hartnell and Douglas F. Rall, Domination in Cartesian Products: Vizing's Conjecture, in *Domination in Graphs—Advanced Topics* edited by Haynes, *et al*, Marcel Dekker, Inc, New York, 1998, 163–189.
- [2] V. G. Vizing, The cartesian product of graphs, *Vyčisl. Sistemy* **9**, 1963, 30–43.