

Infection levels of *Austrodiplostomum compactum* (Digenea, Diplostomidae) metacercariae in *Plagioscion squamosissimus* (Teleostei, Sciaenidae) from the Nova Avanhandava reservoir, São Paulo State, Brazil — [Source link](#)

Jaciara Vanessa Kruger Paes, Edmir Daniel Carvalho, R. J. da Silva

Institutions: Sao Paulo State University

Published on: 01 Sep 2010 - Journal of Helminthology (Cambridge University Press)

Related papers:

- [Diplostomum \(Austrodiplostomum\) compactum \(Lutz, 1928\) \(Platyhelminthes, Digenea\) metacercariae in fish from the floodplain of the Upper Paraná River, Brazil.](#)
- [Parasitology meets ecology on its own terms: Margolis et al. revisited.](#)
- [Austrodiplostomum compactum \(Lutz, 1928\) \(Digenea, diplostomidae\) infecting *Geophagus proximus* Castelnau, 1855 \(Cichlidae, Perciformes\) in the Tietê River, Nova Avanhandava reservoir, municipality of Buritama, São Paulo State, Brazil.](#)
- [Novas ocorrências de metacercária de *Austrodiplostomum compactum* \(Lutz, 1928\) \(Platyhelminthes: Digenea\) parasito de olhos de peixes da Bacia do Rio Paraná](#)
- [Infection by *Austrodiplostomum compactum* metacercariae in fish from the Nova Avanhandava reservoir, Tietê river, São Paulo State, Brazil](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/infection-levels-of-austrodiplostomum-compactum-digenea-14tigw0gby>

Infection levels of *Austrodiplostomum compactum* (Digenea, Diplostomidae) metacercariae in *Plagioscion squamosissimus* (Teleostei, Sciaenidae) from the Nova Avanhandava reservoir, São Paulo State, Brazil

J.V.K. Paes^{1†}, E.D. Carvalho^{2‡} and R.J. da Silva^{3*}

¹Pós-graduação em Ciências Biológicas (Zoologia), Instituto de Biociências, Universidade Estadual Paulista (UNESP) – Botucatu, Distrito de Rubião Júnior s/n°, Botucatu, 18618-00 São Paulo Brazil:

²Departamento de Morfologia, Instituto de Biociências, Universidade Estadual Paulista, Botucatu, São Paulo, Brazil: ³Departamento de Parasitologia, Instituto de Biociências, Universidade Estadual Paulista, Botucatu, São Paulo, Brazil

(Accepted 26 August 2009; First Published Online 5 November 2009)

Abstract

This study was aimed at evaluating the infection levels of *Austrodiplostomum compactum* metacercariae in 378 specimens of *Plagioscion squamosissimus* ('corvina' or 'pescada branca') from the Nova Avanhandava reservoir, low Tietê River, São Paulo State, Brazil. High prevalence, mean intensity of infection and abundance were observed in *P. squamosissimus* during most of the study, with the exception of March 2004. The relative condition factor (Kn) did not differ between parasitized and non-parasitized fish. A statistically significant positive correlation was observed between host standard length and intensity of infection. The analysis of biotic and abiotic variables showed that no abiotic variable correlated with parasitic infection levels. Moreover, *P. squamosissimus* status was demonstrated to be unaffected by *A. compactum* metacercariae infection.

Introduction

Digenetic trematodes of the genus *Austrodiplostomum* Szidat & Nani, 1951 have been reported infecting several vertebrate species (Yamaguti, 1971; Kohn *et al.*, 1995; Ramos-Ramos, 1995). The metacercariae of these parasites can infect the vitreous humour (Scholtz *et al.*, 1995; Silva-Souza, 1998; Amato *et al.*, 2001) and, more rarely, the brains of the fish (intermediate hosts)

(Ostrowski-Nunez, 1982; Silva-Souza, 1998; Amato *et al.*, 2001). The life cycle of this diplostomid is completed in the intestines of birds, such as 'biguá' *Phalacrocorax olivaceus* (Humboldt, 1805) (definitive host) (Rietschel & Werding, 1978) (Ostrowski-Nunez, 1982; Ramos-Ramos, 1995; Flowers *et al.*, 2004; Machado *et al.*, 2005). The genus *Austrodiplostomum* comprises 40 digenetic helminth species, including adults and metacercariae, and can be found in 125 host species (Eiras, 1994; Niewiadomska, 1996) distributed across all continents, but most commonly in Eurasia and North America, and occasionally in South America (Niewiadomska, 1996; Niewiadomska & Laskowski, 2002).

*E-mail: reinaldo@ibb.unesp.br

†E-mail: jvkpaes@gmail.com

‡E-mail: carvalho@ibb.unesp.br

Austrodiplostomum compactum (Lutz, 1928) Dubois, 1970, is widely distributed in the Neotropical region, where several fish species from Brazilian freshwater rivers are parasitized by metacercariae, including: *Cichla ocellaris* Bloch & Schneider, 1801 ('tucunaré' or 'tucunaré açu'); *Cichla monoculus* Spix & Agassiz, 1831 (tucunaré or tucunaré açu); *Crenicichla britskii* Kullander, 1982 ('joaninha'); *Hoplias malabaricus* (Bloch, 1794) ('traíra' or 'lobó'); *Plagioscion squamosissimus* (Heckel, 1840) ('corvina' or 'pescada branca'); and *Satanoperca pappaterra* (Heckel, 1840) ('cará', 'acará' or 'porquinho') from the Itaipu reservoir and the Upper Paraná River floodplain, State of Paraná (Kohn *et al.*, 1995; Silva-Souza, 1998; Pavanelli *et al.*, 2000; Santos *et al.*, 2002; Machado *et al.*, 2005; Yamada *et al.*, 2007); *Geophagus brasiliensis* Quoy & Gaimard, 1824 (cará or acará) from Barra Bonita (middle Tietê River, State of São Paulo) (Novaes *et al.*, 2006); and *P. squamosissimus* from Nova Avanhandava, lower Tietê River, State of São Paulo (Paes *et al.*, 2003) and from the Volta Grande reservoir,

State of Minas Gerais (Martins *et al.*, 1999, 2002). In Venezuela, Cuba, Colombia and Mexico, other metacercariae of the genus *Diplostomum* have also been recorded infecting *Rhamdia quelen* (Quoy & Gaimard, 1824) ('bagre' or 'jundiá'), *Oreochromis aureus* (Steindachner, 1864) ('tilapia' or 'tilapia azul') and *Oreochromis mossambicus* (Peters, 1852) (tilapia or 'tilapia de mozambique') (Pérez-Ponce de Leon *et al.*, 1992; Garcia *et al.*, 1993; Aragort *et al.*, 1997; Aguirre-Macedo *et al.*, 2001).

Plagioscion squamosissimus is a widespread fish species from the Amazonian basin. Because of its economical importance, it was introduced in the Paraná basin, mainly in great reservoirs such as Itaipu, Promissão and Marimbondo (Torloni *et al.*, 1993; Moretto *et al.*, 2008), where it is now the major fish product in the south-east of Brazil (Graça & Pavanelli, 2007; Vidotto & Carvalho, 2007).

Considering the previous occurrence of *A. compactum* metacercariae in the Tietê River (Paes *et al.*, 2003), the aim




Fig. 1. Map of Brazil highlighting São Paulo State and the study area at the mouth of the Santa Bárbara River, in the Nova Avanhandava reservoir (lower Tietê River).

of this study was to determine the prevalence, intensity of infection, abundance, seasonality and host status of *A. compactum* metacercariae infection. In addition, the existence of possible correlations between biotic factors (populational and parasitic) and some abiotic variables were assessed.

Materials and methods

Study area and parasitological procedures

This study was conducted in the Santa Bárbara River (municipality of Buritama, São Paulo State, Brazil), a major tributary of the Nova Avanhandava reservoir (21°07'S and 50°17'W), and the penultimate of the middle Tietê River cascade of reservoirs (fig. 1). Nova Avanhandava is the second largest hydroelectric plant in operation on the Tietê River, with an installed power capacity of 347.40 MW. The reservoir has an area of 210 km², total water volume of 2720 × 10⁶ m³, mean discharge rate of 688 m³/s, mean depth of 13 m and water permanence time of 46 days (Torloni *et al.*, 1993; CESP, 1998; Rodgher *et al.*, 2002).

Plagioscion squamosissimus specimens were collected using nylon monofilament gill nets with mesh sizes of 3–14 cm (not opposite knot lengths) from six sites on the Santa Bárbara River. Nets were placed at 17.00 h and removed by 07.00 h the following day, total exposure 14 h, during the months of July, September, October and December of 2003 and February, March, April and June of 2004. All samples were collected from the same sites with the same network area (standardized effort = 1200 m² of nets per month). The total weight (in grams) and standard length (distance from the anterior extremity of the head to the end of the hipural bone, in centimetres) of each specimen were recorded.

Three to six hours after collection, the eyes of all individuals were analysed under a stereomicroscope. Prevalence (number of hosts infected with one or more individuals of a particular parasite species), mean intensity of infection (mean number of parasites of a particular species found in the infected hosts of a particular species) and mean abundance (total number

of individuals of a particular parasite species in a sample of a particular host divided by the total number of examined hosts of that species) were calculated according to Bush *et al.* (1997). Voucher helminths were deposited in the Coleção Helmintológica do Departamento de Parasitologia, Instituto de Biociências (CHIBB), Universidade Estadual Paulista, municipality of Botucatu, São Paulo State, Brazil.

Data analysis

The relative condition factor (Kn) was employed to determine the status of *P. squamosissimus* infected by *A. compactum* (Andrade-Talmelli *et al.*, 1999; Dias *et al.*, 2004; Yamada *et al.*, 2008). Data on the weight and length of *P. squamosissimus* specimens were pooled to establish the species weight–length relationship (LeCren, 1951). A total of 377 specimens (non-parasitized = 20; parasitized = 357) were used to determine the parameters of the equation:

$$W_{\text{exp}} = L^b \times a,$$

where W_{exp} is the expected weight; L is the fish length; a is the intercept; b is the slope.

Host status was assessed using the 'relative condition factor' (Kn), estimated by dividing the observed weight of an individual by the expected weight for a given length ($\text{Kn} = W/W_{\text{exp}}$). The non-parametric Mann–Whitney statistical test was used to compare Kn between parasitized and non-parasitized fish. Intensity of infection in all samples was compared by the Kruskal–Wallis test, and multiple comparisons were performed by Dunn's method. Significance level was set at 5% in all statistical analyses.

Spearman's rank correlation coefficient (r_s) was used to test the relationship between host length and intensity of infection (Abdallah *et al.*, 2005) in order to check whether the infection process is cumulative (Takemoto *et al.*, 2005). The distribution of mean intensity of infection according to host length in parasitized specimens was also evaluated (Sturges, 1926).

Table 1. Parasitic infection data and abiotic variables monitored by collection during the study period.

Date	CHIBB	Parasitic infection data						Abiotic variables					
		N	N_i	N_m	P	MII ± SE (R)	MA ± SE (R)	Pr	Tw	pH	K	T	DO
Jul 03	511	31	30	635	96.8	21 ± 5.4 (1–129)	20 ± 5.2 (0–129)	9.2	0.8	8.3	154.0	23.4	7.6
Sep 03	512	55	53	1161	96.4	22 ± 3.7 (1–167)	21 ± 3.7 (0–167)	15.1	0.9	8.5	163.4	22.6	8.5
Oct 03	520	53	50	1067	94.3	21 ± 3.7 (1–140)	20 ± 3.5 (0–140)	101.9	1.5	7.8	176.5	25.5	9.3
Dec 03	970	63	57	1124	90.5	20 ± 3.3 (1–159)	18 ± 3.2 (0–159)	1.0	1.2	8.5	149.7	29.0	7.8
Feb 04	459	75	73	1807	97.3	25 ± 3.6 (1–159)	24 ± 3.6 (0–159)	146.7	2.4	6.8	178.9	29.2	6.8
Mar 04	561	38	31	196	81.6	6 ± 1.3 (1–33)	5 ± 1.2 (0–33)	53.3	2.0	7.0	201.8	28.4	6.2
Apr 04	476	40	39	1275	97.5	33 ± 7.2 (1–232)	32 ± 7.1 (0–232)	116.6	2.4	7.2	203.2	27.2	6.1
Jun 04	485	23	23	455	100.0	20 ± 3.4 (1–55)	20 ± 3.4 (1–55)	39.6	1.4	8.1	204.2	22.4	9.0

CHIBB, Coleção Helmintológica of the Departamento de Parasitologia, Instituto de Biociências, Universidade Estadual Paulista; N , number of collected specimens; N_i , number of parasitized specimens; N_m , total number of metacercariae; P , prevalence (%); MII, mean infection intensity; R, range; SE, standard error; MA, mean abundance; Pr, precipitation (mm); Tw, water transparency (m); K , conductivity ($\mu\text{S}/\text{cm}$); T , water temperature ($^{\circ}\text{C}$); DO, dissolved oxygen (mg/l).




Fig. 2. Weight-length relationship in *Plagioscion squamosissimus* specimens ($n = 377$) captured in the study area during the study period. Equation $y = 0.0649x^{2.5732}$; $R^2 = 0.8171$.

Concomitantly, Spearman's correlation test was used to check the existence of correlations between parasitic infection levels and some abiotic variables, such as accumulated monthly precipitation (data provided by the AES Tietê Company), water transparency (measured with a Secchi disk) and pH, electrical conductivity, water temperature and dissolved oxygen, measured with Horiba U-22 (Paes, 2006).

Results

A total of 378 *P. squamosissimus* specimens were analysed. The mean prevalence observed was $94.3 \pm 2.1\%$. The number of metacercariae collected ranged from 1 to 232 parasites. Mean intensity of infection and mean abundance were 21.6 ± 1.5 and 20.2 ± 1.6 , respectively (table 1).

During the study period, the prevalence of *A. compactum* metacercariae infection remained above 90%,




Fig. 3. Change in condition factor values (Kn) in *Plagioscion squamosissimus* around the 1.0 default, according to the total weight of parasitized (white circle; $n = 357$) and non-parasitized (black circle; $n = 20$) specimens.




Fig. 4. Correlation between standard length and abundance of *Austrodiplostomum compactum* from the Nova Avanhandava reservoir, between July 2003 and June 2004. Linear regression $y = 2.8292x - 32.004$; $R^2 = 0.1201$.

except in March 2004, when it was 81.6%. Mean intensity of infection and mean abundance also showed low variation, except in March 2004, when a significant reduction in these parameters was observed ($P \leq 0.001$) (table 1).

Host fish length and weight were 18.5 ± 3.5 (11.5–35) cm and 131.4 ± 91.8 (27.7–920.6) g, respectively. The weight-length relationship (fig. 2) was established and Kn was calculated for parasitized and non-parasitized specimens (fig. 3). No significant difference in Kn was observed between parasitized fish (1.03 ± 0.2 ; 0.44–2.1) and non-parasitized fish (1.0 ± 0.1 ; 0.86–1.09; $T = 2913$; $P = 0.068$).

The number of parasites significantly correlated with host length ($r_s = 0.32$, $P < 0.001$) and host weight ($r_s = 0.36$, $P < 0.001$) (figs 4 and 5). Therefore, a greater intensity of infection (78 parasites/individual) was observed in the larger specimens ($L_s = 26.6$ –31.5 cm) (fig. 6).

The analysis of biotic and abiotic variables (table 1) revealed that no abiotic variable correlated with parasitic infection level ($P > 0.05$).




Fig. 5. Correlation between standard length and abundance of *Austrodiplostomum compactum* from the Nova Avanhandava reservoir, between July 2003 and June 2004. Linear regression $y = 0.1114x + 5.8046$; $R^2 = 0.1264$.




Fig. 6. Mean *Austrodiplostomum compactum* infection intensity and standard deviation in *Plagioscion squamosissimus* according to host length.

Discussion

In this study, a high prevalence of *A. compactum* metacercariae infection in *P. squamosissimus* was observed. This fish species from the Amazon Basin was introduced in São Paulo Power Company (CESP) reservoirs between 1966 and 1973 (Torloni *et al.*, 1993). According to Pojmanska & Chabros (1993), the prevalence of diplostomids in allochthonous fish (species from other Neotropical basins) is higher than that of autochthonous fish (species from the Upper Paraná). This was also observed by Machado *et al.* (2005), who found higher rates of infection in the fish introduced into the floodplain of the upper Paraná River, where *P. squamosissimus* species showed the highest prevalence and intensity of infection. Thus, it is possible that *A. compactum* metacercariae utilize native fish as intermediate hosts, and that their adaptation to these native hosts is the explanation for the low prevalence and intensity of infection in these fish (Machado *et al.*, 2005).

According to Pavanelli *et al.* (2000), the occurrence of *A. compactum* metacercariae in several fish species and in different aquatic ecosystems (rivers, lakes and dams) shows that while this trematode may have been introduced with *P. squamosissimus*, it appears to have found appropriate ecological conditions for its parasitic cycle because, in addition to fish (second intermediate hosts), it also found snails (first intermediate hosts) (Machado *et al.*, 2005; Ruiz & Aguilar, 2005) and birds (definitive host) (Silva-Souza, 1998; Machado *et al.*, 2005) in many of these ecosystems. Moreover, the low level of specificity of the metacercariae associated with the presence of intermediate hosts allows metacercariae to spread and infect other species of autochthonous fish. Specifically, in the Nova Avanhandava reservoir, the intermediate hosts, i.e. gastropod molluscs *Biomphalaria glabrata* (Say, 1818) and *Biomphalaria intermedia* (Paraense & Deslandes, 1962) (França *et al.*, 2007), as well as birds *Phalacrocorax brasilianus* (Gmelin, 1789) (Branco, 2003) and those of the genus *Casmerodius* (J.V.K. Paes, pers. obs.) were also observed.

The analysis of *A. compactum* infection in *P. squamosissimus* showed that infection levels (prevalence, mean intensity of infection and abundance) were high during most parts of the study period, with the exception of March 2004. Dörücü & İspir (2001), studying *Diplostomum* sp. metacercariae infection in *Acanthobrama marmid* Heckel, 1843, also noted a seasonal variation in the intensity of infection. Silva-Souza (1998) observed variation in monthly prevalence. However, rather than a clear seasonal pattern, the author found only a trend to higher *A. compactum* prevalence in *P. squamosissimus* in the driest months. These findings differ from those reported by Santos *et al.* (2002), who found a trend towards an increase in infection levels with the increase in precipitation and temperature.

The relative condition factor, however, did not differ significantly between parasitized and non-parasitized fish, indicating a good general fish status, irrespective of the presence of the parasite. In support of this hypothesis, studies on fish composition have shown that *P. squamosissimus* is dominant in number and biomass among the species recorded in Nova Avanhandava (Paes, 2006; Vidotto & Carvalho, 2007). This indicates that *A. compactum* metacercariae parasitism does not exert a substantially negative influence on the host population. Nonetheless, lower condition factors in *P. squamosissimus* individuals parasitized by *A. compactum* metacercariae as compared to non-parasitized fish were observed in the Tibagi River, Paraná State by Silva-Souza (1998).

An increase in food supply might be a compensatory factor that minimizes the negative effect of parasitism. Indeed, this may have occurred in Nova Avanhandava, where *P. squamosissimus* preferably consumed aquatic insects, especially Odonata and Ephemeroptera (Vidotto, 2005) and Libellulidae and Gomphidae (Ramos *et al.*, 2008), and shrimps (*Macrobrachium* sp.) (Vidotto, 2005; Ramos *et al.*, 2008), which are very abundant in the reservoirs located on the middle and lower Tietê River (Pereira *et al.*, 2002). In the Tibagi River, however, the main food item for this species was fish (*Astyanax altiparanae* Garutti & Britski, 2000 and *P. squamosissimus* young) (Bennemann & Shibatta, 2002), which require a greater effort (and tactics) to catch in comparison with shrimps and aquatic insects.

Differences in the period of introduction and establishment in ecological aquatic ecosystems, as well as different limnological features may also explain the discrepancy between *P. squamosissimus* data obtained in the Tibagi River (Silva-Souza, 1998) and those found in the present study in Nova Avanhandava (Bittencourt-Oliveira, 2002; Shibatta *et al.*, 2002; Yabe & Gimenez, 2002; Paes, 2006; Vidotto & Carvalho, 2007). According to some reports, *P. squamosissimus* was equivocally introduced in the Tietê River during the 1960s (Torloni *et al.*, 1993), while in the Tibagi River, Paraná State, it was introduced more recently, in the early 1990s (Bennemann & Shibatta, 2002). This information suggests that the longer the time of *A. compactum* metacercariae interaction with *P. squamosissimus*, the smaller the negative effects on the host.

In this study, the significant correlation found between the length of parasitized specimens and mean intensity of infection indicates that *A. compactum* metacercariae

infection is a cumulative process. Aragort *et al.* (1997) also found a positive correlation between length and intensity of infection in a study of the parasitic fauna in tilapia (*O. mossambicus* and *O. aureus*) of the Valencia Lake (Venezuela). They observed that intensity of infection was higher in the larger fish (in weight), and suggested that the relationship between intensity of infection and length might be due to the fact that larger specimens have a greater body surface for infection. The present study and that of Aragort *et al.* (1997) corroborate the island biogeography theory (Kuris *et al.*, 1980).

Austrodiplostomum compactum metacercariae most commonly parasitize the vitreous humour, but some metacercariae can also be found in the aqueous humour (Garcia *et al.*, 1993) and the brain of their hosts (Ostrowski-Nunez, 1982; Conroy *et al.*, 1985; Pineda-López, 1985; Osorio-Sarabia *et al.*, 1987; Silva-Souza, 1998). In this study, however, although the brains or other organs of the specimens were not examined, only free metacercariae were found in the vitreous humour, and no eye lesions were observed macroscopically.

According to Martins (1998), digenetic trematodes are more pathogenic when a fish is acting as an intermediate host because the parasite is more aggressive to the host at this metacercaria stage than in its adult forms. Migration to the host tissues in order to reach other infection sites (eyes, brain, among others) can cause injury and, if encysted, tissue changes. The fact that metacercariae cause diseases and weaken hosts is an important ecological characteristic. Weakened fish can be easily preyed and, as a consequence, the parasite cycle can be completed in the predator (bird), the definitive host (Pavanelli *et al.*, 2006).

Evans *et al.* (1976) and Eiras (1994) showed that, depending on host size, 40 metacercariae/eye may cause blindness, cataracts or undermined vision. Garcia *et al.* (1993) found that the presence of *A. compactum* metacercariae in the eyes (aqueous and vitreous humour) and brain of tilapia *O. aureus* and *O. mossambicus* from the Amela Lake in Colima (Mexico) caused injuries in the cornea and conjunctiva. Silva-Souza (1998) did not find any injury or inflammatory reaction resulting from infection in *P. squamosissimus*. The same (no damage, cataracts or blindness) was observed in this study despite the record of up to 136 metacercariae in a single eyeball. It is noteworthy that metacercariae of other species of the Diplostomidae family have caused mortality and decreases in condition factors in catfish *Ictalurus punctatus* (Rafinesque, 1818) at fish farms in the USA (Overstreet & Curran, 2004).

Acknowledgements

The authors are thankful to Laboratório de Biologia e Ecologia de Peixes and Laboratório de Parasitologia de Animais Silvestres (Universidade Estadual Paulista, UNESP – Instituto de Biociências – IB, Campus de Botucatu, São Paulo, Brazil), for the use of their facilities during the conduction of this research; the Post-Graduate Course in Biological Sciences (Zoology) for their support during every stage of this work; Dr Ana Paula Vidotto

and Dr José Luís Costa Novaes (Departamento de Morfologia, IB, UNESP) for their help in the critical reading of the manuscript; Ricardo A. S. Teixeira, Renato Devidé (Departamento de Morfologia, IB, UNESP) and Dr Karina Rodrigues dos Santos (Departamento de Parasitologia, IB, UNESP, Botucatu) for their help in the field and laboratory work; Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq (process n. 133203/2004-2 and 140838/2007-4), Fundação Paulista de Amparo a Pesquisa – FAPESP (process n. 03/11 239-2) and Fundação para o Desenvolvimento da Unesp – FUNDUNESP (process n. 00329/04) for their financial support. This work complies with current Brazilian laws.

References

- Abdallah, V.D., Azevedo, R.K. & Luque, J.L. (2005) Ecologia da comunidade de metazoários parasitos do sairú *Cyphocarax gilbert* (Quoy e Gaimard, 1824) (Characiformes: Curimatidae) do rio Guandu, Estado do Rio de Janeiro, Brasil. *Revista Brasileira de Parasitologia Veterinária* **14**, 154–159.
- Aguirre-Macedo, M.L., Scholz, T., González-Solís, D., Vidal-Martinez, V.M., Posel, P., Arjona-Torres, G., Siu-Estrada, E. & Dumailo, S. (2001) Larval helminths parasitizing freshwater fishes from the Atlantic Coast of Nicaragua. *Comparative Parasitology* **68**, 42–51.
- Amato, S.B., Amato, J.F.R. & Albrecht, M. (2001) Metacercárias livres de diplostomídeos (Digenea, Diplostomidae) em *Loricariichthys anus* (Val., 1840) (Siluriformes, Loricariidae) do Estado do Rio Grande do Sul, Brasil. *Parasitologia al día* **25**, 1–7.
- Andrade-Talmelli, E.F., Fenerich-Verani, N. & Verani, J.R. (1999) Fator de condição relativo (Kn): um critério para selecionar fêmeas de Piabinha, *Brycon insignis* (Steindachner, 1876) (Pisces: Bryconinae), para indução reprodutiva. *Boletim do Instituto de Pesca de São Paulo* **25**, 95–99.
- Aragort, F.W., Leon, E.A., Guillén, A.T., Silva, M. & Balestrini, C. (1997) Fauna parasitaria em tilapias del Lago de Valencia. *Veterinaria Tropical* **22**, 171–187.
- Bennemann, S.T. & Shibatta, A.O. (2002) Dinâmica de uma assembléia de peixes do rio Tibagi. pp. 433–442 in Medri, M.E., Bianchini, E., Shibatta, A.O. & Pimenta, J.A. (Eds) *A bacia do rio Tibagi*. Londrina, UEL.
- Bittencourt-Oliveira, M.C. (2002) A comunidade fitoplanctônica do rio Tibagi: uma abordagem preliminar de sua diversidade. pp. 373–402 in Medri, M.E., Bianchini, E., Shibatta, A.O. & Pimenta, J.A. (Eds) *A bacia do rio Tibagi*. Londrina, UEL.
- Branco, M.B.C. (2003) Diversidade da avifauna aquática nas represas do Médio e Baixo Rio Tietê (SP) e no sistema de lagos do Médio Rio Doce (MG) e sua relação com o estado trófico e a morfometria dos ecossistemas aquáticos Masters dissertation, Universidade Federal de São Carlos – UFSCar, São Carlos (SP).
- Bush, A.O., Lafferty, K.D., Lotz, J.M. & Shostak, A.W. (1997) Parasitology meets ecology on its own terms: Margolis revisited. *The Journal of Parasitology* **83**, 575–583.

- CESP (Companhia Energética de São Paulo)** (1998) *Conservação e manejo nos reservatórios: limnologia, ictiologia e pesca*. Série Divulgação e informação. São Paulo, CESP.
- Conroy, G., Conroy, D.A., Santacana, J.A. & Perdomo, F.** (1985) Diplostomatosis in cultured Venezuelan grey mullets. *Bulletin of the European Association of Fish Pathologists* **5**, 14–16.
- Dias, P.G., Furuya, W.M., Pavanelli, G.C., Machado, M.H. & Takemoto, R.M.** (2004) Carga parasitária de *Rondonia rondoni* Travassos, 1920 (Nematoda, Atrictidae) e fator de condição do armado, *Pterodoras granulosus*, Valenciennes, 1833 (Pisces, Doradidae). *Acta Scientiarum Biology Science* **26**, 151–156.
- Dörücü, M. & İspir, Ü.** (2001) Seasonal variation of *Diplostomum* sp. infection in eyes of *Acanthobrama marmid* Heckel, 1843 in Keban Dam Lake, Elazığ, Turkey. *E.U. Journal of Fisheries and Aquatic Sciences* **18**, 301–305.
- Eiras, J.C.** (1994) *Elementos de Ictioparasitologia*. 339 pp. Porto, Fundação Engenheiro Antônio de Almeida.
- Evans, R.S., Heckmann, R.A. & Palmieri, J.** (1976) Diplostomiasis in Utah. *Utah Academy Proceedings* **53**, 20–25.
- Flowers, J.R., Poore, M.F., Mullen, J.E. & Levy, M.G.** (2004) Digeneans collected from piscivorous birds in North Carolina, USA. *Comparative Parasitology* **71**, 243–244.
- França, R.S., Suriani, A.L. & Rocha, O.** (2007) Composição das espécies de moluscos bentônicos nos reservatórios do baixo rio Tietê (São Paulo, Brasil) com uma avaliação do impacto causado pelas espécies exóticas invasoras. *Revista Brasileira de Zoologia* **24**, 41–51.
- García, M.L.J., Osório-Sarabia, D. & Constantino, F.** (1993) Prevalência de los parasitos y las alteraciones histológicas que producen a las tilapias de la laguna de Amela Tecoman Colima. *Veterinaria México* **24**, 199–205.
- Graça, W.J. & Pavanelli, C.S.** (2007) *Peixes da planície de inundação do alto rio Paraná e áreas adjacentes*. 241 pp. Maringá, EDUEM.
- Kohn, A., Fernandes, B.M.M. & Baptista-Farias, M.F.D.** (1995) Metacercariae of *Diplostomum* (*Austrodiplostomum*) *compactum* (Trematoda, Diplostomidae) in the eyes of *Plagioscion squamosissimus* (Teleostei, Sciaenidae) from the reservoir of the Hydroelectric Power Station of Itaipu, Brazil. *Memórias do Instituto Oswaldo Cruz* **90**, 341–344.
- Kuris, A.M., Blaustein, A.R. & Alió, J.J.** (1980) Hosts as islands. *The American Naturalist* **116**, 570–586.
- LeCren, E.D.** (1951) The length–weight relationship and seasonal cycle in gonad weight and condition in perch *Perca fluviatilis*. *The Journal of Animal Ecology* **20**, 201–219.
- Machado, P.M., Takemoto, R.M. & Pavanelli, G.C.** (2005) *Diplostomum* (*Austrodiplostomum*) *compactum* (Lutz, 1928) (Platyhelminthes, Digenea) metacercariae in fish from the floodplain of the Upper Paraná River, Brazil. *Parasitology Research* **97**, 436–444.
- Martins, M.L.** (1998) *Doenças infecciosas e parasitárias de peixes*. Jaboticabal, Boletim Técnico do Centro de Aqüicultura da UNESP.
- Martins, M.L., Fujimoto, R.Y., Nascimento, A.A. & Moraes, E.R.** (1999) Ocorrência de *Diplostomum* sp. Nordmann, 1832 (Digenea: Diplostomidae) em *Plagioscion squamosissimus* (Heckel, 1840) proveniente do reservatório de Volta Grande, MG, Brasil. *Acta Scientiarum* **21**, 263–266.
- Martins, M.L., Paiva, A.M.F.C., Fujimoto, R.Y., Schalch, S.H.C. & Colombano, N.C.** (2002) Prevalência, sazonalidade e intensidade de infecção por *Diplostomum* (*Austrodiplostomum*) *compactum* Lutz, 1928 (Digenea, Diplostomidae), em peixes do reservatório de Volta Grande, Estado de Minas Gerais, Brasil. *Acta Scientiarum* **24**, 469–474.
- Moretto, E.M., Marciano, F.T., Velludo, M.R., Fenerich-Verani, N., Espindola, E.L.G. & Rocha, O.** (2008) The recent occurrence, establishment and potential impact of *Geophagus proximus* (Cichlidae: Perciformes) in the Tietê river reservoirs: an Amazonian fish species introduced in the Paraná Basin (Brazil). *Biodiversity and Conservation* **17**, 3013–3025.
- Niewiadomska, K.** (1996) The genus *Diplostomum* – taxonomy, morphology and biology. *Acta Parasitologica* **41**, 55–66.
- Niewiadomska, K. & Laskowski, Z.** (2002) Systematic relationships among six species of *Diplostomum* Nordmann, 1832 (Digenea) based on morphological and molecular data. *Acta Parasitologica* **47**, 20–28.
- Novaes, J.C.L., Ramos, I.P., Carvalho, E.D. & Silva, R.J.** (2006) Metacercariae of *Diplostomum compactum* Lutz, 1928 (Trematoda, Diplostomidae) in the eyes of acara *Geophagus brasiliensis* Quoy & Gaimard, 1824 (Teleostei, Cichlidae) from Barra Bonita Reservoir – São Paulo, Brazil. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* **58**, 1229–1231.
- Osorio-Sarabia, V., Pineda-López, R. & Salgado-Maldonado, G.** (1987) Fauna helmintológica de peces dulceacuícolas de Tabasco. Estudio preliminar. *Universidad y Ciencia* **4**, 5–31.
- Ostrowski-Nunez, M.** (1982) The life histories of *Diplostomum* (*Austrodiplostomum*) *compactum* (Lutz, 1928) Dubois, 1970 and *D. (A.) mordax* (Szidat and Nani, 1951) n. comb. in South América. *Zoologischer Anzeiger* **208**, 393–404.
- Overstreet, R.M. & Curran, S.S.** (2004) Defeating diplostomoid dangers in USA catfish aquaculture. *Folia Parasitologica* **51**, 153–165.
- Paes, J.V.K.** (2006) A ictiofauna associada e as condições limnológicas num sistema de piscicultura em tanques-rede no reservatório de Nova Avanhandava (baixo Rio Tietê, SP). Masters dissertation, Universidade Estadual Paulista – UNESP, Botucatu (SP).
- Paes, J.V.K., Santos, K.R., Carvalho, E.D. & Silva, R.J.** (2003) Ocorrência de metacercária de *Diplostomum compactum* (Trematoda, Diplostomidae) parasitando *Plagioscion squamosissimus* (Teleostei, Sciaenidae) proveniente do reservatório de Nova Avanhandava, Buritama, São Paulo. *Arquivos do Instituto Biológico* **70**, 383–387.
- Pavanelli, G.C., Takemoto, R.M., Machado, P.M., Lizama, M.A.P., Tanaka, L.K., Guidelli, G.M., Issac, A., Carvalho, S. & França, J.G.** (2000) *A planície alagável do Alto Rio Paraná: estrutura e processos ambientais*.

- Componentes biológicos: ictioparasitologia. Programa PELD/CNPQ.
- Pavanelli, G.C., Takemoto, R.M., Lizama, M.A.P., Guidelli, G.M., Lacerda, A.C.F., Yamada, F.H., Tavernari, F.C., Moreira, L.H.A., Ceschini, T.L., Bellay, S. & Fernandes, E.S.** (2006) *A planície de inundação do alto rio Paraná: Ictioparasitologia*. Programa PELD/CNPQ.
- Pereira, C.C.G.F., Smith, W.S., Espíndola, E.L.G. & Rocha, O.** (2002) Alterações tróficas nas espécies de peixes em decorrência da construção de reservatórios em cascata no médio e baixo rio Tietê. pp. 29–41 in PPG – CEA. *Recursos Hidroenergéticos: Usos, Impactos e Planejamento Integrado*. São Carlos, Rima Editora.
- Pérez-Ponce de Leon, G., Osório-Saraiba, D. & Garcia-Prieto, L.** (1992) Helminthofauna del 'Juile' *Rhamdia guatemalensis* (Pisces: Pimelodidae), del lago de Catemaco, Veracruz. *Revista da Sociedade Mexicana de História Natural* **43**, 25–31.
- Pineda-López, R.** (1985) Infección por metacercárias (Platyhelminthes: Trematoda) em peixes de agua dulce de Tabasco. *Universidad y Ciencia* **2**, 47–60.
- Pojmanska, T. & Chabros, M.** (1993) Parasites of common carp and three introduced cyprinid fish in pound culture. *Acta Parasitologica* **38**, 101–108.
- Ramos, I.P., Vidotto-Magnoni, A.P. & Carvalho, E.D.** (2008) Influence of cage fish farming on the diet of dominant fish species of a Brazilian reservoir (Tietê River, High Paraná River Basin). *Acta Limnologica Brasiliensia* **20**, 245–252.
- Ramos-Ramos, P.** (1995) Algunos trematodos de vertebrados de la presa Miguel Aleman em Temascal, Oaxaca, México. *Anales del Instituto de Biología, Serie Zoología* **66**, 241–246.
- Rietschel, G. & Werding, B.** (1978) Trematodes of birds from Northern Columbia. *Zeitschrift für Parasitenkunde* **57**, 57–82.
- Rodger, S., Espíndola, E.L.G., Fracácio, R., Rodrigues, M.H., Pereira, R.H.G. & Rocha, O.** (2002) Estudos ecotoxicológicos nos reservatórios em cascata do médio e baixo rio Tietê: uma avaliação dos impactos ambientais. pp. 131–144 in PPG – CEA. *Recursos Hidroenergéticos: Usos, Impactos e Planejamento Integrado*. São Carlos, Rima Editora.
- Ruiz, L.J.R. & Aguilar, J.G.** (2005) Estructura de la comunidad y dinámica poblacional de gasterópodos em una zona enzoótica de fasciolosis em Tabasco, México. *Acta Zoologica Mexicana* **21**, 79–85.
- Santos, R.S., Pimenta, F.D.A., Martins, M.L., Takahashi, H.K. & Marengoni, N.G.** (2002) Metacercárias de *Diplostomum (Austrodiplostomum) compactum* Lutz, 1928 (Digenea, Diplostomidae) em peixes do rio Paraná, Brasil, Prevalência, sazonalidade e intensidade de infecção. *Acta Scientiarum* **24**, 475–480.
- Scholtz, T., Vargas-Vázquez, J., Moravec, F., Vivas-Rodríguez, C. & Mendonza-Franco, E.** (1995) Metacercariae of trematodes of fishes from cenotes (=sinkholes) of the Yucatan Península, México. *Folia Parasitologica* **42**, 173–192.
- Shibatta, O.A., Orsi, M.L., Bennemann, S.T. & Silva-Souza, A.T.** (2002) Diversidade e distribuição de peixes na bacia do rio Tibagi. pp. 403–423 in Medri, M.E., Bianchini, E., Shibatta, A.O. & Pimenta, J.A. (Eds) *A bacia do rio Tibagi*. Londrina, Universidade Estadual de Londrina.
- Silva-Souza, A.T.** (1998) Estudo do parasitismo de *Plagioscion squamosissimus* (Heckel 1940) (Perciformes, Sciaenidae) por *Diplostomum (Austrodiplostomum) compactum* (Lutz, 1928) (Trematoda, Digenea) no rio Tibagi, PR. PhD Thesis, UFSCar, São Carlos.
- Sturges, H.A.** (1926) The choice of a class interval. *Journal of the American Statistical Association* **21**, 65–66.
- Takemoto, R.M., Lizama, M.A.P., Guidelli, G.M., Pavanelli, G.C., Moreira, S.T., Ito, K.F., Lacerda, A.C.F. & Bellay, S.** (2005) Fauna parasitária de peixes em reservatórios. pp. 269–280 in Rodrigues, L., Thomaz, S.M., Agostinho, A.A. & Gomes, L.C. (Eds) *Biocenoses em reservatórios: padrões espaciais e temporais*. São Carlos, Rima Editora.
- Torloni, C.E.C., Corrêa, A.R.A., Carvalho, A.A. Jr, Santos, J.J., Gonçalves, J.L., Gereto, E.J., Cruz, J.A., Moreira, J.A., Silva, D.C., Deus, E.F. & Ferreira, A.S.** (1993) *Produção pesqueira e composição das capturas em reservatórios sob concessão da CESP nos rios Tietê, Paraná e Grande, no período de 1986 a 1991*. 73 pp. Série pesquisa e desenvolvimento No. 1. São Paulo, CESP.
- Vidotto, A.P.** (2005) Estrutura da comunidade de peixes do reservatório de Nova Avanhandava (baixo rio Tietê SP), com ênfase na dinâmica populacional e dieta das espécies introduzidas. Masters dissertation, Universidade Estadual Paulista – UNESP, Botucatu (SP).
- Vidotto, A.P. & Carvalho, E.D.** (2007) Composition and structure of fish community in a stretch of the Santa Bárbara River influenced by Nova Avanhandava Reservoir (low Tietê River, São Paulo State, Brazil). *Acta Limnologica Brasiliensia* **19**, 233–245.
- Yabe, M.J.S. & Gimenez, S.M.N.** (2002) Caracterização físico-química da água da bacia do rio Tibagi. pp. 355–371 in Medri, M.E., Bianchini, E., Shibatta, A.O. & Pimenta, J.A. (Eds) *A bacia do rio Tibagi*. Londrina, UEL.
- Yamada, F.H., Takemoto, R.M. & Pavanelli, G.C.** (2007) Ecological aspects of ectoparasites from the gills of *Satanoperca pappaterra* (Heckel, 1840) (Cichlidae) from the upper Paraná river floodplain, Brazil. *Acta Scientiarum Biology Science* **29**, 331–336.
- Yamada, F.H., Moreira, L.H.A., Ceschini, T.L., Takemoto, R.M. & Pavanelli, G.C.** (2008) Novas ocorrências de metacercárias de *Austrodiplostomum compactum* (Lutz, 1928) (Platyhelminthes: Digenea) parasito de olhos de peixes da bacia do rio Paraná. *Revista Brasileira de Parasitologia Veterinária* **17**, 163–166.
- Yamaguti, S.** (1971) *Synopsis of Digenetic Trematodes of Vertebrates*. Vol. II. 1074 pp. Tokyo, Keigaku Publishing Company.