



Infinite Matrices and Almost Convergence

Ab. Hamid Ganie^{a,b}, Neyaz Ahmad Sheikh^a

^aDepartment of mathematics, National Institute of Technology Srinagar, 190006, India

^bDepartment of mathematics, S.S. M College of Engineering and Technology Pattan -193121, J and K.

Abstract. In the present paper, we characterize $(r^q(u, p) : f_\infty)$, $(r^q(u, p) : f)$ and $(r^q(u, p) : f_0)$; where f_∞ , f and f_0 denotes, respectively, the spaces of almost bounded sequences, almost convergent sequences and almost sequences converging to zero, where the space $r^q(u, p)$ of non-absolute type have recently been introduced by Neyaz and Hamid (see, [13]).

1. Introduction, Background and Notation

Let ω denote the space of all sequences (real or complex). The family under pointwise addition and scalar multiplication forms a linear (vector) space over real or complex numbers. Any subspace of ω is called the sequence space. So the sequence space is the set of scalar sequences (real or complex) which is closed under co-ordinate wise addition and scalar multiplication. Throughout paper \mathbb{N} , \mathbb{R} and \mathbb{C} denotes the set of non-negative integers, the set of real numbers and the set of complex numbers, respectively.

Let X and Y be two sequence spaces and let $A = (a_{nk})$ be an infinite matrix of real or complex numbers a_{nk} , where $n, k \in \mathbb{N}$. Then, the matrix A defines the A -transformation from X into Y , if for every sequence $x = (x_k) \in X$ the sequence $Ax = \{(Ax)_n\}$, the A -transform of x exists and is in Y ; where $(Ax)_n = \sum_k a_{nk} x_k$. For simplicity in notation, here and in what follows, the summation without limits runs from 0 to ∞ . By $A \in (X : Y)$ we mean the characterizations of matrices from X to Y i.e., $A : X \rightarrow Y$. A sequence x is said to be A -summable to l if Ax converges to l which is called as the A -limit of x . For a sequence space X , the matrix domain X_A of an infinite matrix A is defined as

$$X_A = \{x = (x_k) \in \omega : Ax \in X\}. \quad (1)$$

Let l_∞ and c be Banach spaces of bounded and convergent sequences $x = \{x_n\}_{n=0}^\infty$ with supremum norm $\|x\| = \sup_n |x_n|$. Let T denote the shift operator on ω , that is, $Tx = \{x_n\}_{n=1}^\infty$, $T^2x = \{x_n\}_{n=2}^\infty$ and so on. A Banach limit L is defined on l_∞ as a non-negative linear functional such that L is invariant i.e., $L(Sx) = L(x)$ and

2010 Mathematics Subject Classification. Primary 40A05, 46A45; Secondary 46C05

Keywords. Sequence space of non-absolute type, almost convergence, β -duals and matrix transformations

Received: 23 November 2013; Accepted: 04 April 2014

Communicated by Dragan S. Djordjević

Corresponding author: Ab. Hamid Ganie

Email addresses: ashamidg@rediffmail.com (Ab. Hamid Ganie), neyaznit@yahoo.co.in (Neyaz Ahmad Sheikh)

$L(e) = 1, e = (1, 1, 1, \dots)$ (see, [2]).

Lorentz (see, [7]), called a sequence $\{x_n\}$ almost convergent if all Banach limits of $x, L(x)$, are same and this unique Banach limit is called F -limit of x . In his paper, Lorentz proved the following criterion for almost convergent sequences.

A sequence $x = \{x_n\} \in l_\infty$ is almost convergent with F -limit $L(x)$ if and only if

$$\lim_{m \rightarrow \infty} t_{mn}(x) = L(x)$$

where, $t_{mn}(x) = \frac{1}{m} \sum_{j=0}^{m-1} T^j x_n, (T^0 = 0)$ uniformly in $n \geq 0$.

We denote the set of almost convergent sequences by f .

Nanda (see, [11]) has defined a new set of sequences f_∞ as follows:

$$f_\infty = \left\{ x \in l_\infty : \sup_{mn} |t_{mn}(x)| < \infty \right\}.$$

We call f_∞ as the set of all almost bounded sequences.

A infinite matrix $A = (a_{nk})$ is said to be regular (see, [11]) if and only if the following conditions (or Toeplitz conditions) hold:

- (i) $\lim_{n \rightarrow \infty} \sum_{k=0}^{\infty} a_{nk} = 1,$
- (ii) $\lim_{n \rightarrow \infty} a_{nk} = 0, (k = 0, 1, 2, \dots),$
- (iii) $\sum_{k=0}^{\infty} |a_{nk}| < M, (M > 0, j = 0, 1, 2, \dots).$

Let (q_k) be a sequence of positive numbers and let us write, $Q_n = \sum_{k=0}^n q_k$ for $n \in \mathbb{N}$.

Then the matrix $R^q = (r_{nk}^q)$ of the Riesz mean (R, q_n) is given by

$$r_{nk}^q = \begin{cases} \frac{q_k}{Q_n}, & \text{if } 0 \leq k \leq n, \\ 0, & \text{if } k > n. \end{cases}$$

The Riesz mean (R, q_n) is regular if and only if $Q_n \rightarrow \infty$ as $n \rightarrow \infty$ (see, [1, 11]).

Following [1-3, 6-8, 12-14], the sequence space $r^q(u, p)$ is defined as

$$r^q(u, p) = \left\{ x = (x_k) \in \omega : \sum_k \left| \frac{1}{Q_k} \sum_{j=0}^k u_j q_j x_j \right|^{p_k} \right\} < \infty, (0 < p_k \leq H = \sup p_k < \infty)$$

where, $u = (u_k)$ is a sequence such that $u_k \neq 0$ for all $k \in \mathbb{N}$.

With the notation of (1), we may re-define $r^q(u, p)$ as follows:

$$r^q(u, p) = \{l(p)\}_{R_u^q}.$$

Define the sequence $y = (y_k)$, which will be used, by the R_u^q -transform of a sequence $x = (x_k)$, i.e.,

$$y_k = \frac{1}{Q_k} \sum_{j=0}^k u_j q_j x_j. \tag{2}$$

We denote by X^β the β - dual of X and mean the set of all sequences $x = (x_k)$ such that $xy = (x_k y_k) \in cs$ for all $y = (y_k) \in X$.

2. Main Results

We first state some lemmas which are needed in proving the main results.

Lemma 2.1 [13]. Define the sets $D_1(u, p)$ and $D_2(u, p)$ as follows,

$$D_1(u, p) = \left\{ a = (a_k) \in \omega : \sup_k \left| \Delta \left(\frac{a_k}{u_k q_k} \right) Q_k \right|^{p_k} < \infty \text{ and } \sup_k \left| \frac{a_k}{u_k q_k} Q_k \right|^{p_k} < \infty \right\}$$

and

$$D_2(u, p) = \left\{ a = (a_k) \in \omega : \sum_k \left| \Delta \left(\frac{a_k}{u_k q_k} \right) Q_k C^{-1} \right|^{p'_k} < \infty \text{ and } \left\{ \left(\frac{a_k}{u_k q_k} Q_k C^{-1} \right)^{p'_k} \right\} \in l_\infty \right\}.$$

Then,

$$[r^q(u, p)]^\beta = D_1(u, p); (0 < p_k \leq 1) \text{ and } [r^q(u, p)]^\beta = D_2(u, p); (1 < p_k \leq H < \infty).$$

Lemma 2.2 [13]. $r^q(u, p)$ is a complete linear metric space paranormed by g where

$$g(x) = \left[\sum_k \left| \frac{1}{Q_k} \sum_{j=0}^k u_j q_j x_j \right|^{p_k} \right]^{\frac{1}{M}} \text{ with } 0 < p_k \leq H < \infty, M = \max\{1, H\}.$$

Lemma 2.3 [13]. The Riesz sequence space $r^q(u, p)$ of non-absolute type is linearly isomorphic to the space $l(p)$, where $0 < p_k \leq H < \infty$.

Lemma 2.4 [7]. $f \subset f_\infty$.

For simplicity in notation, we shall write

$$t_{mn}(Ax) = \frac{1}{m+1} \sum_{j=0}^m A_{n+j}(x) = \sum_k a(n, k, m) x_k$$

where,

$$a(n, k, m) = \frac{1}{m+1} \sum_{j=0}^m a_{n+j,k}; \quad (n, k, m \in \mathbb{N}).$$

Also,

$$\widehat{a}(n, k, m) = \Delta \left[\frac{a(n, k, m)}{u_k q_k} \right] Q_k$$

where,

$$\Delta \left[\frac{a(n, k, m)}{u_k q_k} \right] Q_k = \left[\frac{a(n, k, m)}{u_k q_k} - \frac{a(n, k+1, m)}{u_{k+1} q_{k+1}} \right] Q_k.$$

Theorem 2.5.(i) Let $1 < p_k \leq H < \infty$ for every $k \in \mathbb{N}$. Then $A \in (r^q(u, p) : f_\infty)$ if and only if there exists an integer $C > 1$ such that

$$\sup_{n, m \in \mathbb{N}} \sum_k |\widehat{a}(n, k, m) C^{-1}|^{p'_k} < \infty, \tag{3}$$

$$\left\{ \left(\frac{a_{nk}}{u_k q_k} Q_k C^{-1} \right)^{p'_k} \right\} \in l_\infty \text{ for all } n \in \mathbb{N}. \tag{4}$$

(ii) Let $0 \leq p_k \leq 1$ for every $k \in \mathbb{N}$. Then $A \in (r^q(u, p) : f_\infty)$ if and only if

$$\sup_{n, k, m \in \mathbb{N}} |\widehat{a}(n, k, m)|^{p_k} < \infty. \tag{5}$$

Proof. Since $f_\infty = l_\infty$, so the proof follows from [Th. 3.1, 13]. \square

Theorem 2.6. Let $1 < p_k \leq H < \infty$ for every $k \in \mathbb{N}$. Then $A \in (r^q(u, p) : f)$ if and only if (3), (4), (5) holds of Theorem 2.5 and there is a sequence of scalars (β_k) such that

$$\lim_{m \rightarrow \infty} \widehat{a}(n, k, m) = \beta_k, \text{ uniformly in } n \in \mathbb{N}. \tag{6}$$

Proof: Sufficiency. Suppose that the conditions (3), (4), (5) and (6) hold and $x \in r^q(u, p)$. Then Ax exists and we have by (6) that $|a(n, k, m) C^{-1}|^{p'_k} \rightarrow |\beta_k C^{-1}|^{p'_k}$ as $m \rightarrow \infty$, uniformly in n and for each $k \in \mathbb{N}$, which leads us with (3) to the inequality

$$\sum_{j=0}^k |\beta_j C^{-1}|^{p'_j} = \lim_{m \rightarrow \infty} \sum_{j=0}^k |\widehat{a}(n, j, m) C^{-1}|^{p'_j} \text{ (uniformly in } n)$$

$$\leq \sup_{n,m \in \mathbb{N}} \sum_{j=0}^k |\widehat{a}(n, j, m)C^{-1}|^{p'_k} < \infty, \tag{7}$$

holding for every $k \in \mathbb{N}$. Since $x \in r^q(u, p)$ by hypothesis and $r^q(u, p) \cong lp$ (by Lemma 2.3 above), we have $y \in l(p)$. Therefore, it follows from (7) that the series $\sum_k \beta_k y_k$ and $\sum_k \widehat{a}(n, k, m)y_k$ converges for each m, n and

$y \in l(p)$. Now, for a given $\epsilon > 0$, choose a fixed $k_0 \in \mathbb{N}$ such that $\left(\sum_{k=k_0+1}^{\infty} |y_k|^{p_k}\right)^{\frac{1}{p_k}} < \epsilon$. Then, there is some

$m_0 \in \mathbb{N}$ such that $\left|\sum_{k=0}^{k_0} [\widehat{a}(n, k, m) - \beta_k]\right| < \epsilon$ for every $m \geq m_0$, uniformly in n . Since (6) holds, it follows that $\left|\sum_{k=0}^{\infty} [\widehat{a}(n, k, m) - \beta_k]\right|$ is arbitrary small. Therefore, $\lim_m \sum_k a(n, k, m)x_k = \lim_m \sum_k \widehat{a}(n, k, m)y_k = \sum_k \beta_k y_k$ uniformly in n . Hence, $Ax \in f$, this proves sufficiency.

Necessity. Suppose that $A \in (r^q(u, p) : f)$. Then, since $f \subset f_\infty$ (by Lemma 2.4 above), the necessities of (3) and (4) are immediately obtained from Theorem 2.3 (above). To prove the necessity of (6), consider the sequence

$$b_n^k(q) = \begin{cases} (-1)^{n-k} \frac{Q_k}{u_n q_n}, & \text{if } k \leq n \leq k + 1, \\ 0, & \text{if } 0 \leq n < k \text{ or } n > k + 1. \end{cases}$$

Since Ax exists and is in f for each $x \in r^q(u, p)$, one can easily see that $Ab^{(k)}(q) = \left\{ \Delta \left(\frac{a_{nk}}{u_k q_k} \right) Q_k \right\}_{n \in \mathbb{N}} \in f$ for all $k \in \mathbb{N}$, which proves the necessity of (6). This concludes the proof. \square

Theorem 2.7. Let $1 < p_k \leq H < \infty$ for every $k \in \mathbb{N}$. Then $A \in (r^q(u, p) : f_0)$ if and only if (3), (4), (5) and (6) holds with $\beta_k \rightarrow 0$ for each $k \in \mathbb{N}$.

Proof. The proof follows from above Theorem 2.6 by taking $\beta_k \rightarrow 0$ for each $k \in \mathbb{N}$.

Corollary 2.8. By taking $u_k = 1$ for all $k \in \mathbb{N}$, we get results obtained by Jalal and Ganie [5].

Acknowledgement. We are very thankful to the referee(s) for the careful reading, valuable suggestions which improved the presentation of the paper.

References

[1] B. Altay and F. Başar, On the paranormed Riesz sequence space of nonabsolute type, Southeast Asian Bulletin of Mathematics 26(2002) 701–715.
 [2] S. Banach, Théories des operations linéaires, Warszawa 1932.
 [3] C. Cakan, B. Altay and M. Mursaleen, The σ -convergence and σ -core of double sequences, Applied Mathematics Letters 19(2006) 1122–1128.
 [4] S. A. Gupkari, Some new sequence space and almost convergence, Filomat 22(2)(2008) 59–64.
 [5] T. Jalal and A. H. Ganie, Almost Convergence and some matrix transformation, Shekhar(New Series)- International Journal of Mathematics 1(1) (2009) 133–138.
 [6] K. Kayaduman and M. Sengnl, The spaces of Cesáro almost convergent sequences and core theorems, Acta Mathematica Scientia 32 B(6)(2012) 2265–2278.
 [7] G. G. Lorentz, A contribution to the theory of divergent series, Acta Mathematica (80)(1948) 167–190.

- [8] M. Mursaleen, A. M. Jarrah and S. A. Mohiuddine, Almost convergence through the generalized de la Vallee-Pousin mean, *Iranian Journal of Science and Technology Transaction A-science* 33(2009) 169–177.
- [9] M. Mursaleen and S.A. Mohiuddine, Double σ -multiplicative matrices, *Journal of Mathematical Analysis and Applications* 327(2007) 991–996.
- [10] M. Mursaleen and E. Savas, Almost regular matrices for double sequences, *Studia Scientiarum Mathematicarum Hungarica* 40(2003) 205–212.
- [11] S. Nanda, Matrix transformations and almost boundedness, *Glasnik Matematicki* 14(34) (1979) 99–107.
- [12] G. M. Petersen, *Regular matrix transformations*. McGraw-Hill Publishing Co. Ltd., London-New York-Toronto 1966.
- [13] N. A. Sheikh and A. H. Ganie, A new paranormed sequence space and some matrix transformations, *Acta Mathematica Academiae Paedagogicae Nyiregyhaziensis* 28(1)(2012) 47–58.
- [14] N. A. Sheikh and A. H. Ganie, A new type of sequence space of non-absolute type and matrix transformation, *WSEAS Transaction Journal of Mathematics* 8(12) (August-2013) 852–589.
- [15] N. A. Sheikh and A. H. Ganie, On the spaces of λ -convergent sequences and almost convergence, *Thai Journal of Mathematics* 11(2)(2013) 393–398.