

INFINITE SUBCLASSES OF RECURSIVELY ENUMERABLE CLASSES

J. B. FLORENCE

P. R. Young [1] has constructed an infinite recursively enumerable (r.e.) class with no proper infinite r.e. subclasses, and has asked if infinite r.e. classes with $m+1$ infinite r.e. subclasses exist for every $m \geq 0$. It can further be asked what is the most general partially ordered set we can represent by the infinite r.e. subclasses of such a class (under inclusion). These questions are answered by the theorem below. The author wishes to thank A. H. Lachlan for his guidance and encouragement. Our construction is based on a formulation of Young's due to Lachlan.

THEOREM. (a) *Let $m \geq 1$, $n \geq 1$. Let $\{F_i | 1 \leq i \leq m+1\}$ be a class of subsets of $\{x | 1 \leq x \leq n\}$ closed to subsets and with*

$$\bigcup \{F_i | 1 \leq i \leq m+1\} = \{x | 1 \leq x \leq n\}.$$

Then there is an infinite class \mathcal{C}^ of r.e. sets and distinct r.e. sets A_1, \dots, A_n not in \mathcal{C}^* such that*

$$\mathcal{C} = \mathcal{C}^* \cup \{A_i | 1 \leq i \leq n\}$$

is an infinite r.e. class with infinite r.e. subclasses

$$\mathcal{C} - \{A_i | i \in F_j\} \quad (1 \leq j \leq m+1).$$

(b) *There is an infinite r.e. class with one infinite r.e. subclass.*

Conversely, any infinite r.e. class with finitely many infinite r.e. subclasses is of one of these forms.

PROOF. We prove the converse first. Let \mathcal{C} be an infinite r.e. class with $m+1$ ($m \geq 1$) infinite r.e. subclasses. If \mathcal{C}_1 is an r.e. subclass and $X \in \mathcal{C} - \mathcal{C}_1$, then $\mathcal{C}_1 \cup \{X\}$ is an r.e. subclass. It follows that each infinite r.e. subclass lacks only finitely many members of \mathcal{C} . Let \mathcal{C}^* be the intersection of the infinite r.e. subclasses, then $\mathcal{C} - \mathcal{C}^*$ is finite, with members A_1, \dots, A_n , say ($n \geq 1$). Now the infinite r.e. subclasses of \mathcal{C} have the form

$$\mathcal{C} - \{A_i | i \in F_j\} \quad (1 \leq j \leq m+1)$$

where the F_j are subsets of $\{x | 1 \leq x \leq n\}$. These are closed to subsets, for if $F \subseteq F_j$

Received by the editors June 13, 1966.

$$\mathfrak{C} - \{A_i \mid i \in F\} = (\mathfrak{C} - \{A_i \mid i \in F_j\}) \cup \{A_i \mid i \in F_j - F\}$$

and the union of two r.e. classes is an r.e. class. Also, by definition of A_1, \dots, A_n ,

$$\cup\{F_i \mid 1 \leq i \leq m + 1\} = \{x \mid 1 \leq x \leq n\}.$$

This completes the proof of the converse.

Note that in (a) $m + 1 \leq 2^n$. We give a construction for the case $m + 1 < 2^n$ and obtain as corollaries the case $m + 1 = 2^n$ and (b), both of which Young has already proved.

Since $m + 1 < 2^n$ there are subsets G_1, \dots, G_l say of $\{x \mid 1 \leq x \leq n\}$ different from all the F_i . For each i, k ($1 \leq i \leq l, 1 \leq k \leq m + 1$) there is a number

$$p(i, k) \in G_i - F_k,$$

since the F_i are closed to subsets. For each k with $1 \leq k \leq m + 1$ there will be a different variation of the construction. We will show that these variations in the construction all give rise to the same $\mathfrak{C}^*, A_1, \dots, A_n$.

At this point we make some informal remarks. \mathfrak{C} will be enumerated in an r.e. sequence

$$\langle A_1, \dots, A_n, V_0, V_1, \dots \rangle.$$

To begin with the sets in the sequence are all disjoint. By "amalgamating" sets we force the following situation: there will be an increasing function $r(i)$ such that the V_i different from each of A_1, \dots, A_n are $V_{r(0)}, V_{r(1)}, \dots$. These will be disjoint from each other and from each of A_1, \dots, A_n . Also if an r.e. set W intersects infinitely many of $V_{r(0)}, V_{r(1)}, \dots$ it intersects them all and

$$1 \leq i \leq l \Rightarrow (Ex)[x \in W \ \& \ (z)(x \in A_z \Leftrightarrow z \in G_i)].$$

Now if we take W to be $\cup \mathfrak{C}_1$ where \mathfrak{C}_1 is an infinite r.e. subclass we get for all i $V_{r(i)} \in \mathfrak{C}_1$ and for all i with $1 \leq i \leq l$ there is $z \in G_i$ with $A_z \in \mathfrak{C}_1$. Thus the only possibilities for \mathfrak{C}_1 are

$$\mathfrak{C} - \{A_z \mid z \in F_k\} \quad (1 \leq k \leq m + 1).$$

$A_z, r(x)$ and $V_{r(x)}$ will be independent of the variation used to get the sequence

$$\langle A_1, \dots, A_n, V_0, V_1, \dots \rangle,$$

but in variation k use of the function p will ensure for no y, z do we have

$$V_y = A_z \text{ with } z \in F_k,$$

and thus an enumeration of $\mathcal{C} - \{A_z | z \in F_k\}$ is easily obtained. The number $r(i)$ is approximated to by a sequence

$$r(i, 0), r(i, 1), \dots$$

Because we care only about r.e. sets W which intersect $V_{r(j)}$ for infinitely many j we are able to allow W to make $r(i, s+1) \neq r(i, s)$ only if $i > e$ (the index of W) and thus to make the sequence constant eventually. Now we proceed with the details.

Variation k. Let $\langle W_e | e \geq 0 \rangle$ be an r.e. sequence enumerating all the r.e. sets. Let $P = \{ (d, e) | d \in W_e \}$. The s th pair will mean the s th member of P to appear in an effective enumeration of P without repetitions. We define:

$$W_e^s = \{ d | (d, e) \text{ is among the first } s \text{ pairs} \}.$$

We define for each $s \geq 0$, by induction on s , sets A_1^s, \dots, A_n^s and a sequence of sets $\langle V_0^s, V_1^s, \dots \rangle$. Let

$$A_x^0 = \{ x - 1 \} \quad (1 \leq x \leq n), \quad V_x^0 = \{ n + x \} \quad (x \geq 0).$$

Assume for induction

- (1) the V_x^s, A_x^s are all finite;
- (2) the A_x^s are all different, in fact $x - 1 \in A_x^s - \cup \{ A_y^s | y \neq x \}$;
- (3) only finitely many of the V_x^s can be the same;
- (4) if V_x^s is different from all the A_j^s then it is disjoint from them all;
- (5) of the V_x^s which differ from all the A_y^s , different ones are disjoint;
- (6) V_0^s is different from all the A_x^s , properties evidently possessed by the A_x^0, V_x^0 .

We can define a function $r(i, s)$ by

$$\begin{aligned} r(0, s) &= 0, \\ r(i + 1, s) &= \mu x [x > r(i, s) \ \& \ V_x^s \neq A_j^s \text{ for } 1 \leq j \leq n \\ &\quad \& \ V_x^s \neq V_{r(i,s)}^s \text{ for } 0 \leq j \leq i]. \end{aligned}$$

Suppose the $(s+1)$ th pair is (d, e) and there is a $j > e$ such that $d \in V_{r(j,s)}^s$. There is at most one such j by (5). Define

$$R(e, s, i, x) = x \in W_e^{s+1} \ \& \ (z) (x \in A_z^s \Leftrightarrow z \in G_i).$$

Case 1. There is i with $1 \leq i \leq l$ and $\sim (Ex)R(e, s, i, x)$. Let i be the least such number. Put

$$\begin{aligned}
 A_x^{s+1} &= A_x^s \cup V_{r(j,s)}^s && \text{if } x \in G_i, \\
 A_x^{s+1} &= A_x^s && \text{if } x \notin G_i, \\
 V_x^{s+1} &= A_y^{s+1} && \text{if } V_x^s = A_y^s \text{ for some } y \text{ (there can be only} \\
 &&& \text{one such } y \text{ by (2)),} \\
 V_x^{s+1} &= A_{p(i,k)}^s \cup V_{r(j,s)}^s && \text{if } V_x^s = V_{r(j,s)}^s, \\
 V_x^{s+1} &= V_x^s && \text{otherwise.}
 \end{aligned}$$

This is where variation k arises.

Case 2. Case 1 does not obtain but there is i with $i < j$ and $W_e^{s+1} \cap V_{r(i,s)}^s = \emptyset$. Let i be the least such number. Put

$$\begin{aligned}
 A_x^{s+1} &= A_x^s && \text{for all } x, \\
 V_x^{s+1} &= V_{r(i,s)}^s \cup V_{r(j,s)}^s && \text{if } V_x^s = V_{r(i,s)}^s \text{ or } V_{r(j,s)}^s, \\
 V_x^{s+1} &= V_x^s && \text{otherwise.}
 \end{aligned}$$

If there is no $j > e$ such that $d \in V_{r(j,s)}^s$ or if neither Case 1 nor Case 2 occurs, put

$$A_x^{s+1} = A_x^s \text{ for all } x; \quad V_x^{s+1} = V_x^s \text{ for all } x.$$

We show that (1), \dots , (6) are preserved. We have $A_x^s \subseteq A_x^{s+1}$, $V_x^s \subseteq V_x^{s+1}$ (so that the A_x^s, V_x^s are all nonempty), $V_x^s = V_y^s \Rightarrow V_x^{s+1} = V_y^{s+1}$, $V_x^s = A_y^s \Rightarrow V_x^{s+1} = A_y^{s+1}$ for all x, y . (1) and (3) are clear.

For (2) we need consider only Case 1, where the result follows by induction hypothesis (4) and the definition of r .

For (4) and (5), consider V_x^{s+1}, V_y^{s+1} different from each other and from all the A_z^{s+1} . Then V_x^s, V_y^s are different from each other and from all the A_z^s , so by induction hypothesis (4), (5) they are disjoint from each other and from all the A_z^s . The desired conclusion is that V_x^{s+1}, V_y^{s+1} are disjoint from each other and from all the A_z^{s+1} . If Case 1 occurs, then neither V_x^s nor V_y^s is equal to $V_{r(j,s)}^s$, for $V_x^s = V_{r(j,s)}^s$ implies that $V_x^{s+1} = A_{p(i,k)}^s \cup V_{r(j,s)}^s = A_{p(i,k)}^{s+1}$. Thus $V_x^{s+1} = V_x^s$ and $V_y^{s+1} = V_y^s$. The two are therefore disjoint. Also V_x^s is disjoint from $V_{r(j,s)}^s$ and from A_z^s for all z , and so V_x^{s+1} is disjoint from A_z^{s+1} for all z . If Case 2 occurs at least one of V_x^s, V_y^s is different from both $V_{r(i,s)}^s$ and $V_{r(j,s)}^s$, or $V_x^{s+1} = V_y^{s+1}$. If both have this property then $V_x^{s+1} = V_x^s$ and $V_y^{s+1} = V_y^s$ and the result follows since $A_z^{s+1} = A_z^s$ for all z . This leaves the case where say $V_x^s = V_{r(i,s)}^s$ and V_y^s differs from both $V_{r(i,s)}^s$ and $V_{r(j,s)}^s$, then $V_y^{s+1} = V_y^s$ and the result follows by induction hypotheses (4), (5).

(6) follows by a similar argument, using the fact that in the construction $j > 0$.

Define:

$$A_x = \cup \{ A_x^s \mid s \geq 0 \} \quad (1 \leq x \leq n),$$

$$V_x = \cup \{ V_x^s \mid s \geq 0 \} \quad (x \geq 0).$$

We can find the members of A_x^s, V_x^s effectively from x, s so

$$\langle A_1, \dots, A_n, V_0, V_1, \dots \rangle$$

is an r.e. sequence enumerating an r.e. class \mathcal{C} . Let $\mathcal{C}^* = \mathcal{C} - \{A_1, \dots, A_n\}$.

For each x and all sufficiently large (s.l.) s ,

$$r(x, s) \text{ is a constant, say } r(x).$$

We prove this by induction on x . $r(0, s) = 0$ for all s , so $r(0) = 0$. We suppose the result holds for all $y \leq x$ and we show it holds for $x + 1$. There is an s_0 such that if $s \geq s_0$ $r(y, s) = r(y)$ for all $y \leq x$. In Case 1 or Case 2

$$r(z, s + 1) = r(z, s) \quad \text{if } z < j,$$

$$r(z, s + 1) = r(z + 1, s) \quad \text{if } z \geq j,$$

for if we divide the members of the sequence

$$V_0^s, V_1^s, \dots$$

which differ from each of A_1^s, \dots, A_n^s into equivalence classes under set equality, $V_{r(z,s)}^s$ is the first member of the $(z + 1)$ th such class, and by (4) and (5) the only effect of either case on the computation of r is the loss of the original $(j + 1)$ th class. So if $r(x + 1, s + 1) \neq r(x + 1, s)$ with $s \geq s_0$, Case 1 or Case 2 occurs with $x + 1 \geq j > e$ and by our induction hypothesis $x + 1 = j$. If Case 1 occurs we have $R(e, t, i, d)$ for all $t > s$ by (4), because d does not belong to $V_{r(j,t)}^s$ for any j . Now $r(x + 1, s + 1) \neq r(x + 1, s)$ can hold for only finitely many $s \geq s_0$ through Case 1—at most l times for each $e < x + 1$, and through Case 2—at most $x + 1$ times for each $e < x + 1$ by our induction hypothesis. Thus $r(x + 1, s) = r(x + 1)$, a constant for all s.l. s .

By (2) A_1^s, \dots, A_n^s are distinguished by the numbers $0, \dots, n - 1$ for all s , so A_1, \dots, A_n are.

We now wish to prove that the $V_{r(u)}$ are distinct, disjoint from each other and from all the A_u . Suppose $x \in V_{r(u)} \cap V_{r(v)}$ with $u \neq v$. Then for all s.l. s $x \in V_{r(u)}^s \cap V_{r(v)}^s$, so all s.l. $s, x \in V_{r(u,s)}^s \cap V_{r(v,s)}^s$, contradict-

ing (5). Similarly $x \in V_{r(u)} \cap A_v$ contradicts (4). The $V_{r(u)}$ are therefore distinct since they are nonempty.

Now we show that for all x , either $V_x = V_{r(u)}$ for some u or $V_x = A_{p(u,k)}$ for some u . It follows by induction on s that if V_x^s is equal to one of the A_y^s then it is equal to $A_{p(u,k)}^s$ for some u . Suppose that there is no u such that $V_x = V_{r(u)}$. There is then a u such that $r(u) < x < r(u+1)$. Consider s s.l. that $r(v, s) = r(v)$ for all $v \leq u+1$. Then $V_x^s = A_{p(z,k)}^s$ for some z , in which case $V_x = A_{p(z,k)}$, or $V_x^s = V_{r(v,s)}^s$ for some $v \leq u$, in which case $V_x = V_{r(v)}$.

By induction on s , $A_z^s, r(x, s)$ and $V_{r(z,s)}^s$ are all independent of k . Thus $A_z, r(x)$ and $V_{r(x)}$ are all independent of k .

Now we show that if W_e intersects infinitely many of $V_{r(0)}, V_{r(1)}, \dots$, then

$$1 \leq i \leq l \Rightarrow (Ex)R(e, i, x), \text{ where we define}$$

$$(7) \quad R(e, i, x) = x \in W_e \ \& \ (z)(x \in A_z \Leftrightarrow z \in G_i);$$

$$(8) \quad i \geq 0 \Rightarrow (Ex)[x \in W_e \cap V_{r(i)}].$$

First we have: if a, t are any given numbers there is $s > t$ and $j > a$ such that the $(s+1)$ th pair is (d, e) and $d \in V_{r(j,s)}^s$. For there are infinitely many $y > a$ such that W_e intersects $V_{r(y)}$. Also these $V_{r(y)}$ are disjoint. So for infinitely many members d of $W_e, d \in V_{r(y)}$ for some $y > a$. So there is $s > t$ such that (d, e) is the $(s+1)$ th pair and $d \in V_{r(y)}$ with $y > a$. Now $d \in V_{r(j,s)}^s$ for some j . Suppose $j < y$. Consider $u > s$ s.l. that $r(y, u) = r(y)$ and $d \in V_{r(y)}^u$. Then $d \in V_{r(y,u)}^u \cap V_{r(j,s)}^u$ so by (4) and (5) $V_{r(y,u)}^u = V_{r(j,s)}^u$, but $r(y, u) > r(j, u) \geq r(j, s)$ contradicting the definition of $r(y, u)$. Thus $j \geq y \geq a$.

Now suppose there is a least $i \leq l$ such that $\sim(Ex)R(e, i, x)$. For each $y < i$ let $d(y)$ be such that $R(e, y, d(y))$. Let t be s.l. that for each $y < i$ $R(e, t, y, d(y))$. Put $a = e$ and let s, j correspond to a, t as above. Since for each $y < i$ we have $R(e, s, y, d(y))$ and $\sim(Ex)R(e, s, i, x)$ Case 1 ensures that $R(e, s+1, i, d)$ and so $R(e, i, d)$, contradiction.

Suppose there is a least i such that $\sim(Ex)[x \in W_e \cap V_{r(i)}]$. Let t be s.l. that $r(i, t) = r(i)$, for each $y < i$ $(Ex)[x \in W_e^{t+1} \cap V_{r(y,t)}^s]$ and $r(y, t) = r(y)$ and for $1 \leq y \leq l$ $R(e, t, y, d(y))(d(y)$ as above). Put $a = \max(e, i)$ and let s, j correspond to a, t as above. Then Case 2 ensures that $d \in W_e^{s+2} \cap V_{r(t,s+1)}^{s+1}$ so $(Ex)[x \in W_e \cap V_{r(i)}]$, contradiction.

Recall that \mathfrak{C} has distinct members

$$A_1, \dots, A_n, V_{r(0)}, V_{r(1)}, \dots$$

and $\mathfrak{C}^* = \{V_{r(0)}, V_{r(1)}, \dots\}$. Suppose \mathfrak{C}_1 is an infinite r.e. subclass. Put $U\mathfrak{C}_1 = W_e$ (it is an r.e. set). W_e intersects infinitely many of the

$V_{r(i)}$ so we can apply (7) and (8). By (8), since the $V_{r(i)}$ are disjoint from each other and from all the A_i , $V_{r(i)} \in \mathcal{C}_1$ for all i . So the only possibilities for \mathcal{C}_1 are

$$\begin{aligned} \mathcal{C} - \{A_z \mid z \in G_i\} & \quad (1 \leq i \leq l) \\ \mathcal{C} - \{A_z \mid z \in F_k\} & \quad (1 \leq k \leq m + 1). \end{aligned}$$

We discount the first possibility. For by (7),

$$x \in W_e \ \& \ (z)(x \in A_z \Leftrightarrow z \in G_i) \quad \text{for some } x.$$

$x \in V_{r(j)}$ for any j so the only members of \mathcal{C} which x belongs to are the $\{A_z \mid z \in G_i\}$. So one of these sets must be in \mathcal{C}_1 .

We complete the proof by showing that

$$\mathcal{C} - \{A_z \mid z \in F_k\} \text{ is r.e.} \quad (1 \leq i \leq m + 1).$$

For consider the construction of \mathcal{C} by variation k , in an r.e. sequence

$$\langle A_1, \dots, A_n, V_0, V_1, \dots \rangle.$$

Since $V_x \neq$ any member of $\mathcal{C}^* \Rightarrow V_x = A_{p(i,k)}$ some i and $p(i, k) \notin F_k$, the r.e. sequence obtained by omitting the z th member of the original one for each $z \in F_k$ enumerates the desired class.

(a) when $m + 1 = 2^n$: Let $n' = n + 1$, $m' = 2^{n+1} - 2$ and the F'_k ($1 \leq k \leq 2^{n+1} - 1$) be all the subsets of $\{x \mid 1 \leq x \leq n + 1\}$ other than the whole set. Let $\mathcal{C}, \mathcal{C}^*$ be constructed for m', n', F'_k as above. Define

$$\mathcal{C}_1 = \mathcal{C}^* \cup \{A_{n+1}\}.$$

Then \mathcal{C}_1 has no proper infinite r.e. subclasses (and this proves (b)) and $\cup \mathcal{C}_1 \subseteq \{x \mid x \geq n\}$. Define

$$\mathcal{C}_2 = \mathcal{C}_1 \cup \{\{i\} \mid 0 \leq i \leq n - 1\}$$

and \mathcal{C}_2 is the required class. For given an infinite r.e. subclass \mathcal{C}_3 of \mathcal{C}_2 , $\mathcal{C}_3 - \{\{i\} \mid 0 \leq i \leq n - 1\}$ is an infinite r.e. subclass of \mathcal{C}_1 and therefore is \mathcal{C}_1 . On the other hand, any combination of the $\{i\}$ can be added to \mathcal{C}_1 .

REFERENCE

1. P. R. Young, *A theorem on recursively enumerable classes and splinters*, Proc. Amer. Math. Soc. 17 (1966), 1050-1056.

SIMON FRASER UNIVERSITY