

Review

Open Access

Inflammatory mechanisms in ischemic stroke: therapeutic approaches

Shaheen E Lakhan*, Annette Kirchgessner and Magdalena Hofer

Address: Global Neuroscience Initiative Foundation, Los Angeles, CA, USA

Email: Shaheen E Lakhan* - slakhan@gnif.org; Annette Kirchgessner - akirchgessner@gnif.org; Magdalena Hofer - lhofer@gnif.org

* Corresponding author

Published: 17 November 2009

Received: 3 August 2009

Journal of Translational Medicine 2009, **7**:97 doi:10.1186/1479-5876-7-97

Accepted: 17 November 2009

This article is available from: <http://www.translational-medicine.com/content/7/1/97>

© 2009 Lakhan et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Acute ischemic stroke is the third leading cause of death in industrialized countries and the most frequent cause of permanent disability in adults worldwide. Despite advances in the understanding of the pathophysiology of cerebral ischemia, therapeutic options remain limited. Only recombinant tissue-plasminogen activator (rt-PA) for thrombolysis is currently approved for use in the treatment of this devastating disease. However, its use is limited by its short therapeutic window (three hours), complications derived essentially from the risk of hemorrhage, and the potential damage from reperfusion/ischemic injury. Two important pathophysiological mechanisms involved during ischemic stroke are oxidative stress and inflammation. Brain tissue is not well equipped with antioxidant defenses, so reactive oxygen species and other free radicals/oxidants, released by inflammatory cells, threaten tissue viability in the vicinity of the ischemic core. This review will discuss the molecular aspects of oxidative stress and inflammation in ischemic stroke and potential therapeutic strategies that target neuroinflammation and the innate immune system. Currently, little is known about endogenous counterregulatory immune mechanisms. However, recent studies showing that regulatory T cells are major cerebroprotective immunomodulators after stroke suggest that targeting the endogenous adaptive immune response may offer novel promising neuroprotectant therapies.

Introduction

Stroke is the third leading cause of death in industrialized countries [1] and the most frequent cause of permanent disability in adults worldwide [2]. Three months following a stroke, 15-30% of stroke survivors are permanently disabled and 20% require institutional care [3]. Deficits can include partial paralysis, difficulties with memory, thinking, language, and movements. In the Western world, over 70% of individuals experiencing a stroke are over 65 years of age. Since life expectancy continues to grow, the absolute number of individuals with stroke will further increase in the future.

The most common cause of stroke is the sudden occlusion of a blood vessel by a thrombus or embolism, resulting in an almost immediate loss of oxygen and glucose to the cerebral tissue. Although different mechanisms are involved in the pathogenesis of stroke, increasing evidence shows that ischemic injury and inflammation account for its pathogenic progression [4]. Cerebral ischemia triggers the pathological pathways of the ischemic cascade and ultimately causes irreversible neuronal injury in the ischemic core within minutes of the onset [5].

However, a much larger volume of brain tissue surrounding this ischemic core, known as the penumbra, can be salvaged if cerebral blood flow is promptly restored. Thus, the original definition of the ischemic penumbra referred to areas of brain that were damaged but not yet dead, offering the promise that if proper therapies could be found, one could rescue brain tissue after stroke and reduce post-stroke disability.

Despite advances in the understanding of the pathophysiology of cerebral ischemia, therapeutic options for acute ischemic stroke remain very limited [2]. Only one drug is approved for clinical use for the thrombolytic treatment of acute ischemic stroke in the US and that is intravenous recombinant tissue plasminogen activator (rt-PA). When delivered within three hours after symptom onset, rt-PA reduces neurological deficits and improves the functional outcome of stroke patients. However, this improvement in recovery is achieved at the expense of an increased incidence in symptomatic intracranial hemorrhage, which occurs in ~6% of patients. Furthermore, since the large majority of patients with acute ischemic stroke do not go to the hospital within three hours of stroke onset most do not receive rt-PA treatment [6]. Consequently, the successful treatment of acute ischemic stroke remains one of the major challenges in clinical medicine.

This review will provide a brief overview of the current understanding of the inflammatory mechanisms involved in an acute ischemic stroke and the neuroprotective agents that can curtail neuroinflammation and potentially show utility in the treatment of stroke. Neuroprotective treatments are therapies that block the cellular, biochemical, and metabolic elaboration of injury during exposure to ischemia. Of the more than 100 neuroprotective agents that reached randomized clinical trials in focal ischemic stroke, none has proven unequivocally efficacious, despite success seen in preceding animal studies [7]. However, the failed trials of the past have greatly increased our understanding of the fundamental biology of ischemic brain injury and have laid a strong foundation for future advance. New anti-inflammatory targets continue to be identified, which is an important area for translational medicine in acute stroke. Overall, the prospects for safe neuroprotective therapies to improve stroke outcome remain promising [8]

Ischemic cascade

Acute ischemic stroke accounts for about 85% of all cases while hemorrhagic stroke is responsible for almost 15% of all strokes. Ischemic stroke results from the sudden decrease or loss of blood circulation to an area of the brain, resulting in a corresponding loss of neurological function. It is a nonspecific term encompassing a heterogeneous group of etiologies including thrombosis, embolism, and relative hypoperfusion. In most cases, the cause is atherothrombosis of large cervical or intracranial arteries, or embolism from the heart.

Within seconds to minutes after the loss of blood flow to a region of the brain, the ischemic cascade is rapidly initiated, which comprises a series of subsequent biochemical events that eventually lead to disintegration of cell membranes and neuronal death at the center/core of the infarction. Ischemic stroke begins with severe focal hypoperfusion, that leads to excitotoxicity and oxidative damage which in turn cause microvascular injury, blood-brain barrier dysfunction and initiate post-ischemic inflammation. These events all exacerbate the initial injury and can lead to permanent cerebral damage (see Figure 1). The amount of permanent damage depends on several factors: the degree and the duration of ischemia and the capability of the brain to recover and repair itself [5].

As a result of residual perfusion from the collateral blood vessels, regions where blood flow drops to approximately 30 ml/100 g/min ischemic cascade progresses at a slower rate. Neuronal cells may tolerate this level of reduced (20-40% of control values) blood flow for several hours from the stroke onset with full recovery of function following restoration of blood flow [9].

In the center of the ischemic region cells undergo anoxic depolarization and they never repolarize. While in the




Figure 1

Ischemic cascade leading to cerebral damage.

Ischemic stroke leads to hypoperfusion of a brain area that initiates a complex series of events. Excitotoxicity, oxidative stress, microvascular injury, blood-brain barrier dysfunction and postischemic inflammation lead ultimately to cell death of neurons, glia and endothelial cells. The degree and duration of ischemia determines the extent of cerebral damage.

penumbral region, the cells can repolarize at the expense of further energy consumption and depolarize again in response to elevated levels of extracellular glutamate and potassium ions. Such repetitive depolarizations called "peri-infarct depolarizations" lead to the increased release of the excitatory neurotransmitter glutamate with resulting excitotoxic cell damage [10]. Ultimately, the severity of functional and structural changes in the brain caused by ischemia will depend on its degree and duration.

Hyperbaric (HBO) and normobaric oxygen (NBO) therapies attempt to increase the partial pressure of oxygen to the tissue and thereby limit the damage caused by hypoperfusion. However, three clinical trials of hyperbaric oxygen therapy failed to show efficacy [11]. Normobaric, high-flow oxygen therapy was shown to cause a transient improvement of clinical deficits and MRI abnormalities in a sub-group of patients with acute ischemic stroke. Further studies are needed to investigate the safety and efficacy of hyperoxia as a stroke therapy [12].

Oxidative stress

Oxidative stress contributes to the pathogenesis of a number of neurological conditions including stroke. Oxidative stress is defined as the condition occurring when the physiological balance between oxidants and antioxidants is disrupted in favor of the former with potential damage for the organism. Oxidative stress leading to ischemic cell death involves the formation of ROS/reactive nitrogen species through multiple injury mechanisms, such as mitochondrial inhibition, Ca^{2+} overload, reperfusion injury, and inflammation [13]. Plenty of ROS are generated during an acute ischemic stroke and there is considerable evidence that oxidative stress is an important mediator of tissue injury in acute ischemic stroke [14]. Brain ischemia generates superoxide (O_2^-), which is the primary radical from which hydrogen peroxide is formed. Hydrogen peroxide is the source of hydroxyl radical (OH^\bullet). Nitric oxide is a water- and lipid-soluble free radical that is produced from L-arginine by three types of nitric oxide synthases (NOS). Ischemia causes an increase in NOS type I and III activity in neurons and vascular endothelium, respectively. At a later stage, elevated NOS type II (iNOS) activity occurs in a range of cells including glia and infiltrating neutrophils. Thus, free radicals are regarded as an important therapeutic target for improving the outcome of an ischemic stroke. Several compounds with significant antioxidant properties including ebselen [15], and resveratrol [16], a natural phytoalexin found in some dietary sources such as grapes and red wine, have been demonstrated to reduce stroke-related brain damage in animal models.

The transcription factor Nrf2

Nuclear factor erythroid-related factor 2 (Nrf2) is a transcription factor that regulates an expansive set of antioxi-

dant genes that act in synergy to remove ROS through sequential enzymatic reactions [17].

Nrf2 gene targets, collectively referred to as phase II genes, are involved in free radical scavenging, detoxification of xenobiotics, and maintenance of redox potential. Nrf2 is normally localized to the cytoplasm, tethered to the regulatory protein, kelch-like erythroid cell-derived protein with CNC homology associated protein 1 (Keap1) (Figure 2). Oxidative stress, or electrophilic agents that mimic oxidative stress, can modify key sulfhydryl group interactions in the Keap-Nrf2 complex, allowing dissociation and nuclear translocation of Nrf2. When activated, Nrf2 specifically targets genes bearing an antioxidant response element (ARE) within their promoters such as heme oxygenase 1, 1-ferritin, and glutathione peroxidase, which maintain redox homeostasis and influence the inflammatory response. Wide ranges of natural and synthetic small molecules are potent inducers of Nrf2 activity. These molecules have been identified from diverse chemical backgrounds including isothiocyanates, which are abundant in cruciferous vegetables, heavy metals, and hydroperoxides.




Figure 2
Nuclear erythroid-related factor 2 (Nrf2) anti-oxidant signaling in acute ischemic stroke. Nrf2 is the principal transcription factor that regulates antioxidant response element (ARE)-mediated expression of phase II detoxifying antioxidant enzymes. Under normal conditions, Nrf2 is sequestered in the cytoplasm by an actin-binding (Kelch-like) protein (Keap1); on exposure of cells to oxidative stress, Nrf2 dissociates from Keap1, translocates into the nucleus, binds to ARE, and transactivates phase II detoxifying and antioxidant genes. Among the spectrum of antioxidant genes controlled by Nrf2 are catalase, superoxide dismutase (SOD), glutathione reductase, and glutathione peroxidase.

Several studies have shown that increasing Nrf2 activity is highly neuroprotective in *in vitro* models that stimulate components of stroke damage, such as oxidative glutamate toxicity, H₂O₂ exposure, and Ca²⁺ overload [18]. Administration of the well characterized Nrf2 inducer, *tert*-butylhydroquinone (tBHQ), a metabolite of the widely used food antioxidant butylated hydroxyanisole, significantly improved sensorimotor and histological outcome in two models of I/R in rats and mice [19]. Within this injury paradigm, Nrf2 activation before stroke was able to salvage the cortical penumbra but not the stroke core. Clear differences in stroke outcome were found as early as 24 hours after reperfusion. Moreover, prophylactic treatment improved functional recovery up to one month after transient MCAO suggesting that previous Nrf2 activation may reduce neuronal cell death during delayed apoptosis and inflammation long after stroke onset.

Conversely, Nrf2-deficient mice are significantly more prone to ischemic brain injury and neurological deficits than WT mice. Deletion of the Nrf2 gene renders animals more susceptible to various stressors mainly because of the failure to induce phase II enzymes. Furthermore, an Nrf2 inducer was able to reverse neuronal cell death induced by the free radical donor *tert*-butylhydroperoxide (t-BuOOH) [19]. The MCAO and reperfusion model is known to induce a transient focal ischemic cascade that uniquely includes a substantial surge of free radical damage.

Ischemia/reperfusion (I/R) injury

The ischemic cascade usually goes on for hours but can last for days, even after restoration of blood circulation. Although reperfusion of ischemic brain tissue is critical for restoring normal function, it can paradoxically result in secondary damage, called ischemia/reperfusion (I/R) injury.

The definitive pathophysiology regarding I/R injury still remains obscure; however, oxidative stress mediators such as reactive oxygen species (ROS) released by inflammatory cells around the I/R injured areas are suggested to play a critical role [20]. The increase in oxygen free radicals triggers the expression of a number of pro-inflammatory genes by inducing the synthesis of transcription factors, including NF- κ B, hypoxia inducible factor 1, interferon regulator factor 1 and STAT3. As a result, cytokines are upregulated in the cerebral tissue and consequently, the expression of adhesion molecules on the endothelial cell surface is induced, including intercellular adhesion molecule 1 (ICAM-1), P-selectin and E-selectin which mediate adhesion of leukocytes to endothelia in the periphery of the infarct [21].

Furthermore, the complement cascade has been shown to play a critical role in I/R injury [22]. In addition to direct cell damage, regional brain I/R induces an inflammatory response involving complement activation and generation of active fragments such as C3a and C5a anaphylatoxins. Expression of C3a and complement 5a receptors was found to be significantly increased after middle cerebral artery occlusion (MCAO) in the mouse indicating an active role of the complement system in cerebral ischemic injury. Complement inhibition resulted in neuroprotection in animal models of stroke [23].

Post-ischemic inflammation

Although for many years the CNS was considered an immune-privileged organ, it is now well accepted that the immune and the nervous system are engaged in bi-directional crosstalk. Moreover, mounting data suggest that in the brain, as in peripheral organs, inflammatory cells participate in tissue remodeling after injury.

Microglial cells are the resident macrophages of the brain and play a critical role as resident immunocompetent and phagocytic cells in the CNS. Ekdahl and colleagues [24] reported an increased number of activated microglial cells up to 16 weeks after two hour MCAO in rats. After activation by ischemia, microglia can transform into phagocytes and they can release a variety of substances many of which are cytotoxic and/or cytoprotective. Microglia may exert neuroprotection by producing neurotrophic molecules such as brain-derived neurotrophic factor (BDNF), insulin-like growth factor I (IGF-I), and several other growth factors. There is substantial evidence that activated microglial cells in response to ischemia have the potential of releasing several pro-inflammatory cytokines such as TNF- α , IL-1 β , and IL-6, as well as other potential cytotoxic molecules including NO, ROS, and prostanoids [25].

Astrocytes, like microglia, are capable of secreting inflammatory factors such as cytokines, chemokines, and NO [26]. Cytokines upregulate the expression of cell adhesion molecules (CAMs). Within four to six hours after ischemia onset, circulating leukocytes adhere to vessel walls and migrate into the brain with subsequent release of additional pro-inflammatory mediators and secondary injury in the penumbra. Neutrophils are the earliest leukocyte subtype to show substantial upregulation in gene expression studies and to infiltrate areas of brain ischemia (see below). Recently, Shichita et al. [27] demonstrated an infiltration of γ dT cells 3 days after the onset of ischemia in a mouse model, along with a production of IL-17 which amplify the inflammatory cascade. IL-23 from infiltrating macrophages appear to produce IL-23 which attracts the infiltrating γ dT cells. Blocking a specific γ dT cell receptor with an antibody effectively reduced three-day infarct volumes, even when treatment was initiated at

24 hours after onset of cerebral ischemia. Targeting these γ dT cells may offer a clinical opportunity with a longer therapeutic window to prevent the secondary inflammatory expansion of cerebral damage after stroke.

The described post-ischemic neuroinflammatory changes lead to dysfunction of the blood-brain barrier, cerebral edema, and neuronal cell death (summarized in Figure 3). Therefore, therapeutic targeting of the neuroinflammatory pathways in acute ischemic stroke has become an important area of research in translational medicine.

Cytokines and brain inflammation

Cytokines are a group of small glycoproteins that are produced in response to an antigen and were originally described as mediators for regulating the innate and adaptive immune systems. Cytokines are thus upregulated in the brain in a variety of diseases, including stroke. In the brain, cytokines are expressed not only in the cells of the immune system, but are also produced by resident brain cells, including neurons and glia [28]. In addition, it has been shown that peripherally derived cytokines are involved in brain inflammation. Thus, peripherally derived mononuclear phagocytes, T lymphocytes, NK

cells and polymorphonuclear leukocytes produce and secrete cytokines and might contribute to inflammation of the CNS [29].

The most studied cytokines related to inflammation in acute ischemic stroke are tumor necrosis factor- α (TNF- α), the interleukins (IL), IL-1 β , IL-6, IL-20, IL-10 and transforming growth factor (TGF)- β . While IL-1 β and TNF- α , appear to exacerbate cerebral injury, TGF- β and IL-10 may be neuroprotective [30,31]. Increased production of pro-inflammatory cytokines and lower levels of the anti-inflammatory IL-10 are related to larger infarctions and poorer clinical outcome.

Elevated IL-1 β mRNA expression occurs within the first 15-30 min after permanent MCAO and elevated IL-1 β protein expression occurs a few hours later and remains elevated for up to 4 days [32]. There are studies that correlate an increase in the levels of IL-1 β after ischemia with worsening of the infarct severity. For example, Yamasaki et al [33] demonstrated that intraventricular injection of recombinant IL-1 β after MCAO increases the formation of brain edema, the volume of the size and the influx of neutrophils. In addition, IL-1 β deficient mice presented smaller infarcts in comparison with wild-type mice [34].

High circulating IL-1 β elevates circulating IL-6, another well known cytokine that is upregulated following cerebral ischemia [35]. Moreover, the serum level of IL-6 correlates with brain infarct volume [36] and is a powerful predictor of early neurological deterioration [37]. On the other hand, Clark et al [38] demonstrated that infarct size and neurological function were not different in animals deficient in IL-6 after transient CNS ischemia. This suggests that IL-6 does not have a direct influence on acute ischemic injury.

IL-20 is induced when IL-1 β modulates p38 MAPK and the NF- κ B pathway. IL-20 in turn induces the production of IL-6. Inhibition of IL-20 by a specific mAb significantly ameliorated the brain ischemic infarction in rats following MCAO [39].

Several approaches are under investigation for managing IL-1 in stroke (Table 1). IL-1 acts via membrane receptors (IL-1R), which can be blocked by a receptor antagonist (IL-1RA). In a randomized trial for acute stroke, IL-1RA readily crossed the blood-brain barrier, was safe to use, and seemed to afford some benefit, particularly for patients with cortical infarcts [40].

IL-10 is an anti-inflammatory cytokine that acts by inhibiting IL-1 and TNF- α , and by suppressing cytokine receptor expression and receptor activation as well. As a consequence, IL-10 could provide neuroprotection in




Figure 3
Postischemic inflammatory response. Excitotoxicity and oxidative stress caused by the initial ischemic event activate microglia and astrocytes which react by secreting cytokines, chemokines and matrix metalloproteases (MMP). These inflammatory mediators lead to an upregulation of cell adhesion molecules on endothelial cells, allowing blood-derived inflammatory cells, mainly neutrophils, to infiltrate the ischemic brain area. Neutrophils themselves also secrete cytokines which cause a further activation of glial cells. These processes all result in neuronal cell death and enhance the damage to the ischemic brain.

Table 1: Clinical studies of agents targeting inflammatory pathways in acute ischemic stroke.

Neuroprotective Agent	Mode of Action	Reference
Recombinant human IL-1RA	Interleukin-1 receptor antagonist	[67]
Enlimomab	Anti-ICAM-1 monoclonal antibody	[68]
Tirilazad	Lipid peroxidation inhibitor	[69]
UK-279, 276	Neutrophil inhibitory factor	[70]
Cerovive (NXY-059)	Nitrone-based free radical trapping agent	[71,72]
Acetaminophen (Paracetamol)	Anti-pyretic effect	[73]
Minocycline	Anti-inflammatory	[74]
Ginsenoside	Ca ²⁺ channel antagonist	[75]
Edaravone MCI-186	Free radical scavenger	[76]
ONO-2506 (Arundic Acid)	Astrocyte modulator	[77]

Adapted from Shah et al., 2009 [78].

acute ischemic stroke. Both central and systemic administration of IL-10 to rats subjected to MCAO significantly reduced infarct size 30 min to three hours post MCAO [30]. In acute ischemic stroke, elevated concentrations of IL-10 in CSF have been found [41]. Moreover, patients with low plasma levels (<6 pg/ml) of IL-10 during the first hours after stroke were three times more likely to have worsening neurological symptoms within 48 hours following the stroke [37]. IL-10 also seems to mediate the reduction in infarct size by regulatory T cells (see below).

Chemokines and brain inflammation

Chemokines, for example, monocyte chemoattractant protein 1, are a class of cytokines that guide the migration of blood borne inflammatory cells, such as neutrophils and macrophages, towards the source of the chemokine. Consequently, they play important roles in cellular communication and inflammatory cell recruitment. Expression of chemokines such as MCP-1, macrophage inflammatory protein-1 α (MIP-1 α), and fractalkine following focal ischemia is thought to have a deleterious effect by increasing leukocyte infiltration [42]. The level of a variety of chemokines has been found to increase in animal models of ischemia and their inhibition or deficiency has been associated with reduced injury [43-45]. Mice without the chemokine receptor CCR2 are protected against ischemia-reperfusion injury [46].

Cellular adhesion molecules

There is increasing evidence that cellular adhesion molecules (CAMs) play an important role in the pathophysiol-

ogy of acute ischemic stroke [21]. CAMs are upregulated in the first days after stroke by various cytokines and are responsible for the adhesion and migration of the leukocytes. Leukocytes roll on the endothelial surface and then adhere to the endothelial cells. The interaction between leukocytes and the vascular endothelium is mediated by three main groups of CAMs: the selectins, the immunoglobulin gene superfamily, and the integrins. Selectins, especially E- and P-selectins are upregulated and mediate leukocyte rolling and recruitment during the early stages of ischemia [47]. Among the immunoglobulin family member, intercellular adhesions molecule-1 (ICAM-1) and vascular cell adhesion molecule-1 have been the most extensively investigated in cerebral ischemia. Within hours after stroke onset, ICAM-1 expression increases upon stimulation by cytokines [48].

Patients with acute ischemic stroke had higher soluble intercellular adhesion molecule-1 (sICAM-1) levels compared to patients without cardiovascular disease. Moreover, sICAM-1 levels were significantly higher in patients who died compared to those who survived [49]. High sICAM-1 levels on admission are associated with early death in ischemic middle-aged stroke patients suggesting a pathogenic role of inflammation in the evolution of ischemic stroke.

A number of animal studies have shown that after transient and permanent focal ischemia the upregulation of CAMs, especially ICAM-1, P- and E-selectin, preceded the invasion of neutrophils into brain. There is ample evi-

dence from animal models of MCAO that expression of CAMs is associated with cerebral infarct size. Thus, genetic ablation of CAMs resulted in reduced infarct size, which could be mimicked by treatment with anti-CAM antibodies [50,51]. Inhibition of leukocyte activation and infiltration into the ischemic cerebral tissue has, therefore, been an important area of neuroprotection research. Thus far, anti-CAM treatment has not been successful in patients with acute ischemic stroke. However, further translational research into the therapeutic targeting of CAM is ongoing.

The spatiotemporal profile of CAMs is still largely unresolved, even though they are crucial for efficient anti-inflammatory therapies. More knowledge of the spatiotemporal profile of CAMs may lead the way to successful application and monitoring of promising anti-inflammatory treatment strategies after stroke.

Matrix metalloproteinases

MMPs are a family of proteolytic enzymes that are responsible for remodeling the extracellular matrix and that can degrade all its constituents. Expression of MMPs in the adult brain is very low to undetectable, but many MMPs are upregulated in the brain in response to injury [52]. Neurons, astrocytes, microglia, and endothelial cells have all been shown to express MMPs after injury. Stroke is associated with a biphasic disruption of the blood brain barrier (BBB) leading to vasogenic edema and hemorrhage and experimental studies have shown that that BBB breakdown and hemorrhage results from the expression and activation of MMPs [53].

MMP-2 and MMP-9 have been implicated in cerebral ischemia. Elevated MMP-9 levels were found in brain tissue and in serum from patients with acute ischemic stroke and in animal models of stroke beginning at 12 h after permanent MCAO [54]. MMP-9 is normally absent and this is the major MMP associated with neuroinflammation. Early (day 1) MMP-9 inhibition reduced infarction of day 14. However, benefit was lost when the treatment was delayed until day 3 and stroke pathology was exacerbated when administration was delayed until day 7 [55]. These studies all suggest that MMP inhibition could have a beneficial effect on the outcome of stroke but the effect will depend on the timing of treatment in relation to the stage of brain injury [55].

Regulatory T lymphocytes

Severe brain ischemia also perturbs innate and adaptive immune cells, resulting in systemic immunodepression that predisposes patients after stroke to life-threatening infections [56]. Postischemic alterations in the immune system might represent a useful immunomodulatory adaptation, preventing autoimmune reactions against CNS antigens after stroke.


Recently, regulatory T lymphocytes (T_{reg}) were shown to play an important role in protecting cells in a mouse model for stroke [57]. Thymus-derived $CD4^+CD25^+Foxp3^+$ T_{reg} cells play a key part in controlling immune responses under physiological conditions and in various systemic and CNS inflammatory diseases [58]. T_{reg} are generated by dendritic or antigen-presenting cells expressing the immunosuppressive mediator indoleamine 2,3-dioxygenase, the first enzyme in the kynurenine pathway, that degrades and converts tryptophan to kynurenine [59]. Interferon- γ and TNF- α which are both present at high levels in the ischemic brain induce IDO in response to chronic immune activation, possibly in microglia [60].

A stroke in mice with no functioning T_{reg} cells in their blood caused much greater damage to the brain and greater disabilities than in animals with functioning T_{reg} cells. T_{reg} cells protect cells by suppressing the harmful activation of the immune system and can thus also prevent autoimmune diseases from developing. IL-10 is a cytokine that is produced by the T_{reg} cells and seems to play an important role during a stroke. Mice with no functioning T_{reg} cells that were injected with IL-10 on the first day following a stroke had markedly less brain damage than mice that did not receive IL-10. On the other hand, the transfer of genetically modified T_{reg} cells unable to produce IL-10 offered no protection [57]. T_{reg} cells producing IL-10 induce IDO suggesting that IL-10 may act upstream by modulating the production of IDO.

Depletion of T_{reg} cells profoundly increased delayed brain damage and deteriorated functional outcome. Absence of T_{reg} cells augmented postischemic activation of resident and invading inflammatory cells including microglia and T cells, the main sources of cerebral TNF- α and IFN- γ , respectively. T_{reg} cells prevent secondary infarct growth by counteracting excessive production of proinflammatory cytokines and by modulating invasion and/or activation of lymphocytes and microglia in the ischemic brain. Liesz et al [57] found that T_{reg} cells antagonize enhanced TNF- α and IFN- γ production, which induce delayed inflammatory brain damage, and that T_{reg} cell-derived secretion of IL-10 is the key mediator of the cerebroprotective effect via suppression of proinflammatory cytokine production. IL-10 potently reduced infarct size in normal mice and prevented delayed lesion growth after T_{reg} cells depletion (Figure 4).

Post-stroke recovery

Patients experiencing a typical large-vessel acute ischemic stroke will lose 120 million neurons each hour. Compared with the normal rate of neuron loss during aging, the ischemic brain will age 3.6 years for every hour the stroke goes untreated. Thus, it is not surprising that the majority of stroke patients exhibit certain levels of motor

**Figure 4**

Regulatory T (T_{reg}) cells protect the brain after stroke. Experiments by Liesz et al. [57] show that T_{reg} cells prevent delayed lesion expansion in an IL-10-dependent manner in a mouse model of acute ischemic stroke. They also reduce the proinflammatory cytokine levels during the early postischemic inflammatory phase. Injection of IL-10 in the brain reduces infarct volume. Reprinted by permission from Macmillan Publishers Ltd: Nature Medicine 15, 138-139 Copyright 2009.

weakness and sensory disturbances [2]. However, over time, most will show a certain degree of functional recovery which may be explained by brain reorganization and brain plasticity.

Brain plasticity refers to the brain's ability to change its structure and function during development, learning, and pathology. For example, within the minutes following ischemia, rapid changes are observed in the number and length of dendritic spines of the neurons in the penumbra region. The initial loss is then followed by the re-establishment of the dendritic spine synapses several months after the initial stroke as part of the functional recovery process [61].

Functional MRI studies have demonstrated that the damaged adult brain is able to reorganize to compensate for motor deficits [62,63]. The main mechanism underlying recovery of motor abilities appears to involve enhanced activity in preexisting networks. Studies in experimental stroke models demonstrate that focal cerebral ischemia promotes neurogenesis in the subventricular zone (SVZ) and subgranular zone (SGZ) of the dentate gyrus and induces SVZ neuroblast migration towards the ischemic boundary. More importantly, stroke-induced neurogenesis has also recently been demonstrated in the adult human brain, even in advanced age patients [64-66]. These findings have led to a hope for a neurorestorative treatment of stroke which aims to manipulate endogenous neurogenesis and thereby enhance brain repair.

Conclusion

In conclusion, in the presented work, we sought to provide a brief overview of the current understanding of

inflammatory mechanisms involved during acute ischemic stroke and neuroprotective agents that can curtail neuroinflammation and could have utility in the treatment of stroke (see Table 1). As discussed, evidence suggests that post-ischemic oxidative stress and inflammation contribute to brain injury and to the expansion of the ischemic lesion. On the other hand, an adequate adaptive immune response after acute brain ischemia also plays an important role in response to ischemic injury as shown by the tremendous potential of T_{reg} cells to prevent secondary infarct growth by counteracting the production of proinflammatory cytokines and by modulating the activation of lymphocytes and microglia in the ischemic brain [57]. These results provide new insights into the immunopathogenesis of acute ischemic stroke and could lead to new approaches that involve immune modulation using T_{reg} cells.

To date, 1,026 drugs have been tested in various animal models, of which 114 underwent clinical evaluation [8]. The greater part of the agents studied until now have failed. Consequently, rt-PA remains the only agent shown to improve stroke outcome in clinical trials, despite the many clinical trials conducted. However, its use is limited by its short therapeutic window (three hours), by its complications derived essentially from the risk of hemorrhage, and by the potential damage by R/I injury. Because of these drawbacks the optimum treatment of cerebral focal ischemia remains one of the major challenges in clinical medicine.

Abbreviations

ARE: Antioxidant response element; BDNF: brain-derived neurotrophic factor; CAM: cell adhesion molecule; IGF-I:

insulin-like growth factor I; IL: interleukin; IL-1R: interleukin-1 membrane receptor; IL-1RA: interleukin-1 receptor antagonist; Keap1: kelch-like erythroid cell-derived protein with CNC homology associated protein 1; MMP: matrix metalloproteinase; iNOS: nitric oxide synthase type II; ICAM-1: intracellular adhesion molecule 1; MCAO: middle cerebral artery occlusion; MCP-1: monocyte chemoattractant protein-1; NOS: nitric oxide synthase; Nrf2: nuclear factor erythroid-related factor 2; ROS: reactive oxygen species; rt-PA: recombinant tissue plasminogen activator; T_{reg}: regulatory T lymphocytes; sICAM-1: soluble intracellular adhesion molecule 1; SOD: superoxide dismutase; t-BuOOH: tert-butylhydroperoxide; tBHQ: tert-butylhydroquinone; TGF: transforming growth factor; TNF- α : tumor necrosis factor- α .

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

All authors participated in the preparation of the manuscript, and read and approved the final manuscript.

Acknowledgements

The authors wish to extend special thanks to GNIF research associates Elissa Hamlat, Julie Aeschliman, and Lorraine Webster for their suggestions and editing support.

References

- Lo EH, Dalkara T, Moskowitz MA: **Mechanisms, challenges and opportunities in stroke.** *Nat Rev Neurosci* 2003, **4**:399-415.
- Donnan GA, Fisher M, Macleod M, Davis SM: **Stroke.** *Lancet* 2008, **371**:1612-1623.
- American Heart Association: *Heart Disease and Stroke Statistics - 2005 Update* Dallas, Texas: American Heart Association; 2005.
- Muir KW, Tyrrell P, Sattar N, Warburton E: **Inflammation and ischaemic stroke.** *Curr Opin Neurol* 2007, **20**:334-342.
- Dirnagl U, Iadecola C, Moskowitz MA: **Pathobiology of ischaemic stroke: an integrated view.** *Trends Neurosci* 1999, **22**:391-397.
- Furlan AJ, Katzan IL, Caplan LR: **Thrombolytic therapy in acute ischemic stroke.** *Curr Treat Options Cardiovasc Med* 2003, **5**:171-180.
- O'Collins VE, Macleod MR, Donnan GA, Horky LL, Worp BH van der, Howells DW: **1,026 experimental treatments in acute stroke.** *Ann Neurol* 2006, **59**:467-477.
- De la Ossa NP, Davalos A: **Neuroprotection in cerebral infarction: the opportunity of new studies.** *Cerebrovasc Dis* 2007, **24**:153-156.
- Hossmann K: **Thresholds of ischemic injury.** In *Cerebrovascular Disease: Pathophysiology Diagnosis and Management* Edited by: Ginsberg MD, Bogousslavsky J. Malden, UK: Blackwell Science; 1988:193-204.
- Hossmann KA: **Periinfarct depolarizations.** *Cerebrovasc Brain Metab Rev* 1996, **8**:195-208.
- Singhal AB: **Oxygen therapy in stroke: past, present, and future.** *Int J Stroke* 2006, **1**:191-200.
- Singhal AB, Benner T, Roccatagliata L, Koroshetz WJ, Schaefer PW, Lo EH, Buonanno FS, Gonzalez RG, Sorensen AG: **A Pilot Study of Normobaric Oxygen Therapy in Acute Ischemic Stroke.** *Stroke* 2005, **36**:797-802.
- Coyle JT, Puttfarcken P: **Oxidative stress, glutamate, and neurodegenerative disorders.** *Science* 1993, **262**:689-695.
- Cuzzocrea S, Riley DP, Caputi AP, Salvemini D: **Antioxidant therapy: a new pharmacological approach in shock, inflammation, and ischemia/reperfusion injury.** *Pharmacol Rev* 2001, **53**:135-159.
- Yamagata K, Ichinose S, Miyashita A, Tagami M: **Protective effects of ebselen, a seleno-organic antioxidant on neurodegeneration induced by hypoxia and reperfusion in stroke-prone spontaneously hypertensive rat.** *Neuroscience* 2008, **153**:428-435.
- Ozkan OV, Yuzbasioglu MF, Ciralik H, Kurutas EB, Yonden Z, Aydin M, Bulbuloglu E, Semerci E, Goksu M, Atli Y, Bakan V, Duran N: **Resveratrol, a natural antioxidant, attenuates intestinal ischemia/reperfusion injury in rats.** *Tohoku J Exp Med* 2009, **218**:251-258.
- Nguyen T, Nioi P, Pickett CB: **The nrf2-antioxidant response element signaling pathway and its activation by oxidative stress.** *J Biol Chem* 2009, **284**:13291-13295.
- Johnson JA, Johnson DA, Kraft AD, Calkins MJ, Jakel RJ, et al.: **The Nrf2-ARE pathway: an indicator and modulator of oxidative stress in neurodegeneration.** *Ann N Y Acad Sci* 2008, **1147**:61-69.
- Shih AY, Li P, Murphy TH: **A small molecule inducible Nrf2-mediated antioxidant response provides effective prophylaxis against cerebral ischemia in vivo.** *J Neurosci* 2005, **25**:10321-10335.
- Wong Ch, Crack PJ: **Modulation of neuro-inflammation and vascular response by oxidative stress following cerebral ischemia-reperfusion injury.** *Curr Med Chem* 2008, **15**:1-14.
- Yilmaz G, Granger DN: **Cell adhesion molecules and ischemic stroke.** *Neurol Res* 2008, **30**:783-93.
- D'Ambrosio AL, Pinsky DJ, Connolly ES: **The role of the complement cascade in ischemia/reperfusion injury: implications for neuroprotection.** *Mol Med* 2001, **7**:367-382.
- Arumugam TV, Woodruff TM, Lathia JD, Selvaraj PK, Mattson MP, Taylor SM: **Neuroprotection in stroke by complement inhibition and immunoglobulin therapy.** *Neuroscience* 2009, **158**:1074-1089.
- Ekdahl CT, Kokaia Z, Lindvall O: **Brain inflammation and adult neurogenesis: the dual role of microglia.** *Neuroscience* 2009, **158**:1021-1029.
- Lucas SM, Rothwell NJ, Gibson RM: **The role of inflammation in CNS injury and disease.** *Br J Pharmacol* 2006, **147**(Suppl 1):S232-40.
- Swanson RA, Ying W, Kauppinen TM: **Astrocyte influences on ischemic neuronal death.** *Curr Mol Med* 2004, **4**:193-205.
- Shichita T, Sugiyama Y, Ooboshi H, Sugimori H, Nakagawa R, Takada I, Iwaki T, Okada Y, Iida M, Cua DJ, Iwakura Y, Yoshimura A: **Pivotal role of cerebral interleukin-17-producing $\gamma\delta$ T cells in the delayed phase of ischemic brain injury.** *Nat Med* 2009, **15**:946-950.
- Barone FC, Feuerstein GZ: **Inflammatory mediators and stroke: new opportunities for novel therapeutics.** *J Cereb Blood Flow Metab* 1999, **19**:819-34.
- Ferrarese C, Mascarucci P, Zoia C, Cavarretta R, Frigo M, Begni B, Sarinella F, Frattola L, De Simoni MG: **Increased cytokine release from peripheral blood cells after acute stroke.** *J Cereb Blood Flow Metab* 1999, **19**:1004-9.
- Zhu Y, Yang GY, Ahlemeyer B, Pang L, Che XM, Culmsee C, Klumpp S, Kriegstein J: **Transforming growth factor-beta 1 increases bad phosphorylation and protects neurons against damage.** *J Neurosci* 2002, **22**:3898-909.
- Spera PA, Ellison JA, Feuerstein GZ, Barone FC: **IL-10 reduces rat brain injury following focal stroke.** *Neurosci Lett* 1998, **251**:189-92.
- Caso JR, Moro MA, Lorenzo P, Lizasoain I, Leza JC: **Involvement of IL-1 β in acute stress induced worsening of cerebral ischemia in rats.** *Eur Neuropsychopharmacol* 2007, **17**:600-607.
- Yamasaki Y, Matsuura N, Shozuhara H, Onodera H, Itoyama Y, Kogure K: **Interleukin-1 as a pathogenetic mediator of ischemic brain damage in rats.** *Stroke* 1995, **26**:676-680.
- Boutin H, LeFeuvre RA, Horai R, Asano M, Iwakura Y, Rothwell NJ: **Role of IL-1 α and IL-1 β in ischemic brain damage.** *J Neurosci* 2001, **21**:5528-5534.
- Clark WM, Rinker LG, Lessov NS, Hazel K, Eckenstein F: **Time course of IL-6 expression in experimental CNS ischemia.** *Neurol Res* 1999, **21**:287-292.
- Acalovschi D, Wiest T, Hartmann M, Farahmi M, Mansmann U, Aufarth GU, Grau AJ, Green FR, Grond-Ginsbach C, Schwanninger M: **Multiple levels of regulation of the interleukin-6 system in stroke.** *Stroke* 2003, **34**:1864-1870.

37. Vila N, Castillo J, Davalos A, Chamorro A: **Proinflammatory cytokines and early neurological worsening in ischemic stroke.** *Stroke* 2001, **31**:2325-2329.
38. Clark WM, Rinker LG, Lessov NS, Hazel K, Hill JK, Stenzel-Poore M, Eckenstein F: **Lack of interleukin-6 expression is not protective against focal central system ischemia.** *Stroke* 2001, **31**:1715-1720.
39. Chen WY, Chang MS: **IL-20 is regulated by hypoxia-inducible factor and up-regulated after experimental ischemic stroke.** *J Immunol* 2009, **182**:5000-5012.
40. Gueorguieva I, Clark SR, McMahon CJ, Scarth S, Rothwell NJ, Tyrrell PJ, Hopkins SJ, Rowland M: **Pharmacokinetic modeling of interleukin-1 receptor antagonist in plasma and cerebrospinal fluid of patients following subarachnoid hemorrhage.** *Br J Clin Pharmacol* 2008, **65**:317-325.
41. Tarkowski E, Rosengren L, Blomstrand C, Wikkelso C, Jensen C, Ekholm S, Tarkowski A: **Intrathecal release of pro- and anti-inflammatory cytokines during stroke.** *Clin Exp Immunol* 1997, **110**:492-499.
42. Kim JS, Gautam SC, Chopp M, Zaloga C, Jones ML, Ward PA, Welch KMA: **Expression of monocyte chemoattractant protein-1 and macrophage inflammatory protein-1 after focal cerebral ischemia in the rat.** *J Neuroimmunol* 1995, **56**:127-134.
43. Soriano S, Amaravadi L, Wang Y, Zhou H, Yu G, Tonra J, Fairchild-Huntress V, Fang Q, Dunmore J, Huszar D: **Mice deficient in fractalkine are less susceptible to cerebral ischemia-reperfusion injury.** *J Neuroimmunol* 2002, **125**:59-65.
44. Hughes PM, Allegrini PR, Rudin M, Perry VH, Mir AK, Wiessner C: **Monocyte chemoattractant protein-1 deficiency is protective in a murine stroke model.** *J Cereb Blood Flow Metab* 2002, **22**:308-17.
45. Kumai Y, Ooboshi H, Takada J, Kamouchi M, Kitazono T, Egashira K, Ibayashi S, Iida M: **Anti-monocyte chemoattractant protein-1 gene therapy protects against focal brain ischemia in hypertensive rats.** *J Cereb Blood Flow Metab* 2004, **24**:1359-1368.
46. Dimitrijevic OB, Stamatovic SM, Keep RF, Andjelkovic AV: **Absence of the chemokine receptor CCR2 protects against cerebral ischemia/reperfusion injury in mice.** *Stroke* 2007, **38**:1345-1353.
47. Zhang R, Chopp M, Zhang Z, Jiang N, Powers C: **The expression of P- and E-selectins in three models of middle cerebral artery occlusion.** *Brain Res* 1998, **785**:207-214.
48. Lindsberg PJ, Carpen O, Paetau A, Karjalainen-Lindsberg ML, Kaste M: **Endothelial ICAM-1 Expression Associated With Inflammatory Cell Response in Human Ischemic Stroke.** *Circulation* 1996, **94**:939-945.
49. Rallidis LS, Zolindaki MG, Vikelis M, Kaliva K, Papadopoulos C, Kremastinos DT: **Elevated soluble intercellular adhesion molecule-1 levels are associated with poor short-term prognosis in middle-aged patients with acute ischaemic stroke.** *Int J Cardiol* 2009, **132**:216-220.
50. Zhang RL, Chopp M, Li Y, et al.: **Anti-ICAM-1 antibody reduces ischemic cell damage after transient middle cerebral artery occlusion in the rat.** *Neurology* 1994, **44**:1747-1751.
51. Goussev AV, Zhang Z, Anderson DC, Chopp M: **P-selectin antibody reduces hemorrhage and infarct volume resulting from MCA occlusion in the rat.** *J Neurosci* 1998, **18**:16-22.
52. Montaner J, Alvarez-Sabin J, Molina C, Angles A, Abilleira S, Arenillas J, Gonzalez MA, Monasterio J: **Matrix metalloproteinase expression after human cardioembolic stroke: temporal profile and relation to neurological impairment.** *Stroke* 2001, **32**:1759-66.
53. Asahi M, Wang X, Mori T, Sumii T, Jung JC, Moskowitz MA, Fini ME, Lo EH: **Effects of matrix metalloproteinase-9 gene knock-out on the proteolysis of blood-brain barrier and white matter components after cerebral ischemia.** *J Neurosci* 2001, **21**:7724-7732.
54. Park KP, Rosell A, Foerch C, Xing C, Kim WJ, Lee S, Opdenakker G, Furie KL, Lo EH: **Plasma and brain matrix metalloproteinase-9 after acute focal cerebral ischemia in rats.** *Stroke* 2009, **40**:2836-2842.
55. Zhao BQ, Wang S, Kim HY, Storrie H, Rosen BR, Mooney DJ, Wang X, Lo EH: **Role of metalloproteinases in delayed cortical responses after stroke.** *Nat Med* 2006, **12**:441-445.
56. Meisel C, Schwab JM, Prass K, Meisel A, Dirnagl U: **Central nervous system injury-induced immune deficiency syndrome.** *Nat Rev Neurosci* 2005, **6**:775-786.
57. Liesz A, Suri-Payer E, Veltkamp C, Doerr H, Sommer C, Rivest S, Giese T, Veltkamp R: **Regulatory T cells are key cerebroprotective immunomodulators in acute experimental stroke.** *Nat Med* 2009, **15**:192-199.
58. O'Garra A, Vieira P: **Regulatory T cells and mechanisms of immune system control.** *Nat Med* 2004, **10**:801-805.
59. Sharma MD, Baban B, Chandler P, Hou DY, Singh N, Yagita H, Azuma M, Blazar BR, Mellor AL, Munn DH: **Plasmacytoid dendritic cells from mouse tumor-draining lymph nodes directly activate mature Tregs via indoleamine 2,3-dioxygenase.** *J Clin Invest* 2007, **117**:2570-2582.
60. O'Connor JC, Andre C, Wang Y, Lawson MA, Szegedi SS, Lestage J, Castanon N, Kelley KW, Dantzer R: **Interferon-gamma and tumor necrosis factor-alpha mediate the upregulation of indoleamine 2,3-dioxygenase and the induction of depressive-like behavior in mice in response to bacillus Calmette-Guerin.** *J Neurosci* 2009, **29**:4200-9.
61. Brown CE, Wong C, Murphy TH: **Rapid morphologic plasticity of peri-infarct dendritic spines after focal ischemic stroke.** *Stroke* 2008, **39**:1286-1291.
62. Calautti C, Baron J-C: **Functional Neuroimaging Studies of Motor Recovery After Stroke in Adults: A Review.** *Stroke* 2003, **34**:1553-1566.
63. Eliassen JC, Boespflug EL, Lamy M, Allendorfer J, Chu WJ, Szaflarski JP: **Brain-mapping techniques for evaluating poststroke recovery and rehabilitation: a review.** *Top Stroke Rehabil* 2008, **15**:427-450.
64. Macas J, Nern C, Plate KH, Momma S: **Increased generation of neuronal progenitors after ischemic injury in the aged adult human forebrain.** *J Neurosci* 2006, **26**:13114-13119.
65. Minger SL, Ekonomou A, Carta EM, Chinoy A, Perry RH, Ballard CG: **Endogenous neurogenesis in the human brain following cerebral infarction.** *Regen Med* 2007, **2**:69-74.
66. Yamashita T, Ninomiya M, Hernandez Acosta P, Garcia-Verdugo JM, Sunabori T, Sakaguchi M, Adachi K, Kojima T, Hirota Y, Kawase T, Araki N, Abe K, Okano H, Sawamoto K: **Subventricular zone-derived neuroblasts migrate and differentiate into mature neurons in the post-stroke adult striatum.** *J Neurosci* 2006, **26**:6627-6636.
67. Emsley HC, Smith CJ, Georgiou RF, Vail A, Hopkins SJ, Rothwell NJ, Tyrrell PJ: **Acute Stroke Investigators: A randomized phase II study of interleukin-1 receptor antagonist in acute stroke patients.** *J Neurol Neurosurg Psychiatry* 2005, **76**:1366-1372.
68. Enlimomab Acute Stroke Trial Investigators: **Use of anti-ICAM-1 therapy in ischemic stroke: results of the Enlimomab Acute Stroke Trial.** *Neurology* 2001, **57**:1428-1434.
69. Bath PM, Iddenden R, Bath FJ, Orgogozo JM, Tirilazad International Steering Committee: **Tirilazad for acute ischaemic stroke.** *The Cochrane Database Syst Rev* 2001, **4**:CD002087.
70. Krams M, Lees KR, Hacke W, Grieve AP, Orgogozo JM, Ford GA, ASTIN Study Investigators: **Acute stroke therapy by inhibition of neutrophils (ASTIN): an adaptive dose-response study of UK-279,276 in acute ischaemic stroke.** *Stroke* 2003, **34**:2543-2548.
71. Lyden PD, Shuaib A, Lees KR, Davalos A, Davis SM, Diener HC, Grotta JC, Ashwood TJ, Hardemark HG, Svensson HH, Rodichok L, Wasiewski WW, Ahlberg G, CHANT Trial Investigators: **Safety and tolerability of NXY-059 for acute intracerebral hemorrhage: the CHANT trial.** *Stroke* 2007, **38**:2262-2269.
72. Shuaib A, Lees KR, Lyden P, Grotta J, Davalos A, Davis SM, Diener HC, Ashwood T, Wasiewski WW, Emeribe U, SAINT II Trials Investigators: **NXY-059 for treatment of acute ischemic stroke.** *N Engl J Med* 2007, **357**:562-571.
73. van Breda EJ, Worp HB van der, van Gemert HM, Algra A, Kappelle LJ, van Gijn J, Koudstaal PJ, Dippel DW, PAIS investigators: **PAIS: paracetamol (acetaminophen) in stroke; protocol for a randomized, double blind clinical trial. [ISRCTN 74418480].** *BMC Cardiovasc Disord* 2005, **5**:24.
74. Lampl Y, Boaz M, Gilad R, Lorberboym M, Dabby R, Rapoport A, Anca-Herszkowitz M, Sadeh M: **Minocycline treatment in acute stroke: an open-label, evaluator-blinded study.** *Neurology* 2007, **69**:1404-10.
75. Liu X, Xia J, Wang L, Song Y, Yang J, Yan Y, Ren H, Zhao G: **Efficacy and safety of ginsenoside-Rd for acute ischaemic stroke: a randomized, double-blind, placebo-controlled, phase II multicenter trial.** *Eur J Neurol* 2009, **5**:569-75.

76. The Edaravone Acute Brain Infarction Study Group: **Effect of a novel free radical scavenger, Edaravone (MCI-186), on acute brain infarction. Randomized, placebo-controlled, double-blind study at multicenters.** *Cerebrovasc Dis* 2003, **15**:222-229.
77. Pettigrew LC, Kasner SE, Albers GW, Gorman M, Grotta JC, Sherman DG, Funakoshi Y, Ishibashi H, Arundic Acid (ONO-2506) Stroke Study Group: **Safety and tolerability of arundic acid in acute ischemic stroke.** *J Neurol Sci* 2006, **251**:50-6.
78. Shah IM, Macrae M, Di Napoli M: **Neuroinflammation and neuro-protective strategies in acute ischaemic stroke - from bench to bedside.** *Curr Mol Med* 2009, **9**:336-354.

Publish with **BioMed Central** and every scientist can read your work free of charge

"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:
http://www.biomedcentral.com/info/publishing_adv.asp

