

Influence of Aeration Speed on Bacterial Colony Forming Unit (CFU) Formation Capacity

Md. Sakil Munna, Sadika Tamanna, Most. Rumana Afrin, Gulshan Ara Sharif, Chaity Mazumder, Konica Sarker Kana, Nusrat Jahan Urmi, Md. Aftab Uddin, Tasmina Rahman, Rashed Noor*

Department of Microbiology, Stamford University Bangladesh, Dhaka, Bangladesh

*Corresponding author: noor.rashed@yahoo.com

Received December 30, 2013; Revised January 10, 2014; Accepted February 09, 2014

Abstract Bacterial homeostasis depends on an array of physical and chemical stimulants. Current investigation assessed the impact of one of such factors, the speed of aeration, on cell viability and culturability of *Escherichia coli*, *Pseudomonas* spp. and *Bacillus* spp. Each of the bacterial strain was incubated at 37°C with a shaking speed of 0, 100 or 200 rotation per minute (rpm) separately up to 72 hours, with a simultaneous monitoring of morphological changes and cell culturability. All bacterial species were found to optimally grow at 100 rpm whereas at 0 rpm growths of *E. coli* and *Pseudomonas* spp. were bit slower compared to that of *Bacillus* spp. The capacity to form colony forming units (CFUs) of *E. coli* and *Pseudomonas* spp. on Luria Burtani (LB) agar plates were observed to be inhibited after 36 hours of growth at 200 rpm; i.e., approximately 3-log reduced CFUs than those formed by *Bacillus* spp. Besides, morphologically impaired cells were observed for the former two bacteria cultivated at 200 rpm. Taken together, it is assumed that the high speed shaking might evolve the oxidative stress endogenously which possibly rendered the cells lose their culturability.

Keywords: *Escherichia coli*, *Pseudomonas* spp., *Bacillus* spp., aeration speed, colony forming units (CFUs), oxidative stress

Cite This Article: Md. Sakil Munna, Sadika Tamanna, Most. Rumana Afrin, Gulshan Ara Sharif, Chaity Mazumder, Konica Sarker Kana, Nusrat Jahan Urmi, Md. Aftab Uddin, Tasmina Rahman, and Rashed Noor, "Influence of Aeration Speed on Bacterial Colony Forming Unit (CFU) Formation Capacity." *American Journal of Microbiological Research*, vol. 2, no. 1 (2014): 47-51. doi: 10.12691/ajmr-2-1-7.

1. Introduction

Study of stress responses in bacteria is indeed an exhilarating vicinity of basic and applied biomedical and microbiological research. Variations in any of the growth influencing factors including nutrient availability, temperature, pH, aeration, redox potential, water activity, solute concentrations, and the culture volume along with the vessel size may affect bacterial growth rate, which is universally known as stress phenomenon [1-6]. Upon entry of into the stationary phase, *Escherichia coli* cells have been shown to lose their ability to reproduce on standard nutrient plates, i.e., cells enter into a viable but non-culturable (VBNC) state, possibly due to the accumulation of reactive oxygen species (ROS) [3,4,8-15]. However, such oxidative stress can be combated by catalase and superoxide dismutase mediated response in *E. coli* [3,4,9,10,11,16,17,18]. Other bacteria including *Pseudomonas* spp. and *Bacillus* spp. have also been reported to elicit responses against such oxidative stress [19-26].

Effects of agitation on bacterial growth kinetics have been studied earlier; however, neither the association between the shaking speed with the advancement of endogenous oxidative stress nor the influence of agitation

speed on cell culturability has been deliberated [27,28,29]. Our previous studies showed three independent aspects in evolving stress: (i) a close proportional relation with the temperature up-shift with the generation of oxidative stress [30,31]; (ii) spontaneous accumulation of ROS generating oxidative stress at the early stationary phase [3] and (iii) the commencement of such stress upon the addition of 3 mM hydrogen peroxide (H₂O₂) [3,32,33]. In all cases, *E. coli* strains were found to deteriorate their viability with a reduced rate of the formation of colony forming units (CFUs) [2,3,6,30,32]. Apart from the external induction by H₂O₂, temperature up-shift or nutrient starvation, one of the critical parameters decreasing cell viability could be the culture shaking speed [27,28,34,35]. An imitation of such stress responsive event with other bacterial strains or species might be of interest to the inclusive insight of the cellular responses. Along these lines, current investigation assessed the effect of aeration speed apart from the external of addition of H₂O₂ on *E. coli*, *Pseudomonas*, and *Bacillus* spp.

2. Materials and Methods

2.1. Bacterial Stain, Medium and Culture Condition

Bacterial strains used in this study were previously preserved on Nutrient Agar (NA) at 4°C [32]. After 2 hours of growth of freshly grown culture (pre culture) in 5 ml nutrient broth (NB), the OD at 600 nm (OD₆₀₀) of the culture broth was adjusted to 0.1. A volume of 30 µL was introduced into 3 different sets of 30 ml of nutrient broth and was incubated at 37°C with shaking at 0, 100 and 200 rotation per minute (rpm). At every 12 hours of intervals cell growth was monitored by measuring OD₆₀₀ and the colony forming units (CFU)/ml was estimated by counting the colonies on nutrient agar (NA) at every 24 hours [4,30,32].

2.2. Microscopy

Samples were collected at every 12 hours of intervals and the shape and arrangement of the bacterial cells were observed by using simple staining (Crystal Violet, Hucker's Solution) under light microscope (Optima Biological Microscope G206, manufactured in Taiwan) at 1000 × magnification [32].

2.3. Spot Test




Figure 1. Assessment of stress response in *Escherichia coli* through the examination of cell turbidity (a), enumeration of CFU (b), observation for the accumulation of cell aggregates in culture media (c), morphological study: arrowheads indicate impaired cells (d) and the demonstration of loss of cell culturability by means of spot test (e). Cells were grown in nutrient broth at 37°C under 0, 100 and 200 rpm as described in Materials and Methods. At the times indicated, all examinations were carried out

As depicted in Figure 1 c, *E. coli* culture medium was observed to be filled with cell aggregates, suggestive of the defective cell lysis phenomenon [3,4,6]. Such defective cells are suggested to consist of a major fraction of VBNC cells, which was further confirmed through the loss of culturability [1,2,3,4,6,9,11]. Besides, flat colonies characteristic of stressed cells [6] were observed for *E. coli* cells after 36 hours and the non-pigmented colonies were observed for *Pseudomonas* spp. upon prolonged under 200 rpm aeration. Thus the decline in CFUs as well as the colony characteristics referred to the stress commencement in *Pseudomonas* cells upon cultivation at 200 rpm. No such impaired phenotypes in *Bacillus* colonies were observed nor was the accumulation of cell aggregates detected in their culture media. Rather a steady

The bacterial suspension was serially diluted in 9 ml nutrient broth up to 10⁻⁴. From each of the 4 dilutions, 5 µL of the bacterial suspension was spotted onto NA plate. After inoculums dried off, plates were incubated at 37°C for 24 hours. Spotting was done after every 12 hours interval [30,32].

3. Results and Discussion

3.1. Growth Retardation of *E. Coli* and *Pseudomonas* spp. Under 200 rpm

All three bacterial species showed a relatively slower growth at 0 rpm compared to those at 100 rpm (Figure 1 a, Figure 1 b, Figure 2 a, Figure 2 b, and Figure 3 a, Figure 3 b). While shaken at the highest speed of aeration, i.e., at 200 rpm, the colony forming units (CFUs) were observed to be significantly reduced after 36 hour in case of *E. coli* and *Pseudomonas* cells. A minor reduction in CFU in *Bacillus* cells was also observed after 48 hour; however, the trend was insignificant compared to the former strains.

growth was monitored all along the incubation period in case of *Bacillus* spp. at 200 rpm (Figure 3 a, Figure 3 b), possibly due to the upregulation of the general stress regulon by the unique stressosome complex as described previously [36].

Earlier studies showed *Pseudomonas* cells to exhibit a reduced growth rate under high oxygen tension [8,23]. However, *P. aeruginosa* is known to express the pqrCBAR genes which in turn may neutralize the oxidants [18,21,37,38]. In our experimental condition, the reduction trend in CFUs by *Pseudomonas* cells after 36 hour might be due to the impairment of such defense mechanism which is in cohort to the previous studies [8,23].




Figure 2. Assessment of stress response in *Pseudomonas* spp. through the examination of cell turbidity (a), ability to form CFUs (b), morphological observation: arrowheads indicate impaired cells (c) and the demonstration of loss of cell culturability through spot test (d). Cells were grown in nutrient broth at 37°C under 0, 100 and 200 rpm as described in Materials and Methods. At the times indicated, examinations were carried out. No accumulation of cell aggregates in culture media was observed




Figure 3. Assessment of stress response in *Bacillus* spp. through the examination of cell turbidity (a), CFU formation (b), morphological observation (c) and the demonstration of loss of cell culturability by means of spot test (d). Cells were grown in nutrient broth at 37°C under 0, 100 and 200 rpm as described in Materials and Methods. At the times indicated, all examinations were carried out. The morphological change at 72 hour of incubation in static condition is shown (e)

3.2. Morphological Changes in Bacterial Cells in Relation to Cell Viability

All three bacterial species showed a relatively slower growth. While at 36 hours of incubation, no significant changes in cell morphology and shape were observed in *E. coli* at all rpm; however, both the static condition and the highest agitation (i.e., 200 rpm) resulted in cell aggregates

with a cessation in regular rod size (Figure 1 d). This is consistent with the observed changes in colony morphology after 36 hours, with a parallel impact on the reduction of culturability (Figure 1 e). Interestingly, under 200 rpm aeration, most of the *Pseudomonas* spp. cells were found to transform to spherical shape after 36 hours, and at 72 hours these cells were observed to be in aggregated condition (Figure 2 c). Like *E. coli* cells,

Pseudomonas cells were also found to lose their culturability after 36 hours at 200 rpm (Figure 2 d). Thus, both in the cases of *E. coli* and *Pseudomonas* cells, morphologically defective cells were observed which were worth to explain the ultimate non-culturability of these strains (Figure 1 e, Figure 2 d) at 72 hours under 200 rpm [3,4,6,9].

One important thing is to ponder that unlike *E. coli*, *Pseudomonas* culture media did not show any accumulation of cell aggregates; however, interestingly the pigmentation in the *Pseudomonas* culture media was found to be eliminated concomitantly with the loss of their culturability. Distinct from *E. coli* and *Pseudomonas* cells, no significant changes in colony phenotype or cell morphology and arrangement were observed for the *Bacillus* spp. cells up to 72 hours upon the variation in the aeration speed. (Figure 3 c). The comparative resistance might be due to the unique cellular survival mechanism imparted by *Bacillus* spp. [39,40,41,42,43].

An interested morphological trait was observed in case of *Bacillus* cells in static condition. At 72 hour under 0 rpm, a significant fraction of cells were observed to be segmented (Figure 3 e). Such event was indicative of the existence of dividing cells resulting in the cellular culturability possible due to the presence of stressosome regulating the secondary messenger levels [36,44,45]. This notion is in cohort with the load of final bacterial load at 72 hour under 200 rpm; i.e., *Bacillus* cells were found to form around 3-log higher CFU/mL than both *E. coli* and *Pseudomonas* spp. (Figure 1, Figure 2 and Figure 3). Another important facet of growth of *Bacillus* spp. lied under the fact that these cells showed a comparatively higher growth than the other two all along at the static condition. Such a growth response irrespective of the aeration speed again could be explained through the stressosomal activities of *Bacillus* spp. [36,44].

4. Conclusion

It's been a long while the aeration speed for bacterial cultivation known to influence the growth rate [46,47,48]. In the current investigation, the preliminary experiments achieved a bit expected result that the excessive aeration caused significant impact on *Escherichia coli* and *Pseudomonas* spp. cells viability and culturability and that the *Bacillus* cells were bit unaffected by the aeration speed. Thus our earlier results relating to the cessation of bacterial culturability caused by upshifting temperature or by externally adding oxidant might recruit the elevated aeration speed as another stress evoking stimulant [3,30,32]. However, the suggestive relation between increased aeration and the commencement of oxidative stress needs to further concluded by the direct measurement of ROS. Besides, the genetic revelation of the phenotypes studied here would unveil the mechanism of cellular survival against intrinsic stress stimuli.

Acknowledgements

Authors are thankful to Kamal Kanta Das, Mrityunjoy Acharjee and Saurab Kishore Munshi for technical assistance, valuable suggestions and careful monitoring

during the experiments. Authors also gratify Stamford University Bangladesh for the financial and logistic supports.

Conflict of Interest

None of the authors have conflict of interest.

References

- [1] Nagamitsu, H., Murata, M., Kosaka, T., Kawaguchi, J., Mori, H., Yamada, M., "Crucial roles of MicA and RybB as vital factors for σ E dependent cell lysis in *Escherichia coli* long-term stationary phase," *J Mol Microbiol Biotechnol*, 23, 227-232, 2013.
- [2] Murata, M., Noor, R., Nagamitsu, H., Tanaka, S., Yamada, M., "Novel pathway directed by σ E to cause cell lysis in *Escherichia coli*," *Genes to Cells*, 17, 234-247, 2012.
- [3] Noor, R., Murata, M., Yamada, M., "Oxidative stress as a trigger for growth phase-specific σ E dependent cell lysis in *Escherichia coli*," *Journal of Molecular Microbiology and Biotechnology*, 17, 177-187, 2009.
- [4] Noor, R., Murata, M., Nagamitsu, H., Klein, G., Rain, S., Yamada, M., "Dissection of sigma-E dependent cell lysis in *Escherichia coli*: roles of RpoE regulators RseA, RseB and periplasmic folding catalyst Ppid," *Genes to cells*, 14, 885-899, 2009.
- [5] Moat, A.G., Foster, J.W., Spector, M.P., *Lipid and Sterol in Microbial Physiology*, Wiley and Sons, New York, 2002.
- [6] Nitta, T., Nagamitsu, H., Murata, M., Izu, H., Yamada, M., "Function of the sigma-E regulon in dead-cell lysis in stationary phase *Escherichia coli*," *J Bacteriol*, 182, 5231-5237, 2000.
- [7] Baez, A., Shiloach, J., "*Escherichia coli* avoids high dissolved oxygen stress by activation of SoxRS and manganese-superoxide dismutase," *Microb Cell Fact*, 12, 12-23, 2013.
- [8] Imlay J.A., "Cellular Defenses against Superoxide and Hydrogen Peroxide," *Annual Review of Biochemistry*, 77, 755-76, 2008.
- [9] Cuny, C., Dukan, L., Frayssse, L., Ballesteros, M., Dukan, S., "Investigations of the first events leading to the loss of culturability during *Escherichia coli* starvation: future non-culturable bacteria form a subpopulation," *J Bacteriol*, 187, 2244-2248, 2005.
- [10] Nystrom, T., "Role of oxidative carbonylation in protein quality control and senescence," *The EMBO Journal*, 24, 1311-1317, 2005.
- [11] Desnues, B., Cuny, C., Gregori, G., Dukan, S., Aguilaniu, H., Nystrom, T., "Differential oxidative damage and expression of stress defence regulons in culturable and nonculturable *Escherichia coli* cells," *EMBO reports*, 4: 400-404, 2003.
- [12] Cabiscol, E., Tamarit, J., Ros, J., "Oxidative stress in bacteria and protein damage by reactive oxygen species," *Internatl Microbiol*, 3, 3-8, 2000.
- [13] Dukan, S., Nystrom, T., "Oxidative stress defense and deterioration of growth arrested *Escherichia coli* cells," *J Biol Chem*, 274, 26027-26032, 1999.
- [14] Dukan, S., Nyström, T., "Bacterial senescence: stasis results in increased and differential oxidation of cytoplasmic proteins leading to developmental induction of the heat shock regulon," *Genes Dev*, 12, 3431-3441, 1998.
- [15] Farr, S.B., Kogoma, T., "Oxidative stress responses in *Escherichia coli* and *Salmonella typhimurium*," *American Society for Microbiology*, 55 (4), 561-585, 1991.
- [16] Hunter, G.J., Hunter, T., "GroESL protects superoxide dismutase (SOD)-deficient cells against oxidative stress and is a chaperone for SOD," *SciRes*, 5, 1719-1729, 2013.
- [17] Chiang, S.M., Schellhorn, H.E., "Regulators of oxidative stress response genes in *Escherichia coli* and their functional conservation in bacteria," *Arch Biochem Biophys*, 525 (2), 161-169, 2012.
- [18] Hunter, T., Bannister, J.V., Hunter, G.J., "Thermo stability of manganese and iron-superoxide dismutases from *Escherichia coli* is determined by the characteristic position of a glutamine residue," *European Journal of Biochemistry*, 269, 5137-5148, 2002.
- [19] Rungrasamee, W., Kelly, C., Ryan, M.J.M., Pablo, J.P., "The PqrR Transcriptional repressor of *Pseudomonas aeruginosa*

- transduces redox signals via an iron containing prosthetic group," *Journal of Bacteriology*, 191, 6709-6721, 2009.
- [20] Zuber, P., "Management of oxidative stress in *Bacillus* spp.," *Annual Review of Microbiology*, 63, 575-597, 2009.
- [21] Rungrassamee, W., Oxidative stress responses in *Escherichia coli* and *Pseudomonas aeruginosa*, Electronic Doctoral Dissertations for University of Massachusetts Amherst, 2008. Available at: <http://scholarworks.umass.edu/dissertations/AAI3325244>. [January 1, 2008].
- [22] Helmann, J.D., Wu, M.F.W., Gaballa, A., Kobel, P.A., Morshedi, M.M., Fawcett, P., Paddon, C., "The global transcriptional response of *Bacillus subtilis* to peroxide stress is coordinated by three transcription factors," *J Bacteriol*, 185, 243-253, 2003.
- [23] Sabra, Eun-Jin, K., Anping, Z., "Physiological response of *Pseudomonas aeruginosa* PA01 to oxidative stress in controlled microaerobic and aerobic cultures," *Microbiology*, 148, 3195-3202, 2002.
- [24] Friedberg, E.C., Walker, G.C., Siede, W., *DNA repair and mutagenesis*, American Society for Microbiology, Washington DC, 1995.
- [25] Givskov, M., Eberl, L., Moller, S., Poulsen, L. K., Molin, S., "Responses to nutrient starvation in *Pseudomonas Putida* KT2442: analysis of general cross-protection, cell shape, and macromolecular content," *J Bacteriol*, 176, 7-14, 1994.
- [26] Inaoka, T., Matsumura, Y., Tsuchido, T., "SodA and manganese are essential for resistance to oxidative stress in growing and sporulating cells of *Bacillus subtilis*," *J Bacteriol*, 181 (6), 1939-1943, 1990.
- [27] Soini, J., Ukkonen, K., Neubauer, P., "High cell density media for *Escherichia coli* are generally designed for aerobic cultivations-consequences for large-scale bioprocesses and shake flask cultures," *Microbial Cell Factories*, 7, 26, 2008.
- [28] Colemana, M.E., Tamplin, M.L., Phillips, J.G., Marmer, B.S., Influence of agitation, inoculum density, pH, and strain on the growth parameters of *Escherichia coli* O157: H7-relevance to risk assessment," *International Journal of Food Microbiology*, 83, 147-160, 2003.
- [29] Conter, A., Gangneux, C., Suzanne, M., Gutierrez, C., "Survival of *Escherichia coli* during long-term starvation: effects of aeration, NaCl, and the rpoS and osmC gene products," *Res Microbiol*, 152, 17-26, 2001.
- [30] Noor, R., Islam, Z., Munshi, S.H., and Rahman, F., "Influence of temperature on *Escherichia coli* growth in different culture media," *Journal of pure and applied microbiology*, 7 (2), 899-904, 2013.
- [31] Yamada, M., Noor, R., Nagamitsu, H., and Murta, M., "The higher temperature, the more oxidative stress and lysis in *Escherichia coli*," *The 3rd International Conference on Fermentation Technology for Value Added Agricultural Products*, Khon Kaen, Thailand, 2009.
- [32] Munna, M.S., Nur, I., Rahman, T., Noor, R., "Influence of exogenous oxidative stress on *Escherichia coli* cell growth, viability and morphology," *American Journal of BioScience*, 1 (4), 59-62, 2013.
- [33] Kabir, M.S., Yamashita, D., Noor, R., Yamada, M., "Effect of σ^S on σ^E -directed cell lysis in *Escherichia coli* early stationary phase," *J Mol Microbiol Biotechnol*, 2004, 8, 189-194.
- [34] Mary, K.B., David, S., Phillips, Janet, O.B., "Shaker agitation rate and orbit affect growth of cultured bacteria," *Thermo Fisher Scientific*, 2011.
- [35] Juergensmeyer, M.A., Nelson, E.S., Juergensmeyer, E.A., "Shaking alone, without concurrent aeration, affects the growth characteristics of *Escherichia coli*," *Letters in Applied Microbiology*, 45, 179-183, 2007.
- [36] Quin, M.B., Berrisford, J.M., Newman, J.A. Baslé, A., Lewis, R.J., Marles, W.J., "The bacterial stressosome: a modular system that has been adapted to control secondary messenger signaling," *Structure*, 20 (2), 350-363, 2012.
- [37] Wilkinson, S.P., Grove, A., "Ligand-responsive transcriptional regulation by members of the MarR family of winged helix proteins," *Curr Issues Mol Biol*, 8, 51-62, 2006.
- [38] Salunkhe, P., Topfer, T., Buer, J., Tummeler, B., "Genome-wide transcriptional profiling of the steady-state response of *Pseudomonas aeruginosa* to hydrogen peroxide," *J Bacteriol*, 187, 2565-2572, 2005.
- [39] Den Besten, H., M., W., Effraimidou, S., Abee, T., "Catalase activity as a biomarker for mild stress induced robustness in *Bacillus weihenstephanensis*," *Appl Environ Microbiol*, 79 (1), 57-62, 2013.
- [40] Huillet, E., Tempelars, M., Andre, L.G., Wanapaisan, P., Bridoux, L., Makhzamis, et al., "P1cRa, a new quorum-sensing regulator from *Bacillus cereus*, play a role in oxidative stress response and cysteine metabolism in stationary phase," *Genetique Microbiennet Environent*, 7 (12), e51047, 2012.
- [41] Pohl, S., Tu, W., Aldridge, P., Gillespie, C., Hahne, H., Mader, U., "Combined proteomic and transcriptomic analysis of the response of *Bacillus anthracis* to oxidative stress," *Proteomics*, 11 (15), 3036-55, 2011.
- [42] Passalacqua, K., Bergman, N., Leejy, Sherman, D., Hanna, P., "The superoxide dismutases of *Bacillus anthracis* do not cooperatively protect against endogenous superoxide stress," *J Bacteriol*, 188 (11), 3837-48, 2006.
- [43] Passalacqua, K., Bergman, N., Leejy, Sherman, D., Hanna, P., "The global transcriptional responses of *Bacillus anthracis* Sterne (34F2) and a delta sodAI mutant to paraquat reveal metal ion homeostasis imbalances during endogenous superoxide stress," *J Bacteriol*, 189 (11), 3996-4013, 2007.
- [44] Hiremath, P.S., Parashuram, B., "Automatic identification and classification of bacilli bacterial cell growth phases," *IJCA Special Issue on Recent Trends in Image Processing and Pattern Recognition*, IJCA Journal, 48-52.
- [45] Setlow, B., Setlow, P., "Role of DNA repair in *Bacillus subtilis* spore resistance," *J Bacteriol*, 178, 3486-3495, 1996.
- [46] Losen, M., Frölich, B., Pohl, M., Büchs, J., "Effect of oxygen limitation and medium composition on *Escherichia coli* fermentation in shake-flask cultures," *Biotechnol Prog*, 20 (4), 1062-1068, 2004.
- [47] Bunch, A.W., "High cell density growth of microorganisms," *Biotechnol Genet Eng Rev*, 12, 535-561, 1994.
- [48] McDaniel, L.E., Bailey, E.G., Zimmerli, A., "Effect of oxygen-supply rates on growth of *Escherichia coli*. II. Comparison of results in shake flasks and 50-liter fermentor," *Appl Microbiol*, 13:115-9, 1965.