

Influence of dopaminergic and serotonergic genes on working memory in healthy subjects

Monika Wilkość^{1*}, Joanna Hauser², Marta Tomaszewska¹, Monika Dmitrzak-Węglarz², Maria Skibińska², Aleksandra Szczepankiewicz², and Alina Borkowska¹

¹Chair and Department of Clinical Neuropsychology, Nicolaus Copernicus University in Torun, Collegium Medicum in Bydgoszcz, Bydgoszcz, Poland, *Email: monikawilkosc@gmail.com; ²Laboratory of Psychiatric Genetics, Department of Psychiatry, Poznań University of Medical Sciences, Poznan, Poland

Working memory is an ability to keep information in short-term memory and manipulate them “on line”. Working memory is also involved in complex frontal executive functions. The role of dopaminergic system in modulating working memory processes in prefrontal cortex is well established. Also the role of serotonergic receptors is postulated. The purpose of this study was to assess the association between the polymorphisms of dopaminergic (DRD1, DRD3, DRD4, COMT) and serotonergic (SERT – serotonin transporter, 5HT2A, 5HT2C) genes’ polymorphisms and performance on WCST in 200 volunteers from the Polish population. We found the association between DRD1, DRD4, COMT and SERT genes polymorphisms and the performance on WCST. The results obtained in the study indicate that dopaminergic and serotonergic genes may play a role in modulating the executive function and working memory processes in healthy subjects. The pattern of this influence may be different in males and females. Moreover, the relationship between the efficacy of prefrontal cognitive function and genes polymorphisms may differ between healthy subjects and schizophrenic patients.

Key words: Wisconsin Card Sorting Test, dopaminergic and serotonergic genes, working memory, healthy subjects

INTRODUCTION

Working memory is an ability to keep information in short-term memory and manipulate them “on line”. Working memory is also involved in complex frontal executive functions (Baddeley 2000, D’Esposito et al. 2000, Fletcher and Henson 2001). Working memory deficit in schizophrenic and bipolar patients, as well as in their first-degree relatives indicates the genetic mechanism underlying its function. Currently it is proposed as a cognitive endophenotypic marker of vulnerability to these illnesses (Borkowska and Rybakowski 2001,

Everett et al. 2001, Gottesman and Gould 2003, Glahn et al. 2004, Zalla et al. 2004, McIntosh et al. 2005).

The role of dopaminergic system in modulating working memory processes in prefrontal cortex is well established (Seamans and Yang 2004). Many studies demonstrated the significance of activity of D1 dopamine receptors in determining the efficacy of the “on-line” processes, whereas interaction of D1 and D2 receptors influences the overall efficacy of working memory (Goldman-Rakic 2000, Goldman-Rakic et al. 2004). The association between the polymorphism of DRD1 receptor gene and the performance on Wisconsin Card Sorting Test (WCST) in schizophrenic patients was demonstrated (Rybakowski et al. 2005). Laszy and coauthors (2005) postulated the role of D3 dopamine receptors in the processes of memory and learning. Szekeres and colleagues (2004) found the correlation between the results on WCST and the polymorphism of DRD3 gene. The experimental study on the dopamine receptor D4 antagonists has demonstrated their beneficial influence on

Correspondence should be addressed to M. Wilkość
Email: monikawilkosc@gmail.com

*Present address of the Author: (A) Department of Individual Differences Psychology, Psychology Institute, Kazimierz Wielki University in Bydgoszcz, Bydgoszcz, Poland; (B) Department of Psychiatry, Nicolaus Copernicus University in Torun, Collegium Medicum in Bydgoszcz, Bydgoszcz, Poland

Received 08 June 2009, accepted 26 February 2010

working memory (Zhang et al. 2004). Also the interaction of DRD4 and NMDA in the prefrontal cortex may play a significant role in determining cognitive efficiency by modulating synaptic plasticity (Wang et al. 2003).

It has been shown that the correlation between the amount of dopamine in the prefrontal cortex and cognitive performance can be represented as an inverted U, which means that overflow as well as deficiency of dopamine may have a negative impact on cognitive functions, particularly on working memory (Arnsten and Li 2005). One of the most studied factors that modulate the dopamine levels in prefrontal cortex is catechol-O-methyltransferase (COMT). The Met variant of this enzyme shows a 3–4 times lower activity leading to the increased dopamine level (Lachman et al. 1996, Chen et al. 2004). Numerous studies investigated the influence of COMT on prefrontal function measured by WCST (Egan et al. 2001, Malhotra et al. 2002, Rosa et al. 2004, Bruder et al. 2005, Rybakowski et al. 2005, 2006, Barnett et al. 2007a). Sambataro and coworkers (2009) demonstrated that the effect of Val108(158)Met polymorphism on activity and functional connectivity of brain regions involved in working memory network increases with age. There is also an evidence for a sexually dimorphic influence of COMT (Sazci et al. 2004, Harrison and Turnbridge 2008). The other polymorphisms of COMT are also postulated to have an influence on prefrontal cognitive functions (Meyer-Lindenberg et al. 2006, Diaz-Asper et al. 2008), as well as interaction between COMT and DRD4 (Alfimova et al. 2007) or DRD2 polymorphisms (Xu et al. 2007).

The role of the serotonin system in the evolution of human cognitive capacities was supported by changes in the supply of this neurotransmitter to the frontal cortex (Raghanti et al. 2008). The serotonin transporter (SERT) plays an essential role in serotonergic neurotransmission as it determines the quality and duration of the serotonin signal in the synaptic cleft. The 5HT receptors play also an important role in brain development and memory processes (Meneses et al. 1997, Meneses 1999, van Kesteren and Spencer 2003) and may be a target for cognitive enhancement (Roth 2004). It was demonstrated that memory performance may be correlated with SERT density in the dorsolateral prefrontal cortex, orbitofrontal cortex, and parietal cortex which are implicated in memory function. (McCann et al. 2008). The 5HT function also modulates dopaminergic activity (Nocjar 2002). Regulatory role of serotonin-dopamine interactions in cognition and the prefrontal

Table I

Results of WCST in relation to –48A/G polymorphism of DRD1 gene (X ± SD)				
DRD1 genotype	Total			
	WCST	A/A	A/G	G/G
%P	7.69 ± 2.4	8.36 ± 3.4	8.26 ± 2.6	
%NP	8.26 ± 4.2	7.52 ± 4.0	7.67 ± 3.3	
%CONC	81.52 ± 7.8	81.52 ± 7.9	81.23 ± 6.5	
CC	6.00 ± 0.0	6.00 ± 0.2	6.00 ± 0.1	
1stCAT	13.78 ± 6.2	13.87 ± 6.5	13.26 ± 5.0	
Females				
DRD1 genotype	Total			
	WCST	A/A	A/G	G/G
%P	8.00 ± 2.9	8.34 ± 3.3	8.30 ± 2.7	
%NP	9.77 ± 4.7	7.84 ± 4.2	7.83 ± 4.2	
%CONC	78.69 ± 8.5	81.48 ± 7.1	80.43 ± 8.0	
CC	6.00 ± 0.0	5.96 ± 0.2	6.00 ± 0.0	
1stCAT	16.00 ± 7.6*	14.11 ± 6.4	12.33 ± 4.1	
Males				
DRD1 genotype	Total			
	WCST	A/A	A/G	G/G
%P	7.30 ± 1.6	8.39 ± 3.4	8.22 ± 2.5	
%NP	6.30 ± 2.4*	7.13 ± 3.7	7.53 ± 2.3	
%CONC	85.21 ± 4.9*	81.57 ± 8.8	81.89 ± 4.9	
CC	6.00 ± 0.0	5.96 ± 0.3	5.97 ± 0.2	
1stCAT	10.9 ± 0.6**	13.59 ± 6.6 [#]	13.97 ± 5.6	

*Difference A/A vs. A/G $P < 0.05$; **A/A vs. A/G $P < 0.01$; [#] A/G vs. G/G

Table II

Results of WCST in relation to polymorphism -521C/T of DRD4 gene ($X \pm SD$)				
DRD4 genotype	Total			
	WCST	C/C	C/T	T/T
%P	7.50 \pm 2.7	8.58 \pm 3.1	8.25 \pm 2.9	
%NP	6.92 \pm 2.9	8.04 \pm 4.1	7.28 \pm 3.6	
%CONC	83.02 \pm 6.0	80.45 \pm 7.7	82.22 \pm 7.2	
CC	6.00 \pm 0.0	5.95 \pm 0.3	5.99 \pm 0.1	
1stCAT	13.08 \pm 5.3	13.7 \pm 5.9	13.74 \pm 6.1	
Females				
WCST	C/C	C/T	T/T	
%P	7.48 \pm 3.3	8.58 \pm 3.3	8.47 \pm 2.7	
%NP	7.67 \pm 3.3	9.00 \pm 4.7	6.75 \pm 3.9 [#]	
%CONC	81.52 \pm 6.5	79.7 \pm 8.17	82.39 \pm 6.9	
CC	6.00 \pm 0.0	5.98 \pm 0.2	5.97 \pm 0.2	
1stCAT	12.91 \pm 5.5	14.33 \pm 6.1	13.42 \pm 5.8	
Males				
WCST	C/C	C/T	T/T	
%P	7.53 \pm 2.0	8.59 \pm 3.1	8 \pm 3.2	
%NP	6.00 \pm 2.1*	7.16 \pm 3.3	7.88 \pm 3.1	
%CONC	84.88 \pm 5.0	81.14 \pm 7.3	82.03 \pm 7.7	
CC	6.00 \pm 0.0	5.93 \pm 0.3	6.00 \pm 0.0	
1stCAT	13.3 \pm 5.4	13.2 \pm 5.7	14.1 \pm 6.5	

*Difference C/C vs. T/T $P < 0.05$; [#]T/T vs. C/T $P < 0.05$

cortex and the striatum as a neuroanatomical substrate for these interactions, are postulated. Experimental data revealed that disruption of serotonin neurotransmission results in a facilitative effect on the processing of mnemonic information by refrontal cortex and the striatum which are under strong, functional DA modulation, while increased serotonin neurotransmission has a detrimental effect on cognitive functions connected with these brain structures (Olvera-Cortés et al. 2008). Neuroanatomical findings in schizophrenic patients revealed an altered 5HT_{2A} serotonergic receptors distribution in prefrontal cortex (Dean 1996, Dean and Hayes 1996). Üçok and colleagues (2007) found a correlation between the results of WCST and polymorphism of 5HT_{2A} receptors' gene.

Relatively few studies examined the relationship between polymorphisms of dopaminergic and serotonergic receptors' genes and prefrontal cognitive functions in healthy subjects.

The purpose of this study was to assess the association between the polymorphisms of dopamine and serotonin systems' genes and working memory in healthy subjects. The association between dopaminergic (DRD1, DRD3, DRD4, COMT) and serotonergic (SERT – serotonin transporter, 5HT_{2A}, 5HT_{2C}) genes' polymorphisms and performance on WCST in the Polish population was assessed.

METHODS

Subjects

The study included 200 healthy volunteers (100 males and 100 females) aged 18–60 (mean 34.4 \pm 11.7) years with no history of any psychiatric disorder, substance abuse or serious somatic illnesses. All subjects were assessed using the Polish version of M.I.N.I Plus scale (Lecrubier et al. 1997) for excluding presence of any serious mental health problems. The Bioethics Committee of Nicolaus Copernicus University Torun, Collegium Medicum in Bydgoszcz, approved the study. Informed consent was received from all participants after the aim and procedure of the study were fully explained to them. All subjects were Caucasians of Polish origin.

Neuropsychological assessment

Working memory evaluation was made using the Heaton and coauthors (1993) computerized version of

WCST. The following domains of WCST were measured: (1) the percentage of perseverative errors (%P) – connected with inability to change reaction due to ignorance of relevant stimuli; (2) the percentage of non-perseverative errors (%NP) – connected with attention inability to avoid distraction; (3) the percentage of conceptual level response (%CONC) – connected with ability to utilize new information and previous experience to create conception; (4) the number of correctly completed categories (CC) – connected with effectiveness of measured cognitive function; (5) the set to complete first category (1stCAT) – connected with speed of formulating logical conception.

Genotyping

Genomic DNA was extracted from 10 ml of ethylene-diaminetetraacetic (EDTA) acid-anticoagulated venous blood using the salting out method (Miller et al. 1988). The polymorphisms of interest were amplified by PCR-RLFP method.

The following polymorphisms were determined using cited methods: -48 A/G of DRD1 (Cichon et al. 1994), Ser9Gly of DRD3 (Lannfelt et al. 1992), -521C/T of DRD4 (Jönsson et al. 2001), Val108(158)Met of COMT (Li et al. 1996), ins/del of SERT (Stoltenberg et al. 2002), T102C of 5HTR2A (Du et al. 2000), C68G (Cys23Ser) of 5HT2C (Lappalainen et al. 1995).

Statistics

For statistical evaluation The Shapiro-Wilk test was used to evaluate the normality of the variables' distribution. In case of normal distribution one-way ANOVA was applied. Otherwise non-parametric tests of Friedman ANOVA and U Mann-Whitney were used. All statistical analyses were performed using Statistica 7.0 package (STATSOFT, Poland). The concordance of genotypes with Hardy-Weinberg equilibrium was assessed using „Utility Programs for Analysis of Genetic Linkage” (Copyright © 1988 J. Ott).

RESULTS

The analysis of -48A/G polymorphism showed that women with A/A genotype obtained significantly worse results comparing to those with G/G one in WCST-1stCAT, while males with A/A and A/G genotypes performed better in this parameter of WCST

Table III

Results of WCST in relation to polymorphism Val108(158)Met of COMT gene (X ± SD)			
COMT genotype	Total		
	Met/Met	Val/Met	Val/Val
WCST			
%P	8.31 ± 3.1	8.1 ± 2.9	8.55 ± 3.2
%NP	7.36 ± 4.2	7.87 ± 3.7	7.36 ± 3.7
%CONC	81.95 ± 7.8	81.2 ± 7.1	81.4 ± 8.1
CC	6.00 ± 0.0	5.95 ± 0.3	6.00 ± 0.0
1stCAT	13.15 ± 4.8	13.7 ± 6.0	13.93 ± 6.6
Females			
WCST			
%P	8.75 ± 3.7	8.35 ± 3.0	8.13 ± 2.9
%NP	9.38 ± 4.1	7.98 ± 4.3	7.42 ± 4.5
%CONC	78.94 ± 8.0	80.62 ± 7.7	81.96 ± 7.5
CC	6.00 ± 0.0	5.97 ± 0.2	6.00 ± 0.0
1stCAT	13.50 ± 3.7	13.97 ± 6.2	13.93 ± 6.6
Males			
WCST			
%P	8.00 ± 2.8	7.78 ± 2.7	9.00 ± 3.5
%NP	5.96 ± 3.7	7.74 ± 2.9**	7.3 ± 2.7*
%CONC	84.04 ± 7.1	81.9 ± 6.4	80.83 ± 8.8
CC	6.00 ± 0.0	5.94 ± 0.3	6.00 ± 0.0
1stCAT	12.9 ± 5.4	13.36 ± 5.7	14.00 ± 6.6

*Difference Met/Met vs. Val/Val $P < 0.05$;

**Met/Met vs. Val/Met $P < 0.005$

Table IV

Results of WCST in relation to polymorphism ins/del of SERT gene ($X \pm SD$)			
SERT genotype	Total		
	s/s	s/l	l/l
WCST			
%P	7.95 \pm 3.6	8.42 \pm 3.1	8.12 \pm 2.7
%NP	8.50 \pm 4.8	7.35 \pm 3.5	7.77 \pm 3.7
%CONC	81.65 \pm 7.9	81.34 \pm 7.1	81.60 \pm 7.3
CC	6.00 \pm 0.0	5.96 \pm 0.2	6.97 \pm 0.2
1stCAT	12.65 \pm 5.4*	12.74 \pm 4.6 ^{##}	15.04 \pm 6.9
Females			
WCST	s/s	s/l	l/l
%P	8.55 \pm 4.3	8.55 \pm 3.0	8.02 \pm 2.9
%NP	10.64 \pm 5.5**	6.89 \pm 3.4	6.28 \pm 4.3
%CONC	78.90 \pm 8.1	81.68 \pm 6.9	80.74 \pm 8.0
CC	6.00 \pm 0.0	5.98 \pm 0.2	5.98 \pm 0.2
1stCAT	14.00 \pm 7.1	12.27 \pm 3.7 [#]	15.07 \pm 6.9
Males			
WCST	s/s	s/l	l/l
%P	7.22 \pm 2.5	8.29 \pm 3.2	8.26 \pm 2.6
%NP	5.89 \pm 1.6	7.85 \pm 3.6	7.03 \pm 2.6
%CONC	85.00 \pm 6.5	80.98 \pm 7.4	82.83 \pm 6.1
CC	6.00 \pm 0.0	5.95 \pm 0.3	5.97 \pm 0.2
1stCAT	11.00 \pm 2.8*	12.24 \pm 5.4	15.00 \pm 7.0

*Difference s/s vs. l/l $P < 0.05$; **s/s vs. l/s $P < 0.05$;

s/l vs. l/l $P < 0.05$; ##s/l vs. l/l $P < 0.01$

compared to G/G. Moreover male homozygotes A performed better in WCST-%NP and %CONC compared to those with G/G genotype (Table I).

The analysis of -521C/T polymorphism of DRD4 demonstrated that males with T/T genotype performed worse than those with C/C one, while in females, T/T genotype was connected with better performance comparing to females with T/C genotype on WCST-%NP (Table II).

Table III shows the results the analysis of Val108(158) Met polymorphism of COMT gene. The correlation with results of WCST in this case was observed only in the male group where genotype Met/Met was connected with significantly better results in WCST-%NP than other genotypes.

The analysis of ins/del polymorphism of SERT gene showed that subjects with l/l genotype obtained worse results in WCST-1stCAT than in other genotypes (Table IV). Moreover, in females, s/s genotype was connected with worse results in WCST-%NP comparing to l/s genotype, while l/s one was related to better results in WCST-1stCAT comparing to l/l. Also in males, l/l genotype was connected with worse results in WCST-1st CAT.

We found no correlation between the polymorphisms: Ser9Gly of DRD3 gene, T102C of 5HTR2A gene, C68G (Cys23Ser) of 5HT2C gene and the results of WCST in healthy subjects.

DISCUSSION

The -48A/G polymorphism of DRD1 has been postulated to modulate transcription activity of this gene (Cichon et al. 1994). Our analysis showed different pattern of influence in males and females. The impact in the male group was more significant and observed in more domains. Women with A/A genotype obtained significantly worse results comparing to those with G/G one in WCST-1stCAT, while males with A/A and A/G genotypes performed better in this parameter of WCST compared to other genotypes. The study of this polymorphism in schizophrenia showed that G/G genotype was connected with the worst results in all domains of WCST (Rybakowski et al. 2005). Also in our study we found that G/G genotype in healthy males was connected with worse results in WCST-%NP, %CONC, 1stCAT, while female group with G/G genotype obtained better results in WCST-1stCAT compared to A/A.

We found no correlation of Ser9Gly polymorphism of DRD3 with results on WCST. DRD3 are located mostly

in limbic structures, hippocampus and hypothalamus (Levesque et al 1992) that may cause their possible larger influence on long-term memory processes than those of working memory. The serine variant of the protein shows a higher affinity for dopamine (Lundstrom and Turpin 1996). The results of previous cognitive studies were ambiguous. Szekeres and coauthors (2004) have found that schizophrenic patients with Ser/Ser genotype performed better in WCST-%P and CC comparing to Ser/Gly one, but work of Rybakowski and others (2005) did not confirm these findings.

The analysis of -521C/T polymorphism of DRD4 gene showed that allele C has a 40% higher transcription activity than allele T (Okuyama et al. 1999). In this study we found the relation between this polymorphism and the results in WCST-%NP. Similarly to the case of DRD1 gene polymorphism, the influence of DRD4 differed according to gender. Males with T/T genotype performed worse than those with C/C one, while in females, T/T genotype was connected with better performance compared to females with T/C genotype on WCST-%NP. In the Polish study, no association between WCST performance and this polymorphism in schizophrenic patients was found (Rybakowski et al. 2005).

Our study demonstrated that Met/Met genotype of Val108(158)Met polymorphism of COMT gene was connected with significantly better results in WCST-%NP compared to other genotypes, but only in males. The previous studies showed that metionine allele of COMT gene is related to better performance on WCST, especially in %P parameter (Egan et al. 2001, Malhotra et al. 2002, Rosa et al. 2004). The metaanalysis of Barnett and colleagues (2007a) have confirmed association of Met homozygotic genotype with better performance in WCST-%P in healthy subjects. They also indicated that the influence of COMT gene might be greater in populations other than Caucasian. The studies postulated the interaction between COMT activity and sex (Gogos et al. 1998). Sazci and coworkers (2004) demonstrated that the influence of Val108(158)Met polymorphism of COMT is more significant in schizophrenic women, while Barnett and others (2007b) showed that COMT genotype effect on IQ is significantly higher in boys than in girls. We found no correlation between this polymorphism and performance on WCST in healthy women. Sambataro and coauthors (2009) showed that Val/Val genotype in older subjects is connected with increased activity in dorsolateral prefrontal cortex and decreased activity in ventrolateral

prefrontal cortex comparing to Met/Met genotype in younger subjects. In schizophrenia, the metionine allele was found to be related to better performance in WCST-CC in study of Bruder and colleagues (2005), while Rybakowski and coworkers (2005) observed the association of Val/Val genotype with better results in WCST-%P in male schizophrenic patients, but with worse results in WCST-%NP in females with schizophrenia. The studies of Bilder and coauthors (2002) and Stefanis and others (2004) showed no correlation between COMT gene's polymorphism and performance on WCST in schizophrenia.

The analysis of serotonergic genes showed the correlation only in case of ins/del polymorphism of SERT. Subjects with l/l genotype obtained worse results in WCST-1stCAT than in other genotypes. Borg and colleagues (2009) have found significant association between SERT genotype and performance in the Wisconsin Card Sorting Test. Subjects with the s allele performed better compared to the l homozygotes. The study of Bosia and others (2010) showed an association between SERT polymorphism and executive functions and inversely with sustained attention. The l/l genotype was connected with better executive and poorer attention performances. We also observed a different pattern of correlation in males and females what indicates that serotonin may have a sexually dimorphic influence on cognition. In females, s/s genotype was connected with worse results in WCST-%NP comparing to l/s genotype, while l/s one was related to better results in WCST-1stCAT comparing to l/l. In males, l/l genotype was connected with worse results in WCST-1st CAT.

We found no correlation between polymorphisms: T102C of 5HT2A and Ser23Gly of 5HT2C genes and performance on WCST. It may indicate lack of influence of these polymorphisms on prefrontal cognitive function in healthy subjects.

CONCLUSIONS

The results obtained in the study indicate that dopaminergic and serotonergic genes may play a role in modulating the executive function and working memory processes in healthy subjects. The pattern of this influence may be different in males and females. Moreover, the relationship between the efficacy of prefrontal cognitive function and genes polymorphisms in healthy subjects may differ from that reported in schizophrenic patients.

ACKNOWLEDGEMENT

The study was supported by grant no. P05B123 26 (Ministry of Science and Higher Education, Poland).

REFERENCES

- Alfimova MV, Golimbet VE, Gritsenko IK (2007) Interaction of dopamine system genes and cognitive functions in patients with schizophrenia and their relatives and in healthy subjects from general population. *Neurosci Behav Physiol* 37: 643–650.
- Arnsten AF, Li BM (2005) Neurobiology of executive functions: catecholamine influences on prefrontal cortical functions. *Biol Psychiatry* 57: 1377–1384.
- Baddeley AD (2000) The episodic buffer: a new component of working memory. *Trends Cogn Sci* 4: 417–423.
- Barnett JH, Jones PB, Robbins TW, Müller U (2007a) Effects of the catechol-O-methyltransferase Val158Met polymorphism on executive function: a meta-analysis of the Wisconsin Card Sort Test in schizophrenia and healthy controls. *Mol Psychiatry* 12: 502–509.
- Barnett JH, Heron J, Ring SM, Golding J, Goldman D, Xu K, Jones PB (2007b) Gender-specific effects of the catechol-O-methyltransferase Val108/158Met polymorphism on cognitive function in children. *Am J Psychiatry* 164: 142–149.
- Bilder RM, Volavka J, Czobor P, Malhotra AK, Kennedy JL, Ni X, Goldman RS, Hoptman MJ, Sheitman B, Lindenmayer JP, Citrome L, McEvoy JP, Kunz M, Chakos M, Cooper TB, Lieberman JA (2002) Neurocognitive correlates of the COMT Val(158)Met polymorphism in chronic schizophrenia. *Biol Psychiatry* 52: 701–707.
- Borg J, Henningson S, Saijo T, Inoue M, Bah J, Westberg L, Lundberg J, Jovanovic H, Andrée B, Nordstrom AL, Halldin C, Eriksson E, Farde L (2009) Serotonin transporter genotype is associated with cognitive performance but not regional 5-HT_{1A} receptor binding in humans. *Int J Neuropsychopharmacol* 12: 783–792.
- Borkowska A, Rybakowski J (2001) Neuropsychological frontal lobe tests indicate that bipolar depressed patients are more impaired than unipolar. *Bipolar Disord* 3: 88–94.
- Bosia M, Anselmetti S, Pirovano A, Ermoli E, Marino E, Bramanti P, Smeraldi E, Cavallaro R (2010) HTTLPR functional polymorphism in schizophrenia: Executive functions vs. sustained attention dissociation. *Prog Neuropsychopharmacol Biol Psychiatry* 34: 81–85.
- Bruder G, Keilp J, Xu H, Shikman M, Schori E, Gorman J, Gilliam TC (2005) Catechol-O-methyltransferase (COMT) genotypes and working memory: associations with differing cognitive operations. *Biol Psychiatry* 58: 901–907.
- Chen J, Lipska BK, Halim M, Ma QD, Matsumoto M, Melhem S, Kolachana BS, Hyde TM, Herman MM, Apud J, Egan MF, Kleinman JE, Weinberger DR (2004) Functional analysis of genetic variation in catechol-O-methyltransferase (COMT): effects on mRNA, protein, and enzyme activity in post mortem human brain. *Am J Hum Genet* 75: 807–821.
- Cichon S, Nothen MM, Erdmann J, Propping P (1994) Detection of four polymorphic sites in the human dopamine D1 receptor gene (DRD1). *Hum Mol Genet* 3: 209.
- D'Esposito M, Postle BR, Rypma B (2000) Prefrontal cortical contribution to working memory: evidence from event-related fMRI studies. *Exp Brain Res* 133: 3–11.
- Dean B (1996) Serotonin 2 receptors and the serotonin transporter in the schizophrenic brain. *Behav Brain Res* 73: 169–175.
- Dean B, Hayes W (1996) Decreased frontal cortical serotonin_{2A} receptors in schizophrenia. *Schizophr Res* 21: 133–139.
- Diaz-Asper CM, Golberg TE, Kolachana BS, Straub RE, Egan MF, Weinberger DR (2008) Genetic variation in catechol-O-methyltransferase: effects on working memory in schizophrenic patients, their siblings, and healthy controls. *Biol Psychiatry* 63: 72–79.
- Du L, Bakish D, Lapierre YD, Ravindran AV, Hrdina PD (2000) Association of polymorphism of serotonin 2A receptor gene with suicidal ideation in major depressive disorder. *Am J Med Genet* 96: 56–60.
- Egan MF, Goldberg TE, Kolachana BS, Callicott JH, Mazzanti CM, Straub RE (2001) Effect of COMT Val 108/158 Met phenotype on frontal lobe function and risk for schizophrenia. *Proc Natl Acad Sci U S A* 98: 6917–6922.
- Everett J, Lavoie K, Gagnon JF, Gosselin N (2001) Performance of patients with schizophrenia on the Wisconsin Card Sorting Test (WCST). *J Psychiatry Neurosci* 26: 123–130.
- Fletcher PC, Henson RN (2001) Frontal lobes and human memory: insights from functional neuroimaging. *Brain* 124: 849–881.
- Glahn DC, Bearden CE, Niendam TA, Escamilla MA (2004) The feasibility of neuropsychological endophenotypes in the search for genes associated with bipolar affective disorder. *Bipolar Disord* 6: 171–182.

- Gogos JA, Morgan M, Luine W, Santha M, Ogawa S, Pfaff D, Karayiorgou M (1998) Catechol-O-Methyltransferase-deficient mice exhibit sexually dimorphic changes in catecholamine levels and behaviour. *Proc Natl Acad Sci U S A* 95: 9991–9996.
- Goldman-Rakic P (2000) D1 receptors in prefrontal cells and circuits. *Brain Res Rev* 31: 295–301.
- Goldman-Rakic PS, Castner SA, Svensson TH, Siever LJ, Williams GV (2004) Targeting the dopamine D1 receptor in schizophrenia: insights for cognitive dysfunction. *Psychopharmacology (Berl)* 174: 3–16.
- Gottesman II, Gould TD (2003) The endophenotype concept in psychiatry: etymology and strategic intentions. *Am J Psychiatry* 160: 636–645.
- Harrison PJ, Tunbridge EM (2008) Catechol-O-methyltransferase (COMT): a gene contributing to sex differences in brain function, and to sexual dimorphism in the predisposition to psychiatric disorders. *Neuropsychopharmacology* 33: 3037–3045.
- Heaton RK, Chelune GJ, Talley JL, Kay GG, Curtis G (1993) Wisconsin Card Sorting Test Manual: Revised and expanded. Psychological Assessment Resources, Odessa, FL.
- Jönsson EG, Ivo R, Forslund K, Mattila-Evenden M, Rylander G, Cichon S, Propping P, Nothen MM, Asberg M, Sedvall GC (2001) No association between a promoter dopamine D(4) receptor gene variant and schizophrenia. *Am J Med Genet* 105: 525–528.
- Lachman HM, Papolos DF, Saito T, Yu YM, Szumlanski CL, Weinshilboum RM (1996) Human catechol-O-methyltransferase pharmacogenetics: description of a functional polymorphism and its potential application to neuropsychiatric disorders. *Pharmacogenetics* 6: 243–250.
- Lannfelt L, Sokoloff P, Martres M, Pilon C, Giros B, Jonsson E, Sedvall G, Schwartz JC (1992) Amino-acid substitution in the dopamine D3 receptor as a useful polymorphism for investigating psychiatric disorders. *Psychiatr Genet* 2: 249–256.
- Lappalainen J, Zhang L, Dean M, Oz M, Ozaki N, Yu DH, Virkkunen M, Weight F, Linnoila M, Goldman D (1995) Identification, expression, and pharmacology of a Cys23-Ser23 substitution in the human 5-HT_{2c} receptor gene (HTR_{2C}). *Genomics* 27: 274–279.
- Laszlo J, Laszlovszky I, Gyertyán I (2005) Dopamine D3 receptor antagonists improve the learning performance in memory-impaired rats. *Psychopharmacology* 179: 567–575.
- Leclercq Y, Sheehan D, Weiller E, Amorim P, Bonora I, Sheehan K, Janavs J, Dunbar G (1997) The MINI International Neuropsychiatric Interview (M.I.N.I.) A Short Diagnostic Structured Interview: Reliability and Validity According to the CIDI. *European Psychiatry* 12: 224–231.
- Levesque D, Diaz J, Pilon C, Martres MP, Giros B, Souil E, Schott D, Morgat JL, Schwartz JC, Sokoloff P (1992) Identification, characterization, and localization of the dopamine D3 receptor in rat brain using 7-[³H]hydroxy-N,N-di-n-propyl-2-aminotetralin. *Proc Natl Acad Sci U S A* 89: 8155–8159.
- Li T, Vallada H, Curtis D, Arranz M, Xu K, Cai G, Deng H, Liu J, Murray R, Liu X, Collier D (1996) Catechol-O-methyltransferase Val158Met polymorphism: frequency analysis in HanChinese subjects and allelic association of the low activity allele with bipolar affective disorder. *Pharmacogenetics* 7: 349–353.
- Lundstrom K, Turpin M (1996) Proposed schizophrenia-related gene polymorphism: expression of the Ser9Gly mutant human dopamine D3 receptor with the Semliki Forest virus system. *Biochem Biophys Res Commun* 225: 1068–1072.
- Malhotra AK, Kestler LJ, Mazzanti C, Bates JA, Goldberg T, Goldman D (2002) A functional polymorphism in the COMT gene and performance on a test of prefrontal cognition. *Am J Psychiatry* 159: 652–654.
- McCann UD, Szabo Z, Vranesic M, Palermo M, Mathews WB, Ravert HT, Dannals RF, Ricaurte GA (2008) Positron emission tomographic studies of brain dopamine and serotonin transporters in abstinent (+/-)3,4-methylenedioxymethamphetamine (“ecstasy”) users: relationship to cognitive performance. *Psychopharmacology (Berl)* 200: 439–450.
- McIntosh AM, Harrison LK, Forrester K, Lawrie SM, Johnstone EC (2005) Neuropsychological impairments in people with schizophrenia or bipolar disorder and their unaffected relatives. *Br J Psychiatry* 186: 378–385.
- Meneses A, Terrón JA, Hong E (1997) Effects of the 5-HT receptor antagonists GR127935 (5-HT_{1B/1D}) and MDL100907 (5-HT_{2A}) in the consolidation of learning. *Behav Brain Res* 89: 217–223.
- Meneses A (1999) Are 5-HT(1B/1D) and 5-HT(2A/2B/2C) receptors involved in learning and memory processes? *IDrugs* 2: 796–801.
- Meyer-Lindenberg A, Nichols T, Callicott JH, Ding J, Kolachana B, Buckholtz J, Mattay VS, Egan M, Weinberger DR (2006) Impact on complex genetic variation in COMT on human brain function. *Mol Psychiatry* 11: 867–877.
- Miller SA, Dykes D, Plesky HF (1988) A simple salting out procedure for extracting DNA from human nucleated cells. *Nucleic Acids Res* 16: 1215.

- Nocjar C (2002) Localization of 5-HT(2A) receptors on dopamine cells in subnuclei of the midbrain A10 cell group. *Neuroscience* 111: 163–176.
- Okuyama Y, Ishiguro H, Toru M, Arinami T (1999) A genetic polymorphism in the promoter region of DRD4 associated with expression and schizophrenia. *Biochem Biophys Res Commun* 258: 292–295.
- Olvera-Cortés ME, Anguiano-Rodríguez P, López-Vázquez MA, Alfaro JM (2008) Serotonin/dopamine interaction in learning. *Prog Brain Res* 172: 567–602.
- Raghanti MA, Stimpson CD, Marcinkiewicz JL, Erwin JM, Hof PR, Sherwood CC (2008) Differences in cortical serotonergic innervation among humans, chimpanzees, and macaque monkeys: a comparative study. *Cereb Cortex* 18: 584–597.
- Rosa A, Peralta V, Cuesta MJ, Zarzuela A, Serrano F, Martínez-Larrea A, Fañanás L (2004) New evidence of association between COMT gene and prefrontal neurocognitive function in healthy individuals from sibling pairs discordant for psychosis. *Am J Psychiatry* 161: 1110–1112.
- Roth B (2004) Serotonin receptors represent highly favorable molecular targets for cognitive enhancement in schizophrenia and other disorders. *Psychopharmacology* 174: 17–24.
- Rybakowski JK, Borkowska A, Czerski PM, Kapelski P, Dmitrzak-Weglarz M, Hauser J (2005) An association study of dopamine receptors polymorphisms and the Wisconsin Card Sorting Test in schizophrenia. *J Neural Transm* 112: 1575–1582.
- Rybakowski JK, Borkowska A, Czerski PM, Dmitrzak-Weglarz M, Skibinska M, Kapelski P, Hauser J (2006) Performance on the Wisconsin Card Sorting Test in schizophrenia and genes of dopaminergic inactivation (COMT, DAT, NET). *Psychiatry Res* 143: 13–19.
- Sambataro F, Reed JD, Murty VP, Das S, Tan HY, Callicott JH, Weinberger DR, Mattay VS (2009) Catechol-O-methyltransferase valine(158)methionine polymorphism modulates brain networks underlying working memory across adulthood. *Biol Psychiatry* 66: 540–548.
- Sazci A, Ergul E, Kucukali I, Kilic G, Kaya G, Kara I (2004) Catechol-O-Methyltransferase gene Val108/158Met polymorphism and susceptibility to schizophrenia: association is more significant in woman. *Brain Res* 132: 51–56.
- Seamans JK, Yang CR (2004) The principal features and mechanisms of dopamine modulation in the prefrontal cortex. *Prog Neurobiol* 74: 1–58.
- Stefanis NC, Van Os J, Avamopoulos D, Smyrnis M, Evdokimidis I, Hantoumi I, Stefanis CN (2004) Variation in catechol-O-methyltransferase val158met genotype associated with schizotypy: a population study of 543 young men. *Biol Psychiatry* 56: 510–515.
- Stoltenberg SF, Twitchell GR, Hanna GL, Cook EH, Fitzgerald HE, Zucker RA, Little KY (2002) Serotonin transporter promoter polymorphism, peripheral indexes of serotonin function, and personality measures in families with alcoholism. *Am J Med Genet* 114: 230–4.
- Szekeres G, Kéri S, Juhász A (2004) Role of dopamine D3 receptor (DRD3) and dopamine transporter (DAT) polymorphism in cognitive dysfunctions and therapeutic response to atypical antipsychotics in patients with schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* 124: 1–5.
- Uçok A, Alpsan H, Cakir S, Saruhan-Direskeneli G (2007) Association of a serotonin receptor 2A gene polymorphism with cognitive functions in patients with schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* 144: 704–707.
- van Kesteren RE, Spencer GE (2003) The role of neurotransmitters in neurite outgrowth and synapse formation. *Rev Neurosci* 14: 217–231.
- Wang X, Zhong P, Gu Z, Yan Z (2003) Regulation of NMDA receptors by dopamine D4 signaling in prefrontal cortex. *J Neurosci* 23: 9852–9861.
- Xu H, Kallendonk CB, Simpson EH, Keilp JG, Bruder GE, Polan HJ, Kandel ER, Gilliam TC (2007) DRD2 C957T polymorphism interacts with the COMT Val158Met polymorphism in human working memory ability. *Schizophr Res* 90: 104–107.
- Zalla T, Joyce C, Szöke A, Schürhoff F, Pillon B, Komano O, Perez-Diaz F, Bellivier F, Alter C, Dubois B, Rouillon F, Houde O, Leboyer M (2004) Executive dysfunctions as potential markers of familial vulnerability to bipolar disorder and schizophrenia. *Psychiatry Res* 121: 207–217.
- Zhang K, Grady CJ, Tsapakis EM, Andersen SL, Tarazi FI, Baldessarini RJ (2004) Regulation of working memory by dopamine D4 receptor in rats. *Neuropsychopharmacology* 29: 1648–1655.