

## **Informed decision making: an annotated bibliography and systematic review**

H Bekker  
JG Thornton  
CM Airey  
JB Connelly  
J Hewison  
MB Robinson

J Lilleyman  
M MacIntosh  
AJ Maule  
S Michie  
AD Pearman



---

**Health Technology Assessment  
NHS R&D HTA Programme**



# Standing Group on Health Technology

## Current members

---

### Chair:

**Professor Sir Miles Irving,**  
Professor of Surgery, University  
of Manchester, Hope Hospital,  
Salford

Professor Martin Buxton,  
Professor of Economics,  
Brunel University

Professor Francis Creed,  
School of Psychiatry  
& Behaviour Sciences,  
University of Manchester

Professor Charles Florey,  
Department of Epidemiology  
& Public Health, Ninewells  
Hospital & Medical School,  
University of Dundee

Professor John Gabbay,  
Director, Wessex Institute for  
Health Research & Development

Professor Sir John  
Grimley Evans,  
Department of  
Geriatric Medicine,  
Radcliffe Infirmary, Oxford

Dr Tony Hope,  
The Medical School,  
University of Oxford

Professor Richard Lilford,  
Regional Director, R&D,  
West Midlands

Dr Jeremy Metters,  
Deputy Chief Medical Officer,  
Department of Health

Professor Maggie Pearson,  
Regional Director of R&D,  
NHS Executive North West

Mr Hugh Ross,  
Chief Executive, The United  
Bristol Healthcare NHS Trust

Professor Trevor Sheldon,  
Director, NHS Centre for  
Reviews & Dissemination,  
University of York

Professor Mike Smith,  
Director, The Research  
School of Medicine,  
University of Leeds

Dr John Tripp,  
Department of Child Health,  
Royal Devon & Exeter  
Healthcare NHS Trust

Professor Tom Walley,  
Department of  
Pharmacological Therapeutics,  
University of Liverpool

Dr Julie Woodin,  
Chief Executive,  
Nottingham Health Authority

Professor Kent Woods  
(**Chair Designate**),  
Regional Director of R&D,  
NHS Executive Trent

## Past members

---

Dr Sheila Adam,  
Department of Health

Professor Angela Coulter,  
Director, King's Fund, London

Professor Anthony Culyer,  
Deputy Vice-Chancellor,  
University of York

Dr Peter Doyle,  
Executive Director, Zeneca Ltd,  
ACOST Committee on Medical  
Research & Health

Professor John Farndon,  
Professor of Surgery,  
University of Bristol

Professor Howard  
Glennester,  
Professor of Social Science  
& Administration, London  
School of Economics &  
Political Science

Mr John H James,  
Chief Executive,  
Kensington, Chelsea &  
Westminster Health Authority

Professor Michael Maisey,  
Professor of Radiological  
Sciences, Guy's, King's & St  
Thomas's School of Medicine  
& Dentistry, London

Mrs Gloria Oates,  
Chief Executive,  
Oldham NHS Trust

Dr George Poste,  
Chief Science & Technology  
Officer, SmithKline Beecham

Professor Michael Rawlins,  
Wolfson Unit of  
Clinical Pharmacology,  
University of Newcastle-  
upon-Tyne

Professor Martin Roland,  
Professor of General Practice,  
University of Manchester

Professor Ian Russell,  
Department of Health Sciences  
& Clinical Evaluation,  
University of York

Dr Charles Swan,  
Consultant Gastroenterologist,  
North Staffordshire  
Royal Infirmary

Details of the membership of the HTA panels, the NCCHTA Advisory Group and the HTA Commissioning Board are given at the end of this report.



**INAHTA**

### **How to obtain copies of this and other HTA Programme reports.**

An electronic version of this publication, in Adobe Acrobat format, is available for downloading free of charge for personal use from the HTA website (<http://www.hta.ac.uk>). A fully searchable CD-ROM is also available (see below).

Printed copies of HTA monographs cost £20 each (post and packing free in the UK) to both public **and** private sector purchasers from our Despatch Agents.

Non-UK purchasers will have to pay a small fee for post and packing. For European countries the cost is £2 per monograph and for the rest of the world £3 per monograph.

You can order HTA monographs from our Despatch Agents:

- fax (with **credit card** or **official purchase order**)
- post (with **credit card** or **official purchase order** or **cheque**)
- phone during office hours (**credit card** only).

Additionally the HTA website allows you **either** to pay securely by credit card **or** to print out your order and then post or fax it.

### **Contact details are as follows:**

HTA Despatch  
c/o Direct Mail Works Ltd  
4 Oakwood Business Centre  
Downley, HAVANT PO9 2NP, UK

Email: [orders@hta.ac.uk](mailto:orders@hta.ac.uk)  
Tel: 02392 492 000  
Fax: 02392 478 555  
Fax from outside the UK: +44 2392 478 555

NHS libraries can subscribe free of charge. Public libraries can subscribe at a very reduced cost of £100 for each volume (normally comprising 30–40 titles). The commercial subscription rate is £300 per volume. Please see our website for details. Subscriptions can only be purchased for the current or forthcoming volume.

### **Payment methods**

#### *Paying by cheque*

If you pay by cheque, the cheque must be in **pounds sterling**, made payable to *Direct Mail Works Ltd* and drawn on a bank with a UK address.

#### *Paying by credit card*

The following cards are accepted by phone, fax, post or via the website ordering pages: Delta, Eurocard, Mastercard, Solo, Switch and Visa. We advise against sending credit card details in a plain email.

#### *Paying by official purchase order*

You can post or fax these, but they must be from public bodies (i.e. NHS or universities) within the UK. We cannot at present accept purchase orders from commercial companies or from outside the UK.

### **How do I get a copy of HTA on CD?**

Please use the form on the HTA website ([www.hta.ac.uk/htacd.htm](http://www.hta.ac.uk/htacd.htm)). Or contact Direct Mail Works (see contact details above) by email, post, fax or phone. *HTA on CD* is currently free of charge worldwide.

---

The website also provides information about the HTA Programme and lists the membership of the various committees.



# **Informed decision making: an annotated bibliography and systematic review**

H Bekker<sup>1</sup>

JG Thornton<sup>2</sup>

CM Airey<sup>3</sup>

JB Connelly<sup>3</sup>

J Hewison<sup>1</sup>

MB Robinson<sup>3</sup>

J Lilleyman<sup>2</sup>

M MacIntosh<sup>2</sup>

AJ Maule<sup>4</sup>

S Michie<sup>5</sup>

AD Pearman<sup>4</sup>

<sup>1</sup> School of Psychology, University of Leeds, UK

<sup>2</sup> Centre for Reproduction Growth and Development,  
University of Leeds, UK

<sup>3</sup> Nuffield Institute, University of Leeds, UK

<sup>4</sup> Business School, University of Leeds, UK

<sup>5</sup> Psychology and Genetics Research Group, UMDS, London, UK

Published February 1999

---

This report should be referenced as follows:

Bekker H, Thornton JG, Airey CM, Connelly JB, Hewison J, Robinson MB, *et al.* Informed decision making: an annotated bibliography and systematic review. *Health Technol Assess* 1999;**3**(1).

*Health Technology Assessment* is indexed in *Index Medicus/MEDLINE* and *Excerpta Medica/EMBASE*. Copies of the Executive Summaries are available from the NCCHTA web site (see overleaf).

# NHS R&D HTA Programme

The overall aim of the NHS R&D Health Technology Assessment (HTA) programme is to ensure that high-quality research information on the costs, effectiveness and broader impact of health technologies is produced in the most efficient way for those who use, manage and work in the NHS. Research is undertaken in those areas where the evidence will lead to the greatest benefits to patients, either through improved patient outcomes or the most efficient use of NHS resources.

The Standing Group on Health Technology advises on national priorities for health technology assessment. Six advisory panels assist the Standing Group in identifying and prioritising projects. These priorities are then considered by the HTA Commissioning Board supported by the National Coordinating Centre for HTA (NCCHTA).

This report is one of a series covering acute care, diagnostics and imaging, methodology, pharmaceuticals, population screening, and primary and community care. It was identified as a priority by the Screening Panel and funded as project number 94/27/03.

The views expressed in this publication are those of the authors and not necessarily those of the Standing Group, the Commissioning Board, the Panel members or the Department of Health. The editors wish to emphasise that funding and publication of this research by the NHS should not be taken as implicit support for the recommendations for policy contained herein. In particular, policy options in the area of screening will be considered by the National Screening Committee. This Committee, chaired by the Chief Medical Officer, will take into account the views expressed here, further available evidence and other relevant considerations.

Reviews in *Health Technology Assessment* are termed 'systematic' when the account of the search, appraisal and synthesis methods (to minimise biases and random errors) would, in theory, permit the replication of the review by others.

Series Editors: Andrew Stevens, Ruairidh Milne and Ken Stein  
Editorial Assistant: Melanie Corris

The editors have tried to ensure the accuracy of this report but cannot accept responsibility for any errors or omissions. They would like to thank the referees for their constructive comments on the draft document.

ISSN 1366-5278

© Crown copyright 1999

Enquiries relating to copyright should be addressed to the NCCHTA (see address given below).

Published by Core Research, Alton, on behalf of the NCCHTA.

Printed on acid-free paper in the UK by The Basingstoke Press, Basingstoke.

---

Copies of this report can be obtained from:

The National Coordinating Centre for Health Technology Assessment,  
Mailpoint 728, Boldrewood,  
University of Southampton,  
Southampton, SO16 7PX, UK.  
Fax: +44 (0) 1703 595 639 Email: [hta@soton.ac.uk](mailto:hta@soton.ac.uk)  
<http://www.soton.ac.uk/~hta>



# Contents

<b>List of abbreviations</b> .....	i	<b>References</b> .....	33
<b>Executive summary</b> .....	iii	<b>Appendix 1</b> References used to develop the search strategy .....	35
<b>1 Introduction</b> .....	1	<b>Appendix 2</b> The principle of the search strategy .....	37
Factors associated with informed decision making .....	1	<b>Appendix 3</b> MEDLINE search strategy .....	39
Measures of informed decision making .....	2	<b>Appendix 4</b> Data extraction coding forms ..	41
Decision-making theory .....	2	<b>Appendix 5</b> Bibliography: studies by study number .....	43
Aims of review .....	3	<b>Appendix 6</b> Bibliography: studies by author .....	67
<b>2 Methods</b> .....	5	<b>Appendix 7</b> Decision details by study design (grouped by health area) .....	81
Search strategy .....	5	<b>Appendix 8</b> Intervention details by study design (grouped by health area in same order as appendix 7) .....	125
Inclusion criteria .....	5	<b>Health Technology Assessment reports published to date</b> .....	151
Data extraction .....	6	<b>Health Technology Assessment panel membership</b> .....	153
Data synthesis .....	8		
<b>3 Results</b> .....	11		
Summaries of extracted information by study .....	11		
Summary by frequency tables .....	11		
Summary by study number listings .....	14		
Synthesising findings .....	14		
<b>4 Discussion</b> .....	29		
Evaluating the methods .....	29		
Overview of results .....	29		
Suggested applications of the review .....	31		
Recommendations for research .....	32		







## List of abbreviations

AA	Alcoholics Anonymous*
ACS	American Cancer Society*
A&E	accident and emergency
BPH	benign prostatic hypertrophy
BSE	breast self examination*
CF	cystic fibrosis*
CPR	cardiopulmonary resuscitation*
CVD	cardiovascular disease*
FOB	faecal occult blood*
GP	general practitioner*
HBM	health belief model*
HCP	healthcare professional
HIV	human immunodeficiency virus
HRT	hormone replacement therapy*
RCT	randomised controlled trial (In this study a, b and c are used to denote RCTs with a low risk of bias, an unknown risk of bias, and a high risk of bias, respectively.)
SLT	social learning theory
STD	sexually transmitted disease
TB	tuberculosis
TRA	theory of reasoned action
VI	verbal information
WI	written information

\* Used only in appendices





## Executive summary

### Background

Everyone makes decisions about their health, and many healthcare interventions aim to encourage this. An informed decision is one where a reasoned choice is made by a reasonable individual using relevant information about the advantages and disadvantages of all the possible courses of action, in accord with the individual's beliefs.

### Objectives

- To provide an unbiased bibliography of controlled studies evaluating interventions that may affect informed patient decision making.
- To classify studies by research design, decision-making theory, type of intervention and health setting.
- To describe the measures of informed patient decision making and other outcomes reported.
- To identify under-reported areas and direct future research.

### Methods

#### Data sources

The electronic databases MEDLINE, BIDS (social science), and PsycLIT were searched for 1991–96. The journals *Medical Decision Making*, *Patient Education and Counseling*, and *Preventive Medicine* were handsearched for 1986–96.

#### Inclusion criteria

Studies were included if they reported the results of a controlled study of any intervention using real patients making a health decision. Specifically, randomised controlled trials (RCTs), non-randomised concurrent studies, historical studies, and same subject 'before and after' studies were included. Health decisions were defined to include any reported health behaviour change as well as explicit decisions. Interventions were defined broadly to include any that could reasonably be expected to affect informed decision making, such as changes in information provision, cost, or service provision. Patients were defined as any individual making a decision about health care. Experimental studies on healthy student volunteers, studies of health

professionals making decisions about another individual's care, and articles not published in English were excluded.

#### Data extraction

This was performed using coding forms by a member of the project group and checked by a team member, with disagreements resolved by discussion. Abstracts were assessed and the articles retrieved if the review criteria were met. Final inclusion decisions were made by the first author of this report and verified by another member of the project group.

Articles were classified by study quality according to the hierarchy of evidence, underlying theory, the domain of health care, and the health decision. The comparison groups, other factors associated with the decision-making process, reported measures, and a summary of the findings were recorded.

#### Data synthesis

Descriptive summaries and qualitative analysis were performed. The health domains and decisions were too diverse for meaningful quantitative meta-analysis.

### Results

Following handsearching and abstract evaluation 825 articles were distributed to the project group members and 547 were subsequently included within the bibliography.

#### Study quality

There were 336 RCTs, 114 non-randomised concurrent studies, 34 historical, and 63 'before and after' same-sample studies. Only 51 of the RCTs were classified as having a low risk of bias. A total of 267 studies claimed to have approached a representative sample of participants, but only 243 reported the number invited to take part. Few studies provided adequate descriptions of the intervention materials.

#### Theoretical context

A total of 206 studies referred to an underlying theory. Of these, 101 referred to theories explain-

ing decision making such as expected utility theory, prospect theory or social cognition models.

### Health domain and the decision

A total of 251 studies were in general medicine, 114 cancer, 108 genitourinary medicine, 61 primary care, 31 paediatrics; 15 mental health, 10 dentistry, 11 surgery, seven genetics, and 31 obstetrics and gynaecology, and midwifery. The decision was classified as a life-style change in 357 studies, a screening decision in 114, a treatment decision in 107, a decision to participate in the consultation in 51, and as another type of decision in 26 studies.

### Interventions

A total of 301 interventions were of information provision itself, 273 varied the delivery of information, 208 provided patient feedback, 94 manipulated information in some other way, 55 prompted active patient participation, and 89 of another intervention altogether.

### Decision-making factors

A total of 512 studies assessed actual rather than hypothetical decisions, 476 involved decisions affecting the participant rather than a third party and in 525 studies the decision was made without time pressure. Only 26 studies explicitly made patients aware of their involvement in the decision-making process.

### Measures assessed

Demographic details were recorded in 515 studies, knowledge in 181, decision-making measures in 169, measures of affect in 69, satisfaction in 60, self-efficacy in 75, personality trait in 20 and other variables in 111 studies.

### Summary result

Only five studies were **theory** driven, assessed measures associated with **informed** decision making, **and** used a low risk of bias design. Although of disparate design these five studies suggest that information and education are relatively ineffective ways of facilitating informed decision making, compared with the context and social influences. Studies reporting manipulation of information, and provision of feedback, were the most likely to report an effect.

## Conclusions

There is a paucity of well-designed, theoretically driven and adequately operationalised research assessing informed patient decision making. Given the small number of high-quality studies and the relatively slow increase in research in this area there is no need for the NHS to revisit this topic as a review for 5 years. Resources should be concentrated on better primary research.

### Recommendations for research

Future primary research should work under an explicit theory of decision making, record process measures to permit evaluation of whether the decision was informed, and if evaluating experimental interventions use randomised trials with a low risk of bias.

A booklet describing the main decision-making theories, and an inventory of suitable outcome measures could be developed to help clinical researchers design appropriate studies.

Complementary systematic reviews would be valuable.

- The effect of interventions on patient preferences. (At least 50 trials were excluded from the present bibliography because no behaviour change was recorded.)
- Observational studies of real patient decision making. Studies using tape-recorded consultations, verbal thinking aloud protocols, and other written or computer-based process-tracing methods will predominate.
- Assessing the effect of additional information, manipulation of information, provision of feedback, and group delivery of information on informed patient decision making.

Primary research is a priority in areas such as genetics, prenatal diagnosis and where decisions are often made by proxy, such as paediatrics and mental health.

Primary research is required to evaluate the following types of interventions:

- decision aids, such as graphical and computer-based devices
- information manipulation, such as decision analysis, prompts, and feedback.

# Chapter I

## Introduction

Everyone makes decisions about their health care, such as whether to brush their teeth, smoke, or take headache pills, and some have to make difficult decisions, such as whether to undergo a prenatal diagnosis test, or choosing between chemotherapy or surgery for cancer. Generally people want to make these decisions for themselves and to be given the information required to make them wisely. Many healthcare interventions aim to encourage this.

However, there is no agreed definition of an informed decision because it depends on the theoretical predisposition of the researcher. For some an informed decision is one where the final choice accords with an underlying theory, such as expected utility theory. For others it is the process that matters, and whether this follows a particular theory, that defines an informed decision. The two approaches cannot easily be reconciled, and the following is a compromise definition: an informed decision is one where a reasoned choice is made by a reasonable individual, using relevant information about the advantages and disadvantages of all the possible courses of action, in accord with the individual's beliefs. In practice individuals rarely make decisions this way. They have a limited processing capacity and overcome these by using modes of thinking and reasoning that simplify the issues. These heuristic modes of thinking determine which aspects are attended to and how they are interpreted.

Several models and theories have been developed to explain how individuals make decisions and how decision making may be facilitated. Some of the factors that affect an individual's ability to make an informed decision and theories of decision making are discussed in more detail below.

### Factors associated with informed decision making

The factors associated with an individual's ability to make an informed health decision can be grouped into those pertaining to a) the decision context, b) the decision maker, and c) other influences. Only the last are directly alterable.

### Decision context

The following features of the health decision suggest that different psychological evaluations are required, which may alter the extent to which an individual makes an informed decision.

- **Type of health decision**, for example, smoking or drinking alcohol; attending a physician appointment; adhering to medication; having a diagnostic test; participating in choice of treatment.
- **Seriousness of the outcome**, for example, choosing to take a headache pill, deciding whether to donate an organ or not.
- **Familiarity with the decision**, for example, deciding whether or not to exercise is more commonly experienced by people than deciding whether or not to have a genetic test.
- **Level of certainty**, for example, declining insulin **always** makes diabetic people ill, whereas declining prenatal screening only carries a **risk** of giving birth to a Down's syndrome child.
- **Health domain**, for example, making choices in medicine, in surgery or in primary care.
- **Recipient**, for example, deciding for yourself or for your child.

### Decision maker

People vary in their preferred degree of involvement in health decisions.<sup>1</sup> Some want all possible information whereas others are happy to rely on a physician's recommendation or a more intuitive, 'gut reaction'. Personality traits, need for control, beliefs about the doctor-patient role, current state of health or illness, and the degree of anxiety or depression and an individual's ability to understand health information vary widely.<sup>2</sup> It is likely that these individual differences will affect the degree to which informed decisions are made.

### Other influences

Individuals are often unable to deal systematically with large amounts of information, and employ heuristics to reduce the processing required. These often result in decisions being made from the context rather than the content of the information. Decisions may change when the same factual information is presented in slightly different ways, for example, by **framing** the information either positively or negatively, expressing figures as

percentages, risk, inverse risk, or verbally.<sup>3,4</sup> The context also affects whether a reasoned decision is made. For example, individuals under time pressure or experiencing extreme affect (e.g. frightened or angry), are more likely to use an heuristic, whereas those who perceive a need to justify the decision outcome or who are only moderately anxious are more likely to use all the information.

Decision aids such as memory prompts, repeating information, reducing the quantity or improving the clarity of information, and presenting information in different mediums may also affect decision making. It is likely that extra relevant prompt information will reduce the **cognitive load** and promote informed decision making. However, too much information may make people rely on an established heuristic and lead to worse decisions.

Finally, extraneous factors such as charging for a service, the availability of transport, the shape of an inhaler or the presence of a no-smoking policy will alter an individual's evaluation of the available information.

## Measures of informed decision making

The most frequently used measures of informed decision making and their limitations are discussed below.

- Noting the **health behaviour** indicates a decision was made but no more than this.
- **Knowledge** assesses the individual's ability to recall information. Although, better-informed decision makers probably recall more, this does not mean the information was used.
- **Utilities** measure individual's values or attitudes towards the choices available rather than the decision outcome. Proponents of expected utility theory argue that a good decision is one consistent with the choice derived from an equation that integrates an individual's values for each outcome with the likelihood of the outcome occurring. As an informed decision should accord with a patient's attitudes, it is likely that an informed decision will be consistent with a decision derived from expected utility theory. However, deciding in accord with personal utilities is not sufficient for a decision to be classified as informed.
- **Affect** measures anxiety, satisfaction and regret, which are often reported. Although reduced affect may be desirable it is not necessarily associated with informed decision making.

For example, moderately raised anxiety is associated with a more systematic evaluation of the information but great anxiety is associated with the application of heuristics.<sup>5</sup> Equally an individual is as likely to be satisfied with health care if the decision was made following a health professional's recommendation or after an in-depth evaluation of all the available information.

Most of these measures focus on the outcomes following the making of the health decision. It is likely that the effects of memory and the need to continue with the same healthcare team will affect their validity. Equally, few of these measures evaluate the decision-making process, that is, the type and way information was used by the individual when making the decision. It is also necessary to ensure that the final decision was based on an unbiased evaluation of all the available information including the risks and alternatives. A number of **process-tracing** techniques are available. These include: verbal protocols; analysis of tape-recorded consultations following the application of a technique like content analysis to the transcripts; and the use of information boards or computer programmes to track the information used.

Perhaps because of these issues of measurement, little is known about the impact of encouraging informed decision making on the individual. There is some evidence to suggest that too much evaluation of the available information leads to reduced post-decision satisfaction.<sup>6</sup> However, the application of heuristics is more likely to lead to poorer outcomes than adequately surveying all the alternatives.<sup>7</sup> Such conflicts can be resolved only in the light of an underlying theory.

## Decision-making theory

There are three broad types of decision-making theory.<sup>8</sup> The first, **normative theory** describes what people ought to do if they wish to be rational decision makers. These axiomatic theories are based on mathematical and statistical proofs.<sup>9</sup> The most important, classical decision theory, assumes that people are perfectly informed, and are familiar with all the alternatives that are available to them, as well as their beliefs and preferences associated with these alternatives. The rational or best course of action is the one that maximises expected utility.<sup>7</sup>

**Descriptive theories**, in contrast, describe how people actually make decisions.<sup>10</sup> At least since

the early 1950s it has been clear that people rarely do this in accord with normative theories.<sup>11</sup> Simon suggested that because of their limited processing capacity, or bounded rationality, individuals adopted more simplistic modes of thinking that are generally good, but can lead to poor decision making.<sup>12,13</sup>

The third type of decision-making theory, **prescriptive theory**, recognises that human beings can be poor decision makers and is concerned with the development of decision aids to help them.<sup>14</sup> Early approaches were based on a modified normative theory, taking into consideration evidence from how people actually make decisions unaided. For example, social cognition models<sup>15</sup> and prospect theory<sup>4</sup> maintain that an individual's behaviour is logically related to their values and beliefs. However, by describing the underlying cognitive processes such as attitudes, biases or schemas, they permit the development of more effective decision aids.

An understanding of informed decision making in patients depends, in part, on knowing how patients currently make decisions (descriptive theory) and how they can be assisted to make better decisions (prescriptive theory). There are many descriptive theories relevant to this issue. One concerns the simplistic modes of thinking (heuristics) used when judging risk and uncertainty,<sup>7,12,16</sup> which can lead to misperceptions of risk and poor decisions. Similarly framing, the way a problem is presented, for example, emphasising gains rather than losses, affects risk-taking propensity (preference for a safe over an equivalent risky option).<sup>4,17</sup> Another sub-optimal way in which people tend to gather and interpret information is illustrated by the confirmation bias, involving the selective processing of information that confirms initial expectations.<sup>7</sup> Both professionals and the lay-population are subject to these simplistic modes of thinking.<sup>18,19</sup>

People's commitments to different courses of action are affected by the procedure used to establish their preferences. They often judge one alternative as more attractive but select another when asked to choose between them.<sup>16</sup> This suggests that preferences are not necessarily a good guide to final choices, and challenges the view that people have stable preferences for decision outcomes.

People adopt a range of different underlying information processing strategies for different problems, with each strategy differentiated in terms of the mental effort required and the

accuracy/quality of the decision.<sup>13</sup> Many factors, such as fatigue, time pressure, stress and other emotional states, and lack of engagement with the problem, induce the use of simpler sub-optimal strategies.<sup>5,20,21</sup> Similarly the school of naturalistic decision making has shown that people make decisions in very different ways when they have expertise and experience to draw on.<sup>23</sup>

Not surprisingly such theories have given rise to many different ways to improve decision making. One approach has involved specifying a **structured sequence** of activities similar to those specified by normative theories, for example, decision analysis and the use of decision trees for risky decisions, or Analytical Hierarchy Process for riskless ones.<sup>14,24</sup> A second approach, **scenario planning**<sup>25</sup> has been developed for strategic decisions involving outcomes that occur only in the medium-to-long term, and which cannot be ascribed probabilities. This involves developing positive and negative scenarios that describe how the future could unfold, and choosing the action that is best, regardless of which does occur. A third approach involves training people to reason more effectively such that they follow rules based on formal models from statistics, logic and probability theory.<sup>26</sup> All such approaches recognise that different types of decision require different types of prescriptive solution.<sup>27</sup>

Taken together, this body of theory provides a rich source of information for judging patient decision making, and the development of evidence-based medicine has provided the impetus for a critical evaluation of decision aids. Interventions aimed at supporting patient decision making should be developed in the context of these bodies of theory, and a primary function of this review is to assess the extent to which the published literature on informed decision making in patients achieves this. In addition, this body of knowledge provides a base for evaluating the existing research on informed decision making, and to determine whether the implicit and explicit models of decision making underlying this research are coherent and appropriate.

## Aims of review

The aim is to provide a comprehensive, annotated bibliography of controlled studies of interventions that might plausibly inform patient healthcare decision making. Studies have been classified by the hierarchy of evidence and by theoretical context. Information about the type of intervention,

health setting, measures reported and main research findings are recorded. The bibliography provides a summary of the evidence describing patient decision making, factors associated with decision making, evaluated interventions and theory-driven research. Such a review will enable identification of under-reported areas and aim to provide answers to the following questions.

- Can decision making be facilitated?
- Do people who have more information make better decisions?
- Does the way in which we present information change the effectiveness of the decision?
- Does the context affect decision making?
- Does the effectiveness of information on decision making vary by medical setting?



# Chapter 2

## Methods

### Search strategy

Studies considered for inclusion within the systematic review were first identified from abstracts generated from one of two sources: electronic databases and handsearches of complete sets of journals. Articles were retrieved if the review criteria were met or the abstract contained insufficient information to assess the review criteria.

### Electronic databases

Over 6 months a search strategy was developed for the electronic databases MEDLINE, PsycLIT, and BIDS (social science). A pilot strategy had been developed for inclusion within the initial protocol from a set of keywords chosen by the project group. Additional keywords were included within the strategy derived from: 17 articles considered relevant to the field of informed patient decision making (appendix 1); and 100 articles considered for inclusion within the review following application of the pilot strategy to MEDLINE and handsearching of the journal *Medical Decision Making*. The keywords of the final strategy were modified and expanded within each of the electronic databases by the research officer experienced in electronic searches and systematic reviews (appendix 2). The final strategy included terms from three categories: decision making field or healthcare users and comparative study designs (appendix 3).

The sensitivity of the search strategy was evaluated by comparing the number of articles identified by handsearching a second journal, *Patient Education and Counseling*. The pilot strategy identified 7% of articles considered for inclusion while the final strategy identified 62%. This sensitivity figure was considered satisfactory as electronic searches identify between 17–82% of relevant articles.<sup>28</sup> However, these efforts to increase sensitivity resulted in significantly reducing the specificity of the final strategy, as 17,860 abstracts were generated for the years 1991–96 alone. Consequently, the application of the strategy was limited to articles published between these years to ensure completion of a review considering the resources available.

### Handsearching

Three journals were selected for handsearching because of their relevance to the healthcare users decision making area: *Medical Decision Making*, *Patient Education and Counseling*, and *Preventive Medicine*. In the handsearch, every abstract within every volume was read and considered for inclusion. All volumes were handsearched for the years 1986–96 inclusive.

### Inclusion criteria

The review inclusion criteria were developed over 4 months, during which over 300 abstracts downloaded from the pilot MEDLINE and PsycLIT search strategies were circulated to all project group members. At monthly meetings, each group member fed back the inclusion or exclusion status of each abstract. When members disagreed a discussion took place until a set of working rules was developed. At the end of this development phase, agreement between group members on the inclusion and exclusion criteria was reached. Nevertheless, in this area there are no pre-defined or obvious boundaries to be used for selection criteria, and some of the cut-off categories may appear arbitrary. For inclusion, each study published within an article had to fulfil the following criteria.

- Study participants must be patients. A patient was defined as any individual making a health decision. Studies using university student participants were excluded. Studies assessing health professional decision making about another individual's care were excluded.
- A study must report a behavioural measure of the health decision made. This measure of reported or observed behaviour assesses decision making, a more inclusive category than informed decisions. This broad health behaviour outcome measure incorporates many decisions including: smoking; adherence to medication; attendance for screening; and choices between treatments like chemotherapy or surgery. The issue of whether to include such lifestyle decisions was debated extensively by the group. Some members considered these to be often affected more by other factors than the inform-

ation and decision process itself; others argued that this is only a qualitative difference from other health decisions and an arbitrary line would have to be drawn if they were to be excluded. We decided to include such studies. The health behaviour included real, in the sense that an event happened, intended or hypothetical outcomes. Studies were excluded if the health behaviour measure assessed the effectiveness of a pharmacological treatment such as one comparing the effects of placebo gum with nicotine gum on smoking cessation. Additionally, studies that did not assess actual or reported behaviour were excluded, such as those that only reported biophysiological measures, knowledge, satisfaction, preferences or utilities, and changes in affect.

- A study must assess an experimental intervention. The definition of an experimental study was interpreted broadly to include any design that evaluated an experimental group with a comparison group. Studies with one of six designs were included: randomised controlled trial (RCT) with a low risk of bias (RCTa); RCT with an unknown risk of bias (RCTb); RCT with a high risk of bias (RCTc); non-randomised concurrent; historical studies or 'before and after' intervention with different samples; and, 'before and after' intervention with the same sample. Studies assessing predictors of a behaviour such as factors associated with breast screening attendance, were excluded. The definition of an intervention was broad to incorporate the many factors that may alter decision making: offering additional information; framing figures in different ways; patient use of a decision aid; comparison of information mediums; offers of free transportation; introduction of fluoride to water supplies; and so on.

- The study must be published in English.

All abstracts generated from the search strategies were assessed for inclusion by the first author of this report. Articles were retrieved if the review criteria were met or the abstract contained insufficient information. Batches of ten articles were sent in rotation to each member of the project group. The inclusion criteria were again assessed by that member of the project group and, if included, a data extraction coding form was completed for each study of each article. Completed forms were returned to the first author of this report and checked. Classification disagreements were resolved by discussion. All review details and extracted information were entered into the project electronic database.

## Data extraction

To ensure consistent information was extracted from each study of each article, a data extraction sheet or coding form was developed over a 6-month period (appendix 4). The categories within the coding form were informed with reference to the decision making and integration of research literature, and the expertise of members of the project group. Each draft coding frame was piloted on articles derived from the search strategies and modifications agreed within the group monthly meetings. Following piloting, several categories were simplified or omitted because it proved impossible to achieve consistency between articles, as the information published was not described or presented in a format compatible with the review's research question.

There follows a brief explanation of the information extracted from each article.

### Study identification number

Each study was allocated a unique number. If more than one study was reported in one article, each study was given a separate identification number and coding form. If a study was published more than once in different articles, this too was treated as a separate study.

### Location of research

This meant the country where the study was undertaken.

### Domain of health care

The area of health in which the decision was made was recorded. Responses within this free-text box were later categorised into 11 medical domains: dentistry; cancer; general medicine and infectious diseases; HIV and sexually transmitted diseases (STDs); mental health; paediatrics; genetics; obstetrics and gynaecology, and midwifery; surgery and accident and emergency (A&E); primary care; other (alternative medicine, marital counselling).

### Participants

The person making the health decision whom may or may not receive the health care was recorded.

### Age groups

Participants were classified as child, adult, or elderly by the reviewer.

### Health status

Participants were classified by the reviewer as well, physically impaired, or mentally impaired by the reviewer. For example, participants attending for health promotion programmes were classified as

‘well’; athletes foot, cancer follow-up, and surgery were classified as ‘physically impaired’; depression, anxiety, and schizophrenia were classified as ‘mentally impaired’. If the decision maker was a spouse or relative, the health status recorded was that of the person making the decision not the recipient of the decision. So, the decision maker carer of a person with Alzheimer’s disease was classified as ‘well’.

### Decision associated factor 1

Decisions were classified as real, hypothetical or intended. Real decisions were those health actions carried out by the participant such as attending for mammography or using an inhaler. Hypothetical and intended decisions were later combined and included decisions made based on an option becoming available in the future, a vignette based on imaginary decisions and intentions about future behaviours. For example, a non-pregnant woman making a decision to have a test for Down’s syndrome in a future pregnancy, or an individual intending to use a condom during the next sexual encounter.

### Decision associated factor 2

Decisions were classified as implicit or explicit according to whether the researchers and decision makers were aware that they were making a health decision.

### Decision associated factor 3

Studies were classified into those where the decision maker was the recipient of the decision outcome or those where the recipient was a third party. For example, a decision to be immunised compared with having a child immunised. Some decisions were more difficult to classify and minor rules were established: decisions made by pregnant women were classified as participant even though decisions about amniocentesis and termination would impact on a foetus; and use of condoms was said to affect both the participant and other.

### Decision associated factor 4

Studies were classified into those where the decision was time restrained, made at the same time as the intervention, and those that were not time restrained, made some while after the intervention. The important difference being whether the participant was given the opportunity to reflect on, and change, the decision.

### Health decision

A free-text box was provided to write the type of health decision made. These decision types were then classified into one of six decision categories:

- to use a health service
- to make a lifestyle change
- to attend for testing (screening and diagnostic)
- to adhere to treatment regimens
- to actively participate within the consultation (e.g. question asking and research)
- other (e.g. organ donation and willingness to pay).

### Theoretical context

This noted the authors’ references to a theoretical framework that informed the study design or intervention. Any type of theory alluded to was noted in the free-text box and subsequently classified into one of the following four categories:

- no theory
- explanation decision-making theory (social cognition models, prospect theory and expected utility theory)
- health-promotion theories (social learning theory and skill acquisition theories)
- other psychological models (illness representation, attribution theory and stress coping models).

The project group did not assess the adequacy of the operationalisation of theories nor infer whether the research was theory driven if the authors did not explicitly mention a theory.

### Measures described

This table was used to record any measure described within the published paper other than the health behaviour outcome measure. As noted in the introduction, several measures have been used to assess informed patient decision making, and others are associated with the process of decision making. A free-text box was used to list all the measures the authors described. Before piloting, the project group aimed to note the type of analysis associated with each measure. However, little consistency was achieved across papers in the type of measures used and the reporting of results. Subsequently, only the description of the type of measures referred to was classified and grouped as follows:

- **demographic variables:** age, sex, level of education, social class, marital status, religion
- **personal history:** sexual orientation, medical history, law infringements, health insurance, reproductive history, physiological assessment
- **decision-making process measures:** attitudes, utilities, reasons, preferences, variables of operationalised models, perception of risk, perception of severity, autonomy, intention, motivation

- **efficacy:** efficacy and self-efficacy
- **knowledge measures**
- **satisfaction:** usefulness of information, satisfaction with information
- **affect:** anxiety, depression, regret, worry
- **personality factors:** need for cognition, need for cognitive closure, locus of control
- **other variables:** perception of well being, perception of disability, social support, alienation, memory, willingness to pay.

### Intervention groups

This table provided a description of the main components of each group assessed within the experimental intervention. It was assumed that a routine health encounter would include the provision of verbal information by a health professional to a patient. Any variation from this norm was noted in the free-text box. For example, comparison group: routine care; Group 1: routine care plus leaflet; Group 2: routine care plus leaflet plus free test. The main components were subsequently categorised within the following groups:

- additional information (leaflet, video, computer, posters)
- group-delivery information (media, classroom based, group counselling)
- information manipulation (framing, directiveness, more/less information)
- patient-prompted information (decision aid, memory prompt, pre-consultation rehearsal questions)
- service change (additional contact by health professional, free service)
- environmental change (introduction of no-smoking policy, shops altering food displays)
- other (incentives, alternative treatment offered).

### Quality of the study 1

Study designs were classified according to the hierarchy of evidence (Cochrane handbook) as follows:

- RCTa (e.g. randomisation from serially numbered, opaque sealed envelopes, third party or computer; low risk of bias)
- RCTb (e.g. unclear how randomised; unknown risk of bias)
- RCTc (e.g. tossed coins or approached patients on alternate days; high risk of bias)
- non-randomised concurrent (e.g. intervention community compared with comparative community)
- non-randomised historic (e.g. assessed uptake of cervical screening before and after remuneration of doctors)

- ‘before and after’ studies with same sample (e.g. assessed cessation smoking before and after a workplace intervention).

### Quality of the study 2

This assessed the level of intervention. Studies were classified into those where the unit of intervention was the individual patient, and those where some other unit of allocation such as hospitals, communities, or patients allocated by health professional.

### Quality of the study 3

Where possible, details of the sample approached were recorded in order to collect details on the representativeness of the sample. First, whether the sample invited to participate included: all available participants; a systematic sample; a volunteer or convenience sample; or, not adequately described. Second, the actual number of participants was recorded: those who were invited to participate, those who participated, and those who were included within the final analysis. The latter was often difficult to extract from the published information.

### Quality of the study 4

A free-text box was provided for reviewers to highlight additional causes for concern, such as inappropriate use of statistics, confounds between group allocation, and so on. However, as there was so much diversity between the types of study designs and application of analyses, the comments within this box became meaningless, and it was subsequently omitted from the final bibliography.

### Quality of the study 5

Within the area of informed patient decision making, the issue of the readability and accessibility of intervention materials is of particular importance when aiming to explain variations in understanding. In an attempt to describe the quality of intervention materials a category was created to document whether or not the intervention materials had been piloted or evaluated for readability. Unfortunately the information pertaining to this extraction was inconsistently reported, if mentioned at all, and was omitted from the final coding of articles.

### Data synthesis

Four new variables were calculated to encourage integration of the research findings with the review objectives.

- The **theoretical context variable** combined information on reference to a theory-driven study with reference to a decision-making measure. Four classifications were created: theory and decision-making measure; theory only; decision-making measure only; neither theory nor decision-making measure.
- The **summary of reported effects variable** was a crude record of whether the authors reported an effect size between the intervention and health behaviour outcome measure: effect reported; no effect reported; mixed effects reported (e.g. different interventions did not show differential effects but an effect over time or some behaviour changes showed an effect and others did not).
- The **single variable for health domain** was created to maximise the efficiency of subsequent chi-square analysis. A few studies classified as being associated with more than one area of health were modified to force this information into one mutually exclusive category.
- The **single variable for health decision** was created to maximise the efficiency of subsequent chi-square analysis. A few studies classified as being associated with more than one type of decision were modified to force this information into one mutually exclusive category.

First, the results were presented as descriptive summaries and listings of articles by study number,

grouped by health domain, decision and theoretical context.

Second, the review identified five studies that were described as 'good' studies because they fulfilled the following criteria: design reported as RCTa; referred to a theory when designing the study or intervention; and assessed at least one decision-making measure. The findings of these studies are discussed in detail with reference to the reviews objectives. However, no meaningful quantitative meta-analysis have been applied as the reported information was too diverse.

Third, non-parametric analyses were used in an attempt to answer the five questions outlined within the aims. The associations between the reported effect of the intervention on the health behaviour and health domain, health decision, and type of intervention were assessed.

### **The trajectory of knowledge**

The number of studies using different types of methodology, using a decision-making theory, and recording an informed decision outcome were analysed by publication year. Results were plotted for the 5 years 1991, 1992, 1993, 1994 and 1995, for which both a full-year computer-based search and handsearch had been done.



# Chapter 3

## Results

A total of 547 reports fulfilled the inclusion criteria and formed the bibliography. They are listed by study number in appendix 5 and alphabetically by author in appendix 6.

As the amount of extracted information is large, there are several ways to synthesise and present the results. The remaining format of this section summarises the information in accordance with the research aim of this review. The bibliography is available in electronic format from JG Thornton, Centre for Reproduction, Growth and Development, 34 Hyde Terrace, Leeds, LS2 9LN.

### Summaries of extracted information by study

Appendices 7 and 8 summarise all the information extracted following application of the coding form. Appendix 7 includes information on: health decisions; domain of health care; health status and age of participants; reported theoretical context; intervention groups; and, reported measures. Appendix 8 includes: level of intervention; sample size; representativeness of sample; and summary of reported results.

Both tables follow the order of: study design (RCTa, RCTb, RCTc, non-randomised concurrent, non-randomised historical, and 'before and after' same sample); theory (did or did not report a theory); and health domain.

### Summary by frequency tables

The following tables describe the number of studies within each of the following coding frame categories: study design; type of intervention; health decision; health area; theory; decision associated factors; reported measures; theoretical context; and,

reported effects. Two tables are reported for each category. The first table shows the number of studies within each level of the category. As this review has referred to the hierarchy of evidence as a means of quality assessment, the second table reports the number of studies within each level of category by study design.

#### Study design

Table 1 describes the number of studies by one of six study designs.

#### Type of intervention

Tables 2 and 3 describe the number of studies using a particular type of intervention such as written leaflet or the provision of free transport. Note that these categories are not mutually exclusive. For example, one study may have assessed a number of written leaflets framed in different ways, or a media intervention compared with leaflet distribution in clinics. For example, 67% of RCTa studies compared the effects of additional information with the health decision whereas only 10% evaluated the impact of patient prompt.

#### Health decision

Tables 4 and 5 describe the number of studies looking at a particular health decision or health behaviour. Note that these categories are not mutually exclusive and a study may be classified more than once. For example, a healthy diet programme may have two stages involving cholesterol screening and change in food choice resulting in both the lifestyle and testing health-promotion categories being recorded. Seventy-one per cent of concurrent studies looked at lifestyle change decisions.

The frequencies for the modified, mutually exclusive health decision variable were: lifestyle change, 275 (50%); attendance testing, 103 (19%); adherence treatment, 92 (17%); utilisation of services, 51 (9%); and, other, 26 (5%).

TABLE 1 Frequency of study design (n = 547)

	RCTa low-risk bias	RCTb unknown bias	RCTc high-risk bias	Concurrent	Historical	Before/after same sample
Number of studies	51	208	77	114	34	63

**TABLE 2** Frequency of intervention type

Type of intervention	Examples	n
Used additional information	Written leaflet, audio-tape, video-tape, computer	301
Manipulated information	Illustrations, framing, graphics	94
Changed delivery of information	Letter, telephone, physicians versus nurse	273
Provided feedback	Demonstrate learnt skills, results of screening test	208
Used patient prompt	Decision aid, memory	55
Used group-delivery information	Media, classroom, community programmes	273
Manipulated environment	No-smoking policy, design inhaler, free service	109
Other	Patient incentives, health professional prompt	89

**TABLE 3** Frequency of studies by type of interventions and design

	RCTa	RCTb	RCTc	Concurrent	Historical	Before/after same sample
Used additional information	34	105	52	58	16	36
Manipulated information	15	50	17	11	0	1
Changed delivery of information	27	111	47	50	10	28
Provided feedback	14	81	33	47	5	28
Used patient prompt	5	19	15	9	2	5
Used group-delivery information	11	48	21	63	14	31
Manipulated environment	12	29	11	30	13	14
Other	4	38	9	20	7	11

**TABLE 4** Frequency of studies by type of health decision

Type of decision	Examples	n
Utilisation of services	Attendance at appointment, use call-out service	51
Health promotion – changes lifestyle	Smoking, drinking, exercise, substance use, safe sex	357
Health promotion – attendance testing	Breast, colorectal, cervical screening	114
Adherence to treatment	Medication, surgery, behavioural therapy	107
Active participation	Question asking within consultation, taking part in research	20
Other	Organ donation, willingness to pay	6

**TABLE 5** Frequency of studies by health decision and study design

	RCTa	RCTb	RCTc	Concurrent	Historical	Before/after same sample
Utilisation of services	8	20	9	7	4	3
Health promotion – changes lifestyle	28	128	47	81	21	52
Health promotion – attendance testing	14	39	18	22	10	11
Adherence to treatment	12	46	14	18	6	11
Active participation	3	11	3	0	1	2
Other	0	3	1	2	0	0



## Health domain

Tables 6 and 7 describe the number of studies within a particular health domain. Note that these categories are not mutually exclusive and a study may be classified more than once. For example, a study looking at postnatal depression may be classified as both midwifery and mental health, or a smoking study may come under both general medicine and cancer. Eighteen per cent of 'before and after' same-sample studies assessed an intervention associated with cancer care or prevention.

The frequencies for the modified, mutually exclusive health domain variable were: general medicine, 189 (35%); cancer, 109 (20%); the

human immunodeficiency virus (HIV) and STDs, 103 (19%); primary care, 55 (10%); obstetrics, 29 (5%); and, other, 62 (11%).

## Theory

Tables 8 and 9 describe the number of studies that referred to a theory to drive the research. These show that 85% of historical studies were not theory driven.

## Decision associated factors

As there was little variation between studies on these four variables, the frequencies are reported within the text, and no 'by design' frequencies are given.

**TABLE 6** Frequencies of studies by health domain

Health domain	Examples	n
Cancer	Breast, testicular, colorectal, cervical	114
Dentistry	Oral hygiene	10
Genetics	Huntington's chorea, cystic fibrosis (CF), sickle cell	7
General medicine	Cardiovascular disease (CVD) risk (smoking, exercise, alcohol), chronic illness (asthma, diabetes)	251
HIV and STD	Condom use, needle exchange	108
Mental health	Depression, schizophrenia, anxiety	15
Obstetrics, gynaecology and midwifery	Prenatal care, breast feeding, infertility	31
Paediatrics	Sudden infant death syndrome	6
Primary care	Coughs, colds, immunisations	61
Surgery and A&E	Urology, ophthalmology	11
Other	Alternative medicine, marital difficulties	13

**TABLE 7** Frequency of studies by health domain and design

	RCTa	RCTb	RCTc	Concurrent	Historical	Before/after same sample
Cancer	13	44	16	22	8	11
Dentistry	0	5	0	2	3	0
Genetics	1	1	2	2	1	0
General medicine	22	104	36	54	8	27
HIV and STD	6	32	12	28	8	22
Mental health	3	8	1	2	0	1
Obstetrics, gynaecology and midwifery	6	8	6	4	3	4
Paediatrics	0	3	0	1	2	0
Primary care	5	30	9	8	3	6
Surgery and A&E	2	3	2	3	1	0
Other	0	2	5	3	2	1

**TABLE 8** Frequency of studies by theory reported

Type of theory	Examples	n
No theory reported		341
Decision-making theory	Prospect theory, framing bias, health belief model (HBM), stages change, theory planned behaviour	101
Health promotion theory	Social learning theory (SLT), skill-based theory	63
Other theories	Attribution, stress-coping model, illness representation	42

**TABLE 9** Frequency studies by theory reported and study design

	RCTa	RCTb	RCTc	Concurrent	Historical	Before/after same sample
No theory reported	36	116	41	67	29	52
Decision-making theory	7	48	20	19	2	5
Health promotion theory	2	26	7	23	1	4
Other theories	6	18	9	5	2	2

- Number of studies that referred to real of hypothetical decision: real decisions, 512 (94%).
- Number of studies that encouraged patients to play an active part or increase awareness of the decision-making process: explicit decision-making process encouraged, 26 (5%).
- Number of studies where the recipient of the decision outcome was the participant or a third party: recipient same as decision maker, 476 (87%).
- Number of studies wither with or without time restraint: no time restraint, 525 (96%).

### Reported measures

Tables 10 and 11 describe all the measures reported by the study authors within the text. Note that these categories are not mutually exclusive. One study may have reported demographic variation, assessed anxiety and described reasons for choosing the final behaviour.

### Theoretical context

This derived variable describes the number of studies that reported a theoretical framework and a decision-making process and outcome measures (Tables 12 and 13). It attempts to provide an informed decision-making variable classification. Only five (10%) of RCTa studies referred to a theory and measured some aspect of 'informed' decision making such as attitudes or reasons.

### Reported effects

Tables 14 and 15 describe the number of studies that reported as association, or not, between the intervention and the health decision outcome. This is a crude measure based on the results

reported by the study authors and no independent verification of the quality of the study has been incorporated.

### Summary by study number listings

To facilitate identification of studies, the study numbers of articles are listed by various categories in Tables 16–20. The study number suffixes (indicating the number of studies within an article) have been omitted for this summary unless required to avoid ambiguity. Study numbers in Tables 16–19 are grouped into the following categories:

- type of intervention
- health decision
- health area
- theoretical context.

Study numbers in Table 20 are all RCT studies classified as most likely to assess informed patient decision making. That is to say, studies that have referred to a decision-making theory and assessed a decision-making process measure. the studies in Table 20 are grouped according to type of intervention.

### Synthesising findings

Two techniques were used in an attempt to synthesise the review findings. First, the integration of research findings from the five good studies that fulfilled the specific criteria. Second, the application of chi-square analysis to the crude measure of reported effect with the following variables:

**TABLE 10** Frequency of studies by measures reported

Measures	Examples	n
Demographics	Sex, age, level of education, marital status, religion	515
Personal history	Medical and family history, physiological assessment	386
Decision making	Attitudes, reasons, preferences, perception of risk, social norms	169
Self-efficacy	Perception of efficacy and self-efficacy	75
Knowledge	Knowledge	181
Satisfaction	Perceived satisfaction and usefulness	60
Affect	Depression, anxiety, worry, relief	69
Trait	Type A, locus control, cognitive closure	20
Other	Social support, coping, health professional measures	111

**TABLE 11** Frequencies of studies by measures reported and study design

	RCTa	RCTb	RCTc	Concurrent	Historical	Before/after same sample
Demographics	51	198	76	106	28	56
Personal history	43	159	52	78	17	37
Decision making	11	63	27	35	9	24
Self-efficacy	6	32	11	19	0	7
Knowledge	12	57	27	49	9	27
Satisfaction	9	24	12	5	18	4
Affect	7	39	10	5	0	6
Trait	2	7	3	6	0	2
Other	9	45	16	29	5	7

**TABLE 12** Frequency of studies by theoretical context

	n
Theory and decision-making measures	96
Theory only	110
Decision-making measures only	73
No theory or decision-making measure	268

**TABLE 13** Frequency of studies by theoretical context and study design

	RCTa	RCTb	RCTc	Concurrent	Historical	Before/after same sample
Theory and decision-making measures	5	47	16	21	1	6
Theory only	10	45	20	26	4	5
Decision-making measures only	6	16	11	14	8	18
No theory or decision-making measures	30	100	30	53	21	34

**TABLE 14** Frequency of studies reporting finding an effect or not

	n
Effect reported	294
No effect reported	173
Mixed effects reported	80

**TABLE 15** Frequency of studies by reported finding an effect or not and design

	RCTa	RCTb	RCTc	Concurrent	Historical	Before/after same sample
Effect reported	29	100	38	61	20	46
No effect reported	15	79	31	33	7	8
Mixed effects reported	7	29	8	20	7	9

design; intervention; health domain; health decision; and, theoretical context.

### Five good studies

The following five studies used an RCT with low risk of bias, reported 'informed' decision-making variables and referred to a theory.

- 197 - 1 Plaskon PP, Fadden MJ. Cancer screening utilisation: is there a role for social work in cancer prevention? *Social Work in Health Care* 1995;**21**(4):59–70.
- 310 - 1 Quaid K, Faden R, Vining E, Freeman J. Informed consent for a prescription drug: impact of disclosed information on patient understanding and medical outcomes. *Patient Education and Counseling* 1990;**15**:249–59.
- 325 - 1 Schultz S. Educational and behavioural strategies related to knowledge of and participation in an exercise program after cardiac positron emission tomography. *Patient Education and Counseling* 1993;**22**:47–57.
- 542 - 1 Simel DL, Feussner JR. A randomised controlled trial comparing quantitative informed consent formats. *Journal of Clinical Epidemiology* 1991;**44**(8):771–7.
- 799 - 1 DiClemente RJ, Wingwood GM. A randomised controlled trial of an HIV sexual risk-reduction intervention for young African-American women. *JAMA* 1995;**274**(16):1271–6.

A formal integrative review of these five studies would not be useful as each study evaluates a different decision, refers to a different theory and considers different interventions. Some participants were well and others physically ill, and some samples were selected systematically and others by convenience (*Table 21*).

Nevertheless, it appears that information and education are not the most effective ways of facilitating informed decision making. One study found that changing the amount of information about risks and side-effects of a drug did not affect patients' adherence to the treatment (Quaid *et al*). In another, varying the quantitative information used to introduce a new medication made no difference to the decision to participate in a research trial in 55% of out-patients (Simel and Feussner), though it did influence the choice of the 45% who cited this information in their decision. Although an education intervention was found to increase exercise amongst patients at risk of heart disease over a 12-week period, this included behavioural planning (Schultz). When other behavioural strategies of goal setting, self-monitoring, positive reinforcement and telephone follow-up were added, frequency of exercise was significantly increased after 6 weeks. The importance of situational factors was demonstrated in the decision of general practice patients to use a kit for colorectal cancer screening. All patients were given educational information, but screening increased from 0% to 51% when patients were given a free kit rather than told to go to the general practice office or to one of three pharmacies to buy a kit (Plaskon and Fadden). The role of cognitive and social, as well as behavioural, factors in influencing decision making is evident in an intervention aimed at reducing HIV sexual risk amongst African-American women (DiClemente and Wingwood). Although an educational intervention did not bring about change, a social skills intervention, based on social cognitive theory and theory of gender and power, increased condom use.

Although these studies are of disparate populations making disparate decisions, they show the importance of the context and social influences on individual's decision making.

**TABLE 16** Study numbers (without suffix unless necessary) by type of intervention

Type of intervention and study design	No. of studies	Study numbers
Additional information – written	258	
RTCa	30	39, 50, 64, 67, 85, 101, 118, 120, 122, 196, 197, 200, 216, 275, 299, 301, 316, 325, 333, 350, 387, 393, 483, 509, 542, 648, 724, 741, 778, 793
RCTb	87	19, 22, 28, 48, 63, 104, 110, 114, 135, 140, 143, 150, 157, 158, 158, 167, 176, 183, 207, 210, 211, 213, 216, 232, 233, 234, 241, 259, 265, 268, 269, 277, 279, 285, 288, 295, 302, 327, 332, 337, 357, 372, 376, 414, 420, 434, 454, 465, 469, 482, 490, 491, 506, 511, 520, 525, 530, 533, 541, 572, 587, 600, 616, 632, 632, 635, 639, 644, 652, 667, 668, 671, 675, 679, 683, 690, 693, 700, 705, 706, 728, 732, 747, 757, 783, 794, 796
RCTc	46	58, 72, 83, 98, 115, 154, 169, 170, 181, 242, 248, 254, 287, 296, 303, 320, 324, 329, 343, 349, 354, 360, 363, 375, 381, 407, 442, 459, 471, 472, 473, 484, 507, 508, 528, 601, 640, 646, 649, 712, 717, 743, 752, 754, 790, 792
Concurrent	53	3, 44, 125, 136, 146, 171, 177, 180, 203, 205, 205, 206, 215, 237, 243, 245, 246, 251, 253, 261, 273, 280, 283, 312, 315, 367, 370, 373, 386, 391, 394, 404, 418, 440, 446, 448, 453, 463, 464, 504, 515, 516, 579, 628, 631, 663, 665, 709, 726, 740, 780, 791, 813
Before/after different samples	16	26, 97, 149, 192, 193, 294, 340, 383, 462, 495, 514, 575, 619, 658, 670, 765
Before/after same sample	26	35, 42, 111, 126, 133, 162, 163, 218, 244, 262, 266, 270, 348, 384, 449, 450, 467, 480, 522, 526, 544, 573, 676, 770, 809, 815
Additional information – audiotape	10	
RCTa	2	117, 581
RCTb	4	264, 371, 465, 488
RCTc	3	154, 472, 754
Concurrent	1	663
Additional information – video-tape	74	
RCTa	3	23, 585, 741
RCTb	27	28, 55.1, 55.2, 155, 207, 241, 289, 341, 376, 424, 438, 454, 458, 491, 502, 533, 616, 635, 639, 644, 705, 728, 742, 758, 776, 789, 794
RCTc	15	37, 181, 254, 287, 293, 310, 329, 354, 484, 528, 640, 646, 696, 720, 762
Concurrent	13	184, 205, 205, 240, 246, 261, 311, 313, 464, 579, 628, 631, 791
Before/after different samples	3	26, 294, 658
Before/after same sample	13	35, 49, 79, 123, 364, 450, 467, 526, 594, 610, 651, 655, 746
Additional information – computer based	7	
RCTb	5	327, 396, 488, 493, 667
Concurrent	1	40
Before/after same sample	1	795
Manipulated information – illustration or graphic	14	
RCTa	2	50, 117
RCTb	5	19, 150, 305, 357, 371
RCTc	4	324, 459, 473, 649
Concurrent	2	177, 448
Before/after same sample	1	723
Manipulated information – framing or quantity	84	
RCTa	14	39, 67, 101, 117, 118, 275, 301, 393, 542, 724, 755, 778, 793, 799
RCTb	47	19, 55, 86, 105, 137, 158, 167, 176, 183, 211, 226, 288, 295, 305, 321, 327, 332, 341, 420, 426, 434, 438, 482, 490, 491, 494, 533, 587, 600, 634, 636, 644, 653, 667, 668, 671, 683, 685, 690, 693, 727, 742, 747, 757, 772, 783, 789
RCTc	14	115, 310, 377, 380, 381, 471, 473, 481, 507, 512, 684, 691, 752, 788
Concurrent	9	136, 257, 280, 386, 453, 586, 748, 760, 822

continued

**TABLE 16 contd** Study numbers (without suffix unless necessary) by type of intervention

Type of intervention and study design	No. of studies	Study numbers
Change in delivery of information – community or hospital	10	
RCTa	1	39
RCTb	3	47, 234, 564
RCTc	3	485, 752, 790
Concurrent	3	586, 748, 775
Change in delivery of information – alter contact with health professional	183	
RCTa	18	52, 64, 117, 196, 200, 202, 325, 333, 350, 585, 606, 638, 648, 678, 724, 778, 781, 799
RCTb	70	1, 4, 10, 22, 63, 102, 135, 139, 140, 147, 157, 165, 174, 207, 211, 212, 241, 249, 259, 263, 268, 269, 276, 277, 285, 289, 291, 304, 328, 336, 341, 353, 365, 369, 403, 439, 447, 458, 465, 479, 491, 494, 503, 511, 525, 527, 530, 533, 553, 554, 576, 616, 622, 629, 632, 632, 635, 636, 639, 659, 693, 702, 707, 728, 742, 766, 776, 783, 807, 814
RCTc	23	58, 80, 98, 169, 170, 181, 320, 329, 354, 375, 407, 411, 442, 470, 472, 528, 601.2, 640, 695, 720, 754, 763, 792
Concurrent	36	71, 82, 89, 95, 136, 171, 177, 194, 205.1, 205.2, 206, 237, 246, 251, 253, 256, 260, 273, 284, 312, 313, 367.2, 373, 416, 453, 456, 464, 504, 529, 555, 570, 579, 628, 726, 740, 791
Before/after different samples	8	26, 239, 340, 462, 575, 619, 623, 674
Before/after same sample	28	35, 38, 49, 84, 111, 123, 162, 218, 244, 262, 270, 367, 449, 480, 521, 544, 621, 650, 651, 654, 723, 729, 746, 770, 798, 809, 815, 821
Change in delivery of information – letter	83	
RCTa	10	39, 41, 52, 101, 216, 275, 299, 316, 378, 648
RCTb	39	110, 143, 158.1, 158.2, 167, 213, 216, 259, 265, 269, 285, 352.1, 352.2, 352.3, 372, 395, 414, 420, 426, 439, 447, 469, 543, 566, 572, 589, 620, 630, 634, 652, 653, 668, 683, 727, 728, 747, 757, 783, 784
RCTc	19	56, 115, 154, 172, 248, 296, 320, 349, 363, 381, 412, 484, 507, 512, 640, 712, 717, 752, 790
Concurrent	14	61, 146, 237, 253, 257, 394, 404, 446, 516, 546, 631, 740, 760, 813
Before/after different samples	1	100
Change in delivery of information – telephone	47	
RCTa	7	52, 196, 202, 299, 325, 638, 714
RCTb	23	63, 139, 143, 168, 233, 265, 288, 304, 352.2, 353, 369, 403, 408, 414, 420, 482, 572, 576, 600, 622, 668, 702, 783
RCTc	9	287, 380, 475, 481, 512, 640, 711, 754, 788
Concurrent	3	257, 394, 446
Before/after different samples	4	26, 239, 514, 619
Before/after same sample	1	218

continued

**TABLE 16 contd** Study numbers (without suffix unless necessary) by type of intervention

Type of intervention and study design	No. of studies	Study numbers
Provision of feedback or skills acquisition	208	
RCTa	14	64, 107, 120, 275, 316, 325, 387, 393, 585, 648, 714, 755, 781, 799
RCTb	81	4, 10, 20, 22, 104, 110, 114, 135, 140, 145, 147, 157, 165, 174, 207, 210, 212, 216, 217, 220, 223, 226, 233, 241, 247, 263, 264, 268, 269, 277, 291, 295, 307, 327, 328, 341, 371, 396, 414, 424, 439, 454, 458, 478, 479, 491, 492, 502, 503, 511, 525, 527, 530, 533, 541, 553, 562, 576, 616, 630, 632, 632, 635, 636, 644, 660, 667, 668, 679, 685, 690, 700, 705, 707, 742, 758, 766, 772, 776, 783, 814
RCTc	33	72, 80, 98, 115, 154, 287, 293, 320, 329, 354, 360, 375, 377, 381, 405, 407, 442, 472, 475, 481, 601, 640, 646, 649, 684, 696, 708, 720, 749, 754, 788, 792, 817
Concurrent	47	3, 17, 18, 89, 125, 136, 177, 184, 222, 240, 246, 251, 253, 273, 283, 284, 311, 313, 315, 335, 345, 367, 370, 373, 385, 386, 394, 404, 416, 440, 448, 453, 486, 504, 515, 555, 628, 631, 663, 726, 731, 740, 759, 760, 791, 813, 823
Before/after different samples	5	26, 149, 619, 623, 765
Before/after same sample	28	35, 42, 46, 49, 84, 99, 123, 132, 133, 159, 162, 270, 355, 361, 364, 384, 449, 450, 480, 521, 522, 526, 594, 621, 654, 676, 688, 729
Patient prompt – use of decision aid	5	
RCTb	2	396, 493
RCTc	1	472
Before/after same sample	2	42, 594
Patient prompt – use of memory or information aid	45	
RCTa	4	85, 333, 378, 387
RCTb	13	157, 269, 372, 420, 447, 469, 482, 506, 576, 700, 707, 783, 794
RCTc	14	115, 154, 172, 254, 303, 343, 360, 381, 481, 601, 640, 691, 712, 763
Concurrent	9	44, 243, 246, 253, 315, 394, 448, 631, 760
Before/after different samples	2	340, 421
Before/after same sample	3	162, 449, 544
Patient prompt – shared health professional/patient decision making	5	
RCTa	1	678
RCTb	4	27, 279, 337, 564
Group delivery of information	163	
RCTa	11	50, 107, 117, 120, 200, 483, 585, 648, 724, 778, 799
RCTb	46	1, 4, 22, 96, 143, 145, 147, 174, 207, 211, 212, 223, 241, 255.2, 263, 264, 269, 277, 285, 328, 341, 414, 424, 438, 478, 479, 489, 491, 492, 494, 530, 533, 553, 564, 616, 632, 632, 635, 636, 659, 660, 705, 736, 766, 807, 814
RCTc	21	58, 72, 80, 98, 254, 303, 320, 354, 375, 405, 407, 442, 485, 640, 649, 684, 696, 720, 743, 749, 792
Concurrent	46	3, 18, 44, 45, 95, 125, 136, 146, 180, 184, 203, 222, 238, 240, 253, 258, 261, 273, 280, 283, 284, 335, 370, 373, 385, 416, 440, 448, 453, 456, 463, 515, 555, 570, 579, 608, 628, 665, 686, 709, 731, 740, 759, 760, 780, 825
Before/after different samples	11	97, 149, 192, 340, 383, 461, 514, 662, 670, 680, 765
Before/after same sample	28	38, 42, 49, 79, 99, 123, 132, 133, 159, 262, 270, 348, 355, 361, 364, 467, 521, 522, 526, 573, 621, 650, 651, 654, 657, 723, 746, 809

continued

**TABLE 16 contd** Study numbers (without suffix unless necessary) by type of intervention

Type of intervention and study design	No. of studies	Study numbers
Change in health service provision	83	
RCTa	9	39, 41, 52, 64, 197, 425, 509, 714, 741
RCTb	21	47, 48, 135, 143, 213, 234, 269, 285, 289, 352.3, 372, 376, 439, 454, 554, 566, 683, 732, 736, 768, 769
RCTc	10	56, 254, 287, 293, 296, 412, 470, 484, 646, 752
Concurrent	22	3, 75, 136, 146, 177, 180, 203, 237, 243, 256, 312, 397, 448, 452, 455, 464, 516, 529, 618, 686, 726, 822
Before/after different samples	8	340, 383, 421, 462, 575, 737, 765, 767.1
Before/after same sample	13	38, 163, 218, 244, 342, 355, 537, 655, 729, 770, 795, 798, 809
Manipulation of the environment	23	
RCTa	2	216, 229
RCTb	1	220
RCTc	2	56, 442
Concurrent	10	18, 75, 180, 206, 215, 235, 251, 284, 390, 464
Before/after different samples	6	142, 149, 423, 462, 550, 563
Before/after same sample	2	270, 537
Group delivery – use of media	70	
RCTb	6	22, 135, 143, 217, 285, 705
RCTc	2	58, 792
Concurrent	38	3, 18, 44, 45, 146, 171, 180, 184, 203, 205.1, 205.2, 206, 215, 222, 237, 238, 243, 246, 251, 258, 280, 311, 312, 323, 335, 367.2, 391, 448, 463, 464, 608, 631, 665, 686, 709, 726, 740, 780
Before/after different samples	14	97, 149, 192, 340, 383, 461, 462, 514, 619, 658, 662, 670, 680, 765
Before/after same sample	10	38, 42, 49, 162, 218, 367, 522, 657, 746, 809
Other – health professional prompts for action	14	
RCTa	1	724
RCTb	1	395
Concurrent	5	205, 404, 455, 791, 822
Before/after different samples	4	34, 192, 401, 413
Before/after same sample	3	466, 522, 809
Other – patient incentives	32	
RCTa	2	64, 200
RCTb	12	22, 102, 135, 156, 165, 216, 269, 352, 424, 644, 652, 814
RCTc	2	106, 115
Concurrent	7	18, 44, 184, 280, 283, 367, 515
Before/after different samples	3	149, 462, 765
Before/after same sample	6	162, 179, 262, 367, 450, 688



**TABLE 17** Study numbers (without suffix unless necessary) by health decision

Health decision and study design	No. of studies	Study numbers
Decision to use a service e.g. attend an appointment, use a call-out service	51	
RCTa	8	196, 202, 333, 425, 606, 638, 678, 714
RCTb	20	10, 20, 27, 28, 167, 302, 337, 369, 372, 376, 426, 439, 494, 511, 520, 554, 706, 733, 769, 796
RCTc	9	83, 106, 172, 248, 320, 470, 481, 717, 788
Concurrent	7	75, 82, 323, 331, 394, 504, 748
Before/after different samples	4	97, 294, 563, 737
Before/after same sample	3	42, 537, 676
Decision to make a lifestyle change (health promotion) e.g. stop smoking, exercise	357	
RCTa	28	23, 50, 64, 85, 101, 107, 118, 120, 196, 200, 216, 229, 299, 316, 325, 333, 350, 378, 393, 581, 585, 648, 724, 741, 755, 781, 793, 799
RCTb	128	1, 4, 22, 24, 27, 32, 47, 55, 55, 63, 86, 102, 104, 110, 135, 139, 140, 143, 145, 147, 156, 157, 158, 158, 165, 174, 183, 207, 210, 211, 212, 213, 216, 217, 220, 223, 226, 233, 234, 241, 247, 249, 255, 259, 263, 264, 265, 268, 269, 276, 277, 279, 288, 289, 291, 295, 305, 307, 308, 321, 327, 328, 336, 341, 356, 365, 371, 372, 376, 395, 396, 398, 402, 403, 408, 414, 424, 426, 454, 457, 458, 465, 478, 489, 490, 491, 492, 503, 511, 525, 533, 541, 553, 562, 564, 572, 574, 576, 600, 616, 620, 632, 632, 633, 635, 636, 639, 644, 659, 660, 668, 679, 685, 690, 705, 707, 732, 733, 736, 742, 758, 766, 768, 772, 776, 783, 807, 814
RCTc	47	72, 80, 98, 115, 154, 170, 181, 242, 254, 287, 293, 296, 310, 320, 324, 329, 360, 363, 375, 405, 407, 442, 459, 471, 472, 473, 475, 484, 485, 512, 528, 601, 601, 640, 646, 649, 684, 696, 708, 720, 749, 752, 754, 762, 777, 792, 817
Concurrent	81	3, 17, 18, 40, 71, 89, 95, 125, 136, 146, 161, 171, 177, 180, 184, 194, 206, 209, 215, 222, 235, 237, 238, 240, 245, 246, 251, 253, 255, 256, 257, 258, 260, 261, 273, 280, 283, 284, 311, 312, 313, 345, 367, 370, 373, 385, 386, 390, 391, 392, 409, 416, 440, 453, 455, 456, 463, 464, 515, 529, 555, 570, 579, 618, 628, 631, 663, 665, 709, 726, 731, 740, 759, 773, 775, 779, 780, 791, 813, 822, 823
Before/after different samples	21	26, 97, 100, 142, 149, 239, 340, 421, 423, 461, 462, 514, 550, 619, 623, 647, 658, 662, 680, 765, 767
Before/after same sample	52	35, 38, 49, 79, 84, 91, 123, 126, 132, 133, 159, 162, 163, 179, 198, 218, 244, 262, 266, 270, 342, 348, 355, 364, 367, 384, 450, 466, 467, 480, 521, 522, 526, 544, 545, 573, 610, 621, 650, 651, 654, 655, 657, 676, 688, 723, 729, 746, 770, 798, 815, 821

continued

**TABLE 17 contd** Study numbers (without suffix unless necessary) by health decision

Health decision and study design	No. of studies	Study numbers
Decision to attend for testing (health promotion) e.g. breast/colorectal screening	114	
RCTa	14	39, 41, 52, 85, 101, 108, 117, 122, 197, 275, 316, 333, 509, 778
RCTb	39	48, 86, 114, 232, 285, 288, 289, 305, 307, 352, 352, 352, 353, 372, 395, 420, 438, 439, 482, 491, 543, 566, 587, 589, 620, 630, 634, 652, 653, 675, 683, 690, 713, 727, 728, 747, 757, 784, 789
RCTc	18	37, 56, 154, 248, 296, 354, 377, 380, 381, 412, 484, 507, 508, 695, 712, 743, 752, 790
Concurrent	22	44, 45, 61, 171, 203, 205, 205, 243, 246, 257, 394, 397, 404, 418, 446, 448, 516, 546, 608, 686, 760, 813
Before/after different samples	10	34, 192, 193, 294, 383, 401, 421, 575, 670, 737
Before/after same sample	11	46, 99, 126, 133, 348, 384, 466, 746, 795, 809, 815
Decision to adhere to treatment regimen e.g. drugs, surgery, behaviour therapy	107	
RCTa	12	53, 202, 216.2, 229, 299, 301, 333, 378, 427, 483, 714, 781
RCTb	46	6.1, 6.2, 6.3, 51, 57, 63, 88, 96, 104, 105, 137, 139, 150, 167, 168, 176, 234, 291, 304, 327, 328, 332, 357, 369, 395, 402, 414, 434, 447, 469, 479, 488, 502, 527, 530, 564, 572, 612, 622, 633, 671, 693, 698, 702, 768, 794
RCTc	14	98, 169, 170, 181, 303, 349, 363, 380, 411, 471, 512, 691, 711, 763
Concurrent	18	18, 71, 82, 95, 124, 146, 313, 315, 323, 331, 335, 452, 455, 486, 586, 618, 775, 825
Before/after different samples	6	193, 239, 413, 563, 674, 767
Before/after same sample	11	42, 74, 342, 361, 449, 466, 522, 544, 594, 655, 676
Decision to take an active part in consultation or participate in research	20	
RCTa	3	67, 387, 542
RCTb	11	155, 216, 434, 488, 493, 506, 600, 629, 667, 700, 816
RCTc	3	343, 472, 601
Before/after different samples	1	495
Before/after same sample	2	111, 594
Other decisions e.g. donate an organ, pay for treatment	6	
RCTb	3	19, 176, 408
RCTc	1	58
Concurrent	2	190, 430

**TABLE 18** Study numbers (without suffix unless necessary) by health domain

Health domain and study design	No. of studies	Study numbers
Cancer	114	
RCTa	13	23, 39, 41, 52, 85, 101, 108, 117, 122, 197, 275, 316, 509
RCTb	44	48, 86, 114, 140, 207, 232, 263, 269, 285, 288, 307, 352.1, 352.2, 352.3, 353, 395, 420, 434, 438, 458, 482, 488, 491, 527, 533, 541, 543, 566, 587, 630, 653, 667, 675, 683, 690, 700, 705, 713, 727, 728, 757, 784, 789, 816
RCTc	16	37, 154, 248, 296, 310, 354, 377, 380, 412, 484, 507, 508, 691, 695, 712, 752
Concurrent	22	44, 61, 136, 171, 180, 203, 205.1, 205.2, 237, 243, 246, 385, 386, 397, 404, 418, 446, 448, 546, 608, 686, 760
Before/after different samples	8	34, 100, 192, 383, 401, 421, 575, 765
Before/after same sample	11	46, 99, 126, 133, 198, 266, 466, 746, 795, 809, 815
Dentistry	10	
RCTb	5	371, 457, 562, 576, 758
Concurrent non-randomised	2	529, 618
Before/after different samples	3	97, 563, 737
Genetics	7	
RCTa	1	778
RCTb	1	20
RCTc	2	743, 790
Concurrent	2	430, 516
Before/after different samples	1	294
General medicine	251	
RCTa	22	50, 53, 64, 67, 118, 120, 196, 200, 216.2, 229, 301, 325, 350, 483, 542, 581, 648, 714, 724, 755, 781, 793
RCTb	104	1, 10, 20, 27, 28, 32, 47, 55.1, 55.2, 57, 63, 88, 102, 104, 105, 110, 137, 139, 140, 143, 145, 147, 155, 157, 158.1, 158.2, 168, 174, 183, 207, 210, 211, 213, 216, 217, 220, 223, 226, 233, 241, 247, 249, 255.2, 259, 263, 264, 265, 268, 269, 276, 277, 279, 291, 295, 304, 305, 308, 321, 327, 328, 332, 336, 341, 356, 365, 372, 376, 395, 396, 398, 402, 414, 426, 439, 447, 469, 489, 490, 502, 503, 527, 530, 541, 564, 574, 600, 612, 620, 622, 629, 644, 652, 659, 668, 671, 679, 693, 702, 707, 769, 772, 783, 807, 814
RCTc	36	56, 72, 80, 115, 169, 170, 242, 254, 287, 303, 310, 320, 329, 343, 360, 375, 381, 405, 411, 442, 459, 471, 472, 473, 475, 481, 485, 601.1, 601.2, 640, 708, 711, 752, 754, 763, 792
Concurrent	54	45, 82, 89, 184, 194, 206, 209, 215, 222, 235, 237, 238, 245, 251, 253, 255, 256, 257, 258, 260, 261, 273, 280, 283, 284, 311, 312, 313, 315, 323, 345, 367, 370, 390, 394, 409, 416, 418, 440, 452, 453, 456, 464, 504, 586, 631, 709, 726, 731, 740, 759, 779, 813, 823
Before/after different samples	8	340, 423, 461, 462, 495, 514, 550, 658
Before/after same sample	27	42, 74, 79, 84, 111, 133, 162, 179, 218, 244, 262, 270, 348, 355, 367, 384, 449, 466, 521, 522, 544, 594, 621, 657, 676, 798, 821

continued

**TABLE 18 contd** Study numbers (without suffix unless necessary) by health domain

Health domain and study design	No. of studies	Study numbers
HIV and STD	108	
RCTa	6	107, 202, 393, 585, 741, 799
RCTb	32	4, 22, 24, 156, 165, 212, 223, 226, 247, 277, 289, 341, 424, 454, 465, 478, 492, 525, 600, 632, 632, 635, 636, 639, 660, 685, 732, 733, 736, 742, 766, 776
RCTc	12	181, 360, 407, 528, 646, 649, 684, 696, 720, 749, 762, 817
Concurrent	28	17, 18, 71, 125, 161, 177, 240, 251, 311, 373, 391, 392, 453, 455, 463, 486, 515, 555, 579, 628, 663, 665, 748, 759, 780, 791, 822, 823
Before/after different samples	8	142, 423, 550, 619, 647, 662, 670, 680
Before/after same sample	22	35, 38, 49, 91, 123, 159, 364, 450, 467, 480, 526, 537, 545, 610, 650, 651, 654, 655, 688, 723, 729, 770
Mental health	15	
RCTa	3	67, 427, 606
RCTb	8	28, 96, 479, 493, 494, 511, 554, 794
RCTc	1	717
Concurrent	2	124, 825
Before/after same sample	1	361
Obstetrics and gynaecology, and midwifery	31	
RCTa	6	196, 200, 425, 678, 724, 778
RCTb	8	135, 137, 403, 616, 634, 698, 733, 768
RCTc	6	83, 106, 115, 293, 329, 788
Concurrent	4	3, 71, 190, 486
Before/after different samples	3	26, 514, 623
Before/after same sample	4	132, 163, 621, 650
Paediatrics	6	
RCTb	3	167, 520, 706
Concurrent	1	335
Before/after different samples	2	193, 239
Primary care	61	
RCTa	5	299, 333, 378, 387, 638
RCTb	30	6.1, 6.2, 6.3, 19, 27, 32, 47, 51, 167, 174, 234, 302, 337, 357, 369, 395, 408, 414, 426, 506, 520, 553, 572, 589, 620, 633, 706, 747, 769, 796
RCTc	9	98, 172, 363, 470, 512, 601, 601, 752, 777
Concurrent	8	95, 146, 190, 257, 331, 418, 775, 813
Before/after different samples	3	193, 413, 767
Before/after same sample	6	62, 342, 348, 466, 537, 573

continued

**TABLE 18 contd** Study numbers (without suffix unless necessary) by health domain

Health domain and study design	No. of studies	Study numbers
Surgery/A&E	11	
RCTa	2	581, 714
RCTb	3	150, 490, 622
RCTc	2	80, 777
Concurrent	3	40, 75, 394
Before/after different samples	1	674
Other area of health care	13	
RCTb	2	176, 288
RCTc	5	58, 170, 293, 324, 349
Concurrent	3	194, 570, 773
Before/after different samples	2	149, 461
Before/after same sample	1	84

**TABLE 19** Study numbers (without suffix unless necessary) by reported theory and measure of informed decision making

Theory, measures and study design	No. of studies	Study numbers
Theory and measure reported	96	
RCTa	5	197, 301, 325, 542, 799
RCTb	47	24, 86, 114, 140, 145, 176, 210, 211, 217, 220, 232, 255.2, 276, 277, 288, 321, 371, 376, 420, 424, 426, 434, 438, 454, 458, 478, 492, 502, 525, 616, 620, 622, 632.1, 632.2, 635, 659, 660, 667, 668, 690, 700, 742, 758, 766, 768, 776, 784
RCTc	16	72, 115, 242, 354, 360, 380, 472, 507, 640, 691, 696, 717, 749, 762, 792, 817
Concurrent	21	89, 180, 238, 240, 253, 255, 313, 345, 370, 373, 391, 416, 453, 515, 516, 608, 663, 709, 759, 760, 822
Before/after different samples	1	26
Before/after same samples	6	46, 126, 198, 364, 526, 594
Theory only	110	
RCTa	10	101, 107, 117, 120, 275, 333, 387, 427, 483, 585
RCTb	45	1, 4, 19, 20, 27, 104, 157, 158.1, 158.2, 165, 167, 207, 223, 226, 233, 241, 247, 264, 269, 291, 295, 305, 307, 332, 357, 403, 457, 469, 490, 493, 503, 506, 541, 576, 600, 633, 636, 644, 685, 707, 757, 789, 794, 814, 816
RCTc	20	56, 58, 80, 98, 169, 181, 248, 254, 287, 324, 329, 375, 405, 442, 484, 508, 528, 601.1, 601.2, 646
Concurrent	26	17, 18, 95, 136, 184, 190, 209, 237, 251, 261, 311, 367, 386, 404, 440, 446, 456, 463, 464, 486, 618, 628, 631, 726, 780, 823
Before/after different samples	4	462, 647, 658, 765
Before/after same sample	5	179, 262, 367, 450, 654
Measure only	73	
RCTa	6	39, 316, 350, 509, 581, 678
RCTb	16	102, 105, 135, 147, 183, 212, 249, 302, 336, 341, 489, 494, 520, 562, 698, 783
RCTc	11	154, 310, 343, 381, 459, 473, 649, 684, 712, 720, 743
Concurrent	14	3, 44, 45, 177, 243, 256, 385, 430, 504, 570, 665, 686, 773, 825
Before/after different samples	8	193, 239, 383, 413, 423, 514, 619, 662
Before/after same sample	18	35, 49, 79, 111, 132, 159, 163, 218, 244, 342, 467, 480, 573, 610, 621, 650, 655, 688

**TABLE 20** Study numbers (without suffix unless necessary) by intervention from informed decision-making RCTs

Intervention and study design	No. of studies	Study numbers
Additional information	45	72, 114, 115, 140, 176, 197, 210, 211, 232, 242, 277, 288, 301, 325, 354, 360, 371, 376, 420, 424, 434, 438, 454, 458, 472, 502, 507, 525, 542, 616, 632.1, 632.2, 635, 640, 667, 668, 690, 696, 700, 717, 742, 758, 762, 776, 792
Additional written information	35	72, 114, 115, 140, 176, 197, 210, 211, 232, 242, 277, 288, 301, 325, 354, 360, 376, 420, 434, 454, 472, 507, 525, 542, 616, 632.1, 632.2, 635, 640, 667, 668, 690, 700, 717, 792,
Manipulation of information	20	86, 115, 176, 211, 288, 301, 321, 380, 420, 426, 434, 438, 507, 542, 667, 668, 690, 691, 742, 799
Provision of feedback/ skills acquisition	40	72, 114, 115, 140, 145, 210, 217, 220, 277, 325, 354, 360, 371, 424, 454, 458, 472, 478, 492, 502, 525, 616, 632.1, 632.2, 635, 640, 660, 667, 668, 690, 696, 700, 742, 749, 758, 766, 776, 792, 799, 817
Use of patient prompts	7	115, 360, 420, 472, 640, 691, 700
Group delivery of information	22	72, 145, 211, 255.2, 277, 354, 424, 438, 478, 492, 616, 632.1, 632.2, 635, 640, 659, 660, 696, 749, 766, 792, 799
Manipulation of the environment	1	220
Incentives	2	115, 424

**TABLE 21** Results information for five RCTa studies assessing informed decision making

	Health decision and intervention	Impact of intervention on health decision	Process measures (other measures)	Impact intervention on process and other measures
197	Utilisation screening kit: 1 – verbal information (VI) 2 – VI + free kit provided	0% uptake VI only 51% uptake VI + kit	Perception of risk of having colorectal cancer (none)	Not used in analysis
301	Adherence to medication: 1 – routine written information (WI) 2 – enhanced risk (WI)	No group differences to adherence drug	Perceived seriousness risks and side-effects (anxiety, knowledge) enhanced WI)	Enhanced WI associated greater perceived seriousness risks (no effect anxiety, greater knowledge)
325	Uptake exercise: 1 – VI + screening results 2 – VI + screening results + encouraged to self-monitor by writing down goals, diary of exercise	No group differences on exercise uptake and maintenance. Both groups increased over time	Perception of health (knowledge)	Not used in analysis (no group differential effect knowledge, increase over time)
542	Consent to participate in research: 1 – WI framed as a drug working 'twice as fast' 2 – WI framed as a drug working 'half as fast'	Participation greater in 'twice as fast' group	Type of information used for decision, reasons participation (none)	'Twice as fast' used more 'quantitative' information, not analysed by group
799	Changes risky sex: 1 – delayed intervention 2 – VI about HIV prevention (culturally) framed (one session) 3 – VI + social skills (five sessions)	Social skills group only associated greater safer sex practices  Note: differential attrition rates	Perception of partner norms (knowledge, perceived self-efficacy)	Social skills group associated changes in perception of partner norms (no group differences knowledge, social skills greater perceived self-efficacy)

**TABLE 22** Eight chi-square analyses for type of intervention by reported health behaviour effect

	Effect (n = 294)	No effect (n = 173)	Mixed effect (n = 80)	Chi-square value (d.f. = 2)	Significance
Additional information	157	93	51	2.89	$p = 0.236$
Manipulation of information	42	31	11	1.30	$p = 0.524$
Change in delivery of information	161	75	37	6.17	$p = 0.046$
Provision of feedback	99	70	39	6.70	$p = 0.035$
Use of patient prompts	33	17	5	1.74	$p = 0.420$
Use of group-delivery information	92	61	35	4.42	$p = 0.110$
Manipulation of environment	11	5	7	5.00	$p = 0.082$
Other	52	23	14	1.65	$p = 0.439$

*d.f.* = degrees of freedom

**TABLE 23** Chi-square analysis for health domain by reported health behaviour effect

	Effect	No effect	Mixed effect	Chi-square value (d.f. = 10)	Significance
General medicine	87	68	34	18.7	$p = 0.044$
Cancer	70	31	8		
HIV	55	27	21		
Primary care	28	19	8		
Obstetrics	15	9	5		
Other	19	19	4		

**TABLE 24** Chi-square analysis for type of health decision by reported health behaviour effect

	Effect	No effect	Mixed effect	Chi-square value (d.f. = 8)	Significance
Lifestyle change	129	93	53	24.1	$p = 0.002$
Attendance testing	66	28	9		
Adherence to treatment	58	27	7		
Utilisation of service	23	17	11		
Other	18	8	0		

## Non-parametric analyses

Non parametric chi-square analyses were carried out between the crude reported effect variable and intervention used, health domain, and health decision.

### Intervention

Because many studies involved more than one manipulation within the intervention, such as written information and delivery of information, individual analyses were carried out for each intervention. All eight chi-square analyses assessing the association between the type of intervention employed and the likelihood of reporting an effect size for the health behaviour described are reported in *Table 22*. As several analyses were carried

out, the results ought to be interpreted with caution.

### Health domain

*Table 23* describes the chi-square analysis for health domain by reported health behaviour effect. It assesses the association between the health domain of the decision and the likelihood of reporting an effect for the health behaviour described.


### Type of health decision

*Table 24* describes the chi-square analysis for the type of health decision by reported health behaviour effect. It assesses the association between the type of decision context with the likelihood of reporting an effect for the health behaviour described.


### The trajectory of knowledge

The number of included studies by year of publication is shown in *Figures 1–4*. The total is shown in *Figure 1*, trials broken down by quality of


randomisation in *Figure 2*, non-randomised studies in *Figure 3* and studies using a decision theory or recording an informed decision measure are shown in *Figure 4*.




**FIGURE 1** Number of studies by year of publication



**FIGURE 2** Number of studies by quality of randomisation (□, RCTa; ○, RCTb; △, RCTc)



**FIGURE 3** Number of studies using a decision theory or recording an informed decision measure (□, decision theory only; ○, informed decision measure only; △, theory and informed decision measure)



**FIGURE 4** Number of observation studies by year (□, concurrent; ○, before/after same sample; △, before/after different samples)



# Chapter 4

## Discussion

This bibliography summarises 547 controlled studies evaluating the effect of a range of interventions on real patient decision making. The strengths are the systematic method of compilation, the quality classifications using the hierarchy of evidence and by theory, the categorisation of all measures recorded, and the classification by health domain and by type of intervention. We have identified under-researched areas for future empirical work and well-studied areas to which quantitative systematic reviews can be applied. The following discussion is divided into four parts: an evaluation of our methods; an overview of the findings; suggested uses of the review; and recommendations for further research.

### Evaluating the methods

#### Systematic identification

The advantage of integrating research by the systematic application of set criteria to all eligible articles is reduced bias. Other types of review tend to be guided by specific research agendas and may preferentially include studies supporting this. Recommendations drawn from this type of systematic review will be based on a fairer representation of the literature. The main disadvantage of a rigid methodology is the systematic exclusion of information from studies that do not fulfil the inclusion criteria.

Three areas of related research were excluded for this reason: studies of patient's preferences for health care; studies assessing changes in knowledge without recording behaviour; and comparative studies describing predictors of behaviour. Studies measuring preferences were excluded because these are only weakly associated with the final treatment choice or health decision. Nevertheless, as the patient is not committed to the final choice, they provide an excellent opportunity to evaluate more theoretically driven interventions.<sup>29-31</sup> Many such studies exist and their integration should be a priority for a future systematic review. Likewise, studies describing only knowledge change were excluded because knowledge alone is not a direct measure of informed patient decision making. Comparative studies describing predictors of behaviour such as the reasons why individual's attend or do not

attend for screening, were omitted because there is no evaluation of an intervention. Reviews in all three of these areas would probably increase understanding of how individuals make decisions, and which measures best assess informed decision making but would not evaluate intervention to facilitate informed patient decision making.

Although, this review systematically applied the pre-defined criteria to a large number of studies ( $n = 17,860$ ), not all the available evidence was evaluated. Resource constraints resulted in three electronic databases being searched for publications from only 5 years, and three journals handsearched for papers from a 10-year period. The complete electronic databases and reference sections from included articles or 'grey literature' were not searched. This deficit will have reduced the number of studies in the review. However, it was clear from the handsearching that patient rather than expert or professional decision making is a more recent area of interest.

### Overview of results

#### Describing the area researched

Although the search strategy effectively generated a high proportion of studies with an RCT design (*Table 1*), less than 10% of studies were classified as having a low risk of bias. Although some studies in which the randomisation procedures were not adequately explained may also have had a low risk of bias, this is disappointing. The number of such studies is rising only slowly and as a proportion of all studies, rising not at all (*Figure 2*).

The majority of interventions involved providing patients with more information (55%) and/or changing the delivery of information (50%). Few evaluated the manipulation of information (17%) or the effectiveness of patient prompts or decision aids (10%). This is disappointing as it is manipulation of information and the application of decision aids rather than additional information that are more likely to be associated with changes in informed patient decision making.<sup>32</sup>

The majority of decisions were concerned with health promotion. For example, decisions

regarding healthy eating or attendance for screening. There were few comparative studies evaluating decision making within the area of adherence to treatment, genetics, or active participation within the consultation. The least-researched areas of health appear to be genetics, obstetrics, surgery, dentistry and mental health. This is surprising as both prenatal and genetic testing particularly emphasise the importance of autonomy and informed patient decision making within their aims. Although there is an established literature evaluating the impact and uptake of such testing<sup>33</sup> few studies have used experimental designs to evaluate facilitation of informed patient decision making. The lack of studies in mental health is also particularly striking and should be remedied.

Over 90% of studies recorded some demographic details of participants. By identifying consistent variations within populations, interventions can be appropriately targeted. However, unless researchers investigate why these variations occur it is unlikely that interventions will be sufficiently informed. The results of *Table 10* suggest that few studies are able to explain any between-participant variations. Less than a third assess decision-making process measures or knowledge and less than an eighth assess affect so that few conclusions can be drawn about either the impact or interaction between emotion and decision making. Finally, only 4% assess any trait variations that may account for individual differences for preferences for amount of information or styles in utilising the information.

Only 18% of studies referred to both a theory to inform their study and assessed measures associated with the decision-making process. This figure is, if anything, over-generous. Any theory referred to within the study was recorded whether or not it was adequately operationalised. Additionally, less than 15% of studies were concerned with other factors that have been associated with the decision-making process such as time pressure and proxy decision making. As so few studies actually employed a theory that may explain variations in patient decision-making behaviour, it is not unreasonable to conclude that few experimental studies have actually evaluated informed patient decision making. The absolute rate of studies using a theory and recording a measure of informed decision making is rising only slowly in proportion to the overall rise in studies generally (*Figure 3*).

Only five studies were identified that were theory driven, assessed measures associated with informed decision-making, and used a randomised trial

design with a low risk of bias. This suggests that future NHS research effort in this area should concentrate on better quality primary research rather than on secondary reviews of existing evidence. As the rate of production of such good quality studies is low (*Figures 2 and 3*), the Health Technology Assessment programme need not repeat this systematic review for at least 5 years.

## Synthesising the review data

### **Can decision making be facilitated?**

The ideal evidence to answer this question could be provided by the five studies that assessed informed patient decision making using an RCT design with low risk of bias (*Table 21*). However, of these only three assessed the association between the intervention and decision making process measures. Although research exists evaluating the application of decision-making techniques to professionals' decision making and explaining patient decision making, it is clear that interventions aimed at facilitating patient informed decision making are under-researched.

### **Do people who have more information make better decisions?**

It is impossible to give a single answer to this apparently simple question because it implies that some decisions may be better than others and suggests that there is an external way of validating whether the decision made was correct. The criteria by which a decision is evaluated depends on the underlying theory. In the absence of theory it is not obvious for example whether an intervention that increases knowledge, alters behaviour, but also causes anxiety, is resulting in a more informed decision.

In this review, we assess whether additional information was associated with the health behaviour, but note that few studies operationalised a theory or aimed to manipulate factors associated with the decision process and outcome. Note also that because there was little consistency in the presentation of findings between studies, only a crude measure of intervention effect was created for this latter analysis. This indicated whether the study reported a change (effect), no change (no effect) or a mixture of change and no change (mixed effect) in the health behaviours. Consequently, the findings from this analysis are tentative.

Interventions, then, usually included two or more factors that may be associated with decision-making measures. Subsequent analysis could not compare the effectiveness of different interventions with health decision, as this classification did not have

mutually exclusive categories. Independent chi-square analysis had to be carried out for each intervention and the likelihood of reporting an effect (*Table 22*). The reporting of an effect did not mean that the decision was any more or less informed. The results of the analysis for interventions using additional information are located on row 1 (*Table 22*).

Half of the 301 studies evaluating additional information reported an effect of information on the decision behaviour. However, studies were no more likely to report an effect than report no or mixed effects. So, additional information may or may not be associated with decision behaviour. A more detailed breakdown of the additional information studies would be needed before conclusions could be drawn. For example, what information medium was used? Was the intervention comparing information with no information or information with another type of intervention? What was the quality of the information? All the study numbers of informed decision making studies using RCT designs (good quality) are listed (*Table 20*) to provide the basis for any reader attempting a quantitative review in these areas. For example, 45 studies were recorded under the additional information intervention. Of these 21 (47%) reported an effect on the decision behaviour, 17 (38%) no effect, and 7 (16%) mixed effects.

#### **Does the way in which we present information change the effectiveness of the decision?**

There were a number of interventions that may be referred to when answering this question: additional intervention; manipulation information; delivery information; group delivery information; and, patient prompts. Those studies that altered the delivery of information were more likely to report an effect. Twenty good quality trials manipulated the delivery of the information (*Table 20*), 11 (55%) reported an effect on decision behaviour. Studies that provided feedback and social skills training were more likely to report mixed effects on behaviour. If the 40 good-quality trials using feedback (*Table 20*) were more closely categorised, some reliable differences between effective and ineffective feedback interventions may be illustrated.

#### **Does the effectiveness of information on decision making vary by medical setting?**

The analysis suggests there is an association between the health domain in which the research is carried out and the likelihood of reporting of an effect (*Table 23*). Studies evaluating cancer related decisions were more likely to report an effect on behaviour. Such a finding may mean that inter-

ventions within the cancer domain are more refined so impacting on the behaviour or that there is something about cancer that interacts with the decision. It is probably not justified to extrapolate findings from studies in cancer to other health areas, and researchers need to be aware that the setting may affect the informed decision making process.

#### **Does the context of the decision affect decision making?**

The analysis suggests an association between the type of health decision and the likelihood of studies reporting an effect (*Table 24*). There is an effect from the type of health decision (*Table 23*) with studies concerned with lifestyle and utilisation of services being less likely to report a treatment effect than others. Similarly studies that showed little or no effect of interventions on lifestyle or service utilisation decisions should not be extrapolated to other types of decisions where more effects may be seen.

## **Suggested applications of the review**

### **Identification of under-researched areas**

Certain health areas and types of decisions have been more extensively researched than others and the studies in *Table 20* provide the basis for a number of systematic reviews. Bekker has embarked on two: a review of the effect of manipulation of information, and a review of the effect of the effect of provision of feedback and skills acquisition. Clinical genetics and prenatal diagnosis are other obvious under-researched areas. Work in these areas should be a priority, not only because patient decision making is widely recognised to be important here, but also because the strong scientific base provides an excellent test bed for improving our understanding of real patient decision making generally.

### **Design of the ideal study**

All studies in this area would benefit from reference to one of the decision-making theories, which were described in the chapter 1, at the design stage. A study evaluating how people make decisions ought to include a measure of the individual's decision-making process such as a description of the information used to make the decision. Knowledge measures alone are not sufficient. Explanatory studies will find a cross-sectional design the most useful, and a process tracing technique applied to the transcript of a consultation will usually provide the best understanding of how decisions were made.

Studies evaluating decision facilitation may use observational methods to formulate the research question, but must eventually use the best-quality experimental design, the RCT. Typically, the comparison should be against routine care, as there are no well-evaluated and clearly effective interventions in this area. The following is a description of an unpublished study carried out by Bekker. It is an example of an ideal study to evaluate the facilitation of informed decision making.

**Question: Does decision analysis increase informed decision making by women – choosing to have a diagnostic test in pregnancy after a screen positive?**

**Theory:** Decision analysis is a decision aid derived from expected utility theory. Expected utility theory, the theory of planned behaviour, and reasoned-based models of decision making informed the evaluation measures.

**Design:** RCT with a low risk of bias – numbered, sealed, opaque envelopes were opened for allocation to either a routine information consultation or a decision-analysis consultation.

**Measures:** The consultation was audio-tape recorded, transcribed and coded. A questionnaire was completed after the decision was made and 4 weeks later. The following variables were assessed:

- informed decision making: study specific coding frame and computer programme applied to transcripts assessing thinking words and information utilised
- reasoning: questionnaire included measures of beliefs, attitudes, perception of risk, and knowledge
- affect: anxiety and decisional conflict assessed by questionnaires, use of emotion-laden words assessed by the computer programme analysis of transcripts
- clinical: consultation length recorded and perceived usefulness of information
- behaviour: decision to be or not be tested
- other: demographic variations, reproductive history and individual differences in information processing assessed by questionnaires.

In summary the ideal study should randomise using a method with a low risk of bias, be designed with reference to decision-making theory, and include measures of the decision-making process and outcome.

## Recommendations for research

- Future primary research should not simply rely on behavioural or psychometric outcomes but should work under a theory of decision making, and record decision process measures that permit evaluation of whether the decision was informed. If evaluating experimental interventions, they must use RCT methods with a low risk of bias.
- A booklet describing the main decision-making theories, and an inventory of suitable questionnaires and interview-based outcome measures could be developed to help clinical health researchers design appropriate studies.
- Complementary systematic reviews of the following topics would be valuable:
  - trials evaluating the effect of interventions on patient preferences. At least 50 such studies were excluded from the present bibliography because no behaviour change was recorded.
  - observational studies furthering the understanding of real patient decision making. Three process-tracing methodologies will predominate: tape-recorded consultations; verbal protocols of thinking aloud techniques; and information boards or computer programmes to document the information referred to by patients.
  - assessing the effect of additional information, manipulation of information, provision of feedback, and group delivery of information on subsequent informed patient decision making.
- Primary research is required in the following health areas:
  - genetics and prenatal diagnosis. This area is a good place to begin because the nature of the information base, with relatively simple clearly-defined options with known risks corresponds closely to the terms that decision theories use.
  - areas where decisions are often made by proxy such as paediatrics and mental health.
- Primary research is required to evaluate the following types of interventions:
  - Decision aids, such as graphical, pictorial, computer and Internet-based devices.
  - Information manipulation, such as decision analysis, patient prompts and provision of feedback, and scenario-based approaches to informing choice of treatments.



## References

1. Ende J, Kazis L, Ash A, Moskowitz MA. Measuring patients' desire for autonomy: decision making and information-seeking preferences among medical patients. *J Gen Intern Med* 1989;**4**:23–30.
2. Ley P. Communicating with patients: improving communication, satisfaction and compliance. UK: Croom Helm, 1988.
3. Tversky A, Kahneman D. Judgement under uncertainty: heuristics and biases. *Science* 1974;**185**:1124–31.
4. Tversky A, Kahneman D. The framing of decisions and the psychology of choice. *Science* 1981;**211**:453–8.
5. Janis I, Mann L. Decision making: a psychological analysis of conflict, choice, and commitment. USA: Free Press, 1977.
6. Wilson TD, Schooler JW. Thinking too much: introspection can reduce the quality of preferences and decisions. *J Pers Soc Psychol* 1991;**60**:181–92.
7. Baron J. Thinking and deciding. 2nd edition. New York: Cambridge University Press, 1994.
8. Kleindorfer PR, Kunreuther HC, Schoemaker PJH. Decision sciences: an integrative perspective. New York: Cambridge University Press, 1993.
9. Savage LJ. The foundations of statistics. New York: Wiley.
10. Goldstein WM, Hogarth RM. Judgment and decision research: some historical context. In: Goldstein WM, Hogarth RM, editors. Research on judgment and decision making: currents, connections and controversies. New York: Cambridge University Press, 1997.
11. Simon HA. A behavioral model of rational choice. *QJ Econ* 1955;**69**:99–118.
12. Kahneman D, Slovic P, Tversky A, editors. Judgment under uncertainty: heuristics and biases. Cambridge: Cambridge University Press, 1982.
13. Payne JW, Bettman JR, Johnson EJ. The adaptive decision maker. Cambridge: Cambridge University Press, 1993.
14. Goodwin P, Wright G. Decision analysis for management judgement. 2nd edition. New York: Wiley, 1997.
15. Conner M, Norman P, editors. Predicting health behaviour: research and practice with social cognition models. Milton Keynes: Open University Press, 1996.
16. Bazerman MH, Tenbrusel AE, Wade-Benzani K. Negotiating with yourself and losing: making decisions with competing internal preferences. *Academy of Management Review* 1998;**23**:225–41.
17. Kuhberger A. Theoretical conceptions of framing effects in risky decisions. In: Ranyard R, Crozier WR, Svenson O, editors. Decision making: cognitive models and explanations. London: Routledge, 1997.
18. McNeil BJ, Pauker SG, Sox HC, Tversky A. On the elicitation of preferences for alternative therapies. *New Engl J Med* 1982;**306**:1259–62.
19. Christensen-Szalanski JJJ, Bushyhead JB. Physicians' use of probabilistic information in real clinical settings. *J Exp Psychol [Hum Percept]* 1981;**7**:928–35.
20. Mann L. Stress, affect and decision making. In: Yates FJ, editor. Risk-taking behavior. New York: Wiley, 1991.
21. Mano H. Risk taking, framing effects and affect. *Organizational Behavior and Human Decision Processes* 1994;**57**:35–58.
22. Svenson O, Maule AJ, editors. Time pressure and stress in human judgment and decision making. New York: Plenum, 1993.
23. Klein GA, Orasnu J, Calderwood R, Zsombok CE, editors. Decision making in action: models and methods. USA: Ablex Publishing Corporation, 1993.
24. Olson DL. Decision aids for selection problems. New York: Springer, 1996.
25. Van der Heijden K. Scenarios: the art of strategic conversation. London: Wiley, 1996.
26. Russo J, Shoemaker PJH. Confident decision making. Piatkus, 1989.
27. Klein GA. An overview of naturalistic decision making applications. In: Zsombok CE, Klein G, editors. Naturalistic decision making. New Jersey: Erlbaum, 1997.
28. Dickeson. In: Cooper HM, editor. Integrating research: a guide for literature review. 2nd edition. London: Sage, 1989.
29. Siegert EA, Clipp EC, Mulhausen P, Kochersberger. Impact of advance directive videotape on patient comprehension and treatment preferences. *Arch Fam Med* 1996;**5**:207–12.

30. Mazur DJ, Hickam. Patients and physicians' interpretations of graphic data displays. *Med Decis Making* 1993;**13**:59–63.
31. Mazur DJ, Merz JF. Older patients' willingness to trade off urologic adverse outcomes for a better chance at five-year survival in the clinical setting of prostate cancer. *J Am Geriatr Soc* 1995;**43**:979–84.
32. O'Connor AM, Llewellyn-Thomas HA, Drake ER. An annotated bibliography of research on shared decision making. National Cancer Institute of Canada 1995. Available from URL: <http://hiru.mcmaster.ca/cpep/sdm/>
33. Marteau TM, Richards M, editors. *The troubled helix*. 1997

# Appendix I

## References used to develop the search strategy


- Bekker H, Modell M, Dennis G, Silver A, Matthew C, Bobrow M, *et al.* Uptake of cystic fibrosis testing in primary care; supply push or demand pull. *BMJ* 1993;**306**:1584–6.
- Clayton EW, Hannig VL, Pfothenhauer JP, *et al.* Teaching about cystic fibrosis carrier screening by using written and video information. *Am J Hum Genet*;57:171–81.
- Field T, Sandberg D, Quetel TA, Garcia R, Rosario M. Effects of ultrasound feedback on pregnancy anxiety, fetal activity, and neonatal outcome. *Obstet Gynecol* 1985;**66**:525–8.
- Freda MC, Andersen HF, Damus K, Merkatz IR. Are there differences in information given to private and public prenatal patients? *Am J Obstet Gynecol* 1993;**169**:155–60.
- Heckerling PS, Verp MS. Amniocentesis or chorionic villus sampling for prenatal genetic testing: a decision analysis. *J Clin Epidemiol* 1991;**44**(7):657–70.
- Lichtenstein S, Newman JR. Empirical scaling of common verbal phrases associated with numerical probabilities. *Psychol Sci* 1987;**9**(10):563–4.
- Livingstone J, Axton RA, Mennie M, Gilfillan A, Brock. A preliminary trial of couple screening for cystic fibrosis; designing an appropriate information leaflet. *Clin Genet* 1993;**43**:57–62.
- Llewellyn-Thomas HA, Thiel EC, Sem FW, Woermke DE. Presenting clinical trial information: a comparison of methods. *Patient Education and Counseling* 1995;**25**:97–107.
- Marteau TM, Kidd J, Michie S, Cook R, Johnston M, Shaw RW. Anxiety knowledge and satisfaction in women receiving false positive results on routine prenatal screening: a randomised controlled trial. *J Psychosom Obstet Gynaecol* 1993;**14**:185–96.
- Mazur DJ, Merz JF. Older patients' willingness to trade off urologic adverse outcomes for a better chance at five-year survival in the clinical setting of prostate cancer. *J Am Geriatr Soc* 1995;**43**:979–84.
- Miedzybrodzka ZH, Hall MH, Mollison J, Templeton A, Russell IT, Dean JC, *et al.* Antenatal screening for carriers of cystic fibrosis: randomised trial of stepwise v couple screening. *BMJ* 1995;**310**:353–7.
- Pauker SP, Pauker SG. The amniocentesis decision: ten years of decision analytic experience. Birth defects; original article series 23(2):151–69. March of Dimes Birth Defects Foundation, 1987.
- Savage R, Armstrong D. Effects of a general practitioners consulting style on patient satisfaction: a controlled study. *BMJ* 1990;**301**:968–70.
- Simes RJ, Tattersall MH, Coates AS, Raghaven D, Solomon HG, Smartt H. Randomised comparison of procedures for obtaining informed consent in clinical trials of treatment for cancer. *BMJ* 1986;**293**:1065–8.
- Schapiro MC, Najman A, Change J, Keeping D, Morrison J, Western JS. Information control and the exercise of power in the obstetrical encounter. *Soc Sci Med* 1983;**17**:139–46.
- Thornton JG, Hewison J, Lilford RJ, Vail A. A randomised trial of three methods of giving information about prenatal testing. *BMJ* 1995;**311**:1127–30.
- Wilkinson C, Jones JM, McBride J. Anxiety caused by abnormal result of cervical smear test: a controlled trial. *BMJ* 1990;**300**:440.





## Appendix 2

### The principle of the search strategy





## Appendix 3

### MEDLINE search strategy

career choice/  
 exp decision making/  
 2 not 1  
 exp communication/  
 exp cybernetics/  
 exp advertising/  
 exp animal communication/  
 5 or 6 or 7  
 4 not 8  
 exp decision support techniques/  
 "neural networks (computer)"/  
 10 not 11  
 informed consent/  
 exp thinking/  
 exp autism/  
 creativeness/  
 exp esthetics/  
 15 or 16 or 17  
 14 not 18  
 models, psychological/  
 exp probability/  
 exp risk/  
 risk-taking/  
 exp psychological theory/  
 exp patient acceptance of health care/  
 exp patient participation/  
 exp knowledge, attitudes, practice/  
 exp truth disclosure/  
 exp treatment refusal/  
 (patient adj3 decision).ti,ab.  
 fee\$ charg\$.ti,ab.  
 willing\$ to pay.ti,ab.  
 or/3,9,12-13,19-32

exp consumer participation/  
 exp consumer satisfaction/  
 exp health education/  
 health status/  
 exp professional-patient relations/  
 exp patient acceptance of health care/  
 exp public opinion/  
 sick role/  
 genetic counseling/  
 exp counseling/  
 psychology, applied/  
 (referral and consultation).ti,ab,sh.  
 exp attitude to health/  
 (patient adj3 decision).ti,ab.  
 or/34-47  
 comparative study/  
 cross-sectional studies/  
 exp data collection/  
 exp statistics/  
 exp evaluation studies/  
 exp feasibility studies/  
 exp cohort studies/  
 exp random allocation/  
 exp clinical trials/  
 exp research/  
 exp "study design (non mesh)"/  
 exp pilot projects/  
 exp intervention studies/  
 (clinical adj2 trial).ti,ab.  
 (control\$ adj2 trial\$).ti,ab.  
 measure\$.ti,ab.  
 or/49-64  
 33 and 48 and 65



# Appendix 4

## Data extraction coding forms

Page 1, Table 1 — Article identification details

<i>Field</i>	<i>Details</i>
Article review number	.....
number studies	.....
project number	.....
number of duplicate	.....
Authors	.....
Title	.....
Journal details	.....
Included/excluded (reasons exclude)	<b>included</b> <b>excluded</b> .....
Database	MEDLINE <input type="checkbox"/> PsycLIT <input type="checkbox"/> BIDS <input type="checkbox"/> handsearch <input type="checkbox"/>
Date downloaded	____ / ____
Date sent to reviewer	____ / ____
Date returned reviewer	____ / ____
Initials reviewer	MA <input type="checkbox"/> HB <input type="checkbox"/> JC <input type="checkbox"/> JH <input type="checkbox"/> JM <input type="checkbox"/> MM <input type="checkbox"/> SM <input type="checkbox"/> AP <input type="checkbox"/> MR <input type="checkbox"/> JT <input type="checkbox"/>

Page 2, Table 1 – Describing health decision and quality study

Summary health decision (free text)		
<p><b>Location research:</b> unknown <input type="checkbox"/> UK <input type="checkbox"/> North America <input type="checkbox"/> S. Africa <input type="checkbox"/> Australia/NZ <input type="checkbox"/> other <input type="checkbox"/> (write ..... .....)</p> <p><b>Domain of health care:</b> Write type of illness or health domain research located (e.g. cervical/breast cancer, asthma, health promotion, cardiovascular risk factors, HIV/condom use, smoking cessation, drug use, tuberculoses, malaria, etc.) ..... .....</p> <p><b>Quality of study:</b> <i>Design of study:</i> randomised (RCT) a <input type="checkbox"/> b <input type="checkbox"/> c <input type="checkbox"/> non-randomised concurrent <input type="checkbox"/> before/after different sample <input type="checkbox"/> before/after same sample <input type="checkbox"/> Other (e.g. matched case-controlled) <input type="checkbox"/> .....</p> <p><b>Quality of study continued...</b> <i>Causes for concern:</i> no/not obvious causes for concern <input type="checkbox"/> yes (please write any concerns) <input type="checkbox"/> ..... .....</p>	<p><b>Participants:</b> unknown <input type="checkbox"/> adult <input type="checkbox"/> well <input type="checkbox"/> pregnant <input type="checkbox"/> physically impaired <input type="checkbox"/> elderly (&gt; 65) <input type="checkbox"/> mentally impaired <input type="checkbox"/> child (&lt; 17) <input type="checkbox"/> non-English speaking <input type="checkbox"/></p> <p><b>Defining type of health decision:</b> Write type of health decision (e.g. screening, diagnostic testing, medication/drug therapy, surgical treatment, behavioural therapy, willingness to pay, research, cessation, etc.) ..... .....</p> <p><b>Quality of study continued...</b> <i>Level of intervention:</i> patient level <input type="checkbox"/> other level <input type="checkbox"/> not rec. <input type="checkbox"/> <i>Sample invited to participate:</i> total 'population' available <input type="checkbox"/> stratified/systematic sample <input type="checkbox"/> volunteer/non-systematic sample <input type="checkbox"/> not adequately described <input type="checkbox"/></p> <p><b>Quality of study continued...</b> <i>Development intervention materials:</i> piloted intervention: no <input type="checkbox"/> yes <input type="checkbox"/> in part <input type="checkbox"/> piloted measures: no <input type="checkbox"/> yes <input type="checkbox"/> in part <input type="checkbox"/> readability score: no <input type="checkbox"/> yes <input type="checkbox"/> in part <input type="checkbox"/></p>	<p><b>The nature of the decision is:</b> A) real <input type="checkbox"/> intended <input type="checkbox"/> hypothetical <input type="checkbox"/> B) implicit <input type="checkbox"/> explicit <input type="checkbox"/> C) affects – participant <input type="checkbox"/> other <input type="checkbox"/> D) made at – time intervention <input type="checkbox"/> later <input type="checkbox"/></p> <p><b>Authors awareness theory:</b> <i>Decision making theory:</i> 0 = no theory referred to Write in the name of theory referred to in the introduction or methods section ... ..... .....</p> <p><b>Quality of study continued...</b> <i>Sample size:</i> total number available ____ not rec. <input type="checkbox"/> total number invited ____ not rec. <input type="checkbox"/> number participated ____ not rec. <input type="checkbox"/> number excluded ____ not rec. <input type="checkbox"/> number in final analysis ____ not rec. <input type="checkbox"/></p> <p><b>Reference to be accessed ...</b></p>

Page 3, Table 1 – Description of intervention groups

Nature of intervention:	explicit information <input type="checkbox"/> change in service <input type="checkbox"/> other <input type="checkbox"/>		
Comparative group	Description of intervention (write ...)	Code	Number in group (n)
Group 1	..... .....	[ ] [ ] [ ] [ ]	
Group 2	..... .....	[ ] [ ] [ ] [ ]	
Group 3	..... .....	[ ] [ ] [ ] [ ]	
Group 4	..... .....	[ ] [ ] [ ] [ ]	
Group 5	..... .....	[ ] [ ] [ ] [ ]	

Page 3, Table 2 – Variables described within article

Variable	Code	Descriptives	Main effect analysis	Interaction analysis
Write variables described within the study (e.g. age, gender, attitudes, social support)				
Alongside each variable, tick whether the analysis of the variable was descriptive, main effect or interaction.				

Page 4, Table 1 – Description health decision outcome

Description (write...)	Continuous/categorical	Validated, piloted, study specific	Description validated measure (write):
		val <input type="checkbox"/> pil <input type="checkbox"/> study <input type="checkbox"/>	
		val <input type="checkbox"/> pil <input type="checkbox"/> study <input type="checkbox"/>	

Page 4, Table 2 – Results evaluating the association of the intervention and decision outcome

Variable	Categorical level/group e.g. adhere 1 not adhere 2	Group 1 not available or 'n' or mean (sd) n	Group 2 not available or 'n' or mean (sd) n	Group 3 not available or 'n' or mean (sd) n	Group 4 not available or 'n' or mean (sd) n	Group 5 not available or 'n' or mean (sd) n

sd = standard deviation

Page 4, Table 3 –Text box summarising the results of the intervention

Write summary results of intervention on decision outcome and author's main outcome measures.

## Appendix 5

### Bibliography: studies by study number

- 1 - 1** Lindholm LH, Ekblom T, Dash C, Isacson A, Schersten B. Changes in cardiovascular risk factors by combined pharmacological and nonpharmacological strategies: the main results of the CELL Study. *Journal of Internal Medicine* 1996;**240**(1):13–22.
- 3 - 1** Davies-Adetugbo AA. Promotion of breast feeding in the community: impact of health education programme in rural communities in Nigeria. *Journal of Diarrhoeal Diseases Research* 1996;**14**(1):5–11.
- 4 - 1** Hoffman JA, Caudill BD, Koman JJ 3rd, Luckey JW, Flynn PM, Mayo DW. Psychosocial treatments for cocaine abuse. 12-month treatment outcomes. *Journal of Substance Abuse Treatment* 1996;**13**(1):3–11.
- 6 - 1** El-Chaar GM, Mardy G, Wehlou K, Rubin LG. Randomized, double blind comparison of brand and generic antibiotic suspensions: II. A study of taste and compliance in children. *Pediatric Infectious Disease Journal* 1996;**15**(1):18–22.
- 6 - 2** El-Chaar GM, Mardy G, Wehlou K, Rubin LG. Randomized, double blind comparison of brand and generic antibiotic suspensions: II. A study of taste and compliance in children. *Pediatric Infectious Disease Journal* 1996;**15**(1):18–22.
- 6 - 3** El-Chaar GM, Mardy G, Wehlou K, Rubin LG. Randomized, double blind comparison of brand and generic antibiotic suspensions: II. A study of taste and compliance in children. *Pediatric Infectious Disease Journal* 1996;**15**(1):18–22.
- 10 - 1** Toseland RW, O'Donnell JC, Engelhardt JB, Hendlar SA, Richie JT, Jue D. Outpatient geriatric evaluation and management. Results of a randomized trial. *Medical Care* 1996;**34**(6):624–40.
- 17 - 1** Smith MU, Katner HP. Quasi-experimental evaluation of three AIDS prevention activities for maintaining knowledge, improving attitudes, and changing risk behaviors of high school seniors. *AIDS Education and Prevention* 1995;**7**(5):391–402.
- 18 - 1** Goldberg L, Elliot DL, Clarke GN, MacKinnon DP, Zoref L, Moe E, Green C, Wolf SL. The Adolescents Training and Learning to Avoid Steroids (ATLAS) prevention program. Background and results of a model intervention. *Archives of Pediatrics and Adolescent Medicine* 1996;**150**(7):713–21.
- 19 - 1** Mazur DJ, Hickam DH. Five-year survival curves: how much data are enough for patient–physician decision making in general surgery? *European Journal of Surgery* 1996;**162**(2):101–4.
- 20 - 1** Gil KM, Wilson JJ, Edens JL, Webster DA, Abrams MA, Orringer E, Grant M, Clark WC, Janal MN. Effects of cognitive coping skills training on coping strategies and experimental pain sensitivity in African American adults with sickle cell disease. *Health Psychology* 1996;**15**(1):3–10.
- 22 - 1** Pauw J, Ferrie J, Rivera Villegas R, Medrano Martinez J, Gorter A, Egger M. A controlled HIV/AIDS-related health education programme in Managua, Nicaragua. *AIDS* 1996;**10**(5):537–44.
- 23 - 1** Hagopian GA. The effects of informational audiotapes on knowledge and self-care behaviors of patients undergoing radiation therapy. *Oncology Nursing Forum* 1996;**23**(4):697–700.
- 24 - 1** Gordon CM, Carey MP. Alcohol's effects on requisites for sexual risk reduction in men: an initial experimental investigation. *Health Psychology* 1996;**15**(1):56–60.
- 26 - 1** Hartley BM, O'Connor ME. Evaluation of the 'Best Start' breast-feeding education program. *Archives of Pediatrics and Adolescent Medicine* 1996;**150**(8):868–71.
- 27 - 1** Maisiak R, Austin J, Heck L. Health outcomes of two telephone interventions for patients with rheumatoid arthritis or osteoarthritis. *Arthritis and Rheumatism* 1996;**39**(8):1391–9.
- 28 - 1** Mercer BS. A randomized study of the efficacy of the PROPATH Program for patients with Parkinson disease. *Archives of Neurology* 1996;**53**(9):881–4.
- 32 - 1** Agras WS, Berkowitz RI, Arnow BA, Telch CF, Marnell M, Henderson J, Morris Y, Wilfley DE. Maintenance following a very-low-calorie diet. *Journal of Consulting and Clinical Psychology* 1996;**64**(3):610–3.
- 34 - 1** Foley EC, D'Amico F, Merenstein JH. Five-year follow-up of a nurse-initiated intervention to improve mammography recommendation. *Journal of the American Board of Family Practice* 1995;**8**(6):452–6.
- 35 - 1** O'Hara P, Messick BJ, Fichtner RR, Parris D. A peer-led AIDS prevention program for students in an alternative school. *Journal of School Health* 1996;**66**(5):176–82.

- 37 - 1 Yancey AK, Tanjasiri SP, Klein M, Tunder J. Increased cancer screening behavior in women of color by culturally sensitive video exposure. *Preventive Medicine* 1995;**24**(2):142-8.
- 38 - 1 Bearss N, Santelli JS, Papa P. A pilot program of contraceptive continuation in six school-based clinics. *Journal of Adolescent Health* 1995;**17**(3):178-83.
- 39 - 1 Bowman J, Sanson-Fisher R, Boyle C, Pope S, Redman S. A randomised controlled trial of strategies to prompt attendance for a Pap smear. *Journal of Medical Screening* 1995;**2**(4):211-8.
- 40 - 1 Rabin JM, McNett J, Badlani GH. 'Compu-Void II': the computerized voiding diary. *Journal of Medical Systems* 1996;**20**(1):19-34.
- 41 - 1 Turner KM, Wilson BJ, Gilbert FJ. Improving breast screening uptake: persuading initial non-attenders to attend. *Journal of Medical Screening* 1994;**1**(3):199-202.
- 42 - 1 Gillies J, Barry D, Crane J, Jones D, MacLennan L, Pearce N, Reid J, Toop L. A community trial of a written self management plan for children with asthma. Asthma Foundation of NZ Children's Action. *New Zealand Medical Journal* 1996;**109**(1015):30-3.
- 44 - 1 Gardiner JC, Mullan PB, Rosenman KD, Zhu Z, Swanson GM. Mammography usage and knowledge about breast cancer in a Michigan farm population before and after an educational intervention. *Journal of Cancer Education* 1995;**10**(3):155-62.
- 45 - 1 Heath GW, Fuchs R, Croft JB, Temple SP, Wheeler FC. Changes in blood cholesterol awareness: final results from the South Carolina Cardiovascular Disease Prevention Project. *American Journal of Preventive Medicine* 1995;**11**(3):190-6.
- 46 - 1 Kash KM, Holland JC, Osborne MP, Miller DG. Psychological counseling strategies for women at risk of breast cancer. *Monographs - National Cancer Institute* 1995;(17):73-9.
- 47 - 1 Lave JR, Ives DG, Traven ND, Kuller LH. Participation in health promotion programs by the rural elderly. *American Journal of Preventive Medicine* 1995;**11**(1):46-53.
- 48 - 1 Straton JA, Sutherland GJ, Hyndman JC. Cervical cancer screening for hospital inpatients: report of an intervention study. *Australian Journal of Public Health* 1995;**19**(3):288-93.
- 49 - 1 Sunwood J, Brenman A, Escobedo J, Philpott T, Allman K, Mueller J, Jaeger J, Brown LK, Cole FS. School-based AIDS education for adolescents. *Journal of Adolescent Health* 1995;**16**(4):309-15.
- 50 - 1 Reid C, McNeil JJ, Williams F, Powles J. Cardiovascular risk reduction: a randomized trial of two health promotion strategies for lowering risk in a community with low socioeconomic status. *Journal of Cardiovascular Risk* 1995;**2**(2):155-63.
- 51 - 1 Ellerbeck E, Khallaf N, el Ansary KS, Moursi S, Black R. Caretaker compliance with different antibiotic formulations for treatment of childhood pneumonia. *Journal of Tropical Pediatrics* 1995;**41**(2):103-8.
- 52 - 1 Mohler PJ. Enhancing compliance with screening mammography recommendations: a clinical trial in a primary care office. *Family Medicine* 1995;**27**(2):117-21.
- 53 - 1 Friedman RH, Kazis LE, Jette A, Smith MB, Stollerman J, Torgerson J, Carey K. A telecommunications system for monitoring and counseling patients with hypertension. Impact on medication adherence and blood pressure control. *American Journal of Hypertension* 1996;**9**(4 Pt 1):285-92.
- 55 - 1 Sussman S, Parker VC, Lopes C, Crippens DL, Elder P, Scholl D. Empirical development of brief smoking prevention videotapes which target African-American adolescents. *International Journal of the Addictions* 1995;**30**(9):1141-64.
- 55 - 2 Sussman S, Parker VC, Lopes C, Crippens DL, Elder P, Scholl D. Empirical development of brief smoking prevention videotapes which target African-American adolescents. *International Journal of the Addictions* 1995;**30**(9):1141-64.
- 56 - 1 Christensen B. Payment and attendance at general practice preventive health examinations. *Family Medicine* 1995;**27**(8):531-4.
- 57 - 1 Detry JM, Block P, De Backer G, Degaute JP. Patient compliance and therapeutic coverage: comparison of amlodipine and slow release nifedipine in the treatment of hypertension. The Belgian Collaborative Study Group. *European Journal of Clinical Pharmacology* 1995;**47**(6):477-81.
- 58 - 1 Sanner MA, Hedman H, Tufveson G. Evaluation of an organ-donor-card campaign in Sweden. *Clinical Transplantation* 1995;**9**(4):326-33.
- 61 - 1 Bjorge T, Gunbjorud AB, Haugen OA, Skare GB, Trope C, Thoresen SO. Mass screening for cervical cancer in Norway: evaluation of the pilot project. *Cancer Causes and Control* 1995;**6**(6):477-84.
- 63 - 1 Cherkin DC, Deyo RA, Street JH, Hunt M, Barlow W. Pitfalls of patient education. Limited success of a program for back pain in primary care. *Spine* 1996;**21**(3):345-55.



- 64 - 1 Burton LC, Paglia MJ, German PS, Shapiro S, Damiano AM. The effect among older persons of a general preventive visit on three health behaviors: smoking, excessive alcohol drinking, and sedentary lifestyle. The Medicare Preventive Services Research Team. *Preventive Medicine* 1995;24(5):492-7.
- 67 - 1 Griffith CH 3rd, Wilson JF, Emmett KR, Ramsbottom-Lucier M, Rich EC. Knowledge and experience with Alzheimer's disease. Relationship to resuscitation preference. *Archives of Family Medicine* 1995;4(9):780-4.
- 71 - 1 Elk R, Schmitz J, Manfredi L, Rhoades H, Andres R, Grabowski J. Cessation of cocaine use during pregnancy: a preliminary comparison. *Addictive Behaviors* 1994;19(6):697-702.
- 72 - 1 Berman BA, Gritz ER, Braxton-Owens H, Nisenbaum R. Targeting adult smokers through a multi-ethnic public school system. *Journal of Cancer Education* 1995;10(2):91-101.
- 74 - 1 Conget I, Jansa M, Vidal M, Vidal J, Manzanares JM, Gomis R. Effects of an individual intensive educational control program for insulin-dependent diabetic subjects with poor metabolic control. *Diabetes Research and Clinical Practice* 1995;27(3):189-92.
- 75 - 1 Della Valle CJ, Levitz CL, Bora FW Jr. Health care utilization and attitudes toward health insurance. A comparison of privately insured and medical assistance or uninsured patients. *American Journal of Orthopedics* 1995;24(6):483-7.
- 79 - 1 Fardy PS, White RE, Clark LT, Amodio G, Hurster MH, McDermott KJ, Magel JR. Health promotion in minority adolescents: a Healthy People 2000 pilot study. *Journal of Cardiopulmonary Rehabilitation* 1995;15(1):65-72.
- 80 - 1 Oldenburg B, Martin A, Greenwood J, Bernstein L, Allan R. A controlled trial of a behavioral and educational intervention following coronary artery bypass surgery. *Journal of Cardiopulmonary Rehabilitation* 1995;15(1):39-46.
- 82 - 1 Boulet LP, Boutin H, Cote J, Leblanc P, Laviolette M. Evaluation of an asthma self-management education program. *Journal of Asthma* 1995;32(3):199-206.
- 83 - 1 Binstock MA, Wolde-Tsadik G. Alternative prenatal care. Impact of reduced visit frequency, focused visits and continuity of care. *Journal of Reproductive Medicine* 1995;40(7):507-12.
- 84 - 1 Kuthy S, Grap MJ, Penn L, Henderson V. After the party's over: evaluation of a drinking and driving prevention program. *Journal of Neuroscience Nursing* 1995;27(5):273-7.
- 85 - 1 Ferris DG, Golden NH, Petry LJ, Litaker MS, Nackenson M, Woodward LD. Effectiveness of breast self-examination prompts on oral contraceptive packaging. *Journal of Family Practice* 1996;42(1):43-8.
- 86 - 1 Navarro AM, Senn KL, Kaplan RM, McNicholas L, Campo MC, Roppe B. Por La Vida intervention model for cancer prevention in Latinas. *Monographs - National Cancer Institute* 1995;(18):137-45.
- 88 - 1 Williford SL, Johnson DF. Impact of pharmacist counseling on medication knowledge and compliance. *Military Medicine* 1995;160(11):561-4.
- 89 - 1 Boudreau F, Godin G, Pineau R, Bradet R. Health risk appraisal in an occupational setting and its impact on exercise behavior. *Journal of Occupational and Environmental Medicine* 1995;37(9):1145-50.
- 91 - 1 DiScenza S, Nies M, Jordan C. Effectiveness of counseling in the health promotion of HIV-positive clients in the community. *Public Health Nursing* 1996;13(3):209-16.
- 95 - 1 Simeoni E, Bauman A, Stenmark J, O'Brien J. Evaluation of a community arthritis program in Australia: dissemination of a developed program. *Arthritis Care and Research* 1995;8(2):102-7.
- 96 - 1 McFarlane WR, Lukens E, Link B, Dushay R, Deakins SA, Newmark M, Dunne EJ, Horen B, Toran J. Multiple-family groups and psycho-education in the treatment of schizophrenia. *Archives of General Psychiatry* 1995;52(8):679-87.
- 97 - 1 Bian JY, Zhang BX, Rong WS. Evaluating the social impact and effectiveness of four-year 'Love Teeth Day' campaign in China. *Advances in Dental Research* 1995;9(2):130-3.
- 98 - 1 Taal E, Riemsma RP, Brus HL, Seydel ER, Rasker JJ, Wiegman O. Group education for patients with rheumatoid arthritis. *Patient Education and Counseling* 1993;20(2-3):177-87.
- 99 - 1 Forsyth MC, Fulton DL, Lane DS, Burg MA, Krishna M. Changes in knowledge, attitudes and behavior of women participating in a community outreach education program on breast cancer screening. *Patient Education and Counseling* 1992;19(3):241-50.
- 100 - 1 Maurer WJ. Breast cancer screening complacency and compliance. *Wisconsin Medical Journal* 1995;94(6):305-6.
- 101 - 1 Kendall C, Hailey BJ. The relative effectiveness of three reminder letters on making and keeping mammogram appointments. *Behavioral Medicine* 1993;19(1):29-34.

- 102 - 1** Glasgow RE, Hollis JF, Ary DV, Boles SM. Results of a year-long incentives-based worksite smoking-cessation program. *Addictive Behaviors* 1993;**18**(4):455–64.
- 104 - 1** Boehm S, Schlenk EA, Raleigh E, Ronis D. Behavioral analysis and behavioral strategies to improve self-management of type II diabetes. *Clinical Nursing Research* 1993;**2**(3):327–44.
- 105 - 1** Murray MD, Birt JA, Manatunga AK, Darnell JC. Medication compliance in elderly outpatients using twice-daily dosing and unit-of-use packaging. *Annals of Pharmacotherapy* 1993;**27**(5):616–21.
- 106 - 1** Stevens-Simon C, O'Connor P, Bassford K. Incentives enhance postpartum compliance among adolescent prenatal patients. *Journal of Adolescent Health* 1994;**15**(5):396–9.
- 107 - 1** Caceres CF, Rosasco AM, Mandel JS, Hearst N. Evaluating a school-based intervention for STD/AIDS prevention in Peru. *Journal of Adolescent Health* 1994;**15**(7):582–91.
- 108 - 1** Park SI, Saxe JC, Weesner RE. Does use of the Coloscreen Self-Test improve patient compliance with fecal occult blood screening? *American Journal of Gastroenterology* 1993;**88**(9):1391–4.
- 110 - 1** Fries JF, Bloch DA, Harrington H, Richardson N, Beck R. Two-year results of a randomized controlled trial of a health promotion program in a retiree population: the Bank of America Study. *American Journal of Medicine* 1993;**94**(5):455–62.
- 111 - 1** Holley JL, Nespor S, Rault R. The effects of providing chronic hemodialysis patients written material on advance directives. *American Journal of Kidney Diseases* 1993;**22**(3):413–8.
- 114 - 1** Champion V, Huster G. Effect of interventions on stage of mammography adoption. *Journal of Behavioral Medicine* 1995;**18**(2):169–87.
- 115 - 1** Anderson AS, Campbell DM, Shepherd R. The influence of dietary advice on nutrient intake during pregnancy. *British Journal of Nutrition* 1995;**73**(2):163–77.
- 117 - 1** Weinrich SP, Weinrich MC, Stromborg MF, Boyd MD, Weiss HL. Using elderly educators to increase colorectal cancer screening. *Gerontologist* 1993;**33**(4):491–6.
- 118 - 1** McIntosh NA, Clark NM, Howatt WF. Reducing tobacco smoke in the environment of the child with asthma: a cotinine-assisted, minimal-contact intervention. *Journal of Asthma* 1994;**31**(6):453–62.
- 120 - 1** Shope JT, Kloska DD, Dielman TE, Maharg R. Longitudinal evaluation of an enhanced alcohol misuse prevention study (AMPS) curriculum for grades six-eight. *Journal of School Health* 1994;**64**(4):160–6.
- 122 - 1** Verne J, Kettner J, Mant D, Farmer A, Mortenson N, Northover J. Self-administered faecal occult blood tests do not increase compliance with screening for colorectal cancer: results of a randomized controlled trial. *European Journal of Cancer Prevention* 1993;**2**(4):301–5.
- 123 - 1** Remafedi G. Cognitive and behavioral adaptations to HIV/AIDS among gay and bisexual adolescents. *Journal of Adolescent Health* 1994;**15**(2):142–8.
- 124 - 1** Weiden P, Rapkin B, Zygmunt A, Mott T, Goldman D, Frances A. Postdischarge medication compliance of inpatients converted from an oral to a depot neuroleptic regimen. *Psychiatric Services* 1995;**46**(10):1049–54.
- 125 - 1** Magura S, Kang SY, Shapiro JL. Outcomes of intensive AIDS education for male adolescent drug users in jail. *Journal of Adolescent Health* 1994;**15**(6):457–63.
- 126 - 1** Kurtz ME, Kurtz JC, Given B, Given CC. Promotion of breast cancer screening in a work site population. *Health Care for Women International* 1994;**15**(1):31–42.
- 132 - 1** Kasemsarn P, Ngarmpiyasakul C, Phongpanich S, Pulkasisri N. Baby-friendly hospital: how to sustain? *Journal of the Medical Association of Thailand* 1995;**78**(7):362–8.
- 133 - 1** Sobczyk W, Hazel N, Reed CD, Ciarroccki B, Cohen S, Varga D. Health Promotion Schools of Excellence: a model program for Kentucky and the nation. *Journal of the Kentucky Medical Association* 1995;**93**(4):142–7.
- 135 - 1** Sciacca JP, Dube DA, Phipps BL, Ratliff MI. A breast feeding education and promotion program: effects on knowledge, attitudes, and support for breast feeding. *Journal of Community Health* 1995;**20**(6):473–90.
- 136 - 1** Girgis A, Sanson-Fisher RW, Tripodi DA, Golding T. Evaluation of interventions to improve solar protection in primary schools. *Health Education Quarterly* 1993;**20**(2):275–87.
- 137 - 1** Taddio A, Ito S, Einarson TR, Leeder JS, Koren G. Effect of counseling on maternal reporting of adverse effects in nursing infants exposed to antibiotics through breast milk. *Reproductive Toxicology* 1995;**9**(2):153–7.
- 139 - 1** Kirkman MS, Weinberger M, Landsman PB, Samsa GP, Shortliffe EA, Simel DL, Feussner JR. A telephone-delivered intervention for patients with NIDDM. Effect on coronary risk factors. *Diabetes Care* 1994;**17**(8):840–6.

- 140 - 1** Gritz ER, Carr CR, Rapkin D, Abemayor E, Chang LJ, Wong WK, Belin TR, Calcaterra T, Robbins KT, Chonkich G, Beumer J, Ward PH. Predictors of long-term smoking cessation in head and neck cancer patients. *Cancer Epidemiology, Biomarkers and Prevention* 1993;**2**(3):261-70.
- 142 - 1** Ingold FR, Toussirt M. Transmission of HIV among drug addicts in three French cities: implications for prevention. *Bulletin on Narcotics* 1993;**45**(1):117-34.
- 143 - 1** McFall SL, Michener A, Rubin D, Flay BR, Mermelstein RJ, Burton D, Jelen P, Warnecke RB. The effects and use of maintenance newsletters in a smoking cessation intervention. *Addictive Behaviors* 1993;**18**(2):151-8.
- 145 - 1** Santi SM, Cargo M, Brown KS, Best JA, Cameron R. Dispositional risk factors for smoking-stage transitions: a social influences program as an effect modifier. *Addictive Behaviors* 1994;**19**(3):269-85.
- 146 - 1** Ohmit SE, Furumoto-Dawson A, Monto AS, Fasano N. Influenza vaccine use among an elderly population in a community intervention. *American Journal of Preventive Medicine* 1995;**11**(4):271-6.
- 147 - 1** Lloyd HM, Paisley CM, Mela DJ. Barriers to the adoption of reduced-fat diets in a UK population. *Journal of the American Dietetic Association* 1995;**95**(3):316-22.
- 149 - 1** Morris BA, Trimble NE, Fendley SJ. Increasing bicycle helmet use in the community. Measuring response to a wide-scale, 2-year effort. *Canadian Family Physician* 1994;**40**:1126-31.
- 150 - 1** Delp C, Jones J. Communicating information to patients: the use of cartoon illustrations to improve comprehension of instructions. *Academic Emergency Medicine* 1996;**3**(3):264-70.
- 154 - 1** Richardson JL, Mondrus GT, Danley K, Deapen D, Mack T. Impact of a mailed intervention on annual mammography and physician breast examinations among women at high risk of breast cancer. *Cancer Epidemiology, Biomarkers and Prevention* 1996;**5**(1):71-6.
- 155 - 1** Shafir MS, Silversides C, Waters I, MacRury K, Frank JW, Becker LA. Patient consent to observation. Responses to requests for written consent in an academic family practice unit. *Canadian Family Physician* 1995;**41**:1367-72.
- 156 - 1** Rowan-Szal G, Joe GW, Chatham LR, Simpson DD. A simple reinforcement system for methadone clients in a community-based treatment program. *Journal of Substance Abuse Treatment* 1994;**11**(3):217-23.
- 157 - 1** Greenberg RA, Strecher VJ, Bauman KE, Boat BW, Fowler MG, Keyes LL, Denny FW, Chapman RS, Stedman HC, LaVange LM, Glover LH, Haley NJ, Loda FA. Evaluation of a home-based intervention program to reduce infant passive smoking and lower respiratory illness. *Journal of Behavioral Medicine* 1994;**17**(3):273-90.
- 158 - 1** Strecher VJ, Kreuter M, Den Boer DJ, Kobrin S, Hospers HJ, Skinner CS. The effects of computer-tailored smoking cessation messages in family practice settings. *Journal of Family Practice* 1994;**39**(3):262-70.
- 158 - 2** Strecher VJ, Kreuter M, Den Boer DJ, Kobrin S, Hospers HJ, Skinner CS. The effects of computer-tailored smoking cessation messages in family practice settings. *Journal of Family Practice* 1994;**39**(3):262-70.
- 159 - 1** Miller RL. Assisting gay men to maintain safer sex: an evaluation of an AIDS service organization's safer sex maintenance program. *AIDS Education and Prevention* 1995;**7**(5 Suppl): 48-63.
- 161 - 1** Muller O, Sarangbin S, Ruxrungtham K, Sittitrai W, Phanuphak P. Sexual risk behaviour reduction associated with voluntary HIV counselling and testing in HIV infected patients in Thailand. *AIDS Care* 1995;**7**(5):567-72.
- 162 - 1** Stock LZ, Milan MA. Improving dietary practices of elderly individuals: the power of prompting feedback, and social reinforcement. *Journal of Applied Behavior Analysis* 1993;**26**(3):379-87.
- 163 - 1** Finney JW, Miller KM, Adler SP. Changing protective and risky behaviours to prevent child-to-parent transmission of cytomegalovirus. *Journal of Applied Behaviour Analysis* 1993;**26**:471-2.
- 165 - 1** Deren S, Davis WR, Beardsley M, Tortu S, Clatts M. Outcomes of a risk-reduction intervention with high-risk populations: the Harlem AIDS project. *AIDS Education and Prevention* 1995;**7**(5):379-90.
- 167 - 1** Campbell JR, Szilagyi PG, Rodewald LE, Doane C, Roghmann KJ. Patient-specific reminder letters and pediatric well-child-care show rates. *Clinical Pediatrics* 1994;May:268-72.
- 168 - 1** Schectman G, Hiatt J, Hartz A. Telephone contacts do not improve adherence to niacin or bile acid sequestrant therapy. *Annals of Pharmacotherapy* 1994;**28**:29-35.
- 169 - 1** Lipton HL, Bird JA. The impact of clinical pharmacists' consultations on geriatric patients' compliance and medical care use: a randomized controlled trial. *Gerontologist* 1994;**34**(3):307-15.
- 170 - 1** Vineis P, Ronco G, Ciccone G, Gogliani F. Home injuries in children: a population-based intervention trial. *Epidemiology* 1994;**5**(3):349-51.

- 171 - 1** Biger C, Epstein LM, Hagoel L, Tamir A, Robinson E. An Evaluation of an education programme, for prevention and early diagnosis of malignancy in Israel. *European Journal of Cancer Prevention* 1994;**3**:305–12.
- 172 - 1** Moser SE. Effectiveness of post card appointment reminders. *Family Practice Research Journal* 1994;**14**:281–8.
- 174 - 1** Avila P, Hovell MF. Physical activity training for weight loss in Latinas: a controlled trial. *International Journal of Obesity* 1994;**18**:476–82.
- 176 - 1** Malenka DJ, Baron JA, Johanson S, Wahrenberger JW, Ross JM. The framing effect of relative and absolute risk. *Journal General Internal Medicine* 1993;**8**:543–8.
- 177 - 1** Birkel RC, Golaszewski T, Koman JJ, Singh BK, Catan V, Souply K. Findings from the horizontes acquired immune deficiency syndrome education project: the impact of indigenous outreach workers as change agents for injection drug users. *Health Education Quarterly* 1993;**20**:523–38.
- 179 - 1** Meltzer SB, Hovell MF, Meltzer EO, Atkins CJ, de Peyster A. Reduction of secondary smoke exposure in asthmatic children: parent counseling. *Journal of Asthma* 1993;**30**(5):391–400.
- 180 - 1** Hill D, White V, Marks R, Borland R. Changes in sun-related attitudes and behaviours, and reduced sunburn prevalence in a population at high risk of melanoma. *European Journal of Cancer Prevention* 1993;**2**(6):447–56.
- 181 - 1** Hanna KM. Effect of nurse-client transaction on female adolescent's oral contraceptive adherence. *IMAGE: Journal of Nursing Scholarship* 1993;**25**:285–90.
- 183 - 1** Shore ER. Outcomes of a primary prevention project for business and professional women. *Journal of Studies on Alcohol* 1994;**55**(6):657–9.
- 184 - 1** Kaufman JS, Jason LA, Sawlski LM, Halpert JA. A comprehensive multi-media program to prevent smoking among black students. *Journal of Drug Education* 1994;**24**(2):95–108.
- 190 - 1** Moos MK, Bangdiwala SI, Meibohm AR, Cefalo RC. The impact of a preconceptional health promotion program on intendedness of pregnancy. *American Journal of Perinatology* 1996;**13**(2):103–8.
- 192 - 1** Gilles MT, Crewe S, Granites IN, Coppola A. A community-based, cervical screening program in a remote Aboriginal community in the Northern Territory. *Australian journal of Public Health* 1995;**19**:477–81.
- 193 - 1** Bazeley P, Kemp L. Increasing attendance at immunisation clinics: lessons from a trial program that failed. *Australian Journal of Public Health* 1995;**19**(5):459–64.
- 194 - 1** Martinez R, Levine DW, Martin R, Altman DG. Effect of integration of injury control information into a high school physics course. *Annals of Emergency Medicine* 1996;**27**(2):216–24.
- 196 - 1** Bullock LF, Wells JE, Duff GB, Hornblow AR. Telephone support for pregnant women: outcome in late pregnancy. *New Zealand Medical Journal* 1995;**108**(1012):476–8.
- 197 - 1** Plaskon PP, Fadden MJ. Cancer screening utilization: is there a role for social work in cancer prevention? *Social Work in Health Care* 1995;**21**(4):59–70.
- 198 - 1** Robinson JK, Rademaker AW. Skin cancer risk and sun protection learning by helpers of patients with nonmelanoma skin cancer. *Preventive Medicine* 1995;**24**(4):333–41.
- 200 - 1** Simkin-Silverkin L, Wing RR, Hansen DH, Klem ML, Pasagian-Macauley A, Meilahn EN, Kuller LH. Prevention of Cardiovascular risk factor elevations in healthy premenopausal women. *Preventive Medicine* 1995;**24**:509–17.
- 202 - 1** Gariti P, Alterman AI, Holub-Beyer E, Volpicelli JR, Prentice N, O'Brien CP. Effects of an appointment reminder call on patient show rates. *Journal of Substance Abuse Treatment* 1995;**12**(3):207–12.
- 203 - 1** Urban N, Taplin SH, Taylor VM, Peacock S, Anderson G, Conrad D, Etzioni R, White E, Montano DE, Mahloch J, Majer K. Community organization to promote breast cancer screening among women ages 50–75. *Preventive Medicine* 1995;**24**(5):477–84.
- 205 - 1** Clover K, Redman S, Forbes J, Sanson-Fisher R, Callaghan T. Two sequential randomized trials of community participation to recruit women for mammographic screening. *Preventive Medicine* 1996;**25**:126–34.
- 205 - 2** Clover K, Redman S, Forbes J, Sanson-Fisher R, Callaghan T. Two sequential randomized trials of community participation to recruit women for mammographic screening. *Preventive Medicine* 1996;**25**:126–34.
- 206 - 1** Hoffmeister H, Mensink GBM, Stolzenberg H, Hoeltz J, Kreuter H, Laaser U, Nussel E, Hulleman K-D, Troschke J. Reduction of coronary heart disease risk factors in the German Cardiovascular Prevention study. *Preventive Medicine* 1996;**25**:135–45.
- 207 - 1** Schinke SP, Singer B, Cole K, Contento I. Reducing cancer risk among native American adolescents. *Preventive Medicine* 1996;**25**:146–55.
- 209 - 1** Calfas K, Long B, Sallis J, Wooten W, Pratt M, Patrick K. A controlled trial of physician counseling to promote the adoption of physical activity. *Preventive Medicine* 1996;**25**:225–33.

- 210 - 1** Brug J, Steenhuis I, Assema P, de Vreis H. The impact of a computer-tailored nutrition intervention. *Preventive Medicine* 1996;**25**:236–42.
- 211 - 1** Voorhees C, Stillman F, Swank R, Heagerty P, Levine D, Becker D. Heart, body and soul: Impact of church based smoking cessation interventions on readiness to quit. *Preventive Medicine* 1996;**25**:277–85.
- 212 - 1** Clayton R, Cattarello A, Johnstone B. The effectiveness of Drug Abuse Resistance Education (Project DARE): 5-year follow-up results. *Preventive Medicine* 1996;**25**:307–18.
- 213 - 1** Morgan G, Noll E, Orleans T, Rimer B, Amfoh K, Bonney G. Reaching midlife and older smokers: tailored interventions for routine medical care. *Preventive Medicine* 1996;**25**:346–54.
- 215 - 1** Cheadle A, Psaty B, Diehr P, Koepsell T, Wagner E, Curry S, Kristal A. Evaluating community-based nutrition programs: comparing grocery store and individual-level survey measures of program impact. *Preventive Medicine* 1996;**24**:71–9.
- 216 - 1** Schmid T, Jeffery R, Hellerstedt W. Direct mail recruitment to home based smoking and weight control programs: a comparison of strategies. *Preventive Medicine* 1989;**18**:503–17.
- 216 - 2** Schmid T, Jeffery R, Hellerstedt W. Direct mail recruitment to home based smoking and weight control programs: a comparison of strategies. *Preventive Medicine* 1989;**18**:503–17.
- 217 - 1** Flay B, Miller T, Hedeker D, Siddiqui O, Britton C, Brannon B, Johnson C, Hansen W, Sussman S, Dent C. The television, school and family smoking prevention and cessation project. *Preventive Medicine* 1995;**24**:29–40.
- 218 - 1** Mudde A, Vries H, Dolders M. Evaluation of a Dutch community-based smoking cessation intervention. *Preventive Medicine* 1995;**24**:61–70.
- 220 - 1** Edmundson E, Parcel G, Feldman H, Elder J, Perry C, Johnson C, Williston B, Stone E, Yang M, Lytle L, Webber L. The effects of the Child and Adolescent Trial for Cardiovascular Health upon psychosocial determinants of diet and physical activity behavior. *Preventive Medicine* 1996;**25**:442–54.
- 222 - 1** O'Loughlin J, Renaud L, Paradis G, Meshefedjian G. Screening school personnel for cardiovascular disease risk factors: short term impact on behavior and perceived role as promoters of heart health. *Preventive Medicine* 1996;**25**:660–7.
- 223 - 1** Palinkas L, Atkins C, Millers C, Ferreira D. Social skills training for drug prevention in high-risk female adolescents. *Preventive Medicine* 1996;**25**:692–701.
- 226 - 1** Hansen W, Graham J. Preventing alcohol, marijuana, and cigarette use among adolescents: peer pressure resistance training versus establishing conservative norms. *Preventive Medicine* 1991;**20**:414–30.
- 229 - 1** Hughes J, Wadland W, Fenwick J, Lewis J, Bickel W. Effect of cost on the self-administration and efficacy of nicotine gum: a preliminary study. *Preventive Medicine* 1991;**20**:486–96.
- 232 - 1** Bastani R, Marcus AC, Maxwell AE, Das IP, Yan KX. Evaluation of an intervention to increase mammography screening in Los Angeles. *Preventive Medicine* 1994;**23**(1):83–90.
- 233 - 1** Mayer J, Jermanovich A, Wright B, Elder J, Drew J, Williams S. Changes in health behaviors of older adults: the San Diego Medicare preventive health project. *Preventive Medicine* 1994;**23**:127–33.
- 234 - 1** Ives DG, Lave JR, Traven ND, Kuller LH. Impact of Medicare reimbursement on influenza vaccination rates in the elderly. *Preventive Medicine* 1994;**23**(2):134–41.
- 235 - 1** Brenner H, Fleischle B. Smoking regulations at the workplace and smoking behavior: a study from Southern Germany. *Preventive Medicine* 1994;**23**:230–4.
- 237 - 1** Van Assema P, Steenbakkens M, Kok G, Erikson M, de Vreis H. Results of the Dutch community project 'Healthy Bergeyk'. *Preventive Medicine* 1994;**23**:394–401.
- 238 - 1** Murray DM, Prokhorov AV, Harty KC. Effects of a statewide antismoking campaign on mass media messages and smoking beliefs. *Preventive Medicine* 1994;**23**(1):54–60.
- 239 - 1** Ponsonby A, Dwyer T, Kasl S, Cochrane J, Newman N. An assessment of the impact of public health activities to reduce the prevalence of the prone sleeping position during infancy: the Tasmanian cohort study. *Preventive Medicine* 1994;**23**:402–8.
- 240 - 1** Main DS, Iverson DC, McGloin J, Banspach SW, Collins JL, Rugg DL, Kolbe LJ. Preventing HIV infection among adolescents: evaluation of a school-based education program. *Preventive Medicine* 1994;**23**(4):409–17.
- 241 - 1** McAuley E, Courneya K, Rudolph D, Lox C. Enhancing exercise adherence in middle-aged males and females. *Preventive Medicine* 1994;**23**:498–506.
- 242 - 1** Pallonen UE, Leskinen L, Prochaska JO, Willey CJ, Kaariainen R, Salonen JT. A 2-year self-help smoking cessation manual intervention among middle-aged Finnish men: an application of the transtheoretical model. *Preventive Medicine* 1994;**23**(4):507–14.

- 243 - 1 Zapka J, Costanza M, Harris D, Hosmer D, Stoddard A, Barth R, Gaw V. Impact of a breast cancer screening community intervention. *Preventive Medicine* 1993;**22**:34–53.
- 244 - 1 Lewis B, Lynch W. The effect of physician advice on exercise behavior. *Preventive Medicine* 1993;**22**:110–21.
- 245 - 1 Orleans C, Rotberg H, Quade D, Leeds P. A hospital quit-smoking consult service: clinical report and intervention guidelines. *Preventive Medicine* 1990;**19**:198–212.
- 246 - 1 Worden J, Solomon L, Flynn B, Costanza M, Foster B, Dorwaldt A, Weaver S. A community-wide program in breast self-examination training and maintenance. *Preventive Medicine* 1990;**19**:254–69.
- 247 - 1 Graham J, Johnson C, Hansen W, Flay B, Gee M. Drug use prevention programs, gender and ethnicity: evaluation of three seventh-grade project SMART cohorts. *Preventive Medicine* 1990;**19**:305–13.
- 248 - 1 Paskett E, White E, Carter W, Chu E. Improving follow-up after an abnormal Pap smear: a randomised controlled trial. *Preventive Medicine* 1990;**19**:630–41.
- 249 - 1 Secker-Walker R, Flynn B, Solomon L, Vacek P, Bronson D. Predictors of smoking behavior change 6 and 18 months after individual counseling during periodic health examinations. *Preventive Medicine* 1990;**19**:675–85.
- 251 - 1 Pentz M, MacKinnon D, Dwyer J, Wang E, Hansen W, Flay B, Johnson C. Longitudinal effects of the Midwestern Prevention Project on regular and experimental smoking in adolescents. *Preventive Medicine* 1989;**18**:304–21.
- 253 - 1 Crockett S, Mullis R, Perry C, Luepker R. Parent education in youth-directed nutrition interventions. *Preventive Medicine* 1989;**18**:475–91.
- 254 - 1 Hollis J, Lichtenstein E, Mount K, Vogt T, Stevens V. Nurse-assisted smoking counseling in medical settings: minimising demands on physicians. *Preventive Medicine* 1991;**20**:497–507.
- 255 - 1 Murray D, Perry C, Griffin G, Harty K, Jacobs D, Schmid L, Daly K, Pallonen U. Results from a statewide approach to adolescent tobacco use prevention. *Preventive Medicine* 1992;**21**:449–72.
- 255 - 2 Murray D, Perry C, Griffin G, Harty K, Jacobs D, Schmid L, Daly K, Pallonen U. Results from a statewide approach to adolescent tobacco use prevention. *Preventive Medicine* 1992;**21**:449–72.
- 256 - 1 Hebert J, Kristeller J, Ockene J, Landon J, Luippold R, Goldberg R, Kalan K. Patient characteristics and the effect of three physician-delivered smoking interventions. *Preventive Medicine* 1992;**21**:557–73.
- 257 - 1 Ives D, Kuller L, Schultz R, Traven N, Lave J. Comparison of recruitment strategies and associated disease prevalence for health promotion in rural elderly. *Preventive Medicine* 1992;**21**:582–91.
- 258 - 1 Winkleby M, Fortmann S, Rockhill B. Trends in cardiovascular disease risk factors by educational level: the Stanford five-city project. *Preventive Medicine* 1992;**21**:592–601.
- 259 - 1 Nilssen O. The Tromso Study: identification of and a controlled intervention on a population of early stage risk drinkers. *Preventive Medicine* 1991;**20**:518–28.
- 260 - 1 Ockene J, Hymowitz N, Lagus J, Shaten J. Comparison of smoking behavior change for SI and UC Study Groups. *Preventive Medicine* 1991;**20**:518–28.
- 261 - 1 Lionis C, Kafatos A, Vlachonikolis J, Vakaki M, Tzortzi M, Petraki A. The effects of a health education intervention program among Cretan adolescents. *Preventive Medicine* 1991;**20**(6):685–99.
- 262 - 1 Olsen G, Lacy S, Sprafka J, Arceneaux T, Potts T, Kravat B, Gondek M, Bond G. A 5-year evaluation of a smoking cessation incentive program for chemical employees. *Preventive Medicine* 1990;**19**:774–84.
- 263 - 1 Henderson MM, Kushi LH, Thompson DJ, Gorbach SL, Clifford CK, Insull W, Moskowitz M, Thompson RS. Feasibility of a randomised trial of a low-fat diet for the prevention of breast cancer: dietary compliance in the women's health trial vanguard study. *Preventive Medicine* 1990;**19**:115–33.
- 264 - 1 Murray DM, Kurth C, Mullis R, Jeffery RW. Cholesterol reduction through low intensity interventions: results from the Minnesota Heart Health Program. *Preventive Medicine* 1990;**19**:181–9.
- 265 - 1 Walker R, Heller R, Redman S, O'Connell D, Boulton J. Reduction of ischemic heart disease risk markers in the teenage children of heart attack patients. *Preventive Medicine* 1992;**21**:616–29.
- 266 - 1 Robinson J. Compensation strategies in sun protection behaviors by a population with nonmelanoma skin cancer. *Preventive Medicine* 1992;**21**:754–65.
- 268 - 1 Knutsen SE, Knutsen R. The Tromso Survey: the Family Intervention study—the effect of intervention on some coronary risk factors and dietary habits, a 6-year follow-up. *Preventive Medicine* 1991;**20**(2):197–212.

- 269 - 1** Ho EE, Atwood JR, Benedict J, Ritenbaugh C, Sheehan ET, Abrams C, Alberts D, Meyskens FL, Jr. A community-based feasibility study using wheat bran fiber supplementation to lower colon cancer risk. *Preventive Medicine* 1991;**20**(2):213–25.
- 270 - 1** Edmunds M, Conner H, Jones C, Gorayeb R, Waranch H. Evaluation of a multicomponent group smoking cessation program. *Preventive Medicine* 1991;**20**:774–84.
- 273 - 1** Winkleby M, Fortmann S, Rockhill B. Cigarette smoking trends in adolescents and young adults: the Stanford five-city project. *Preventive Medicine* 1993;**22**:325–34.
- 275 - 1** Curry SJ, Taplin SH, Anderman C, Barlow WE, McBride C. A randomized trial of the impact of risk assessment and feedback on participation in mammography screening. *Preventive Medicine* 1993;**22**(3):350–60.
- 276 - 1** Klepp KI, Tell GS, Vellar OD. Ten-year follow-up of the Oslo Youth Study Smoking Prevention Program. *Preventive Medicine* 1993;**22**(4):453–62.
- 277 - 1** Bell RM, Ellickson PL, Harrison ER. Do drug prevention effects persist into high school? How project ALERT did with ninth graders. *Preventive Medicine* 1993;**22**(4):463–83.
- 279 - 1** Alexandrov A, Maslennikova G, Kulikov S, Propirnij G, Perova N. Primary prevention of cardiovascular disease: 3-year intervention results in boys of 12 years of age. *Preventive Medicine* 1992;**21**:53–62.
- 280 - 1** Korhonen H, Niemensivu H, Piha T, Koskela K, Wiio J, Johnson C, Puska P. National TV smoking cessation program and contest in Finland. *Preventive Medicine* 1992;**21**:74–87.
- 283 - 1** Maheu M, Gevirtz R, Sallis J, Schneider N. Competition/co-operation in worksite smoking cessation using nicotine gum. *Preventive Medicine* 1989;**18**:867–76.
- 284 - 1** Shipley R, Orleans C, Wilbur C, Piserchia P, McFadden D. Effect of the Johnson and Johnson LIVE FOR LIFE program on employee smoking. *Preventive Medicine* 1988;**17**:25–34.
- 285 - 1** Faivre J, Arveux P, Milan C, Durand G, Lamour J, Bedenne L. Participation in mass screening for colorectal cancer: results of screening and rescreening from the Burgundy study. *European Journal of Cancer Prevention* 1991;**1**(1):49–55.
- 287 - 1** Stevens VJ, Glasgow RE, Hollis JF, Lichtenstein E, Vogt TM. A smoking-cessation intervention for hospital patients. *Medical Care* 1993;**31**(1):65–72.
- 288 - 1** Weinstein ND, Sandman PM, Roberts NE. Perceived susceptibility and self-protective behavior: a field experiment to encourage home radon testing. *Health Psychology* 1991;**10**(1):25–33.
- 289 - 1** Rickert VI, Gottlieb AA, Jay MS. Is AIDS education related to condom acquisition? *Clinical Pediatrics* 1992;**31**(4):205–10.
- 291 - 1** Clark N, Feldman C, Evans D, Duzey O, Levison M, Wasilewski Y, Kaplan D, Rips J, Mellins R. Managing better: children, parents and asthma. *Patient Education and Counseling* 1986;**8**:27–38.
- 293 - 1** Tietge N, Bender S, Scutchfield FD. Influence of teaching techniques on infant car seat use. *Patient Education and Counseling* 1987;**9**:167–75.
- 294 - 1** Keenan J, Kastner T, Nathanson R, Richardson N, Hinton J, Cress DA. A statewide public and professional education program on fragile X syndrome. *Mental Retardation* 1992;**30**(6):355–61.
- 295 - 1** Mann K, Sullivan P. Effect of task-centred instructional programs on hypertensives' ability to achieve and maintain reduced dietary sodium intake. *Patient Education and Counseling* 1987;**19**:53–72.
- 296 - 1** Bejes C, Marvel MK. Attempting the improbable: offering colorectal cancer screening to all appropriate patients. *Family Practice Research Journal* 1992;**12**(1):83–90.
- 299 - 1** Rosser WW, Hutchison BG, McDowell I, Newell C. Use of reminders to increase compliance with tetanus booster vaccination. *Canadian Medical Association Journal* 1992;**146**(6):911–17.
- 301 - 1** Quaid K, Faden R, Vining E, Freeman J. Informed consent for a prescription drug: impact of disclosed information on patient understanding and medical outcomes. *Patient Education and Counseling* 1990;**15**:249–59.
- 302 - 1** Hansen BWL. A randomized controlled trial on the effect of an information booklet for young families in Denmark. *Patient Education and Counseling* 1990;**16**:147–50.
- 303 - 1** Hawe P, Higgins G. Can medication education improve the drug compliance of the elderly? Evaluation of an In Hospital program. *Patient Education and Counseling* 1990;**16**:151–60.
- 304 - 1** Salleras Sanmarti L, Alcaide Megias J, Altet Gomez MN, Canela Soler J, Navas Alcala E, Sune Puigbo MR, Serra Majem L. Evaluation of the efficacy of health education on the compliance with antituberculosis chemoprophylaxis in school children. A randomized clinical trial [published erratum appears in *Tubercle Lung Disease* 1993;**74**(3):217]. *Tubercle and Lung Disease* 1993;**74**(1):28–31.
- 305 - 1** Baumann LJ, Zimmerman R, Leventhal H. An experiment in common sense: education at blood pressure screening. *Patient Education and Counseling* 1989;**14**:53–67.

- 307 - 1** Lierman LM, Young HM, Powell-Cope G, Georgiadou F, Benoliel JQ. Effects of education and support on breast self-examination in older women. *Nursing Research* 1994;**43**(3):158–63.
- 308 - 1** Ogutu RO, Oloo AJ, Ekissa WS, Genga IO, Mulaya N, Githure JI. The effect of participatory school health programme on the control of malaria. *East African Medical Journal* 1992;**69**(6):298–302.
- 310 - 1** Sutton S, Hallett R. Smoking intervention in the workplace using videotapes and nicotine chewing gum. *Preventive Medicine* 1988;**17**:48–59.
- 311 - 1** Hansen WB, Anderson Johnson C, Flay B, Graham J, Sobel J. Affective and social influences to the prevention of multiple substance abuse among seventh grade students: results from project SMART. *Preventive Medicine* 1988;**17**:135–54.
- 312 - 1** Pietinen P, Nissinen A, Vartiainen E, Tuomilehto A, Uusitalo U, Ketola A, Moisio S, Puska P. Dietary changes in the North Karelia Project (1972–1982). *Preventive Medicine* 1988;**17**:183–93.
- 313 - 1** de Weerd I, Visser A, Kok G, van der Veen E. Randomized controlled evaluation of an education program for insulin treated patients with diabetes: effects on psychosocial variables. *Patient Education and Counseling* 1989;**14**:191–215.
- 315 - 1** Ware GJ, Holford NH, Davison JG, Harris RG. Unit dose calendar packaging and elderly patient compliance. *New Zealand Medical Journal* 1991;**104**(924):495–7.
- 316 - 1** Lee CY. A randomized controlled trial to motivate worksite fecal occult blood testing. *Yonsei Medical Journal* 1991 Jun;**32**(2):131–8.
- 320 - 1** Roca-Cusachs A, Sort D, Altimira J, Bonet R, Guilera E, Monmany J, Nolla J. The impact of a patient education programme in the control of hypertension. *Journal of Human Hypertension* 1991;**5**(5):437–41.
- 321 - 1** Welsh MC, Labbe EE, Delayney D. Cognitive strategies and personality variables in adherence to exercise. *Psychological Reports* 1991;**68**(3 Pt 2):1327–35.
- 323 - 1** Oladepo O, Okunade A, Brieger W, Oshiname F, Ajuwon A. Outcome of 2 patient education methods on recruitment and compliance with Ivermectin in the treatment of onchocerciasis. *Patient Education and Counseling* 1996;**29**(3):237–45.
- 324 - 1** Cardenas M, Simons-Morton B. The effect of anticipatory guidance on mothers' self-efficacy and behavioral intentions to prevent burns caused by hot tap water. *Patient Education and Counseling* 1993;**21**:117–23.
- 325 - 1** Schultz S. Educational and behavioral strategies related to knowledge of and participation in an exercise program after cardiac positron emission tomography. *Patient Education and Counseling* 1993;**22**:47–57.
- 327 - 1** Huss K, Salerno M, Huss RW. Computer-assisted reinforcement of instruction: effects on adherence in adult atopic asthmatics. *Research in Nursing and Health* 1991;**14**(4):259–67.
- 328 - 1** Glasgow R, Toobert D, Hampson S, Brown J, Lewinsohn P, Donnelly J. Improving self care among older patients with Type II Diabetes: The 'Sixty Something...' study. *Patient Education and Counseling* 1992;**19**:61–74.
- 329 - 1** Price JH, Krol RA, Desmond SM, Losh DP, Roberts SM, Snyder FF. Comparison of three antismoking interventions among pregnant women in an urban setting: a randomized trial. *Psychological Reports* 1991;**68**(2):595–604.
- 331 - 1** Joffe MD, Luberti A. Effect of emergency department immunization on compliance with primary care. *Pediatric Emergency Care* 1994;**10**(6):317–19.
- 332 - 1** Malloy TR, Wigton RS, Meeske J, Tape TG. The influence of treatment descriptions on advance medical directive decisions. *Journal of the American Geriatrics Society* 1992;**40**(12):1255–60.
- 333 - 1** Fabacher D, Josephson K, Pietruszka F, Linderborn K, Morley JE, Rubenstein LZ. An in-home preventive assessment program for independent older adults: a randomized controlled trial. *Journal of the American Geriatrics Society* 1994;**42**(6):630–8.
- 335 - 1** Chirwa B, Briega W, Ramakrishna J. Evaluating health education for oral rehydration therapy at a rural Nigerian clinic: Part II. *Patient Education and Counseling* 1988;**11**:203–13.
- 336 - 1** Marcus A, Wheeler R, Cullen J, Crane L. Quasi experimental evaluation of the Los Angeles Know Your Body program: knowledge, beliefs, and self reported behaviors. *Preventive Medicine* 1987;**16**:803–15.
- 337 - 1** Rutten G, Beek M, van Eijk J. Effects of systematic patient education about cough on the consulting behaviour of a general practice population. *Patient Education and Counseling* 1993;**22**:127–32.
- 340 - 1** Marin B, Perez-Stable E, Marin G, Hauck W. Effects of a community intervention to change smoking behavior among Hispanics. *Am J Preventive Medicine* 1994;**10**(6):340–7.
- 341 - 1** Saxon AJ, Calsyn DA. Alcohol use and high-risk behavior by intravenous drug users in an AIDS education paradigm. *Journal of Studies on Alcohol* 1992;**53**(6):611–18.



- 342 - 1** Thomas DR, Winsted B, Koontz C. Improving neglected influenza vaccination among healthcare workers in long-term care. *Journal of the American Geriatrics Society* 1993;**41** (9):928–30.
- 343 - 1** Sachs GA, Stocking CB, Miles SH. Empowerment of the older patient? A randomized, controlled trial to increase discussion and use of advance directives. *Journal of the American Geriatrics Society* 1992;**40** (3):269–73.
- 345 - 1** Goldberg D, Hoffman A, Farinha M, Marder D, Tinson-Mitchem L, Burton D, Smith E. Physician delivery of smoking-cessation advice based on the stages-of-change model. *American Journal of Preventive Medicine* 1994;**10**:267–74.
- 348 - 1** Cook B, Noteloviz M, Rector C, Krischer J. An osteoporosis patient education and screening program: results and implications. *Patient Education and Counseling* 1991;**17**:135–45.
- 349 - 1** Huws R. Non-attendances at a marital and sexual difficulties clinic: a controlled intervention study. *International Journal of Social Psychiatry* 1992;**38** (4):304–8.
- 350 - 1** Gilbert J, Wilson D, Singer J, Lindsey E, Willms D, Best J, Taylor D. A family physician smoking cessation program: and evaluation of the role of follow up visits. *American Journal of Preventive Medicine* 1992;**8**:91–5.
- 352 - 1** Mayer J, Clapp E, Bartholomew S, Elder J. Facility-based inreached strategies to promote annual mammograms. *American Journal of Preventive Medicine* 1994;**10** (6):353–6.
- 352 - 2** Mayer J, Clapp E, Bartholomew S, Elder J. Facility-based inreached strategies to promote annual mammograms. *American Journal of Preventive Medicine* 1994;**10** (6):353–6.
- 352 - 3** Mayer J, Clapp E, Bartholomew S, Elder J. Facility-based inreached strategies to promote annual mammograms. *American Journal of Preventive Medicine* 1994;**10** (6):353–6.
- 353 - 1** Calle E, Miracle-Mcmahill H, Moss R, Heath C. Personal contact from friends to increase mammography usage. *American Journal of Preventive Medicine* 1994;**10** (6):361–6.
- 354 - 1** Rimer BK, Resch N, King E, Ross E, Lerman C, Boyce A, Kessler H, Engstrom PF. Multistrategy health education program to increase mammography use among women ages 65 and older. *Public Health Reports* 1992;**107** (4):369–80.
- 355 - 1** King A, Carl F, Birkel L, Haskell W. Increasing exercise among blue collar employees: the tailoring of worksite programs to meet specific needs. *Preventive Medicine* 1988;**17**:357–65.
- 356 - 1** Evers S, Bass M, Donner A, McWhinney I. Lack of impact of salt restriction advice on hypertensive patients. *Preventive Medicine* 1987;**16**:213–20.
- 357 - 1** Schwarz-Lookinland S, McKeever L, Saputo M. Compliance with antibiotic regimens in Hispanic mothers. *Patient Education and Counseling* 1989;**13**:171–82.
- 360 - 1** Werch CE, Young M, Clark M, Garrett C, Hooks S, Kersten C. Effects of a take-home drug prevention program on drug-related communication and beliefs of parents and children. *Journal of School Health* 1991;**61** (8):346–50.
- 361 - 1** Elder AD, Zultowsky D. Using written communication with patient groups to promote behavioral change in chronic mentally ill patients. *Hospital and Community Psychiatry* 1991;**42** (3):302–4.
- 363 - 1** Oeffinger KC, Roaten SP, Hitchcock MA, Oeffinger PK. The effect of patient education on pediatric immunization rates. *Journal of Family Practice* 1992;**35** (3):288–93.
- 364 - 1** Jemmott LS, Jemmott JB, 3rd. Increasing condom-use intentions among sexually active black adolescent women. *Nursing Research* 1992;**41** (5):273–9.
- 365 - 1** White E, Hurlich M, Thompson RS, Woods MN, Henderson MM, Urban N, Kristal A. Dietary changes among husbands of participants in a low-fat dietary intervention. *American Journal of Preventive Medicine* 1991;**7** (5):319–25.
- 367 - 1** Elder JP, McGraw SA, Rodrigues A, Lasater TM, Ferreira A, Kendall L, Peteron G, Carleton R. Evaluation of two community-wide smoking cessation contests. *Preventive Medicine* 1987;**16**:221–34.
- 367 - 2** Elder JP, McGraw SA, Rodrigues A, Lasater TM, Ferreira A, Kendall L, Peteron G, Carleton R. Evaluation of two community-wide smoking cessation contests. *Preventive Medicine* 1987;**16**:221–34.
- 369 - 1** Weinberger M, Tierney WM, Booher P, Katz BP. The impact of increased contact on psychosocial outcomes in patients with osteoarthritis: a randomized, controlled trial. *Journal of Rheumatology* 1991;**18** (6):849–54.
- 370 - 1** Shope JT, Dielman TE, Butchart AT, Campanelli PC, Kloska DD. An elementary school-based alcohol misuse prevention program: a follow-up evaluation. *Journal of Studies on Alcohol* 1992;**53** (2):106–21.
- 371 - 1** Knapp LG. Effects of type of value appealed to and valence of appeal on children's dental health behavior. *Journal of Pediatric Psychology* 1991;**16** (6):675–86.
- 372 - 1** Gans KM, Lapane KL, Lasater TM, Carleton RA. Effects of intervention on compliance to referral and lifestyle recommendations given at cholesterol screening programs. *American Journal of Preventive Medicine* 1994;**10** (5):275–82.

- 373 - 1** Eisen M, Zellman GL, McAlister AL. A Health Belief Model-Social Learning Theory approach to adolescents' fertility control: findings from a controlled field trial. *Health Education Quarterly* 1992;**19**(2):249-62.
- 375 - 1** Leslie M, Schuster P. The effect of contingency contracting on adherence and knowledge of exercise regimens. *Patient Education and Counseling* 1993;**21**:51-60.
- 376 - 1** Lichtenstein E, Hollis J. Patient referral to a smoking cessation program: who follows through? *Journal of Family Practice* 1992;**34**(6):739-44.
- 377 - 1** Ward JE, Boyle K, Redman S, Sanson-Fisher RW. Increasing women's compliance with opportunistic cervical cancer screening: a randomized trial. *American Journal of Preventive Medicine* 1991;**7**(5):285-91.
- 378 - 1** Spaulding SA, Kugler JP. Influenza immunization: the impact of notifying patients of high-risk status. *Journal of Family Practice* 1991;**33**(5):495-8.
- 380 - 1** Lerman C, Hanjani P, Caputo C, Miller S, Delmoor E, Nolte S, Engstrom P. Telephone counseling improves adherence to colposcopy among lower-income minority women. *Journal of Clinical Oncology* 1992;**10**(2):330-3.
- 381 - 1** Fitzgerald ST, Gibbens S, Agnew J. Evaluation of referral completion after a workplace cholesterol screening program. *American Journal of Preventive Medicine* 1991;**7**(6):335-40.
- 383 - 1** Mayer JA, Kossman MK, Miller LC, Crooks CE, Slymen DJ, Lee CD, Jr. Evaluation of a media-based mammography program. *American Journal of Preventive Medicine* 1992;**8**(1):23-9.
- 384 - 1** Baier CA, Grodzin CJ, Port JD, Leksas L, Tancredi DJ. Coronary risk factor behavior change in hospital personnel following a screening program. *American Journal of Preventive Medicine* 1992;**8**(2):115-22.
- 385 - 1** Stevens MM, Freeman DH, Jr, Mott LA, Youells FE, Linsey SC. Smokeless tobacco use among children: the New Hampshire Study. *American Journal of Preventive Medicine* 1993;**9**(3):160-7.
- 386 - 1** Spatz TS. Improving breast self-examination training by using the 4MAT instructional model. *Journal of Cancer Education* 1991;**6**(3):179-83.
- 387 - 1** McCann S, Weinman J. Empowering the patient in the consultation: a pilot study. *Patient Education and Counseling* 1996;**27**(3):227-34.
- 390 - 1** Heirich MA, Foote A, Erfurt JC, Konopka B. Work-site physical fitness programs. Comparing the impact of different program designs on cardiovascular risks. *Journal of Occupational Medicine* 1993;**35**(5):510-7.
- 391 - 1** Piotrow PT, Kincaid DL, Hindin MJ, Lettenmaier CL, Kuseka I, Silberman T, Zinanga A, Chikara F, Adamchak DJ, Mbizvo MT, Lynn W, Kumah OM, Kim Y-M. Changing men's attitudes and behavior: the Zimbabwe Male Motivation Project. *Studies in Family Planning* 1992;**23**(6 Pt 1):365-75.
- 392 - 1** Koenig MA, Rob U, Khan MA, Chakraborty J, Fauveau V. Contraceptive use in Matlab, Bangladesh in 1990: levels, trends, and explanations. *Studies in Family Planning* 1992;**23**(6 Pt 1):352-64.
- 393 - 1** Mansfield CJ, Conroy ME, Emans SJ, Woods ER. A pilot study of AIDS education and counseling of high-risk adolescents in an office setting. *Journal of Adolescent Health* 1993;**14**(2):115-19.
- 394 - 1** Newcomb PA, Klein R, Massoth KM. Education to increase ophthalmologic care in older onset diabetes patients: indications from the Wisconsin Epidemiologic Study of Diabetic Retinopathy. *Journal of Diabetes and its Complications* 1992;**6**(4):211-17.
- 395 - 1** Ornstein SM, Garr DR, Jenkins RG, Rust PF, Arnon A. Computer-generated physician and patient reminders. Tools to improve population adherence to selected preventive services. *Journal of Family Practice* 1991;**32**(1):82-90.
- 396 - 1** Turnin MC, Beddok RH, Clottes JP, Martini PF, Abadie RG, Buisson JC, Soule-Dupuy C, Bonneau M, Camare R, Anton JP, Chrismet CY, Farreny H, Bayard F, Tauber J-PJ. Telematic expert system Diabeto. New tool for diet self-monitoring for diabetic patients. *Diabetes Care* 1992;**15**(2):204-12.
- 397 - 1** Ansell D, Lacey L, Whitman S, Chen E, Phillips C. A nurse-delivered intervention to reduce barriers to breast and cervical cancer screening in Chicago inner city clinics. *Public Health Reports* 1994;**109**(1):104-11.
- 398 - 1** Wing RR, Anglin K. Effectiveness of a behavioral weight control program for blacks and whites with NIDDM. *Diabetes Care* 1996;**19**(5):409-13.
- 401 - 1** Kohatsu N, Cramer E, Bohnstedt M. Use of a clinician reminder system for screening mammography in a public health clinic. *American Journal of Preventive Medicine* 1994;**10**(6):348-52.
- 402 - 1** Ojehagen A, Berglund M. Acceptance, attrition, and outcome in an outpatient treatment programme for alcoholics. A comparison between a randomized and a non-randomized process-outcome study. *European Archives of Psychiatry and Clinical Neuroscience* 1992;**242**(2-3):82-4.
- 403 - 1** Chen CH. Effects of home visits and telephone contacts on breastfeeding compliance in Taiwan. *Maternal-Child Nursing Journal* 1993;**21**(3):82-90.

- 404 - 1 Rimer BK, Ross E, Balslem A, Engstrom PF. The effect of a comprehensive breast screening program on self-reported mammography use by primary care physicians and women in a health maintenance organization. *Journal of the American Board of Family Practice* 1993;**6**(5):443–51.
- 405 - 1 Doi SC, DiLorenzo TM. An evaluation of a tobacco use education-prevention program: a pilot study. *Journal of Substance Abuse* 1993;**5**(1):73–8.
- 407 - 1 Colon HM, Robles RR, Freeman D, Matos T. Effects of a HIV risk reduction education program among injection drug users in Puerto Rico. *Puerto Rico Health Sciences Journal* 1993;**12**(1):27–34.
- 408 - 1 Stehr-Green PA, Dini EF, Lindegren ML, Patriarca PA. Evaluation of telephoned computer-generated reminders to improve immunization coverage at inner-city clinics. *Public Health Reports* 1993;**108**(4):426–30.
- 409 - 1 Devine CM, Olson CM, Frongillo EA, Jr. Impact of the Nutrition for Life program on junior high students in New York State. *Journal of School Health* 1992;**62**(8):381–5.
- 411 - 1 Barnwell MD, Chitkara R, Lamberta F. Tuberculosis prevention project. *Journal of the National Medical Association* 1992;**84**(12):1014–18.
- 412 - 1 Freedman JD, Mitchell CK. A simple strategy to improve patient adherence to outpatient fecal occult blood testing. *Journal of General Internal Medicine* 1994;**9**(8):462–4.
- 413 - 1 Merkel PA, Caputo GC. Evaluation of a simple office-based strategy for increasing influenza vaccine administration and the effect of differing reimbursement plans on the patient acceptance rate. *Journal of General Internal Medicine* 1994;**9**(12):679–83.
- 414 - 1 Baum JG, Clark HB, Sandler J. Preventing relapse in obesity through posttreatment maintenance systems: comparing the relative efficacy of two levels of therapist support. *Journal of Behavioral Medicine* 1991;**14**(3):287–302.
- 416 - 1 Clifford PA, Tan S-Y, Gorsuch RL. Efficacy of a self-directed behavioural health change program: weight, body composition, cardiovascular fitness, blood pressure, health risk and psychosocial mediating variables. *Journal of Behavioural Medicine* 1991;**14**:303–23.
- 418 - 1 Sumner W. An evaluation of readable preventive health messages. *Family Medicine* 1991;**23**(6):463–6.
- 420 - 1 Myers RE, Ross EA, Wolf TA, Balslem A, Jepson C, Millner L. Behavioral interventions to increase adherence in colorectal cancer screening. *Medical Care* 1991;**29**(10):1039–50.
- 421 - 1 Gastrin G. Preliminary results of primary screening for breast cancer with the Mama Program. *Sozial- und Präventivmedizin* 1993;**38**(5):280–7.
- 423 - 1 Joseph AM. Nicotine treatment at the Drug Dependency Program of the Minneapolis VA Medical Center. A researcher's perspective. *Journal of Substance Abuse Treatment* 1993;**10**(2):147–52.
- 424 - 1 Levy SR, Perhats C, Weeks K, Handler AS, Zhu C, Flay BR. Impact of a school-based AIDS prevention program on risk and protective behavior for newly sexually active students. *Journal of School Health* 1995;**65**(4):145–51.
- 425 - 1 Sikorski J, Wilson J, Clement S, Das S, Smeeton N. A randomised controlled trial comparing two schedules of antenatal visits: the antenatal care project. *British Medical Journal* 1996;**312**:546–53.
- 426 - 1 Norman P. Applying the health belief model to the prediction of attendance at health checks in general practice. *British Journal of Clinical Psychology* 1995;**34**:461–70.
- 427 - 1 Kemp R, Hayward P, Applewhaite G, Everitt B, David A. Compliance therapy in psychotic patients: randomised controlled trial. *British Medical Journal* 1996;**312**(7027):345–9.
- 430 - 1 Shiloh S, Reznik H, Bat-Miriam-Katznelson M, Goldman B. Pre-marital genetic counselling to consanguineous couples: attitudes, beliefs and decisions among counselled, noncounselled and unrelated couples in Israel. *Social Science and Medicine* 1995;**41**(9):1301–10.
- 434 - 1 Llewellyn-Thomas HA, McGreal MJ, Thiel EC. Cancer patients' decision making and trial-entry preferences: the effects of "framing" information about short-term toxicity and long-term survival. *Medical Decision Making* 1995;**15**(1):4–12.
- 438 - 1 Banks SM, Salovey P, Greener S, Rothman AJ, Moyer A, Beauvais J, Epel E. The effects of message framing on mammography utilization. *Health Psychology* 1995;**14**(2):178–84.
- 439 - 1 Urban N, Taplin S, Taylor V, Peacock S, Anderson G, Conrad D, Etzioni R, White E, Montano D, Mahloch J, Majer K. Community organisation to promote breast cancer screening among women ages 50–75. *Preventive Medicine* 1995;**24**:477–84.
- 440 - 1 Carleton RA, Lasater TM, Assaf AR, Feldman HA, McKinlay S. The Pawtucket Heart Health Program: community changes in cardiovascular risk factors and projected disease risk. *American Journal of Public Health* 1995;**85**(6):777–85.

- 442 - 1 Glasgow RE, Terborg JR, Hollis JF, Severson HH, Boles SM. Take heart: results from the initial phase of a work-site wellness program. *American Journal of Public Health* 1995;**85**(2):209–16.
- 446 - 1 Trock B, Rimer BK, King E, Balslem A, Cristinzio CS, Engstrom PF. Impact of an HMO-based intervention to increase mammography utilization. *Cancer Epidemiology, Biomarkers and Prevention* 1993;**2**(2):151–6.
- 447 - 1 Saunders LD, Irwig LM, Gear JS, Ramushu DL. A randomized controlled trial of compliance improving strategies in Soweto hypertensives. *Medical Care* 1991;**29**(7):669–78.
- 448 - 1 Weinrich SP, Weinrich MC, Boyd MD, Atwood J, Cervenka B. Teaching older adults by adapting for aging changes. *Cancer Nursing* 1994;**17**(6):494–500.
- 449 - 1 Smith NA, Seale JP, Ley P, Mellis CM, Shaw J. Better medication compliance is associated with improved control of childhood asthma. *Monaldi Archives for Chest Disease* 1994;**49**(6):470–4.
- 450 - 1 Rhodes F, Wolitski RJ, Thornton-Johnson S. An experiential program to reduce AIDS risk among female sex partners of injection-drug users. *Health and Social Work* 1992;**17**(4):261–72.
- 452 - 1 Jin BW, Kim SC, Mori T, Shimao T. The impact of intensified supervisory activities on tuberculosis treatment. *Tubercle and Lung Disease* 1993;**74**(4):267–72.
- 453 - 1 St. Pierre TL, Kaltreider DL, Mark MM, Aikin KJ. Drug prevention in a community setting: a longitudinal study of the relative effectiveness of a three-year primary prevention program in boys and girls clubs across the nation. *American Journal of Community Psychology* 1992;**20**(6):673–706.
- 454 - 1 Nyamathi AM, Leake B, Flaskerud J, Lewis C, Bennett C. Outcomes of specialized and traditional AIDS counseling programs for impoverished women of color. *Research in Nursing and Health* 1993;**16**(1):11–21.
- 455 - 1 Winter L, Breckenmaker LC. Tailoring family planning services to the special needs of adolescents. *Family Planning Perspectives* 1991;**23**(1):24–30.
- 456 - 1 Rose MA. Evaluation of a peer-education program on heart disease prevention with older adults. *Public Health Nursing* 1992;**9**(4):242–7.
- 457 - 1 Horowitz LG. Comparing shower-based oral hygiene with traditional and electric tooth-brushing. *Clinical Preventive Dentistry* 1992;**14**(6):11–16.
- 458 - 1 Mermelstein RJ, Riesenber LA. Changing knowledge and attitudes about skin cancer risk factors in adolescents. *Health Psychology* 1992;**11**(6):371–6.
- 459 - 1 Davis SW, Cummings KM, Rimer BK, Sciandra R, Stone JC. The impact of tailored self-help smoking cessation guides on young mothers. *Health Education Quarterly* 1992;**19**(4):495–504.
- 461 - 1 Hingson R, McGovern T, Howland J, Heeren T, Winter M, Zakocs R. Reducing alcohol-impaired driving in Massachusetts: the Saving Lives Program. *American Journal of Public Health* 1996;**86**(6):791–7.
- 462 - 1 Brownson RC, Smith CA, Pratt M, Mack NE, Jackson-Thompson J, Dean CG, Dabney S, Wilkerson JC. Preventing cardiovascular disease through community-based risk reduction: the Bootheel Heart Health Project. *American Journal of Public Health* 1996;**86**(2):206–13.
- 463 - 1 Kegeles SM, Hays RB, Coates TJ. The Mpowerment Project: a community-level HIV prevention intervention for young gay men. *American Journal of Public Health* 1996;**86**(8 Pt 1):1129–36.
- 464 - 1 Young DR, Haskell WL, Taylor CB, Fortmann SP. Effect of community health education on physical activity knowledge, attitudes, and behavior. The Stanford Five-City Project. *American Journal of Epidemiology* 1996;**144**(3):264–74.
- 465 - 1 Quirk ME, Godkin MA, Schwenzfeier E. Evaluation of two AIDS prevention interventions for inner-city adolescent and young adult women. *American Journal of Preventive Medicine* 1993;**9**(1):21–6.
- 466 - 1 Garr DR, Ornstein SM, Jenkins RG, Zemp LD. The effect of routine use of computer-generated preventive reminders in a clinical practice. *American Journal of Preventive Medicine* 1993;**9**(1):55–61.
- 467 - 1 Brown LK, Barone VJ, Fritz GK, Cebollero P, Nassau JH. AIDS education: the Rhode Island experience. *Health Education Quarterly* 1991;**18**(2):195–206.
- 469 - 1 Sclar DA, Chin A, Skaer TL, Okamoto MP, Nakahiro RK, Gill MA. Effect of health education in promoting prescription refill compliance among patients with hypertension. *Clinical Therapeutics* 1991;**13**(4):489–95.
- 470 - 1 Karofsky PS, Rice RL, Hoornstra LL, Slater CJ, Kessinich CA, Goode JR. The effect of the initial family interview on a pediatric practice. *Clinical Pediatrics* 1991;**30**(5):290–4.
- 471 - 1 Sanchez-Craig M, Spivak K, Davila R. Superior outcome of females over males after brief treatment for the reduction of heavy drinking: replication and report of therapist effects. *British Journal of Addiction* 1991;**86**(7):867–76.

- 472 - 1 Rost KM, Flavin KS, Cole K, McGill JB. Change in metabolic control and functional status after hospitalization. Impact of patient activation intervention in diabetic patients. *Diabetes Care* 1991;**14**(10):881–9.
- 473 - 1 Lando HA, Pirie PL, McGovern PG, Pechacek TF, Swim J, Loken B. A comparison of self-help approaches to smoking cessation. *Addictive Behaviors* 1991;**16**(5):183–93.
- 475 - 1 Tu KS, McDaniel G, Gay JT. Diabetes self-care knowledge, behaviors, and metabolic control of older adults – the effect of a posteducational follow-up program. *Diabetes Educator* 1993;**19**(1):25–30.
- 478 - 1 Kirby D, Barth RP, Leland N, Fetro JV. Reducing the risk: impact of a new curriculum on sexual risk-taking. *Family Planning Perspectives* 1991;**23**(6):253–63.
- 479 - 1 Xiang M, Ran M, Li S. A controlled evaluation of psychoeducational family intervention in a rural Chinese community. *British Journal of Psychiatry* 1994;**165**(4):544–8.
- 480 - 1 Slap GB, Plotkin SL, Khalid N, Michelman DF, Forke CM. A human immunodeficiency virus peer education program for adolescent females. *Journal of Adolescent Health* 1991;**12**(6):434–42.
- 481 - 1 Tanke ED, Leirer VO. Automated telephone reminders in tuberculosis care. *Medical Care* 1994;**32**(4):380–9.
- 482 - 1 Myers RE, Balslem AM, Wolf TA, Ross EA, Millner L. Adherence to continuous screening for colorectal neoplasia. *Medical Care* 1993;**31**(6):508–19.
- 483 - 1 Kimberlin CL, Berardo DH, Pendergast JF, McKenzie LC. Effects of an education program for community pharmacists on detecting drug-related problems in elderly patients. *Medical Care* 1993;**31**(5):451–68.
- 484 - 1 Marcus AC, Crane LA, Kaplan CP, Reading AE, Savage E, Gunning J, Bernstein G, Berek JS. Improving adherence to screening follow-up among women with abnormal Pap smears: results from a large clinic-based trial of three intervention strategies. *Medical Care* 1992;**30**(3):216–30.
- 485 - 1 Walsh DC, Hingson W, Merrigan DM, Levenson SM. Treating the employed alcoholic: which interventions work? Special Focus: Alcohol and the workplace. *Alcohol Health and Research World* 1992;**16**(2):140–8.
- 486 - 1 Donnelly BW, Davis BJ. A review of the Chance to Grow Project: a care project for pregnant and parenting adolescents. *Child and Adolescent Social Work Journal* 1994;**11**(6):493–506.
- 488 - 1 Llewellyn-Thomas HA, Thiel EC, Sem FW, Woermke DE. Presenting clinical trial information: a comparison of methods. *Patient Education and Counseling* 1995;**25**(2):97–107.
- 489 - 1 Dobs AS, Masters RB, Rajaram L, Stillman FA, Wilder LB, Margolis S, Becker DM. A comparison of education methods and their impact on behavioral change in patients with hyperlipidemia. *Patient Education and Counseling* 1994;**24**(2):157–64.
- 490 - 1 Butler GS, Hurley CA, Buchanan KL, Smith-VanHorne J. Prehospital education: effectiveness with total hip replacement surgery patients. *Patient Education and Counseling* 1996;**29**:189–97.
- 491 - 1 Alcoe SY, Gilbey VJ, McDermot RSR, Wallace DG. The effects of teaching breast self-examination: reported confidence and frequency of practise over a six-year period. *Patient Education and Counseling* 1993;**21**:117–23.
- 492 - 1 Kipke MD, Boyer C, Hein K. An evaluation of an AIDS risk reduction education and skills training (ARREST) program. *Journal of Adolescent Health* 1993;**14**(7):533–9.
- 493 - 1 Brennan PF, Moore SM, Smyth KA. The effects of a special computer network on caregivers of persons with Alzheimer's disease. *Nursing Research* 1995;**44**(3):166–72.
- 494 - 1 Mittelman MS, Ferris SH, Steinberg G, Shulman E, Mackell TA, Ambinder A, Cohen J. An intervention that delays institutionalization of Alzheimer's disease patients: Treatment of spouse-caregivers. *Gerontologist* 1993;**33**(6):730–40.
- 495 - 1 Robinson MK, DeHaven MJ, Koch KA. Effects of the Patient Self-Determination Act on patient knowledge and behavior. *Journal of Family Practice* 1993;**37**(4):363–8.
- 502 - 1 Pichert JW, Snyder GM, Kinzer CK, Boswell EJ. Problem-solving anchored instruction about sick days for adolescents with diabetes. *Patient Education and Counseling* 1994;**23**(2):115–24.
- 503 - 1 Wierenga ME. Life-style modification for weight control to improve diabetes health status. *Patient Education and Counseling* 1994;**23**(1):33–40.
- 504 - 1 Mesters I, van Nunen M, Crebolder H, Meertens R. Education of parents about paediatric asthma: effects of a protocol on medical consumption. *Patient Education and Counseling* 1995;**25**:131–6.
- 506 - 1 Tabak ER. Encouraging patient question asking: a clinical trial. *Patient Education and Counseling* 1988;**12**:37–49.
- 507 - 1 Drossaert CHC, Boer H, Seydel ER. Health education to repeat participation in the Dutch breast cancer screening programmes: evaluation of a leaflet tailored to previous participant. *Patient Education and Counseling* 1996;**28**:121–31.

- 508 - 1 Paskett ED, Phillips KC, Miller ME. Improving compliance among women with abnormal Papanicolaou smears. *Obstetrics and Gynecology* 1995;**86**(3):353–9.
- 509 - 1 Kiefe CI, McKay SV, Halevy A, Brody BA. Is cost a barrier to screening mammography for low-income women receiving Medicare benefits? A randomized trial. *Archives of Internal Medicine* 1994;**154**(11):1217–24.
- 511 - 1 Montgomery EB, Jr, Lieberman A, Singh G, Fries JF. Patient education and health promotion can be effective in Parkinson's disease: a randomized controlled trial. PROPATH Advisory Board. *American Journal of Medicine* 1994;**97**(5):429–35.
- 512 - 1 Linkins RW, Dini EF, Watson G, Patriarca PA. A randomized trial of the effectiveness of computer-generated telephone messages in increasing immunization visits among preschool children. *Archives of Pediatrics and Adolescent Medicine* 1994;**148**(9):908–14.
- 514 - 1 Campion P, Owen L, McNeill A, McGuire C. Evaluation of a mass media campaign on smoking and pregnancy. *Addiction* 1994;**89**(10):1245–54.
- 515 - 1 Klepp KI, Ndeki SS, Seha AM, Hannan P, Lyimo BA, Msuya MH, Irema MN, Schreiner A. AIDS education for primary school children in Tanzania: an evaluation study. *AIDS* 1994;**8**(8):1157–62.
- 516 - 1 Tambor ES, Bernhardt BA, Chase GA, Faden RR, Geller G, Hofman KJ, Holtzman NA. Offering cystic fibrosis carrier screening to an HMO population: factors associated with utilization. *American Journal of Human Genetics* 1994;**55**(4):626–37.
- 520 - 1 Hansen BL. A health education booklet for young families – its need, use and effect. *Patient Education and Counseling* 1995;**25**:137–42.
- 521 - 1 Garcia R, Suarez R. Diabetes education in the elderly: a 5-year follow-up of an interactive approach. *Patient Education and Counseling* 1996;**29**:87–97.
- 522 - 1 Town GI, Hodges ID, Wilkie AT, Toop LJ, Graham P, Drennan CJ. A community-wide promotion of asthma self-management in New Zealand. *Patient Education and Counseling* 1995;**26**:219–24.
- 525 - 1 Baker A, Kochan N, Dixon J, Heather N, Wodak A. Controlled evaluation of a brief intervention for HIV prevention among injecting drug users not in treatment. *AIDS Care* 1994;**6**(5):559–70.
- 526 - 1 Rotheram-Borus MJ, Rosario M, Reid H, Koopman C. Predicting patterns of sexual acts among homosexual and bisexual youths. *American Journal of Psychiatry* 1995;**152**(4):588–95.
- 527 - 1 Simmons MS, Nides MA, Rand CS, Wise RA, Tashkin DP. Trends in compliance with bronchodilator inhaler use between follow-up visits in a clinical trial. *Chest* 1996;**109**(4):963–8.
- 528 - 1 Siegal HA, Falck RS, Carlson RG, Wang J. Reducing HIV needle risk behaviors among injection-drug users in the Midwest: an evaluation of the efficacy of standard and enhanced interventions. *AIDS Education and Prevention* 1995;**7**(4):308–19.
- 529 - 1 Nordblad A, Suominen-Taipale L, Murtomaa H, Vartiainen E, Koskela K. Smart Habit Xylitol campaign, a new approach in oral health promotion. *Community Dental Health* 1995;**12**(4):230–4.
- 530 - 1 Allen RM, Jones MP, Oldenburg B. Randomised trial of an asthma self-management programme for adults. *Thorax* 1995;**50**(7):731–8.
- 533 - 1 Alcoe SY, Gilbey VJ, McDermot RSR, Wallace DG. The practise of breast self examination over a six-year period following teaching. *Patient Education and Counseling* 1995;**25**:183–96.
- 537 - 1 Kissinger P, Clark R, Rice J, Kutzen H, Morse A, Brandon W. Evaluation of a program to remove barriers to public health care for women with HIV infection. *Southern Medical Journal* 1995;**88**(11):1121–5.
- 541 - 1 Chlebowski RT, Blackburn GL, Buzzard IM, Rose DP, Martino S, Khandekar JD, York RM, Jeffery RW, Elashoff RM, Wynder EL. Adherence to a dietary fat intake reduction program in postmenopausal women receiving therapy for early breast cancer. The Women's Intervention Nutrition Study. *Journal of Clinical Oncology* 1993;**11**(11):2072–80.
- 542 - 1 Simel DL, Feussner JR. A randomized controlled trial comparing quantitative informed consent formats. *Journal of Clinical Epidemiology* 1991;**44**(8):771–7.
- 543 - 1 Lancaster G, Elton P. Does the offer of cervical screening with breast screening encourage older women to have a cervical smear test? *Journal of Epidemiology and Community Health* 1992;**46**(5):523–7.
- 544 - 1 Muhlhauser I, Richter B, Kraut D, Weske G, Worth H, Berger M. Evaluation of a structured treatment and teaching programme on asthma. *Journal of Internal Medicine* 1991;**230**(2):157–64.
- 545 - 1 Anonymous. HIV prevention through case management for HIV-infected persons—selected sites, United States, 1989–1992. *MMWR – Morbidity and Mortality Weekly Report* 1993;**42**(23):448–9, 455–6.

- 546 - 1 Palm BT, Kant AC, van den Bosch WJ, Vooijs GP, van Weel C. Preliminary results of a general practice based call system for cervical cancer screening in The Netherlands. *British Journal of General Practice* 1993;**43**(377):503-6.
- 550 - 1 Joseph AM, Nichol KL, Anderson H. Effect of treatment for nicotine dependence on alcohol and drug treatment outcomes. *Addictive Behaviours* 1993;**18**(6):635-44.
- 553 - 1 Minor MA, Brown JD. Exercise maintenance of persons with arthritis after participation in a class experience. *Health Education Quarterly* 1993;**20**(1):83-95.
- 554 - 1 Christianson JB, Lurie N, Finch M, Moscovice IS, Hartley D. Use of community-based mental-health programs by HMOS – evidence from a MEDICAID demonstration. *American Journal of Public Health* 1992;**82**(6):790-6.
- 555 - 1 Michaud PA, Hausser D. Swiss teenagers, AIDS and sexually transmitted diseases – presentation and evaluation of a preventive exhibition. *Health Education Research* 1992;**7**(1):79-86.
- 562 - 1 Buischi YA, Axelsson P, Oliveira LB, Mayer MP, Gjermo P. Effect of two preventive programs on oral health knowledge and habits among Brazilian schoolchildren. *Community Dentistry and Oral Epidemiology* 1994;**22**(1):41-6.
- 563 - 1 Clarke JR, Bradnock G, Hamburger R. The uptake and completion of dental treatment using a mobile clinic in central Birmingham, UK. *Community Dental Health* 1992;**9**(2):181-5.
- 564 - 1 Walsh DC, Hingson RW, Merrigan DM, Levenson SM, Cupples LA, Heeren T, Coffman GA, Becker CA, Barker TA, Hamilton SK, McGuire TG, Kelly CA. A randomized trial of treatment options for alcohol-abusing workers. *New England Journal of Medicine* 1991;**325**(11):775-82.
- 566 - 1 Miller MF, Wong JG. Reducing financial barriers enhances the return rate of stool Hemocult packets. *American Journal of the Medical Sciences* 1993;**306**(2):98-100.
- 570 - 1 Tenn L, Dewis ME. An evaluation of a Canadian peer-driven injury prevention programme for high-risk adolescents. *Journal of Advanced Nursing* 1996;**23**(2):329-37.
- 572 - 1 Ferson MJ, Fitzsimmons G, Christie D, Woollett H. School health nurse interventions to increase immunisation uptake in school entrants. *Public Health* 1995;**109**:25-109.
- 573 - 1 Kamolratanakul P, Ungtavorn P, Israsena S, Sakulramrung R. The influence of dissemination of information on the changes of knowledge, attitude and acceptance of hepatitis B vaccination among hospital personnel in Chulalongkorn Hospital. *Public Health* 1994;**108**(1):49-53.
- 574 - 1 Hurt RD, Offord KP, Lauger GG, Marusic Z, Fagerstrom KO, Enright PL, Scanlon PD. Cessation of long-term nicotine gum use – a prospective, randomized trial. *Addiction* 1995;**90**(3):407-13.
- 575 - 1 Swaddiwudhipong W, Chaovakiratipong C, Nguntra P, Mahasakpan P, Lerdlukanavong P, Koonchote S. Effect of a mobile unit on changes in knowledge and use of cervical cancer screening among rural Thai women. *International Journal of Epidemiology* 1995;**24**(3):493-8.
- 576 - 1 Weinstein R, Tosolin F, Ghilardi L, Zanardelli E. Psychological intervention in patients with poor compliance. *Journal of Clinical Periodontology* 1996;**23**(3 Pt 2):283-8.
- 579 - 1 Archibald CP, Chan RK, Wong ML, Goh A, Goh CL. Evaluation of a safe-sex intervention programme among sex workers in Singapore. *International Journal of STD and AIDS* 1994;**5**(4):268-72.
- 581 - 1 Myles PS, Hendrata M, Layher Y, Williams NJ, Hall JL, Moloney JT, Powell J. Double-blind, randomized trial of cessation of smoking after audiotape suggestion during anaesthesia. *British Journal of Anaesthesia* 1996;**76**(5):694-8.
- 585 - 1 Stanton BF, Li X, Galbraith J, Feigelman S, Kaljee L. Sexually transmitted diseases, human immunodeficiency virus, and pregnancy prevention. Combined contraceptive practices among urban African-American early adolescents. *Archives of Pediatrics and Adolescent Medicine* 1996;**150**(1):17-24.
- 586 - 1 Kohn MR, Arden MR, Vasilakis J, Shenker IR. Directly observed preventive therapy. Turning the tide against tuberculosis. *Archives of Pediatrics and Adolescent Medicine* 1996;**150**(7):727-9.
- 587 - 1 Wolf AM, Nasser JF, Wolf AM, Schorling JB. The impact of informed consent on patient interest in prostate-specific antigen screening. *Archives of Internal Medicine* 1996;**156**(12):1333-6.
- 589 - 1 Norman P, Fitter M. Patients' views on health screening. *Family Practice* 1991;**8**:129-32.
- 594 - 1 Sheppard S, Coulter A, Farmer A. Using interactive videos in general practice to inform patients about treatment choices: a pilot study. *Family Practice* 1995;**12**:443-7.
- 600 - 1 Spoth R, Redmond C. Effective recruitment of parents into family-focused prevention research: a comparison of two strategies. *Psychology and Health* 1994;**9**:353-70.
- 601 - 1 Sander RW, Holloway RL, Eliason C, Marbella AM, Murphy B, Yuen S. Patient-initiated prevention discussions: two interventions to stimulate patients to initiate prevention discussions. *Journal Family Practice* 1996;**43**:468-74.

- 601 - 2** Sander RW, Holloway RL, Eliason C, marbella AM, Murphy B, Yuen S. Patient-initiated prevention discussions: two interventions to stimulate patients to initiate prevention discussions. *Journal Family Practice* 1996;**43**:468–74.
- 606 - 1** Van Heeringen C, Jannes S, Buylaert W, Henderick H, De Bacquer D, Van Remoortel J. The management of non-compliance with referral to out-patient after-care among attempted suicide patients: a controlled intervention study. *Psychological Medicine* 1995;**25**(5):963–70.
- 608 - 1** Turnbull D, Irwig L, Simpson JM, Donnelly N. The psychosocial impact of implementing a mammography screening campaign in an Australian community. *Social Science and Medicine* 1994;**39**(4):543–51.
- 610 - 1** Skinner D, Metcalf CA, Seager JR, de Swardt JS, Laubscher JA. An evaluation of an education programme on HIV infection using puppetry and street theatre. *AIDS Care* 1991;**3**(3):317–29.
- 612 - 1** Ferguson KA, Ono T, Lowe AA, Keenan SP, Fleetham JA. A randomised crossover study of an oral appliance vs nasal-continuous positive airway pressure in the treatment of mild-moderate obstructive sleep apnea. *Chest* 1996;**109**:1269–75.
- 616 - 1** Rossiter JC. The effect of a culture-specific education program to promote breastfeeding among Vietnamese women in Sydney. *International Journal of Nursing Studies* 1994;**31**(4):369–79.
- 618 - 1** Arinen SS, Sintonen H. The choice of dental-care sector by young- adults before and after subsidization reform in Finland. *Social Science and Medicine* 1994;**39**(2):291–7.
- 619 - 1** Dubois-Arber F, Masur JB, Hausser D, Zimmermann E, Paccaud F. Evaluation of AIDS prevention among homosexual and bisexual men in Switzerland. *Social Science and Medicine* 1993;**37**(12):1539–44.
- 620 - 1** Norman P. Predicting the uptake of health checks in general practice: invitation methods and patients' health beliefs. *Social Science and Medicine* 1993;**37**(1):53–9.
- 621 - 1** Guptill KS, Esrey SA, Oni GA, Brown KH. Evaluation of a face-to-face weaning food intervention in Kwara State, Nigeria: knowledge, trial, and adoption of a home-prepared weaning food. *Social Science and Medicine* 1993;**36**(5):665–72.
- 622 - 1** Jones SL, Jones PK, Katz J. Compliance in acute and chronic patients receiving a health belief model intervention in the emergency department. *Social Science and Medicine* 1991;**32**(10):1183–9.
- 623 - 1** Bruce N, Griffioen A. Usefulness of a non-experimental study design in the evaluation of service developments for infant feeding in a general hospital. *Social Science and Medicine* 1995;**40**(8):1109–16.
- 628 - 1** Rotheram-Borus MJ, Koopman C, Haignere C, Davies M. Reducing HIV sexual risk behaviors among runaway adolescents. *JAMA* 1991;**266**(9):1237–41.
- 629 - 1** Meier DE, Fuss BR, O'Rourke D, Baskin SA, Lewis M, Morrison RS. Marked improvement in recognition and completion of health-care proxies – a randomized controlled trial of counseling by hospital patient representatives. *Archives of Internal Medicine* 1996;**156**(11):1227–32.
- 630 - 1** Sharp DJ, Peters TJ, Bartholomew J, Shaw A. Breast screening: a randomised controlled trial in UK general practice of three interventions designed to increase uptake. *Journal of Epidemiology and Community Health* 1996;**50**(1):72–6.
- 631 - 1** Fortmann SP, Taylor CB, Flora JA, Winkleby MA. Effect of community health education on plasma cholesterol levels and diet: the Stanford Five-City Project. *American Journal of Epidemiology* 1993;**137**(10):1039–55.
- 632 - 1** Sorensen JL, London J, Heitzmann C, Gibson DR, Morales ES, Dumontet R, Acree M. Psychoeducational group approach: HIV risk reduction in drug users. *AIDS Education and Prevention* 1994;**6**(2):95–112.
- 632 - 2** Sorensen JL, London J, Heitzmann C, Gibson DR, Morales ES, Dumontet R, Acree M. Psychoeducational group approach: HIV risk reduction in drug users. *AIDS Education and Prevention* 1994;**6**(2):95–112.
- 633 - 1** Black ME, Ploeg J, Walter SD, Hutchinson BG, Scott EA, Chambers LW. The impact of a public health nurse intervention on influenza vaccine acceptance. *American Journal of Public Health* 1993;**83**(12):1751–3.
- 634 - 1** Torgerson DJ, Donaldson C, Garton MJ, Reid DM, Russell IT. Recruitment methods for screening programmes: the price of high compliance. *Health Economics* 1993;**2**(1):55–8.
- 635 - 1** Malow RM, West JA, Corrigan SA, Pena JM, Cunningham SC. Outcome of psychoeducation for HIV risk reduction. *AIDS Education and Prevention* 1994;**6**(2):113–25.
- 636 - 1** McCusker J, Stoddard AM, Zapka JG, Lewis BF. Behavioral outcomes of AIDS educational interventions for drug users in short-term treatment. *American Journal of Public Health* 1993;**83**(10):1463–6.



- 638 - 1** Wasson J, Gaudette C, Whaley F, Sauvigne A, Baribeau P, Welch HG. Telephone care as a substitute for routine clinic follow-up. *JAMA* 1992;**267**(13):1788–93.
- 639 - 1** Ashworth CS, DuRant RH, Gaillard G, Rountree J. An experimental evaluation of an AIDS educational intervention for WIC mothers. *AIDS Education and Prevention* 1994;**6**(2):154–62.
- 640 - 1** Elder JP, Wildey M, de Moor C, Sallis JF, Jr, Eckhardt L, Edwards C, Erickson A, Golbeck A, Hovell M, Johnston D, Levitz MD, Molgaard L, Young R, Vito D, Woodruff SI. The long-term prevention of tobacco use among junior high school students: classroom and telephone interventions. *American Journal of Public Health* 1993;**83**(9):1239–44.
- 644 - 1** Gomel M, Oldenburg B, Simpson JM, Owen N. Work-site cardiovascular risk reduction: a randomized trial of health risk assessment, education, counseling, and incentives. *American Journal of Public Health* 1993;**83**(9):1231–8.
- 646 - 1** Nyamathi AM, Flaskerud J, Bennett C, Leake B, Lewis C. Evaluation of two AIDS education programs for impoverished Latina women. *AIDS Education and Prevention* 1994;**6**(4):296–309.
- 647 - 1** Kelly JA, St Lawrence JS, Stevenson LY, Hauth AC, Kalichman SC, Diaz YE, Brasfield TL, Koob JJ, Morgan MG. Community AIDS/HIV risk reduction: the effects of endorsements by popular people in three cities. *American Journal of Public Health* 1992;**82**(11):1483–9.
- 648 - 1** Jorenby DE, Smith SS, Fiore MC, Hurt RD, Offord KP, Croghan IT, Hays JT, Lewis SF Baker TB. Varying nicotine patch dose and type of smoking cessation counseling. *JAMA* 1995;**274**(17):1347–52.
- 649 - 1** Bellingham K, Gillies P. Evaluation of an AIDS education programme for young adults. *Journal of Epidemiology and Community Health* 1993;**47**(2):134–8.
- 650 - 1** Berrier J, Sperling R, Preisinger J, Evans V, Mason J, Walther V. HIV/AIDS education in a prenatal clinic: an assessment. *AIDS Education and Prevention* 1991;**3**:100–17.
- 651 - 1** Allen S, Serufulira A, Bogaerts J, Van de Perre P, Nsengumuremyi F, Lindan C, Carael M, Wolf W, Coates T, Hulley S. Confidential HIV testing and condom promotion in Africa. Impact on HIV and gonorrhoea rates. *JAMA* 1992;**268**(23):3338–43.
- 652 - 1** Maiman LA, Hildreth NG, Cox C, Greenland P. Improving referral compliance after public cholesterol screening. *American Journal of Public Health* 1992;**82**(6):804–9.
- 653 - 1** Peeters PH, Beckers CG, Hogervorst JM, Collette HJ. Effect on breast cancer screening response in The Netherlands of inviting women for an additional scientific investigation. *Journal of Epidemiology and Community Health* 1994;**48**(2):175–7.
- 654 - 1** Flowers JV, Booraem C, Miller TE, Iverson AE, Copeland J, Furtado K. Comparison of the results of a standardized AIDS prevention program in three geographic locations. *AIDS Education and Prevention* 1991;**3**:189–96.
- 655 - 1** King R, Estey J, Allen S, Kegeles S, Wolf W, Valentine C, Serufulira A. A family planning intervention to reduce vertical transmission of HIV in Rwanda. *AIDS* 1995;**9**(Suppl 1):S45–51.
- 657 - 1** Warnecke RB, Langenberg P, Wong SC, Flay BR, Cook TD. The second Chicago televised smoking cessation program: a 24-month follow-up. *American Journal of Public Health* 1992;**82**(6):835–40.
- 658 - 1** O'Loughlin J, Paradis G, Kishchuk N, Gray-Donald K, Renaud L, Fines P, Barnett T. Coeur en sante St-Henri—a heart health promotion programme in Montreal, Canada: design and methods for evaluation. *Journal of Epidemiology and Community Health* 1995;**49**(5):495–502.
- 659 - 1** Shattuck AL, White E, Kristal AR. How women's adopted low-fat diets affect their husbands. *American Journal of Public Health* 1992;**82**(9):1244–50.
- 660 - 1** Walter HJ, Vaughan RD. AIDS risk reduction among a multiethnic sample of urban high school students. *JAMA* 1993;**270**(6):725–30.
- 662 - 1** Langer LM, Zimmerman RS, Hendershot EF, Singh M. Effect of Magic Johnson's HIV status on HIV-related attitudes and behaviors of an STD clinic population. *AIDS Education and Prevention* 1992;**4**(4):295–307.
- 663 - 1** Aplasca MR, Siegel D, Mandel JS, Santana-Arciaga RT, Paul J, Hudes ES, Monzon OT, Hearst N. Results of a model AIDS prevention program for high school students in the Philippines. *AIDS* 1995;**9**(Suppl 1):S7–13.
- 665 - 1** Kelly JA, St Lawrence JS, Diaz YE, Stevenson LY, Hauth AC, Brasfield TL, Kalichman SC, Smith JE, Andrew ME. HIV risk behavior reduction following intervention with key opinion leaders of population: an experimental analysis. *American Journal of Public Health* 1991;**81**(2):168–71.
- 667 - 1** Street RL, Jr, Voigt B, Geyer C, Jr, Manning T, Swanson GP. Increasing patient involvement in choosing treatment for early breast cancer. *Cancer* 1995;**76**(11):2275–85.

- 668 - 1 Velicer WF, Prochaska JO, Bellis JM, Diclemente CC, Rossi JS, Fava JL, Steiger JH. An expert system intervention for smoking cessation. *Addictive Behaviours* 1993;18(3):269-90.
- 670 - 1 Ross JD, Scott GR. The association between HIV media campaigns and number of patients coming forward for HIV antibody testing. *Genitourinary Medicine* 1993;69(3):193-5.
- 671 - 1 Phatouros CC, Blake MP. How much now to tell? Patients' attitudes to an information sheet prior to angiography and angioplasty. *Australasian Radiology* 1995;39(2):135-9.
- 674 - 1 Banahan I, Quenby S, Stewart H, Farquharson R. Preliminary evaluation of the effectiveness of a preoperative clinic for gynaecological surgery. *British Journal of Hospital Medicine* 1994;52(10):535-8.
- 675 - 1 Robinson MH, Pye G, Thomas WM, Hardcastle JD, Mangham CM. Haemoccult screening for colorectal cancer: the effect of dietary restriction on compliance. *European Journal of Surgical Oncology* 1994;20(5):545-8.
- 676 - 1 D'Souza W, Burgess C, Ayson M, Crane J, Pearce N, Beasley R. Trial of a 'credit card' asthma self-management plan in a high-risk group of patients with asthma. *Journal of Allergy and Clinical Immunology* 1996;97(5):1085-92.
- 678 - 1 Rowley MJ, Hensley MJ, Brinsmead MW, Wlodarczyk JH. Continuity of care by a midwife team versus routine care during pregnancy and birth: a randomised trial. *Medical Journal of Australia* 1995;163(6):289-93.
- 679 - 1 Harrell JS, McMurray RG, Bangdiwala SI, Frauman AC, Gansky SA, Bradley CB. Effects of a school-based intervention to reduce cardiovascular disease risk factors in elementary-school children: the Cardiovascular Health in Children (CHIC) study. *Journal of Pediatrics* 1996;128(6):797-805.
- 680 - 1 van Haastrecht HJ, van den Hoek JA, Coutinho RA. Evidence for a change in behaviour among heterosexuals in Amsterdam under the influence of AIDS. *Genitourinary Medicine* 1991;67(3):199-206.
- 683 - 1 King J, Fairbrother G, Thompson C, Morris DL. Colorectal cancer screening: optimal compliance with postal faecal occult blood test. *Australian and New Zealand Journal of Surgery* 1992;62(9):714-9.
- 684 - 1 McCusker J, Stoddard AM, Zapka JG, Zorn M. Use of condoms by heterosexually active drug abusers before and after AIDS education. *Sexually Transmitted Diseases* 1993;20(2):81-8.
- 685 - 1 Winter L, Goldy AS. Effects of prebehavioral cognitive work on adolescents' acceptance of condoms. *Health Psychology* 1993;12(4):308-12.
- 686 - 1 Fletcher SW, Harris RP, Gonzalez JJ, Degnan D, Lannin DR, Strecher VJ, Pilgrim C, Quade D, Earp JA, Clark RL. Increasing mammography utilization: a controlled study. *Journal of the National Cancer Institute* 1993;85(2):112-20.
- 688 - 1 Kamenga M, Ryder RW, Jingu M, Mbuyi N, Mbu L, Behets F, Brown C, Heyward WL. Evidence of marked sexual behavior change associated with low HIV-1 seroconversion in 149 married couples with discordant HIV-1 serostatus: experience at an HIV counselling center in Zaire. *AIDS* 1991;5(1):61-7.
- 690 - 1 Champion VL. Results of a nurse-delivered intervention on proficiency and nodule detection with breast self-examination. *Oncology Nursing Forum* 1995;22(5):819-24.
- 691 - 1 Whelan TJ, Levine MN, Gafni A, Lukka H, Mohide EA, Patel M, Streiner DL. Breast irradiation postlumpectomy: development and evaluation of a decision instrument. *Journal of Clinical Oncology* 1995;13(4):847-53.
- 693 - 1 Esposito L. The effects of medication education on adherence to medication regimens in an elderly population. *Journal of Advanced Nursing* 1995;21(5):935-43.
- 695 - 1 Hoare T, Thomas C, Biggs A, Booth M, Bradley S, Friedman E. Can the uptake of breast screening by Asian women be increased? A randomized controlled trial of a linkworker intervention. *Journal of Public Health Medicine* 1994;16(2):179-85.
- 696 - 1 O'Donnell L, San Doval A, Duran R, O'Donnell CR. The effectiveness of video-based interventions in promoting condom acquisition among STD clinic patients. *Sexually Transmitted Diseases* 1995;22(2):97-103.
- 698 - 1 Doren M, Reuther G, Minne HW, Schneider HP. Superior compliance and efficacy of continuous combined oral estrogen-progestogen replacement therapy in postmenopausal women. *American Journal of Obstetrics and Gynecology* 1995;173(5):1446-51.
- 700 - 1 Butow PN, Dunn SM, Tattersall MH, Jones QJ. Patient participation in the cancer consultation: evaluation of a question prompt sheet. *Annals of Oncology* 1994;5(3):199-204.
- 702 - 1 Cargill JM. Medication compliance in elderly people: influencing variables and interventions. *Journal of Advanced Nursing* 1992;17(4):422-6.
- 705 - 1 Hughes BR, Altman DG, Newton JA. Melanoma and skin cancer: evaluation of a health education programme for secondary schools. *British Journal of Dermatology* 1993;128(4):412-17.

- 706 - 1** Usherwood TP. Development and randomized controlled trial of a booklet of advice for parents. *British Journal of General Practice* 1991;**41** (343):58–62.
- 707 - 1** DeBusk RF, Miller NH, Superko HR, Dennis CA, Thomas RJ, Lew HT, Berger WE, 3rd, Heller RS, Rompf J, Gee D, Kraemer HC, Bandura A, Ghandour G, Clark M, Shah RV, Fisher L, Barr-Taylor C. A case-management system for coronary risk factor modification after acute myocardial infarction. *Annals of Internal Medicine* 1994;**120**(9):721–9.
- 708 - 1** Robertson I, Phillips A, Mant D, Thorogood M, Fowler G, Fuller A, Yudkin P, Woods M. Motivational effect of cholesterol measurement in general practice health checks. *British Journal of General Practice* 1992;**42**(364):469–72.
- 709 - 1** Osler M, Jespersen NB. The effect of a community-based cardiovascular disease prevention project in a Danish municipality. *Danish Medical Bulletin* 1993;**40**(4):485–9.
- 711 - 1** Scivoletto S, De Andrade AG, Castel S. The effect of a 'recall-system' in the treatment of alcoholic patients. *British Journal of Addiction* 1992;**87**(8):1185–8.
- 712 - 1** Schapira DV, Kumar NB, Clark RA, Yag C. Mammography screening credit card and compliance. *Cancer* 1992;**70**(2):509–12.
- 713 - 1** Stewart DE, Buchegger PM, Lickrish GM, Sierra S. The effect of educational brochures on follow-up compliance in women with abnormal Papanicolaou smears. *Obstetrics and Gynecology* 1994;**83**(4):583–5.
- 714 - 1** Nelson EW, Van Cleve S, Swartz MK, Kessen W, McCarthy PL. Improving the use of early follow-up care after emergency department visits. A randomized trial. *American Journal of Diseases of Children* 1991;**145**(4):440–4.
- 717 - 1** Webster A. The effect of pre-assessment information on clients' satisfaction, expectations and attendance at a mental health day centre. *British Journal of Medical Psychology* 1992;**65**(Pt 2):89–93.
- 720 - 1** Zapka JG, Stoddard AM, McCusker J. Social network, support and influence: relationships with drug use and protective AIDS behavior. *AIDS Education and Prevention* 1993;**5**(4):352–66.
- 723 - 1** Fox LJ, Bailey PE, Clarke-Martinez KL, Coello M, Ordonez FN, Barahona F. Condom use among high-risk women in Honduras: evaluation of an AIDS prevention program. *AIDS Education and Prevention* 1993;**5**(1):1–10.
- 724 - 1** Windsor RA, Lowe JB, Perkins LL, Smith-Yoder D, Artz L, Crawford M, Amburgy K, Boyd NR, Jr. Health education for pregnant smokers: its behavioral impact and cost benefit. *American Journal of Public Health* 1993;**83**(2):201–6.
- 726 - 1** Luepker RV, Murray DM, Jacobs DR, Jr, Mittelman MB, Bracht N, Carlaw R, Crow R, Elmer P, Finnegan J, Folsom AR, Grimm R, Hannan PJ, Jeffrey R, Laubo H, Govern P, Mullis R, Perry CL, Pechacek T, Pirie P, Sprafka M, Weisbrod R, Blackburn H. Community education for cardiovascular disease prevention: risk factor changes in the Minnesota Heart Health Program. *American Journal of Public Health* 1994;**84**(9):1383–93.
- 727 - 1** Hurley SF, Huggins RM, Jolley DJ, Reading D. Recruitment activities and sociodemographic factors that predict attendance at a mammographic screening program. *American Journal of Public Health* 1994;**84**(10):1655–8.
- 728 - 1** McAvoy BR, Raza R. Can health education increase uptake of cervical smear testing among Asian women? *British Medical Journal* 1991;**302**(6780):833–6.
- 729 - 1** Stephens RC, Feucht TE, Roman SW. Effects of an intervention program on AIDS-related drug and needle behavior among intravenous drug users. *American Journal of Public Health* 1991;**81**(5):568–71.
- 731 - 1** Nutbeam D, Macaskill P, Smith C, Simpson JM, Catford J. Evaluation of two school smoking education programmes under normal classroom conditions. *British Medical Journal* 1993;**306**(6870):102–7.
- 732 - 1** Grosskurth H, Mosha F, Todd J, Mwijarubi E, Klokke A, Senkoro K, Mayaud P, Changalucha J, Nicoll A, ka-Gina G, Newell J, Mugeye K, Mabeg D, Hayes R. Impact of improved treatment of sexually transmitted diseases on HIV infection in rural Tanzania: randomised controlled trial. *Lancet* 1995;**346**(8974):530–6.
- 733 - 1** Black MM, Nair P, Kight C, Wachtel R, Roby P, Schuler M. Parenting and early development among children of drug-abusing women: effects of home intervention. *Pediatrics* 1994;**94**(4 Pt 1):440–8.
- 736 - 1** Calsyn DA, Saxon AJ, Freeman G, Jr, Whittaker S. Ineffectiveness of AIDS education and HIV antibody testing in reducing high-risk behaviors among injection drug users. *American Journal of Public Health* 1992;**82**(4):573–5.
- 737 - 1** Wilson TG, Jr, Hale S, Temple R. The results of efforts to improve compliance with supportive periodontal treatment in a private practice. *Journal of Periodontology* 1993;**64**(4):311–14.
- 740 - 1** Rossouw JE, Jooste PL, Chalton DO, Jordaan ER, Langenhoven ML, Jordaan PC, Steyn M, Swanepoel AS, Rossouw LJ. Community-based intervention: the Coronary Risk Factor Study (CORIS). *International Journal of Epidemiology* 1993;**22**(3):428–38.

- 741 - 1** Wenger NS, Linn LS, Epstein M, Shapiro MF. Reduction of high-risk sexual behavior among heterosexuals undergoing HIV antibody testing: a randomized clinical trial. *American Journal of Public Health* 1991;**81**(12):1580-5.
- 742 - 1** Hobfoll SE, Jackson AP, Lavin J, Britton PJ, Shepherd JB. Reducing inner-city women's AIDS risk activities: a study of single, pregnant women. *Health Psychology* 1994;**13**(5):397-403.
- 743 - 1** Miedzybrodzka ZH, Hall MH, Mollison J, Templeton A, Russell IT, Dean JC, Kelly KF, Marteau TM, Haites NE. Antenatal screening for carriers of cystic fibrosis: randomised trial of stepwise v couple screening. *British Medical Journal* 1995;**310**(6976):353-7.
- 746 - 1** Kernohan EE. Evaluation of a pilot study for breast and cervical cancer screening with Bradford's minority ethnic women; a community development approach, 1991-93. *British Journal of Cancer* 1996;**74**(Suppl 29):S42-6.
- 747 - 1** Garton MJ, Torgerson DJ, Donaldson C, Russell IT, Reid DM. Recruitment methods for screening programmes: trial of a new method within a regional osteoporosis study. *British Medical Journal* 1992;**305**(6845):82-4.
- 748 - 1** Smith S, Robinson J, Hollyer J, Bhatt R, Ash S, Shaunak S. Combining specialist and primary health care teams for HIV positive patients: retrospective and prospective studies. *British Medical Journal* 1996;**312**(7028):416-20.
- 749 - 1** Jemmott JB, 3rd, Jemmott LS, Fong GT. Reductions in HIV risk-associated sexual behaviors among black male adolescents: effects of an AIDS prevention intervention [published erratum appears in *American Journal of Public Health* 1992;**82**(5):684]. *American Journal of Public Health* 1992;**82**(3):372-7.
- 752 - 1** Mant D, Fuller A, Northover J, Astrop P, Chivers A, Crockett A, Clements S, Lawrence M. Patient compliance with colorectal cancer screening in general practice. *British Journal of General Practice* 1992;**42**(354):18-20.
- 754 - 1** Shannon BM, Tershakovec AM, Martel JK, Achterberg CL, Cortner JA, Smiciklas-Wright HS, Stallings VA, Stolley PD. Reduction of elevated LDL-cholesterol levels of 4- to 10-year-old children through home-based dietary education. *Pediatrics* 1994;**94**(6 Pt 1):923-7.
- 755 - 1** Hanlon P, McEwen J, Carey L, Gilmour H, Tannahill C, Tannahill A, Kelly M. Health checks and coronary risk: further evidence from a randomised controlled trial. *British Medical Journal* 1995;**311**(7020):1609-13.
- 757 - 1** Lerman C, Ross E, Boyce A, Gorchov PM, McLaughlin R, Rimer B, Engstrom P. The impact of mailing psychoeducational materials to women with abnormal mammograms. *American Journal of Public Health* 1992;**82**(5):729-30.
- 758 - 1** Tedesco LA, Keffer MA, Davis EL, Christersson LA. Effect of a social cognitive intervention on oral health status, behavior reports, and cognitions. *Journal of Periodontology* 1992;**63**(7):567-75.
- 759 - 1** Mellanby AR, Phelps FA, Crichton NJ, Tripp JH. School sex education: an experimental programme with educational and medical benefit. *British Medical Journal* 1995;**311**(7002):414-7.
- 760 - 1** Aiken LS, West SG, Woodward CK, Reno RR, Reynolds KD. Increasing screening mammography in asymptomatic women: evaluation of a second-generation, theory-based program. *Health Psychology* 1994;**13**(6):526-38.
- 762 - 1** Kalichman SC, Kelly JA, Hunter TL, Murphy DA, Tyler R. Culturally tailored HIV-AIDS risk-reduction messages targeted to African-American urban women: impact on risk sensitization and risk reduction. *Journal of Consulting and Clinical Psychology* 1993;**61**(2):291-5.
- 763 - 1** Lowe CJ, Raynor DK, Courtney EA, Purvis J, Teale C. Effects of self medication programme on knowledge of drugs and compliance with treatment in elderly patients. *British Medical Journal* 1995;**310**(6989):1229-31.
- 765 - 1** Lombard D, Neubauer TE, Canfield D, Winett RA. Behavioral community intervention to reduce the risk of skin cancer. *Journal of Applied Behavior Analysis* 1991;**24**(4):677-86.
- 766 - 1** St Lawrence JS, Brasfield TL, Jefferson KW, Alleyne E, O'Bannon RE, 3rd, Shirley A. Cognitive-behavioral intervention to reduce African American adolescents' risk for HIV infection. *Journal of Consulting and Clinical Psychology* 1995;**63**(2):221-37.
- 767 - 1** Reading R, Colver A, Openshaw S, Jarvis S. Do interventions that improve immunisation uptake also reduce social inequalities in uptake? *British Medical Journal* 1994;**308**(6937):1142-4.
- 768 - 1** Dungy CI, Christensen-Szalanski J, Losch M, Russell D. Effect of discharge samples on duration of breast-feeding. *Pediatrics* 1992;**90**(2 Pt 1):233-7.
- 769 - 1** Dale J, Lang H, Roberts JA, Green J, Glucksman E. Cost effectiveness of treating primary care patients in accident and emergency: a comparison between general practitioners, senior house officers, and registrars. *British Medical Journal* 1996;**312**(7042):1340-4.
- 770 - 1** Colon HM, Sahai H, Robles RR, Matos TD. Effects of a community outreach program in HIV risk behaviors among injection drug users in San Juan, Puerto Rico: an analysis of trends. *AIDS Education and Prevention* 1995;**7**(3):195-209.

- 772 - 1** Miller WR, Benefield RG, Tonigan JS. Enhancing motivation for change in problem drinking: a controlled comparison of two therapist styles. *Journal of Consulting and Clinical Psychology* 1993;**61**(3):455–61.
- 773 - 1** Pendergrast RA, Ashworth CS, DuRant RH, Litaker M. Correlates of children's bicycle helmet use and short-term failure of school-level interventions. *Pediatrics* 1992;**90**(3):354–8.
- 775 - 1** Li J, Taylor B. Comparison of immunisation rates in general practice and child health clinics. *British Medical Journal* 1991;**303**(6809):1035–8.
- 776 - 1** Kelly JA, Murphy DA, Washington CD, Wilson TS, Koob JJ, Davis DR, Ledezma G, Davantes B. The effects of HIV/AIDS intervention groups for high-risk women in urban clinics. *American Journal of Public Health* 1994;**84**(12):1918–22.
- 777 - 1** West S, Munoz B, Lynch M, Kayongoya A, Chilangwa Z, Mmbaga BB, Taylor HR. Impact of face-washing on trachoma in Kongwa, Tanzania. *Lancet* 1995;**345**(8943):155–8.
- 778 - 1** Thornton JG, Hewison J, Lilford RJ, Vail A. A randomised trial of three methods of giving information about prenatal testing. *British Medical Journal* 1995;**311**(7013):1127–30.
- 779 - 1** Swaddiwudhipong W, Chaovakiratipong C, Nguntra P, Khumklam P, Silarug N. A Thai monk: an agent for smoking reduction in a rural population. *International Journal of Epidemiology* 1993;**22**(4):660–5.
- 780 - 1** Santelli JS, Celentano DD, Rozsenich C, Crump AD, Davis MV, Polacsek M, Augustyn M, Rolf J, McAlister AL, Burwell L. Interim outcomes for a community-based program to prevent perinatal HIV transmission. *AIDS Education and Prevention* 1995;**7**(3):210–20.
- 781 - 1** Lahdensuo A, Haahtela T, Herrala J, Kava T, Kiviranta K, Kuusisto P, Peramaki E, Poussa T, Saarelainen S, Svahn T. Randomised comparison of guided self management and traditional treatment of asthma over one year. *British Medical Journal* 1996;**312**(7033):748–52.
- 783 - 1** Orleans CT, Schoenbach VJ, Wagner EH, Quade D, Salmon MA, Pearson DC, Fiedler J, Porter CQ, Kaplan BH. Self-help quit smoking interventions: effects of self-help materials, social support instructions, and telephone counseling. *Journal of Consulting and Clinical Psychology* 1991;**59**(3):439–48.
- 784 - 1** Sutton S, Bickler G, Sancho-Aldridge J, Saidi G. Prospective study of predictors of attendance for breast screening in inner London. *Journal of Epidemiology and Community Health* 1994;**48**(1):65–73.
- 788 - 1** Serwint JR, Wilson MH, Duggan AK, Mellits ED, Baumgardner RA, DeAngelis C. Do postpartum nursery visits by the primary care provider make a difference? *Pediatrics* 1991;**88**(3):444–9.
- 789 - 1** Rothman AJ, Salovey P, Turvey C, Fishkin SA. Attributions of responsibility and persuasion: increasing mammography utilization among women over 40 with an internally oriented message. *Health Psychology* 1993;**12**(1):39–47.
- 790 - 1** Bekker H, Modell M, Denniss G, Silver A, Mathew C, Bobrow M, Marteau T. Uptake of cystic fibrosis testing in primary care: supply push or demand pull? *British Medical Journal* 1993;**306**(6892):1584–6.
- 791 - 1** Bhave G, Lindan CP, Hudes ES, Desai S, Wagle U, Tripathi SP, Mandel JS. Impact of an intervention on HIV, sexually transmitted diseases, and condom use among sex workers in Bombay, India. *AIDS* 1995;**9**(Suppl 1):S21–30.
- 792 - 1** Flynn BS, Worden JK, Secker-Walker RH, Badger GJ, Geller BM, Costanza MC. Prevention of cigarette smoking through mass media intervention and school programs. *American Journal of Public Health* 1992;**82**(6):827–34.
- 793 - 1** Anonymous. Effectiveness of a nicotine patch in helping people stop smoking: results of a randomised trial in general practice. Imperial Cancer Research Fund General Practice Research Group. *British Medical Journal* 1993;**306**(6888):1304–8.
- 794 - 1** Katon W, Von Korff M, Lin E, Walker E, Simon GE, Bush T, Robinson P, Russo J. Collaborative management to achieve treatment guidelines. Impact on depression in primary care. *JAMA* 1995;**273**(13):1026–31.
- 795 - 1** Reid GS, Robertson AJ, Bissett C, Smith J, Waugh N, Halkerston R. Cervical screening in Perth and Kinross since introduction of the new contract. *British Medical Journal* 1991;**303**(6800):447–50.
- 796 - 1** Rutten G, Van Eijk J, Beek M, Van der Velden H. Patient education about cough: effect on the consulting behaviour of general practice patients. *British Journal of General Practice* 1991;**41**(348):289–92.
- 798 - 1** Gibbins RL, Riley M, Brimble P. Effectiveness of programme for reducing cardiovascular risk for men in one general practice. *British Medical Journal* 1993;**306**(6893):1652–6.
- 799 - 1** DiClemente RJ, Wingood GM. A randomized controlled trial of an HIV sexual risk-reduction intervention for young African-American women. *JAMA* 1995;**274**(16):1271–6.
- 807 - 1** Maxwell AE, Hunt IF, Bush MA. Effects of a social support group, as an adjunct to diabetes training, on metabolic control and psychosocial outcomes. *Diabetes Educator* 1992;**18**(4):303–9.

- 809 - 1** Zapka JG, Harris DR, Hosmer D, Costanza ME, Mas E, Barth R. Effect of a community health center intervention on breast cancer screening among Hispanic American women. *Health Services Research* 1993;**28**(2):223–35.
- 813 - 1** Leigh JP, Richardson N, Beck R, Kerr C, Harrington H, Parcell CL, Fries JF. Randomized controlled study of a retiree health promotion program. The Bank of American Study. *Archives of Internal Medicine* 1992;**152**(6):1201–6.
- 814 - 1** Marcus BH, Stanton AL. Evaluation of relapse prevention and reinforcement interventions to promote exercise adherence in sedentary females. *Research Quarterly for Exercise and Sport* 1993;**64**(4):447–52.
- 815 - 1** Rodriguez JG, Velez M, Serrano E, Casado MP. Adolescent student's compliance with testicular self examination. *Boletin – Asociacion Medica de Puerto Rico* 1995;**87**(3–4):49–53.
- 816 - 1** Aaronson NK, Visserpol E, Leenhouts GHMW, Muller MJ, Vanderschot ACM, Vandam FSAM, Keus RB, Koning CCE, Huinink WWT, Vandongen JA, Dubbelman R. Telephone-based nursing intervention improves the effectiveness of the informed consent process in cancer clinical-trials. *Journal of Clinical Oncology* 1996;**14**(3):984–96.
- 817 - 1** McCusker J, Stoddard AM, Zapka JG, Morrison CS, Zorn M, Lewis BF. AIDS education for drug abusers: evaluation of short-term effectiveness. *American Journal of Public Health* 1992;**82**(4):533–40.
- 821 - 1** Williams LR, Ekers MA, Collins PS, Lee JF. Vascular rehabilitation: benefits of a structured exercise/risk modification program. *Journal of Vascular Surgery* 1991;**14**(3):320–6.
- 822 - 1** Kirby D, Waszak C, Ziegler J. Six school-based clinics: their reproductive health services and impact on sexual behavior. *Family Planning Perspectives* 1991;**23**(1):6–16.
- 823 - 1** Anonymous. Effectiveness of a health education curriculum for secondary school students – United States, 1986–1989. *MMWR – Morbidity and Mortality Weekly Report* 1991;**40**(7):113–16.
- 825 - 1** Guimon J. The use of group programs to improve medication compliance in patients with chronic diseases. *Patient Education and Counseling* 1995;**26**:189–93.

# Appendix 6

## Bibliography: studies by author

(Study numbers appear at the end of each reference)

- Aaonson NK, Visserpol E, Leenhouts GHMW, Muller MJ, Vanderschot ACM, Vandam FSAM, Keus RB, Koning CCE, Huinink WWT, Vandonden JA, Dubbelman R. *Journal of Clinical Oncology* 1996;**14**(3):984–96. **816 - 1**
- Agras WS, Berkowitz RI, Arnow BA, Telch CF, Marnell M, Henderson J, Morris Y, Wilfley DE. *Journal of Consulting and Clinical Psychology* 1996;**64**(3):610–13. **32 - 1**
- Aiken LS, West SG, Woodward CK, Reno RR, Reynolds KD. *Health Psychology* 1994;**13**(6):526–38. **760 - 1**
- Alcoe SY, Gilbey VJ, McDermot RSR, Wallace DG. *Patient Education and Counseling* 1993;**21**:117–23. **491 - 1**
- Alcoe SY, Gilbey VJ, McDermot RSR, Wallace DG. *Patient Education and Counseling* 1995;**25**:183–96. **533 - 1**
- Alexandrov A, Maslennikova G, Kulikov S, Propirnij G, Perova N. *Preventive Medicine* 1992;**21**:53–62. **279 - 1**
- Allen RM, Jones MP, Oldenburg B. *Thorax* 1995;**50**(7):731–8. **530 - 1**
- Allen S, Serufulira A, Bogaerts J, Van de Perre P, Nsengumuremyi F, Lindan C, Carael M, Wolf W, Coates T, Hulley S. *JAMA* 1992;**268**(23):3338–43. **651 - 1**
- Anderson AS, Campbell DM, Shepherd R. *British Journal of Nutrition* 1995;**73**(2):163–77. **115 - 1**
- Anonymous. *British Medical Journal* 1993;**306**(6888):1304–8. **793 - 1**
- Anonymous. *MMWR – Morbidity and Mortality Weekly Report* 1993;**42**(23):448–9, 455–6. **545 - 1**
- Anonymous. *MMWR – Morbidity and Mortality Weekly Report* 1991;**40**(7):113–16. **823 - 1**
- Ansell D, Lacey L, Whitman S, Chen E, Phillips C. *Public Health Reports* 1994;**109**(1):104–11. **397 - 1**
- Aplasca MR, Siegel D, Mandel JS, Santana-Arciaga RT, Paul J, Hudes ES, Monzon OT, Hearst N. *AIDS* 1995;**9** Suppl 1:S7–13. **663 - 1**
- Archibald CP, Chan RK, Wong ML, Goh A, Goh CL. *International Journal of STD and AIDS* 1994;**5**(4):268–72. **579 - 1**
- Arinen SS, Sintonen H. *Social Science and Medicine* 1994;**39**(2):291–7. **618 - 1**
- Ashworth CS, DuRant RH, Gaillard G, Rountree J. *AIDS Education and Prevention* 1994;**6**(2):154–62. **639 - 1**
- Avila P, Hovell MF. *International Journal of Obesity* 1994;**18**:476–82. **174 - 1**
- Baier CA, Grodzin CJ, Port JD, Leksas L, Tancredi DJ. *American Journal of Preventive Medicine* 1992;**8**(2):115–22. **384 - 1**
- Baker A, Kochan N, Dixon J, Heather N, Wodak A. *AIDS Care* 1994;**6**(5):559–70. **525 - 1**
- Banahan I, Quenby S, Stewart H, Farquharson R. *British Journal of Hospital Medicine* 1994;**52**(10):535–8. **674 - 1**
- Banks SM, Salovey P, Greener S, Rothman AJ, Moyer A, Beauvais J, Epel E. *Health Psychology* 1995;**14**(2):178–84. **438 - 1**
- Barnwell MD, Chitkara R, Lamberta F. *Journal of the National Medical Association* 1992;**84**(12):1014–18. **411 - 1**
- Bastani R, Marcus AC, Maxwell AE, Das IP, Yan KX. *Preventive Medicine* 1994;**23**(1):83–90. **232 - 1**
- Baum JG, Clark HB, Sandler J. *Journal of Behavioral Medicine* 1991;**14**(3):287–302. **414 - 1**
- Baumann LJ, Zimmerman R, Leventhal H. *Patient Education and Counseling* 1989;**14**:53–67. **305 - 1**
- Bazeley P, Kemp L. *Australian Journal of Public Health* 1995;**19**(5):459–64. **193 - 1**
- Bearss N, Santelli JS, Papa P. *Journal of Adolescent Health* 1995;**17**(3):178–83. **38 - 1**
- Bejes C, Marvel MK. *Family Practice Research Journal* 1992;**12**(1):83–90. **296 - 1**
- Bekker H, Modell M, Denniss G, Silver A, Mathew C, Bobrow M, Marteau T. *British Medical Journal* 1993;**306**(6892):1584–6. **790 - 1**
- Bell RM, Ellickson PL, Harrison ER. *Preventive Medicine* 1993;**22**(4):463–83. **277 - 1**
- Bellingham K, Gillies P. *Journal of Epidemiology and Community Health* 1993;**47**(2):134–8. **649 - 1**
- Berman BA, Gritz ER, Braxton-Owens H, Nisenbaum R. *Journal of Cancer Education* 1995;**10**(2):91–101. **72 - 1**
- Berrier J, Sperling R, Preisinger J, Evans V, Mason J, Walther V. *AIDS Education and Prevention* 1991;**3**:100–17. **650 - 1**
- Bhave G, Lindan CP, Hudes ES, Desai S, Wagle U, Tripathi SP, Mandel JS. *AIDS* 1995;**9**(Suppl 1):S21–30. **791 - 1**
- Bian JY, Zhang BX, Rong WS. *Advances in Dental Research* 1995;**9**(2):130–3. **97 - 1**
- Biger C, Epstein LM, Hagoel L, Tamir A, Robinson E. *European Journal of Cancer Prevention* 1994;**3**:305–12. **171 - 1**

- Binstock MA, Wolde-Tsadiq G. *Journal of Reproductive Medicine* 1995;**40**(7):507-12. **83 - 1**
- Birkel RC, Golaszewski T, Koman JJ, Singh BK, Catan V, Souply K. *Health Education Quarterly* 1993;**20**:523-38. **177 - 1**
- Bjorge T, Gunbjorud AB, Haugen OA, Skare GB, Trope C, Thoresen SO. *Cancer Causes and Control* 1995;**6**(6):477-84. **61 - 1**
- Black ME, Ploeg J, Walter SD, Hutchinson BG, Scott EA, Chambers LW. *American Journal of Public Health* 1993;**83**(12):1751-3. **633 - 1**
- Black MM, Nair P, Kight C, Wachtel R, Roby P, Schuler M. *Pediatrics* 1994;**94**(4 Pt 1):440-8. **733 - 1**
- Boehm S, Schlenk EA, Raleigh E, Ronis D. *Clinical Nursing Research* 1993;**2**(3):327-44. **104 - 1**
- Boudreau F, Godin G, Pineau R, Bradet R. *Journal of Occupational and Environmental Medicine* 1995;**37**(9):1145-50. **89 - 1**
- Boulet LP, Boutin H, Cote J, Leblanc P, Laviolette M. *Journal of Asthma* 1995;**32**(3):199-206. **82 - 1**
- Bowman J, Sanson-Fisher R, Boyle C, Pope S, Redman S. *Journal of Medical Screening* 1995;**2**(4):211-18. **39 - 1**
- Brennan PF, Moore SM, Smyth KA. *Nursing Research* 1995;**44**(3):166-72. **493 - 1**
- Brenner H, Fleischle B. *Preventive Medicine* 1994;**23**:230-4. **235 - 1**
- Brown LK, Barone VJ, Fritz GK, Cebollero P, Nassau JH. *Health Education Quarterly* 1991;**18**(2):195-206. **467 - 1**
- Brownson RC, Smith CA, Pratt M, Mack NE, Jackson-Thompson J, Dean CG, Dabney S, Wilkerson JC. *American Journal of Public Health* 1996;**86**(2):206-13. **462 - 1**
- Bruce N, Griffioen A. *Social Science and Medicine* 1995;**40**(8):1109-16. **623 - 1**
- Brug J, Steenhuis I, Assema P, de Vreis H. *Preventive Medicine* 1996;**25**:236-42. **210 - 1**
- Buischi YA, Axelsson P, Oliveira LB, Mayer MP, Gjermo P. *Community Dentistry and Oral Epidemiology* 1994;**22**(1):41-6. **562 - 1**
- Bullock LF, Wells JE, Duff GB, Hornblow AR. *New Zealand Medical Journal* 1995;**108**(1012):476-8. **196 - 1**
- Burton LC, Paglia MJ, German PS, Shapiro S, Damiano AM. *Preventive Medicine* 1995;**24**(5):492-7. **64 - 1**
- Butler GS, Hurley CA, Buchanan KL, Smith-VanHorne J. *Patient Education and Counseling* 1996;**29**:189-97. **490 - 1**
- Butow PN, Dunn SM, Tattersall MH, Jones QJ. *Annals of Oncology* 1994;**5**(3):199-204. **700 - 1**
- Caceres CF, Rosasco AM, Mandel JS, Hearst N. *Journal of Adolescent Health* 1994;**15**(7):582-91. **107 - 1**
- Calfas K, Long B, Sallis J, Wooten W, Pratt M, Patrick K. *Preventive Medicine* 1996;**25**:225-33. **209 - 1**
- Calle E, Miracle-Mcmahill H, Moss R, Heath C. *Am J Preventive Medicine* 1994;**10**(6):361-6. **353 - 1**
- Calsyn DA, Saxon AJ, Freeman G, Jr, Whittaker S. *American Journal of Public Health* 1992;**82**(4):573-5. **736 - 1**
- Campbell JR, Szilagyi PG, Rodewald LE, Doane C, Roghmann KJ. *Clinical Pediatrics* 1994;May:268-72. **167 - 1**
- Campion P, Owen L, McNeill A, McGuire C. *Addiction* 1994;**89**(10):1245-54. **514 - 1**
- Cardenas M, Simons-Morton B. *Patient Education and Counseling* 1993;**21**:117-23. **324 - 1**
- Cargill JM. *Journal of Advanced Nursing* 1992;**17**(4):422-6. **702 - 1**
- Carleton RA, Lasater TM, Assaf AR, Feldman HA, McKinlay S. *American Journal of Public Health* 1995;**85**(6):777-85. **440 - 1**
- Champion V, Huster G. *Journal of Behavioral Medicine* 1995;**18**(2):169-87. **114 - 1**
- Champion VL. *Oncology Nursing Forum* 1995;**22**(5):819-24. **690 - 1**
- Cheadle A, Psaty B, Diehr P, Koepsell T, Wagner E, Curry S, Kristal A. *Preventive Medicine* 1996;**24**:71-9. **215 - 1**
- Chen CH. *Maternal Child Nursing Journal* 1993;**21**(3):82-90. **403 - 1**
- Cherkin DC, Deyo RA, Street JH, Hunt M, Barlow W. *Spine* 1996;**21**(3):345-55. **63 - 1**
- Chirwa B, Briega W, Ramakrishna J. *Patient Education and Counseling* 1988;**11**:203-13. **335 - 1**
- Chlebowski RT, Blackburn GL, Buzzard IM, Rose DP, Martino S, Khandekar JD, York RM, Jeffery RW, Elashoff RM, Wynder EL. *Journal of Clinical Oncology* 1993;**11**(11):2072-80. **541 - 1**
- Christensen B. *Family Medicine* 1995;**27**(8):531-4. **56 - 1**
- Christianson JB, Lurie N, Finch M, Moscovice S, Hartley D. *American Journal of Public Health* 1992;**82**(6):790-6. **554 - 1**
- Clark N, Feldman C, Evans D, Duzey O, Levison M, Wasilewski Y, Kaplan D, Rips J, Mellins R. *Patient Education and Counseling* 1986;**8**:27-38. **291 - 1**
- Clarke JR, Bradnock G, Hamburger R. *Community Dental Health* 1992;**9**(2):181-5. **563 - 1**
- Clayton R, Cattarello A, Johnstone B. *Preventive Medicine* 1996;**25**:307-18. **212 - 1**
- Clifford PA, Tan S-Y, Gorsuch RL. *Journal of Behavioural Medicine* 1991;**14**:303-23. **416 - 1**
- Clover K, Redman S, Forbes J, Sanson-Fisher R, Callaghan T. *Preventive Medicine* 1996;**25**:126-34. **205 - 1**
- Clover K, Redman S, Forbes J, Sanson-Fisher R, Callaghan T. *Preventive Medicine* 1996;**25**:126-34. **205 - 2**
- Colon HM, Robles RR, Freeman D, Matos T. *Puerto Rico Health Sciences Journal* 1993;**12**(1):27-34. **407 - 1**



- Colon HM, Sahai H, Robles RR, Matos TD. *AIDS Education and Prevention* 1995;**7**(3):195–209. **770 - 1**
- Conget I, Jansa M, Vidal M, Vidal J, Manzanares JM, Gomis R. *Diabetes Research and Clinical Practice* 1995;**27**(3):189–92. **74 - 1**
- Cook B, Noteloviz M, Rector C, Krischer J. *Patient Education and Counseling* 1991;**17**:135–45. **348 - 1**
- Crockett S, Mullis R, Perry C, Luepker R. *Preventive Medicine* 1989;**18**:475–91. **253 - 1**
- Curry SJ, Taplin SH, Anderman C, Barlow WE, McBride C. *Preventive Medicine* 1993;**22**(3):350–60. **275 - 1**
- D'Souza W, Burgess C, Ayson M, Crane J, Pearce N, Beasley R. *Journal of Allergy and Clinical Immunology* 1996;**97**(5):1085–92. **676 - 1**
- Dale J, Lang H, Roberts JA, Green J, Glucksman E. *British Medical Journal* 1996;**312**(7042):1340–4. **769 - 1**
- Davies-Adetugbo AA. *Journal of Diarrhoeal Diseases Research* 1996;**14**(1):5–11. **3 - 1**
- Davis SW, Cummings KM, Rimer BK, Sciandra R, Stone JC. *Health Education Quarterly* 1992;**19**(4):495–504. **459 - 1**
- de Weerd I, Visser A, Kok G, van der Veen E. *Patient Education and Counseling* 1989;**14**:191–215. **313 - 1**
- DeBusk RF, Miller NH, Superko HR, Dennis CA, Thomas RJ, Lew HT, Berger WE, 3rd, Heller RS, Rompf J, Gee D, Kraemer HC, Bandura A, Ghandour G, Clark M, Shah RV, Fisher L, Barr-Taylor C. *Annals of Internal Medicine* 1994;**120**(9):721–9. **707 - 1**
- Della Valle CJ, Levitz CL, Bora FW, Jr. *American Journal of Orthopedics* 1995;**24**(6):483–7. **75 - 1**
- Delp C, Jones J. *Academic Emergency Medicine* 1996;**3**(3):264–70. **150 - 1**
- Deren S, Davis WR, Beardsley M, Tortu S, Clatts M. *AIDS Education and Prevention* 1995;**7**(5):379–90. **165 - 1**
- Detry JM, Block P, De Backer G, Degaute JP. *European Journal of Clinical Pharmacology* 1995;**47**(6):477–81. **57 - 1**
- Devine CM, Olson CM, Frongillo EA, Jr. *Journal of School Health* 1992;**62**(8):381–5. **409 - 1**
- DiClemente RJ, Wingood GM. *JAMA* 1995;**274**(16):1271–6. **799 - 1**
- DiScenza S, Nies M, Jordan C. *Public Health Nursing* 1996;**13**(3):209–16. **91 - 1**
- Dobs AS, Masters RB, Rajaram L, Stillman FA, Wilder LB, Margolis S, Becker DM. *Patient Education and Counseling* 1994;**24**(2):157–64. **489 - 1**
- Doi SC, DiLorenzo TM. *Journal of Substance Abuse* 1993;**5**(1):73–8. **405 - 1**
- Donnelly BW, Davis BJ. *Child and Adolescent Social Work Journal* 1994;**11**(6):493–506. **486 - 1**
- Doren M, Reuther G, Minne HW, Schneider HP. *American Journal of Obstetrics and Gynecology* 1995;**173**(5):1446–51. **698 - 1**
- Drossaert CHC, Boer H, Seydel ER. *Patient Education and Counseling* 1996;**28**:121–131. **507 - 1**
- Dubois-Arber F, Masur JB, Hausser D, Zimmermann E, Paccaud F. *Social Science and Medicine* 1993;**37**(12):1539–44. **619 - 1**
- Dungy CI, Christensen-Szalanski J, Losch M, Russell D. *Pediatrics* 1992;**90**(2 Pt 1):233–7. **768 - 1**
- Edmunds M, Conner H, Jones C, Gorayeb R, Waranch H. *Preventive Medicine* 1991;**20**:774–84. **270 - 1**
- Edmundson E, Parcel G, Feldman H, Elder J, Perry C, Johnson C, Williston B, Stone E, Yang M, Lytle L, Webber L. *Preventive Medicine* 1996;**25**:442–54. **220 - 1**
- Eisen M, Zellman GL, McAlister AL. *Health Education Quarterly* 1992;**19**(2):249–62. **373 - 1**
- El-Chaar GM, Mardy G, Wehlou K, Rubin LG. *Pediatric Infectious Disease Journal* 1996;**15**(1):18–22. **6 - 1**
- El-Chaar GM, Mardy G, Wehlou K, Rubin LG. *Pediatric Infectious Disease Journal* 1996;**15**(1):18–22. **6 - 2**
- El-Chaar GM, Mardy G, Wehlou K, Rubin LG. *Pediatric Infectious Disease Journal* 1996;**15**(1):18–22. **6 - 3**
- Elder AD, Zultowsky D. *Hospital and Community Psychiatry* 1991;**42**(3):302–4. **361 - 1**
- Elder JP, McGraw SA, Rodrigues A, Lasater TM, Ferreira A, Kendall L, Peteron G, Carleton R. *Preventive Medicine* 1987;**16**:221–34. **367 - 1**
- Elder JP, McGraw SA, Rodrigues A, Lasater TM, Ferreira A, Kendall L, Peteron G, Carleton R. *Preventive Medicine* 1987;**16**:221–34. **367 - 2**
- Elder JP, Wildey M, de Moor C, Sallis JF, Jr, Eckhardt L, Edwards C, Erickson A, Golbeck A, Hovell M, Johnston D, Levitz MD, Molgaard L, Young R, Vito D, Woodruff SI. *American Journal of Public Health* 1993;**83**(9):1239–44. **640 - 1**
- Elk R, Schmitz J, Manfredi L, Rhoades H, Andres R, Grabowski J. *Addictive Behaviors* 1994;**19**(6):697–702. **71 - 1**
- Ellerbeck E, Khallaf N, el Ansary KS, Moursi S, Black R. *Journal of Tropical Pediatrics* 1995;**41**(2):103–8. **51 - 1**
- Esposito L. *Journal of Advanced Nursing* 1995;**21**(5):935–43. **693 - 1**
- Evers S, Bass M, Donner A, McWhinney I. *Preventive Medicine* 1987;**16**:213–20. **356 - 1**
- Fabacher D, Josephson K, Pietruszka F, Linderborn K, Morley JE, Rubenstein LZ. *Journal of the American Geriatrics Society* 1994;**42**(6):630–8. **333 - 1**
- Faivre J, Arveux P, Milan C, Durand G, Lamour J, Bedenne L. *European Journal of Cancer Prevention* 1991;**1**(1):49–55. **285 - 1**
- Fardy PS, White RE, Clark LT, Amodio G, Hurster MH, McDermott KJ, Magel JR. *Journal of Cardiopulmonary Rehabilitation* 1995;**15**(1):65–72. **79 - 1**
- Ferguson KA, Ono T, Lowe AA, Keenan SP, Fleetham JA. *Chest* 1996;**109**:1269–75. **612 - 1**

- Ferris DG, Golden NH, Petry LJ, Litaker MS, Nackenson M, Woodward LD. *Journal of Family Practice* 1996;**42**(1):43–8. **85 - 1**
- Ferson MJ, Fitzsimmons G, Christie D, Woollett H. *Public Health* 1995;**109**:25–109. **572 - 1**
- Finney JW, Miller KM, Adler SP. *Journal of Applied Behaviour Analysis* 1993;**26**:471–2. **163 - 1**
- Fitzgerald ST, Gibbens S, Agnew J. *American Journal of Preventive Medicine* 1991;**7**(6):335–40. **381 - 1**
- Flay B, Miller T, Hedeker D, Siddiqui O, Britton C, Brannon B, Johnson C, Hansen W, Sussman S, Dent C. *Preventive Medicine* 1995;**24**:29–40. **217 - 1**
- Fletcher SW, Harris RP, Gonzalez JJ, Degnan D, Lannin DR, Strecher VJ, Pilgrim C, Quade D, Earp JA, Clark RL. *Journal of the National Cancer Institute* 1993;**85**(2):112–20. **686 - 1**
- Flowers JV, Booraem C, Miller TE, Iverson AE, Copeland J, Furtado K. *AIDS Education and Prevention* 1991;**3**:189–96. **654 - 1**
- Flynn BS, Worden JK, Secker-Walker RH, Badger GJ, Geller BM, Costanza MC. *American Journal of Public Health* 1992;**82**(6):827–34. **792 - 1**
- Foley EC, D'Amico F, Merenstein JH. *Journal of the American Board of Family Practice* 1995;**8**(6):452–6. **34 - 1**
- Forsyth MC, Fulton DL, Lane DS, Burg MA, Krishna M. *Patient Education and Counseling* 1992;**19**(3):241–50. **99 - 1**
- Fortmann SP, Taylor CB, Flora JA, Winkleby MA. *American Journal of Epidemiology* 1993;**137**(10):1039–55. **631 - 1**
- Fox LJ, Bailey PE, Clarke-Martinez KL, Coello M, Ordonez FN, Barahona F. *AIDS Education and Prevention* 1993;**5**(1):1–10. **723 - 1**
- Freedman JD, Mitchell CK. *Journal of General Internal Medicine* 1994;**9**(8):462–4. **412 - 1**
- Friedman RH, Kazis LE, Jette A, Smith MB, Stollerman J, Torgerson J, Carey K. *American Journal of Hypertension* 1996;**9**(4 Pt 1):285–92. **53 - 1**
- Fries JF, Bloch DA, Harrington H, Richardson N, Beck R. *American Journal of Medicine* 1993;**94**(5):455–62. **110 - 1**
- Gans KM, Lapane KL, Lasater TM, Carleton RA. *American Journal of Preventive Medicine* 1994;**10**(5):275–82. **372 - 1**
- Garcia R, Suarez R. *Patient Education and Counseling* 1996;**29**:87–97. **521 - 1**
- Gardiner JC, Mullan PB, Rosenman KD, Zhu Z, Swanson GM. *Journal of Cancer Education* 1995;**10**(3):155–62. **44 - 1**
- Gariti P, Alterman AI, Holub-Beyer E, Volpicelli JR, Prentice N, O'Brien CP. *Journal of Substance Abuse Treatment* 1995;**12**(3):207–12. **202 - 1**
- Garr DR, Ornstein SM, Jenkins RG, Zemp LD. *American Journal of Preventive Medicine* 1993;**9**(1):55–61. **466 - 1**
- Garton MJ, Torgerson DJ, Donaldson C, Russell IT, Reid DM. *British Medical Journal* 1992;**305**(6845):82–4. **747 - 1**
- Gastrin G. *Sozial- und Preventivmedizin* 1993;**38**(5):280–7. **421 - 1**
- Gibbins RL, Riley M, Brimble P. *British Medical Journal* 1993;**306**(6893):1652–6. **798 - 1**
- Gil KM, Wilson JJ, Edens JL, Webster DA, Abrams MA, Orringer E, Grant M, Clark WC, Janal MN. *Health Psychology* 1996;**15**(1):3–10. **20 - 1**
- Gilbert J, Wilson D, Singer J, Lindsey E, Willms D, Best J, Taylor D. *Am J Preventive Medicine* 1992;**8**:91–5. **350 - 1**
- Gilles MT, Crewe S, Granites IN, Coppola A. *Australian Journal of Public Health* 1995;**19**:477–81. **192 - 1**
- Gillies J, Barry D, Crane J, Jones D, MacLennan L, Pearce N, Reid J, Toop L. *New Zealand Medical Journal* 1996;**109**(1015):30–3. **42 - 1**
- Girgis A, Sanson-Fisher RW, Tripodi DA, Golding T. *Health Education Quarterly* 1993;**20**(2):275–87. **136 - 1**
- Glasgow R, Toobert D, Hampson S, Brown J, Lewinsohn P, Donnelly J. *Patient Education and Counseling* 1992;**19**:61–74. **328 - 1**
- Glasgow RE, Hollis JF, Ary DV, Boles SM. *Addictive Behaviors* 1993;**18**(4):455–64. **102 - 1**
- Glasgow RE, Terborg JR, Hollis JF, Severson HH, Boles SM. *American Journal of Public Health* 1995;**85**(2):209–16. **442 - 1**
- Goldberg D, Hoffman A, Farinha M, Marder D, Tinson-Mitchem L, Burton D, Smith E. *American Journal of Preventive Medicine* 1994;**10**:267–74. **345 - 1**
- Goldberg L, Elliot DL, Clarke GN, MacKinnon DP, Zoref L, Moe E, Green C, Wolf SL. *Archives of Pediatrics and Adolescent Medicine* 1996;**150**(7):713–21. **18 - 1**
- Gomel M, Oldenburg B, Simpson JM, Owen N. *American Journal of Public Health* 1993;**83**(9):1231–8. **644 - 1**
- Gordon CM, Carey MP. *Health Psychology* 1996;**15**(1):56–60. **24 - 1**
- Graham J, Johnson C, Hansen W, Flay B, Gee M. *Preventive Medicine* 1990;**19**:305–13. **247 - 1**
- Greenberg RA, Strecher VJ, Bauman KE, Boat BW, Fowler MG, Keyes LL, Denny FW, Chapman RS, Stedman HC, LaVange LM, Glover LH, Haley NJ, Loda FA. *Journal of Behavioral Medicine* 1994;**17**(3):273–90. **157 - 1**
- Griffith CH, 3rd, Wilson JF, Emmett KR, Ramsbottom-Lucier M, Rich EC. *Archives of Family Medicine* 1995;**4**(9):780–4. **67 - 1**
- Gritz ER, Carr CR, Rapkin D, Abemayor E, Chang LJ, Wong WK, Belin TR, Calcaterra T, Robbins KT, Chonkich G, Beumer J, Ward PH. *Cancer Epidemiology, Biomarkers and Prevention* 1993;**2**(3):261–70. **140 - 1**
- Grosskurth H, Mosha F, Todd J, Mwijarubi E, Klokke A, Senkoro K, Mayaud P, Changalucha J, Nicoll A, ka-Gina G, Newell J, Mugeye K, Mabeg D, Hayes R. *Lancet* 1995;**346**(8974):530–6. **732 - 1**
- Guimon J. *Patient Education and Counseling* 1995;**26**:189–93. **825 - 1**

- Guptill KS, Esrey SA, Oni GA, Brown KH. *Social Science and Medicine* 1993;**36**(5):665–72. **621 - 1**
- Hagopian GA. *Oncology Nursing Forum* 1996;**23**(4):697–700. **23 - 1**
- Hanlon P, McEwen J, Carey L, Gilmour H, Tannahill C, Tannahill A, Kelly M. *British Medical Journal* 1995;**311**(7020):1609–13. **755 - 1**
- Hanna KM. IMAGE. *Journal of Nursing Scholarship*;25:285–90. **181 - 1**
- Hansen BL. *Patient Education and Counseling* 1995;**25**:137–142. **520 - 1**
- Hansen BWL. *Patient Education and Counseling* 1990;**16**:147–50. **302 - 1**
- Hansen W, Graham J. *Preventive Medicine* 1991;**20**:414–30. **226 - 1**
- Hansen WB, Anderson Johnson C, Flay B, Graham J, Sobel J. *Preventive Medicine* 1988;**17**:135–54. **311 - 1**
- Harrell JS, McMurray RG, Bangdiwala SI, Frauman AC, Gansky SA, Bradley CB. *Journal of Pediatrics* 1996;**128**(6):797–805. **679 - 1**
- Hartley BM, O'Connor ME. *Archives of Pediatrics and Adolescent Medicine* 1996;**150**(8):868–71. **26 - 1**
- Hawe P, Higgins G. *Patient Education and Counseling* 1990;**16**:151–60. **303 - 1**
- Heath GW, Fuchs R, Croft JB, Temple SP, Wheeler FC. *American Journal of Preventive Medicine* 1995;**11**(3):190–6. **45 - 1**
- Hebert J, Kristeller J, Ockene J, Landon J, Luippold R, Goldberg R, Kalan K. *Preventive Medicine* 1992;**21**:557–73. **256 - 1**
- Heirich MA, Foote A, Erfurt JC, Konopka B. *Journal of Occupational Medicine* 1993;**35**(5):510–7. **390 - 1**
- Henderson MM, Kushi LH, Thompson DJ, Gorbach SL, Clifford CK, Insull W, Moskowitz M, Thompson RS. *Preventive Medicine* 1990;**19**:115–33. **263 - 1**
- Hill D, White V, Marks R, Borland R. *European Journal of Cancer Prevention* 1993;**2**(6):447–56. **180 - 1**
- Hingson R, McGovern T, Howland J, Heeren T, Winter M, Zakocs R. *American Journal of Public Health* 1996;**86**(6):791–7. **461 - 1**
- Ho EE, Atwood JR, Benedict J, Ritenbaugh C, Sheehan ET, Abrams C, Alberts D, Meyskens FL, Jr. *Preventive Medicine* 1991;**20**(2):213–25. **269 - 1**
- Hoare T, Thomas C, Biggs A, Booth M, Bradley S, Friedman E. *Journal of Public Health Medicine* 1994;**16**(2):179–85. **695 - 1**
- Hobfoll SE, Jackson AP, Lavin J, Britton PJ, Shepherd JB. *Health Psychology* 1994;**13**(5):397–403. **742 - 1**
- Hoffman JA, Caudill BD, Koman JJ, 3rd, Luckey JW, Flynn PM, Mayo DW. *Journal of Substance Abuse Treatment* 1996;**13**(1):3–11. **4 - 1**
- Hoffmeister H, Mensink GBM, Stolzenberg H, Hoeltz J, Kreuter H, Laaser U, Nussel E, Hullemann K-D, Troschke J. *Preventive Medicine* 1996;**25**:135–45. **206 - 1**
- Holley JL, Nespor S, Rault R. *American Journal of Kidney Diseases* 1993;**22**(3):413–18. **111 - 1**
- Hollis J, Lichtenstein E, Mount K, Vogt T, Stevens V. *Preventive Medicine* 1991;**20**:497–507. **254 - 1**
- Horowitz LG. *Clinical Preventive Dentistry* 1992;**14**(6):11–16. **457 - 1**
- Hughes BR, Altman DG, Newton JA. *British Journal of Dermatology* 1993;**128**(4):412–17. **705 - 1**
- Hughes J, Wadland W, Fenwick J, Lewis J, Bickel W. *Preventive Medicine* 1991;**20**:486–96. **229 - 1**
- Hurley SF, Huggins RM, Jolley DJ, Reading D. *American Journal of Public Health* 1994;**84**(10):1655–8. **727 - 1**
- Hurt RD, Offord KP, Lauger GG, Marusic Z, Fagerstrom KO, Enright PL, Scanlon PD. *Addiction* 1995;**90**(3):407–13. **574 - 1**
- Huss K, Salerno M, Huss RW. *Research in Nursing and Health* 1991;**14**(4):259–67. **327 - 1**
- Huws R. *International Journal of Social Psychiatry* 1992;**38**(4):304–8. **349 - 1**
- Ingold FR, Toussirt M. *Bulletin on Narcotics* 1993;**45**(1):117–34. **142 - 1**
- Ives D, Kuller L, Schultz R, Traven N, Lave J. *Preventive Medicine* 1992;**21**:582–91. **257 - 1**
- Ives DG, Lave JR, Traven ND, Kuller LH. *Preventive Medicine* 1994;**23**(2):134–41. **234 - 1**
- Jemmott JB, 3rd, Jemmott LS, Fong GT. *American Journal of Public Health* 1992;**82**(3):372–7. **749 - 1**
- Jemmott LS, Jemmott JB, 3rd. *Nursing Research* 1992;**41**(5):273–9. **364 - 1**
- Jin BW, Kim SC, Mori T, Shimao T. *Tubercle and Lung Disease* 1993;**74**(4):267–72. **452 - 1**
- Joffe MD, Luberti A. *Pediatric Emergency Care* 1994;**10**(6):317–19. **331 - 1**
- Jones SL, Jones PK, Katz J. *Social Science and Medicine* 1991;**32**(10):1183–9. **622 - 1**
- Jorenby DE, Smith SS, Fiore MC, Hurt RD, Offord KP, Croghan IT, Hays JT, Lewis SF, Baker TB. *JAMA* 1995;**274**(17):1347–52. **648 - 1**
- Joseph AM. *Journal of Substance Abuse Treatment* 1993;**10**(2):147–52. **423 - 1**
- Joseph AM, Nichol KL, Anderson H. *Addictive Behaviors* 1993;**18**(6):635–44. **550 - 1**
- Kalichman SC, Kelly JA, Hunter TL, Murphy DA, Tyler R. *Journal of Consulting and Clinical Psychology* 1993;**61**(2):291–5. **762 - 1**
- Kamenga M, Ryder RW, Jingu M, Mbuyi N, Mbu L, Behets E, Brown C, Heyward WL. *AIDS* 1991;**5**(1):61–7. **688 - 1**

- Kamolratanakul P, Ungtavorn P, Israsena S, Sakulramrung R. *Public Health* 1994;**108**(1):49–53. **573 - 1**
- Karofsky PS, Rice RL, Hoornstra LL, Slater CJ, Kessinich CA, Goode JR. *Clinical Pediatrics* 1991;**30**(5):290–4. **470 - 1**
- Kasemsarn P, Ngarmpiyasakul C, Phongpanich S, Pulkasisri N. *Journal of the Medical Association of Thailand* 1995;**78**(7):362–8. **132 - 1**
- Kash KM, Holland JC, Osborne MP, Miller DG. *Monographs – National Cancer Institute* 1995;(17):73–9. **46 - 1**
- Katon W, Von Korff M, Lin E, Walker E, Simon GE, Bush T, Robinson P, Russo J. *JAMA* 1995;**273**(13):1026–31. **794 - 1**
- Kaufman JS, Jason LA, Sawlski LM, Halpert JA. *Journal of Drug Education* 1994;**24**(2):95–108. **184 - 1**
- Keenan J, Kastner T, Nathanson R, Richardson N, Hinton J, Cress DA. *Mental Retardation* 1992;**30**(6):355–61. **294 - 1**
- Kegeles SM, Hays RB, Coates TJ. *American Journal of Public Health* 1996;**86**(8 Pt 1):1129–36. **463 - 1**
- Kelly JA, Murphy DA, Washington CD, Wilson TS, Koob JJ, Davis DR, Ledezma G, Davantes B. *American Journal of Public Health* 1994;**84**(12):1918–22. **776 - 1**
- Kelly JA, St Lawrence JS, Diaz YE, Stevenson LY, Hauth AC, Brasfield TL, Kalichman SC, Smith JE, Andrew ME. *American Journal of Public Health* 1991;**81**(2):168–71. **665 - 1**
- Kelly JA, St Lawrence JS, Stevenson LY, Hauth AC, Kalichman SC, Diaz YE, Brasfield TL, Koob JJ, Morgan MG. *American Journal of Public Health* 1992;**82**(11):1483–9. **647 - 1**
- Kemp R, Hayward P, Applewhaite G, Everitt B, David A. *British Medical Journal* 1996;**312**(7027):345–9. **427 - 1**
- Kendall C, Hailey BJ. *Behavioral Medicine* 1993;**19**(1):29–34. **101 - 1**
- Kernohan EE. *British Journal of Cancer* 1996;**74**(Suppl 29):S42–6. **746 - 1**
- Kiefe CI, McKay SV, Halevy A, Brody BA. *Archives of Internal Medicine* 1994;**154**(11):1217–24. **509 - 1**
- Kimberlin CL, Berardo DH, Pendergast JF, McKenzie LC. *Medical Care* 1993;**31**(5):451–68. **483 - 1**
- King A, Carl F, Birkel L, Haskell W. *Preventive Medicine* 1988;**17**:357–65. **355 - 1**
- King J, Fairbrother G, Thompson C, Morris DL. *Australian and New Zealand Journal of Surgery* 1992;**62**(9):714–19. **683 - 1**
- King R, Estey J, Allen S, Kegeles S, Wolf W, Valentine C, Serufilira A. *AIDS* 1995;**9**(Suppl 1):S45–51. **655 - 1**
- Kipke MD, Boyer C, Hein K. *Journal of Adolescent Health* 1993;**14**(7):533–9. **492 - 1**
- Kirby D, Barth RP, Leland N, Fetro JV. *Family Planning Perspectives* 1991;**23**(6):253–63. **478 - 1**
- Kirby D, Waszak C, Ziegler J. *Family Planning Perspectives* 1991;**23**(1):6–16. **822 - 1**
- Kirkman MS, Weinberger M, Landsman PB, Samsa GP, Shortliffe EA, Simel DL, Feussner JR. *Diabetes Care* 1994;**17**(8):840–6. **139 - 1**
- Kissinger P, Clark R, Rice J, Kutzen H, Morse A, Brandon W. *Southern Medical Journal* 1995;**88**(11):1121–5. **537 - 1**
- Klepp KI, Ndeki SS, Seha AM, Hannan P, Lyimo BA, Msuya MH, Irema MN, Schreiner A. *AIDS* 1994;**8**(8):1157–62. **515 - 1**
- Klepp KI, Tell GS, Vellar OD. *Preventive Medicine* 1993;**22**(4):453–62. **276 - 1**
- Knapp LG. *Journal of Pediatric Psychology* 1991;**16**(6):67–86. **371 - 1**
- Knutsen SF, Knutsen R. *Preventive Medicine* 1991;**20**(2):197–212. **268 - 1**
- Koenig MA, Rob U, Khan MA, Chakraborty J, Fauveau V. *Studies in Family Planning* 1992;**23**(6 Pt 1):352–64. **392 - 1**
- Kohatsu N, Cramer E, Bohnstedt M. *American Journal of Preventive Medicine* 1994;**10**(6):348–52. **401 - 1**
- Kohn MR, Arden MR, Vasilakis J, Shenker IR. *Archives of Pediatrics and Adolescent Medicine* 1996 Jul;**150**(7):727–9. **586 - 1**
- Korhonen H, Niemensivu H, Piha T, Koskela K, Wiio J, Johnson C, Puska P. *Preventive Medicine* 1992;**21**:74–87. **280 - 1**
- Kurtz ME, Kurtz JC, Given B, Given CC. *Health Care for Women International* 1994;**15**(1):31–42. **126 - 1**
- Kuthy S, Grap MJ, Penn L, Henderson V. *Journal of Neuroscience Nursing* 1995;**27**(5):273–7. **84 - 1**
- Lahdensuo A, Haahtela T, Herrala J, Kava T, Kiviranta K, Kuusisto P, Peramaki E, Poussa T, Saarelainen S, Svahn T. *British Medical Journal* 1996;**312**(7033):748–52. **781 - 1**
- Lancaster G, Elton P. *Journal of Epidemiology and Community Health* 1992;**46**(5):523–7. **543 - 1**
- Lando HA, Pirie PL, McGovern PG, Pechacek TF, Swim J, Loken B. *Addictive Behaviors* 1991;**16**(5):183–93. **473 - 1**
- Langer LM, Zimmerman RS, Hendershot EF, Singh M. *AIDS Education and Prevention* 1992;**4**(4):295–307. **662 - 1**
- Lave JR, Ives DG, Traven ND, Kuller LH. *American Journal of Preventive Medicine* 1995;**11**(1):46–53. **47 - 1**
- Lee CY. *Yonsei Medical Journal* 1991;**32**(2):131–8. **316 - 1**
- Leigh JP, Richardson N, Beck R, Kerr C, Harrington H, Parcell CL, Fries JF. *Archives of Internal Medicine* 1992;**152**(6):1201–6. **813 - 1**
- Lerman C, Hanjani P, Caputo C, Miller S, Delmoor E, Nolte S, Engstrom P. *Journal of Clinical Oncology* 1992;**10**(2):330–3. **380 - 1**
- Lerman C, Ross E, Boyce A, Gorchov PM, McLaughlin R, Rimer B, Engstrom P. *American Journal of Public Health* 1992;**82**(5):729–30. **757 - 1**

- Leslie M, Schuster P. *Patient Education and Counseling* 1993;**21**:51–60. **375 - 1**
- Levy SR, Perhats C, Weeks K, Handler AS, Zhu C, Flay BR. *Journal of School Health* 1995;**65**(4):145–51. **424 - 1**
- Lewis B, Lynch W. *Preventive Medicine* 1993;**22**:110–21. **244 - 1**
- Li J, Taylor B. *British Medical Journal* 1991;**303**(6809):1035–8. **775 - 1**
- Lichtenstein E, Hollis J. *Journal of Family Practice* 1992;**34**(6):739–44. **376 - 1**
- Lierman LM, Young HM, Powell-Cope G, Georgiadou F, Benoliel JQ. *Nursing Research* 1994;**43**(3):158–63. **307 - 1**
- Lindholm LH, Ekblom T, Dash C, Isacson A, Schersten B. *Journal of Internal Medicine* 1996;**240**(1):13–22. **1 - 1**
- Linkins RW, Dini EF, Watson G, Patriarca PA. *Archives of Pediatrics and Adolescent Medicine* 1994;**148**(9):908–14. **512 - 1**
- Lionis C, Kafatos A, Vlachonikolis J, Vakaki M, Tzortzi M, Petraki A. *Preventive Medicine* 1991;**20**(6):685–99. **261 - 1**
- Lipton HL, Bird JA. *Gerontologist* 1994;**34**(3):307–15. **169 - 1**
- Llewellyn-Thomas HA, McGreal MJ, Thiel EC. *Medical Decision Making* 1995;**15**(1):4–12. **434 - 1**
- Llewellyn-Thomas HA, Thiel EC, Sem FW, Woermke DE. *Patient Education and Counseling* 1995;**25**(2):97–107. **488 - 1**
- Lloyd HM, Paisley CM, Mela DJ. *Journal of the American Dietetic Association* 1995;**95**(3):316–22. **147 - 1**
- Lombard D, Neubauer TE, Canfield D, Winett RA. *Journal of Applied Behavior Analysis* 1991;**24**(4):677–86. **765 - 1**
- Lowe CJ, Raynor DK, Courtney EA, Purvis J, Teale C. *British Medical Journal* 1995;**310**(6989):1229–31. **763 - 1**
- Luepker RV, Murray DM, Jacobs DR, Jr, Mittelmark MB, Bracht N, Carlaw R, Crow R, Elmer P, Finnegan J, Folsom AR, Grimm R, Hannan PJ, Jeffrey R, Lanbo H, Govern P, Mullis R, Perry CL, Pechacek T, Pirie P, Sprafka M, Weisbrod R, Blackburn H. *American Journal of Public Health* 1994;**84**(9):1383–93. **726 - 1**
- Magura S, Kang SY, Shapiro JL. *Journal of Adolescent Health* 1994;**15**(6):457–63. **125 - 1**
- Maheu M, Gevirtz R, Sallis J, Schneider N. *Preventive Medicine* 1989;**18**:867–76. **283 - 1**
- Maiman LA, Hildreth NG, Cox C, Greenland P. *American Journal of Public Health* 1992;**82**(6):804–9. **652 - 1**
- Main DS, Iverson DC, McGloin J, Banspach SW, Collins JL, Rugg DL, Kolbe LJ. *Preventive Medicine* 1994;**23**(4):409–17. **240 - 1**
- Maisiak R, Austin J, Heck L. *Arthritis and Rheumatism* 1996;**39**(8):1391–9. **27 - 1**
- Malenka DJ, Baron JA, Johanson S, Wahrenberger JW, Ross JM. *Journal General Internal Medicine* 1993;**8**:543–8. **176 - 1**
- Malloy TR, Wigton RS, Meeske J, Tape TG. *Journal of the American Geriatrics Society* 1992;**40**(12):1255–60. **332 - 1**
- Malow RM, West JA, Corrigan SA, Pena JM, Cunningham SC. *AIDS Education and Prevention* 1994;**6**(2):113–25. **635 - 1**
- Mann K, Sullivan P. *Patient Education and Counseling* 1987;**19**:53–72. **295 - 1**
- Mansfield CJ, Conroy ME, Emans SJ, Woods ER. *Journal of Adolescent Health* 1993;**14**(2):115–9. **393 - 1**
- Mant D, Fuller A, Northover J, Astrop P, Chivers A, Crockett A, Clements S, Lawrence M. *British Journal of General Practice* 1992;**42**(354):18–20. **752 - 1**
- Marcus A, Wheeler R, Cullen J, Crane L. *Preventive Medicine* 1987;**16**:803–15. **336 - 1**
- Marcus AC, Crane LA, Kaplan CP, Reading AE, Savage E, Gunning J, Bernstein G, Berek JS. *Medical Care* 1992;**30**(3):216–30. **484 - 1**
- Marcus BH, Stanton AL. *Research Quarterly for Exercise and Sport* 1993;**64**(4):447–52. **814 - 1**
- Marin B, Perez-Stable E, Marin G, Hauck W. *American Journal of Preventive Medicine* 1994;**10**(6):340–7. **340 - 1**
- Martinez R, Levine DW, Martin R, Altman DG. *Annals of Emergency Medicine* 1996;**27**(2):216–24. **194 - 1**
- Maurer WJ. *Wisconsin Medical Journal* 1995;**94**(6):305–6. **100 - 1**
- Maxwell AE, Hunt IF, Bush MA. *Diabetes Educator* 1992;**18**(4):303–9. **807 - 1**
- Mayer J, Clapp E, Bartholomew S, Elder J. *American Journal of Preventive Medicine* 1994;**10**(6):353–6. **352 - 1**
- Mayer J, Clapp E, Bartholomew S, Elder J. *American Journal of Preventive Medicine* 1994;**10**(6):353–6. **352 - 2**
- Mayer J, Clapp E, Bartholomew S, Elder J. *American Journal of Preventive Medicine* 1994;**10**(6):353–6. **352 - 3**
- Mayer J, Jermanovich A, Wright B, Elder J, Drew J, Williams S. *Preventive Medicine* 1994;**23**:127–33. **233 - 1**
- Mayer JA, Kossman MK, Miller LC, Crooks CE, Slymen DJ, Lee CD, Jr. *American Journal of Preventive Medicine* 1992;**8**(1):23–9. **383 - 1**
- Mazur DJ, Hickam DH. *European Journal of Surgery* 1996;**162**(2):101–4. **19 - 1**
- McAuley E, Courneya K, Rudolph D, Lox C. *Preventive Medicine* 1994;**23**:498–506. **241 - 1**
- McAvoy BR, Raza R. *British Medical Journal* 1991;**302**(6780):833–6. **728 - 1**
- McCann S, Weinman J. *Patient Education and Counseling* 1996;**27**(3):227–34. **387 - 1**
- McCusker J, Stoddard AM, Zapka JG, Lewis BF. *American Journal of Public Health* 1993;**83**(10):1463–6. **636 - 1**
- McCusker J, Stoddard AM, Zapka JG, Morrison CS, Zorn M, Lewis BF. *American Journal of Public Health* 1992;**82**(4):533–40. **817 - 1**

- McCusker J, Stoddard AM, Zapka JG, Zorn M. *Sexually Transmitted Diseases* 1993;**20**(2):81–8. **684 - 1**
- McFall SL, Michener A, Rubin D, Flay BR, Mermelstein RJ, Burton D, Jelen P, Warnecke RB. *Addictive Behaviors* 1993;**18**(2):151–8. **143 - 1**
- McFarlane WR, Lukens E, Link B, Dushay R, Deakins SA, Newmark M, Dunne EJ, Horen B, Toran J. *Archives of General Psychiatry* 1995;**52**(8):679–87. **96 - 1**
- McIntosh NA, Clark NM, Howatt WF. *Journal of Asthma* 1994;**31**(6):453–62. **118 - 1**
- Meier DE, Fuss BR, Orourke D, Baskin SA, Lewis M, Morrison RS. *Archives of Internal Medicine* 1996;**156**(11):1227–32. **629 - 1**
- Mellanby AR, Phelps FA, Crichton NJ, Tripp JH. *British Medical Journal* 1995;**311**(7002):414–17. **759 - 1**
- Meltzer SB, Hovell MF, Meltzer EO, Atkins CJ, de Peyster A. *Journal of Asthma* 1993;**30**(5):391–400. **179 - 1**
- Mercer BS. *Archives of Neurology* 1996;**53**(9):881–4. **28 - 1**
- Merkel PA, Caputo GC. *Journal of General Internal Medicine* 1994;**9**(12):679–83. **413 - 1**
- Mermelstein RJ, Riesenber LA. *Health Psychology* 1992;**11**(6):371–6. **458 - 1**
- Mesters I, van Nunen M, Crebolder H, Meertens R. *Patient Education and Counseling* 1995;**25**:131–6. **504 - 1**
- Michaud PA, Hausser D. *Health Education Research* 1992;**7**(1):79–86. **555 - 1**
- Miedzybrodzka ZH, Hall MH, Mollison J, Templeton A, Russell IT, Dean JC, Kelly KF, Marteau TM, Haites NE. *British Medical Journal* 1995;**310**(6976):353–7. **743 - 1**
- Miller MF, Wong JG. *American Journal of the Medical Sciences* 1993;**306**(2):98–100. **566 - 1**
- Miller RL. *AIDS Education and Prevention* 1995;**7**(Suppl 5):48–63. **159 - 1**
- Miller WR, Benefield RG, Tonigan JS. *Journal of Consulting and Clinical Psychology* 1993;**61**(3):455–61. **772 - 1**
- Minor MA, Brown JD. *Health Education Quarterly* 1993;**20**(1):83–95. **553 - 1**
- Mittelman MS, Ferris SH, Steinberg G, Shulman E, Mackell TA, Ambinder A, Cohen J. *Gerontologist* 1993;**33**(6):730–40. **494 - 1**
- Mohler PJ. *Family Medicine* 1995;**27**(2):117–21. **52 - 1**
- Montgomery EB, Jr, Lieberman A, Singh G, Fries JF. *American Journal of Medicine* 1994;**97**(5):429–35. **511 - 1**
- Moos MK, Bangdiwala SI, Meibohm AR, Cefalo RC. *American Journal of Perinatology* 1996;**13**(2):103–8. **190 - 1**
- Morgan G, Noll E, Orleans T, Rimer B, Amfoh K, Bonney G. *Preventive Medicine* 1996;**25**:346–54. **213 - 1**
- Morris BA, Trimble NE, Fendley SJ. *Canadian Family Physician* 1994;**40**:1126–31. **149 - 1**
- Moser SE. *Family Practice Research Journal* 1994;**14**:281–8. **172 - 1**
- Mudde A, Vries H, Dolders M. *Preventive Medicine* 1995;**24**:61–70. **218 - 1**
- Muhlhauser I, Richter B, Kraut D, Weske G, Worth H, Berger M. *Journal of Internal Medicine* 1991;**230**(2):157–64. **544 - 1**
- Muller O, Sarangbin S, Ruxrungtham K, Sittitrai W, Phanuphak P. *AIDS Care* 1995;**7**(5):567–72. **161 - 1**
- Murray D, Perry C, Griffin G, Harty K, Jacobs D, Schmid L, Daly K, Pallonen U. *Preventive Medicine* 1992;**21**:449–72. **255 - 1**
- Murray D, Perry C, Griffin G, Harty K, Jacobs D, Schmid L, Daly K, Pallonen U. *Preventive Medicine* 1992;**21**:449–72. **255 - 2**
- Murray DM, Kurth C, Mullis R, Jeffery RW. *Preventive Medicine* 1990;**19**:181–9. **264 - 1**
- Murray DM, Prokhorov AV, Harty KC. *Preventive Medicine* 1994;**23**(1):54–60. **238 - 1**
- Murray MD, Birt JA, Manatunga AK, Darnell JC. *Annals of Pharmacotherapy* 1993;**27**(5):616–21. **105 - 1**
- Myers RE, Balslem AM, Wolf TA, Ross EA, Millner L. *Medical Care* 1993;**31**(6):508–19. **482 - 1**
- Myers RE, Ross EA, Wolf TA, Balslem A, Jepson C, Millner L. *Medical Care* 1991;**29**(10):1039–50. **420 - 1**
- Myles PS, Hendrata M, Layher Y, Williams NJ, Hall JL, Moloney JT, Powell J. *British Journal of Anaesthesia* 1996;**76**(5):694–8. **581 - 1**
- Navarro AM, Senn KL, Kaplan RM, McNicholas L, Campo MC, Roppe B. *Monographs – National Cancer Institute* 1995;(18):137–45. **86 - 1**
- Nelson EW, Van Cleve S, Swartz MK, Kessen W, McCarthy PL. *American Journal of Diseases of Children* 1991;**145**(4):440–4. **714 - 1**
- Newcomb PA, Klein R, Massoth KM. *Journal of Diabetes and its Complications* 1992;**6**(4):211–17. **394 - 1**
- Nilssen O. *Preventive Medicine* 1991;**20**:518–28. **259 - 1**
- Nordblad A, Suominen-Taipale L, Murtomaa H, Vartiainen E, Koskela K. *Community Dental Health* 1995;**12**(4):230–4. **529 - 1**
- Norman P. *Social Science and Medicine* 1993;**37**(1):53–9. **620 - 1**
- Norman P. *British Journal of Clinical Psychology* 1995;**34**:461–70. **426 - 1**
- Norman P, Fitter M. *Family Practice* 1991;**8**:129–32. **589 - 1**
- Nutbeam D, Macaskill P, Smith C, Simpson JM, Catford J. *British Medical Journal* 1993;**306**(6870):102–7. **731 - 1**
- Nyamathi AM, Flaskerud J, Bennett C, Leake B, Lewis C. *AIDS Education and Prevention* 1994;**6**(4):296–309. **646 - 1**
- Nyamathi AM, Leake B, Flaskerud J, Lewis C, Bennett C. *Research in Nursing and Health* 1993;**16**(1):11–21. **454 - 1**

- O'Donnell L, San Doval A, Duran R, O'Donnell CR. *Sexually Transmitted Diseases* 1995;**22**(2):97-103. **696 - 1**
- O'Hara P, Messick BJ, Fichtner RR, Parris D. *Journal of School Health* 1996;**66**(5):176-82. **35 - 1**
- O'Loughlin J, Paradis G, Kishchuk N, Gray-Donald K, Renaud L, Fines P, Barnett T. *Journal of Epidemiology and Community Health* 1995;**49**(5):495-502. **658 - 1**
- O'Loughlin J, Renaud L, Paradis G, Meshfedjian G. *Preventive Medicine* 1996;**25**:660-7. **222 - 1**
- Ockene J, Hymowitz N, Lagus J, Shaten J. *Preventive Medicine* 1991;**20**:518-28. **260 - 1**
- Oeffinger KC, Roaten SP, Hitchcock MA, Oeffinger PK. *Journal of Family Practice* 1992;**35**(3):288-93. **363 - 1**
- Ogututu RO, Oloo AJ, Ekissa WS, Genga IO, Mulaya N, Githure JL. *East African Medical Journal* 1992;**69**(6):298-302. **308 - 1**
- Ohmit SE, Furumoto-Dawson A, Monto AS, Fasano N. *American Journal of Preventive Medicine* 1995;**11**(4):271-6. **146 - 1**
- Ojehagen A, Berglund M. *European Archives of Psychiatry and Clinical Neuroscience* 1992;**242**(2-3):82-4. **402 - 1**
- Oladepo O, Okunade A, Brieger W, Oshiname F, Ajuwon A. *Patient Education and Counseling* 1996;**29**(3):237-45. **323 - 1**
- Oldenburg B, Martin A, Greenwood J, Bernstein L, Allan R. *Journal of Cardiopulmonary Rehabilitation* 1995;**15**(1):39-46. **80 - 1**
- Olsen G, Lacy S, Sprafka J, Arceneaux T, Potts T, Kravat B, Gondek M, Bond G. *Preventive Medicine* 1990;**19**:774-84. **262 - 1**
- Orleans C, Rotberg H, Quade D, Leeds P. *Preventive Medicine* 1990;**19**:198-212. **245 - 1**
- Orleans CT, Schoenbach VJ, Wagner EH, Quade D, Salmon MA, Pearson DC, Fiedler J, Porter CQ, Kaplan BH. *Journal of Consulting and Clinical Psychology* 1991;**59**(3):439-48. **783 - 1**
- Ornstein SM, Garr DR, Jenkins RG, Rust PF, Arnon A. *Journal of Family Practice* 1991;**32**(1):82-90. **395 - 1**
- Osler M, Jespersen NB. *Danish Medical Bulletin* 1993;**40**(4):485-9. **709 - 1**
- Palinkas L, Atkins C, Millers C, Ferreira D. *Preventive Medicine* 1996;**25**:692-701. **223 - 1**
- Pallonen UE, Leskinen L, Prochaska JO, Willey CJ, Kaariainen R, Salonen JT. *Preventive Medicine* 1994;**23**(4):507-14. **242 - 1**
- Palm BT, Kant AC, van den Bosch WJ, Vooijs GP, van Weel C. *British Journal of General Practice* 1993;**43**(377):503-6. **546 - 1**
- Park SI, Saxe JC, Weesner RE. *American Journal of Gastroenterology* 1993;**88**(9):1391-4. **108 - 1**
- Paskett E, White E, Carter W, Chu E. *Preventive Medicine* 1990;**19**:630-41. **248 - 1**
- Paskett ED, Phillips KC, Miller ME. *Obstetrics and Gynecology* 1995;**86**(3):353-9. **508 - 1**
- Pauw J, Ferrie J, Rivera Villegas R, Medrano Martinez J, Gorter A, Egger M. *AIDS* 1996;**10**(5):537-44. **22 - 1**
- Peeters PH, Beckers CG, Hogervorst JM, Collette HJ. *Journal of Epidemiology and Community Health* 1994;**48**(2):175-7. **653 - 1**
- Pendergrast RA, Ashworth CS, DuRant RH, Litaker M. *Pediatrics* 1992;**90**(3):354-8. **773 - 1**
- Pentz M, MacKinnon D, Dwyer J, Wang E, Hansen W, Flay B, Johnson C. *Preventive Medicine* 1989;**18**:304-21. **251 - 1**
- Phatouros CC, Blake MP. *Australasian Radiology* 1995;**39**(2):135-9. **671 - 1**
- Pichert JW, Snyder GM, Kinzer CK, Boswell EJ. *Patient Education and Counseling* 1994;**23**(2):115-24. **502 - 1**
- Pietinen P, Nissinen A, Vartiainen E, Tuomilehto A, Uusitalo U, Ketola A, Moiso S, Puska P. *Preventive Medicine* 1988;**17**:183-93. **312 - 1**
- Piotrow PT, Kincaid DL, Hindin MJ, Lettenmaier CL, Kuseka I, Silberman T, Zinanga A, Chikara F, Adamchak DJ, Mbizvo MT, Lynn W, Kumak CM, Kim Y-M. *Studies in Family Planning* 1992;**23**(6 Pt 1):365-75. **391 - 1**
- Plaskon PP, Fadden MJ. *Social Work in Health Care* 1995;**21**(4):59-70. **197 - 1**
- Ponsonby A, Dwyer T, Kasl S, Cochrane J, Newman N. *Preventive Medicine* 1994;**23**:402-8. **239 - 1**
- Price JH, Krol RA, Desmond SM, Losh DP, Roberts SM, Snyder FF. *Psychological Reports* 1991;**68**(2):595-604. **329 - 1**
- Quaid K, Faden R, Vining E, Freeman J. *Patient Education and Counseling* 1990;**15**:249-59. **301 - 1**
- Quirk ME, Godkin MA, Schwenzfeier E. *American Journal of Preventive Medicine* 1993;**9**(1):21-6. **465 - 1**
- Rabin JM, McNett J, Badlani GH. *Journal of Medical Systems* 1996;**20**(1):19-34. **40 - 1**
- Reading R, Colver A, Openshaw S, Jarvis S. *British Medical Journal* 1994;**308**(6937):1142-4. **767 - 1**
- Reid C, McNeil JJ, Williams F, Powles J. *Journal of Cardiovascular Risk* 1995;**2**(2):155-63. **50 - 1**
- Reid GS, Robertson AJ, Bissett C, Smith J, Waugh N, Halkerston R. *British Medical Journal* 1991;**303**(6800):447-50. **795 - 1**
- Remafedi G. *Journal of Adolescent Health* 1994;**15**(2):142-8. **123 - 1**
- Rhodes F, Wolitski RJ, Thornton-Johnson S. *Health and Social Work* 1992;**17**(4):261-72. **450 - 1**
- Richardson JL, Mondrus GT, Danley K, Deapen D, Mack T. *Cancer Epidemiology, Biomarkers and Prevention* 1996;**5**(1):71-6. **154 - 1**

- Rickert VI, Gottlieb AA, Jay MS. *Clinical Pediatrics* 1992;**31**(4):205–10. **289 - 1**
- Rimer BK, Resch N, King E, Ross E, Lerman C, Boyce A, Kessler H, Engstrom PF. *Public Health Reports* 1992;**107**(4):369–80. **354 - 1**
- Rimer BK, Ross E, Balslem A, Engstrom PF. *Journal of the American Board of Family Practice* 1993;**6**(5):443–51. **404 - 1**
- Robertson I, Phillips A, Mant D, Thorogood M, Fowler G, Fuller A, Yudkin P, Woods M. *British Journal of General Practice* 1992;**42**(364):469–72. **708 - 1**
- Robinson J. *Preventive Medicine* 1992;**21**:754–65. **266 - 1**
- Robinson JK, Rademaker AW. *Preventive Medicine* 1995;**24**(4):333–41. **198 - 1**
- Robinson MH, Pye G, Thomas WM, Hardcastle JD, Mangham CM. *European Journal of Surgical Oncology* 1994;**20**(5):545–8. **675 - 1**
- Robinson MK, DeHaven MJ, Koch KA. *Journal-of-Family-Practice* 1993;**37**(4):363–8. **495 - 1**
- Roca-Cusachs A, Sort D, Altimira J, Bonet R, Guilera E, Monmany J, Nolla J. *Journal of Human Hypertension* 1991;**5**(5):437–41. **320 - 1**
- Rodriguez JG, Velez M, Serrano E, Casado MP. *Boletin – Asociacion Medica de Puerto Rico* 1995;**87**(3–4):49–53. **815 - 1**
- Rose MA. *Public Health Nursing* 1992;**9**(4):242–7. **456 - 1**
- Ross JD, Scott GR. *Genitourinary Medicine* 1993;**69**(3):193–5. **670 - 1**
- Rosser WW, Hutchison BG, McDowell I, Newell C. *Canadian Medical Association Journal* 1992;**146**(6):911–17. **299 - 1**
- Rossiter JC. *International Journal of Nursing Studies* 1994;**31**(4):369–79. **616 - 1**
- Rossouw JE, Jooste PL, Chalton DO, Jordaan ER, Langenhoven ML, Jordaan PC, Steyn M, Swanepoel AS, Rossouw LJ. *International Journal of Epidemiology* 1993;**22**(3):428–38. **740 - 1**
- Rost KM, Flavin KS, Cole K, McGill JB. *Diabetes Care* 1991;**14**(10):881–9. **472 - 1**
- Rotheram-Borus MJ, Koopman C, Haignere C, Davies M. *JAMA* 1991;**266**(9):1237–41. **628 - 1**
- Rotheram-Borus MJ, Rosario M, Reid H, Koopman C. *American Journal of Psychiatry* 1995;**152**(4):588–95. **526 - 1**
- Rothman AJ, Salovey P, Turvey C, Fishkin SA. *Health Psychology* 1993;**12**(1):39–47. **789 - 1**
- Rowan-Szal G, Joe GW, Chatham LR, Simpson DD. *Journal of Substance Abuse Treatment* 1994;**11**(3):217–23. **156 - 1**
- Rowley MJ, Hensley MJ, Brinsmead MW, Wlodarczyk JH. *Medical Journal of Australia* 1995;**163**(6):289–93. **678 - 1**
- Rutten G, Beek M, van Eijk J. *Patient Education and Counseling* 1993;**22**:127–32. **337 - 1**
- Rutten G, Van Eijk J, Beek M, Van der Velden H. *British Journal of General Practice* 1991;**41**(348):289–92. **796 - 1**
- Sachs GA, Stocking CB, Miles SH. *Journal of the American Geriatrics Society* 1992;**40**(3):269–73. **343 - 1**
- Salleras Sanmarti L, Alcaide Megias J, Altet Gomez MN, Canela Soler J, Navas Alcala E, Sune Puigbo MR, Serra Majem L. *Tubercle and Lung Disease* 1993;**74**(1):28–31. **304 - 1**
- Sanchez-Craig M, Spivak K, Davila R. *British Journal of Addiction* 1991;**86**(7):867–76. **471 - 1**
- Sander RW, Holloway RL, Eliason C, marbella AM, Murphy B, Yuen S. *Journal Family Practice* 1996;**43**:468–74. **601 - 1**
- Sander RW, Holloway RL, Eliason C, marbella AM, Murphy B, Yuen S. *Journal Family Practice* 1996;**43**:468–74. **601 - 2**
- Sanner MA, Hedman H, Tufveson G. *Clinical Transplantation* 1995;**9**(4):326–33. **58 - 1**
- Santelli JS, Celentano DD, Rozsenich C, Crump AD, Davis MV, Polacsek M, Augustyn M, Rolf J, McAlister AL, Burwell L. *AIDS Education and Prevention* 1995;**7**(3):210–20. **780 - 1**
- Santi SM, Cargo M, Brown KS, Best JA, Cameron R. *Addictive Behaviors* 1994;**19**(3):269–85. **145 - 1**
- Saunders LD, Irwig LM, Gear JS, Ramushu DL. *Medical Care* 1991;**29**(7):669–78. **447 - 1**
- Saxon AJ, Calsyn DA. *Journal of Studies on Alcohol* 1992;**53**(6):611–18. **341 - 1**
- Schapira DV, Kumar NB, Clark RA, Yag C. *Cancer* 1992;**70**(2):509–12. **712 - 1**
- Schectman G, Hiatt J, Hartz A. *The Annals of Pharmacotherapy* 1994;**28**:29–35. **168 - 1**
- Schinke SP, Singer B, Cole K, Contento I. *Preventive Medicine* 1996;**25**:146–55. **207 - 1**
- Schmid T, Jeffery R, Hellerstedt W. *Preventive Medicine* 1989;**18**:503–17. **216 - 1**
- Schmid T, Jeffery R, Hellerstedt W. *Preventive Medicine* 1989;**18**:503–17. **216 - 2**
- Schultz S. *Patient Education and Counseling* 1993;**22**:47–57. **325 - 1**
- Schwarz-Lookinland S, McKeever L, Saputo M. *Patient Education and Counseling* 1989;**13**:171–82. **357 - 1**
- Sciacca JP, Dube DA, Phipps BL, Ratliff MI. *Journal of Community Health* 1995;**20**(6):473–90. **135 - 1**
- Scivoletto S, De Andrade AG, Castel S. *British Journal of Addiction* 1992;**87**(8):1185–8. **711 - 1**
- Sclar DA, Chin A, Skaer TL, Okamoto MP, Nakahiro RK, Gill MA. *Clinical Therapeutics* 1991;**13**(4):489–95. **469 - 1**
- Secker-Walker R, Flynn B, Solomon L, Vacek P, Bronson D. *Preventive Medicine* 1990;**19**:675–85. **249 - 1**



- Serwint JR, Wilson MH, Duggan AK, Mellits ED, Baumgardner RA, DeAngelis C. *Pediatrics* 1991;**88**(3):444-9. **788 - 1**
- Shafir MS, Silversides C, Waters I, MacRury K, Frank JW, Becker LA. *Canadian Family Physician* 1995;**41**:1367-72. **155 - 1**
- Shannon BM, Tershakovec AM, Martel JK, Achterberg CL, Cortner JA, Smiciklas-Wright HS, Stallings VA, Stolley PD. *Pediatrics* 1994;**94**(6 Pt 1):923-7. **754 - 1**
- Sharp DJ, Peters TJ, Bartholomew J, Shaw A. *Journal of Epidemiology and Community Health* 1996;**50**(1):72-6. **630 - 1**
- Shattuck AL, White E, Kristal AR. *American Journal of Public Health* 1992;**82**(9):1244-50. **659 - 1**
- Sheppard S, Coulter A, Farmer A. *Family Practice* 1995;**12**:443-7. **594 - 1**
- Shiloh S, Reznik H, Bat-Miriam-Katznelson M, Goldman B. *Social Science and Medicine* 1995;**41**(9):1301-10. **430 - 1**
- Shipley R, Orleans C, Wilbur C, Piserchia P, McFadden D. *Preventive Medicine* 1988;**17**:25-34. **284 - 1**
- Shope JT, Dielman TE, Butchart AT, Campanelli PC, Kloska DD. *Journal of Studies on Alcohol* 1992;**53**(2):106-21. **370 - 1**
- Shope JT, Kloska DD, Dielman TE, Maharg R. *Journal of School Health* 1994;**64**(4):160-6. **120 - 1**
- Shore ER. *Journal of Studies on Alcohol* 1994;**55**(6):657-9. **183 - 1**
- Siegal HA, Falck RS, Carlson RG, Wang J. *AIDS Education and Prevention* 1995;**7**(4):308-19. **528 - 1**
- Sikorski J, Wilson J, Clement S, Das S, Smeeton N. *British Medical Journal* 1996;**312**:546-53. **425 - 1**
- Simel DL, Feussner JR. *Journal of Clinical Epidemiology* 1991;**44**(8):771-7. **542 - 1**
- Simeoni E, Bauman A, Stenmark J, O'Brien J. *Arthritis Care and Research* 1995;**8**(2):102-7. **95 - 1**
- Simkin-Silverkin L, Wing RR, Hansen DH, Klem ML, Pasagian-Macauley A, Meilahn EN, Kuller LH. *Preventive Medicine* 1995;**24**:509-17. **200 - 1**
- Simmons MS, Nides MA, Rand CS, Wise RA, Tashkin DP. *Chest* 1996;**109**(4):963-8. **527 - 1**
- Skinner D, Metcalf CA, Seager JR, de Swardt JS, Laubscher JA. *AIDS Care* 1991;**3**(3):317-29. **610 - 1**
- Slap GB, Plotkin SL, Khalid N, Michelman DF, Forke CM. *Journal of Adolescent Health* 1991;**12**(6):434-42. **480 - 1**
- Smith MU, Katner HP. *AIDS Education and Prevention* 1995;**7**(5):391-402. **17 - 1**
- Smith NA, Seale JP, Ley P, Mellis CM, Shaw J. *Monaldi Archives for Chest Disease* 1994;**49**(6):470-4. **449 - 1**
- Smith S, Robinson J, Hollyer J, Bhatt R, Ash S, Shaunak S. *British Medical Journal* 1996;**312**(7028):416-20. **748 - 1**
- Sobczyk W, Hazel N, Reed CD, Ciarroccki B, Cohen S, Varga D. *Journal of the Kentucky Medical Association* 1995;**93**(4):142-7. **133 - 1**
- Sorensen JL, London J, Heitzmann C, Gibson DR, Morales ES, Dumontet R, Acree M. *AIDS Education and Prevention* 1994;**6**(2):95-112. **632 - 1**
- Sorensen JL, London J, Heitzmann C, Gibson DR, Morales ES, Dumontet R, Acree M. *AIDS Education and Prevention* 1994;**6**(2):95-112. **632 - 2**
- Spatz TS. *Journal of Cancer Education* 1991;**6**(3):179-83. **386 - 1**
- Spaulding SA, Kugler JP. *Journal of Family Practice* 1991;**33**(5):495-8. **378 - 1**
- Spoth R, Redmond C. *Psychology and Health* 1994;**9**:353-70. **600 - 1**
- St Lawrence JS, Brasfield TL, Jefferson KW, Alleyne E, O'Bannon RE, 3rd, Shirley A. *Journal of Consulting and Clinical Psychology* 1995;**63**(2):221-37. **766 - 1**
- St Pierre TL, Kaltreider DL, Mark MM, Aikin KJ. *American Journal of Community Psychology* 1992;**20**(6):673-706. **453 - 1**
- Stanton BF, Li X, Galbraith J, Feigelman S, Kaljee L. *Archives of Pediatrics and Adolescent Medicine* 1996;**150**(1):17-24. **585 - 1**
- Stehr-Green PA, Dini EF, Lindegren ML, Patriarca PA. *Public Health Reports* 1993;**108**(4):426-30. **408 - 1**
- Stephens RC, Feucht TE, Roman SW. *American Journal of Public Health* 1991;**81**(5):568-71. **729 - 1**
- Stevens MM, Freeman DH, Jr, Mott LA, Youells FE, Linsey SC. *American Journal of Preventive Medicine* 1993;**9**(3):160-7. **385 - 1**
- Stevens VJ, Glasgow RE, Hollis JF, Lichtenstein E, Vogt TM. *Medical Care* 1993;**31**(1):65-72. **287 - 1**
- Stevens-Simon C, O'Connor P, Bassford K. *Journal of Adolescent Health* 1994;**15**(5):396-9. **106 - 1**
- Stewart DE, Buchegger PM, Lickrish GM, Sierra S. *Obstetrics and Gynecology* 1994;**83**(4):583-5. **713 - 1**
- Stock LZ, Milan MA. *Journal of Applied Behavior Analysis* 1993;**26**(3):379-87. **162 - 1**
- Straton JA, Sutherland GJ, Hyndman JC. *Australian Journal of Public Health* 1995;**19**(3):288-93. **48 - 1**
- Strecher VJ, Kreuter M, Den Boer DJ, Kobrin S, Hospers HJ, Skinner CS. *Journal of Family Practice* 1994;**39**(3):262-70. **158 - 1**
- Strecher VJ, Kreuter M, Den Boer DJ, Kobrin S, Hospers HJ, Skinner CS. *Journal of Family Practice* 1994;**39**(3):262-70. **158 - 2**
- Street RL, Jr, Voigt B, Geyer C, Jr, Manning T, Swanson GP. *Cancer* 1995;**76**(11):2275-85. **667 - 1**
- Sumner W. *Family Medicine* 1991;**23**(6):463-6. **418 - 1**

- Sunwood J, Brenman A, Escobedo J, Philpott T, Allman K, Mueller J, Jaeger J, Brown LK, Cole FS. *Journal of Adolescent Health* 1995;**16**(4):309–15. **49 - 1**
- Sussman S, Parker VC, Lopes C, Crippens DL, Elder P, Scholl D. *International Journal of the Addictions* 1995;**30**(9):1141–64. **55 - 1**
- Sussman S, Parker VC, Lopes C, Crippens DL, Elder P, Scholl D. *International Journal of the Addictions* 1995;**30**(9):1141–64. **55 - 2**
- Sutton S, Bickler G, Sancho-Aldridge J, Saidi G. *Journal of Epidemiology and Community Health* 1994;**48**(1):65–73. **784 - 1**
- Sutton S, Hallett R. *Preventive Medicine* 1988;**17**:48–59. **310 - 1**
- Swaddiwudhipong W, Chaovakiratipong C, Nguntr P, Khumklam P, Silarug N. *International Journal of Epidemiology* 1993;**22**(4):660–5. **779 - 1**
- Swaddiwudhipong W, Chaovakiratipong C, Nguntr P, Mahasakpan P, Lerdlukanavong P, Koonchote S. *International Journal of Epidemiology* 1995;**24**(3):493–8. **575 - 1**
- Taal E, Riemsma RP, Brus HL, Seydel ER, Rasker JJ, Wiegman O. *Patient Education and Counseling* 1993;**20**(2–3):177–87. **98 - 1**
- Tabak ER. *Patient Education and Counseling* 1988;**12**:37–49. **506 - 1**
- Taddio A, Ito S, Einarson TR, Leeder JS, Koren G. *Reproductive Toxicology* 1995;**9**(2):153–7. **137 - 1**
- Tambor ES, Bernhardt BA, Chase GA, Faden RR, Geller G, Hofman KJ, Holtzman NA. *American Journal of Human Genetics* 1994;**55**(4):626–37. **516 - 1**
- Tanke ED, Leirer VO. *Medical Care* 1994;**32**(4):380–9. **481 - 1**
- Tedesco LA, Keffer MA, Davis EL, Christersson LA. *Journal of Periodontology* 1992;**63**(7):567–75. **758 - 1**
- Tenn L, Dewis ME. *Journal of Advanced Nursing* 1996;**23**(2):329–37. **570 - 1**
- Thomas DR, Winsted B, Koontz C. *Journal of the American Geriatrics Society* 1993;**41**(9):928–30. **342 - 1**
- Thornton JG, Hewison J, Lilford RJ, Vail A. *British Medical Journal* 1995;**311**(7013):1127–30. **778 - 1**
- Tietge N, Bender S, Scutchfield FD. *Patient Education and Counseling* 1987;**9**:167–75. **293 - 1**
- Torgerson DJ, Donaldson C, Garton MJ, Reid DM, Russell IT. *Health Economics* 1993;**2**(1):55–8. **634 - 1**
- Toseland RW, O'Donnell JC, Engelhardt JB, Hendler SA, Richie JT, Jue D. *Medical Care* 1996;**34**(6):624–40. **10 - 1**
- Town GI, Hodges ID, Wilkie AT, Toop LJ, Graham P, Drennan CJ. *Patient Education and Counseling* 1995;**26**:219–24. **522 - 1**
- Trock B, Rimer BK, King E, Balslem A, Cristinzio CS, Engstrom PF. *Cancer Epidemiology, Biomarkers and Prevention* 1993;**2**(2):151–6. **446 - 1**
- Tu KS, McDaniel G, Gay JT. *Diabetes Educator* 1993;**19**(1):25–30. **475 - 1**
- Turnbull D, Irwig L, Simpson JM, Donnelly N. *Social Science and Medicine* 1994;**39**(4):543–51. **608 - 1**
- Turner KM, Wilson BJ, Gilbert FJ. *Journal of Medical Screening* 1994;**1**(3):199–202. **41 - 1**
- Turnin MC, Beddok RH, Clottes JP, Martini PF, Abadie RG, Buisson JC, Soule-Dupuy C, Bonneau M, Camare R, Anton JP, Chrismet CY, Farreny H, Bayard F, Tauber J-PJ. *Diabetes Care* 1992;**15**(2):204–12. **396 - 1**
- Urban N, Taplin S, Taylor V, Peacock S, Anderson G, Conrad D, Etzioni R, White E, Montano D, Mahloch J, Majer K. *Preventive Medicine* 1995;**24**:477–84. **439 - 1**
- Urban N, Taplin SH, Taylor VM, Peacock S, Anderson G, Conrad D, Etzioni R, White E, Montano DE, Mahloch J, Majer K. *Preventive Medicine* 1995;**24**(5):477–84. **203 - 1**
- Usherwood TP. *British Journal of General Practice* 1991;**41**(343):58–62. **706 - 1**
- van Assema P, Steenbakkens M, Kok G, Erikson M, de Vreis H. *Preventive Medicine* 1994;**23**:394–401. **237 - 1**
- van Haastrecht HJ, van den Hoek JA, Coutinho RA. *Genitourinary Medicine* 1991;**67**(3):199–206. **680 - 1**
- van Heeringen C, Jannes S, Buylaert W, Henderick H, de Bacquer D, van Remoortel J. *Psychological Medicine* 1995;**25**(5):963–70. **606 - 1**
- Velicer WF, Prochaska JO, Bellis JM, Dicelmente CC, Rossi JS, Fava JL, Steiger JH. *Addictive Behaviors* 1993;**18**(3):269–90. **668 - 1**
- Verne J, Kettner J, Mant D, Farmer A, Mortenson N, Northover J. *European Journal of Cancer Prevention* 1993;**2**(4):301–5. **122 - 1**
- Vineis P, Ronco G, Ciccone G, Gogliani F. *Epidemiology* 1994;**5**(3):349–51. **170 - 1**
- Voorhees C, Stillman F, Swank R, Heagerty P, Levine D, Becker D. *Preventive Medicine* 1996;**25**:277–85. **211 - 1**
- Walker R, Heller R, Redman S, O'Connell D, Boulton J. *Preventive Medicine* 1992;**21**:616–29. **265 - 1**
- Walsh DC, Hingson RW, Merrigan DM, Levenson SM. *Alcohol Health and Research World* 1992;**16**(2):140–8. **485 - 1**
- Walsh DC, Hingson RW, Merrigan DM, Levenson SM, Cupples LA, Heeren T, Coffman GA, Becker CA, Barker TA, Hamilton SK, McGuire TG, Kelly CA. *New England Journal of Medicine* 1991;**325**(11):775–82. **564 - 1**
- Walter HJ, Vaughan RD. *JAMA* 1993;**270**(6):725–30. **660 - 1**
- Ward JE, Boyl K, Redman S, Sanson-Fisher RW. *American Journal of Preventive Medicine* 1991;**7**(5):285–91. **377 - 1**

- Ware GJ, Holfard NH, Davison JG, Harris RG. *New Zealand Medical Journal* 1991;**104**(924):495-7. **315 - 1**
- Warnecke RB, Langenberg P, Wong SC, Flay BR, Cook TD. *American Journal of Public Health* 1992;**82**(6):835-40. **657 - 1**
- Wasson J, Gaudette C, Whaley F, Sauvigne A, Baribeau P, Welch HG. *JAMA* 1992;**267**(13):1788-93. **638 - 1**
- Webster A. *British Journal of Medical Psychology* 1992; **65**(Pt 2):89-93. **717 - 1**
- Weiden P, Rapkin B, Zygmunt A, Mott T, Goldman D, Frances A. *Psychiatric Services* 1995;**46**(10):1049-54. **124 - 1**
- Weinberger M, Tierney WM, Booher P, Katz BP. *Journal of Rheumatology* 1991;**18**(6):849-54. **369 - 1**
- Weinrich SP, Weinrich MC, Boyd MD, Atwood J, Cervenka B. *Cancer Nursing* 1994;**17**(6):494-500. **448 - 1**
- Weinrich SP, Weinrich MC, Stromborg MF, Boyd MD, Weiss HL. *Gerontologist* 1993;**33**(4):491-6. **117 - 1**
- Weinstein ND, Sandman PM, Roberts NE. *Health Psychology* 1991;**10**(1):25-33. **288 - 1**
- Weinstein R, Tosolin F, Ghilardi L, Zanardelli E. *Journal of Clinical Periodontology* 1996;**23**(3 Pt 2):283-8. **576 - 1**
- Welsh MC, Labbe EE, Delayney D. *Psychological Reports* 1991;**68**(3 Pt 2):1327-35. **321 - 1**
- Wenger NS, Linn LS, Epstein M, Shapiro MF. *American Journal of Public Health* 1991;**81**(12):1580-5. **741 - 1**
- Werch CE, Young M, Clark M, Garrett C, Hooks S, Kersten C. *Journal of School Health* 1991;**61**(8):346-50. **360 - 1**
- West S, Munoz B, Lynch M, Kayongoya A, Chilangwa Z, Mmbaga BB, Taylor HR. *Lancet* 1995;**345**(8943):155-8. **777 - 1**
- Whelan TJ, Levine MN, Gafni A, Lukka H, Mohide EA, Patel M, Streiner DL. *Journal of Clinical Oncology* 1995;**13**(4):847-53. **691 - 1**
- White E, Hurlich M, Thompson RS, Woods MN, Henderson MM, Urban N, Kristal A. *American Journal of Preventive Medicine* 1991;**7**(5):319-25. **365 - 1**
- Wierenga ME. *Patient Education and Counseling* 1994;**23**(1):33-40. **503 - 1**
- Williams LR, Ekers MA, Collins PS, Lee JF. *Journal of Vascular Surgery* 1991;**14**(3):320-6. **821 - 1**
- Williford SL, Johnson DF. *Military Medicine* 1995;**160**(11):561-4. **88 - 1**
- Wilson TG, Jr, Hale S, Temple R. *Journal of Periodontology* 1993;**64**(4):311-14. **737 - 1**
- Windsor RA, Lowe JB, Perkins LL, Smith-Yoder D, Artz L, Crawford M, Amburgy K, Boyd NR, Jr. *American Journal of Public Health* 1993;**83**(2):201-6. **724 - 1**
- Wing RR, Anglin K. *Diabetes Care* 1996;**19**(5):409-13. **398 - 1**
- Winkleby M, Fortmann S, Rockhill B. *Preventive Medicine* 1992;**21**:592-601. **258 - 1**
- Winkleby M, Fortmann S, Rockhill B. *Preventive Medicine* 1993;**22**:325-34. **273 - 1**
- Winter L, Breckenmaker LC. *Family Planning Perspectives* 1991;**23**(1):24-30. **455 - 1**
- Winter L, Goldy AS. *Health Psychology* 1993;**12**(4):308-12. **685 - 1**
- Wolf AM, Nasser JF, Wolf AM, Schorling JB. *Archives of Internal Medicine* 1996;**156**(12):1333-6. **587 - 1**
- Worden J, Solomon L, Flynn B, Costanza M, Foster B, Dorwaldt A, Weaver S. *Preventive Medicine* 1990;**19**:254-69. **246 - 1**
- Xiang M, Ran M, Li S. *British Journal of Psychiatry* 1994;**165**(4):544-8. **479 - 1**
- Yancey AK, Tanjasiri SP, Klein M, Tunder J. *Preventive Medicine* 1995;**24**(2):142-8. **37 - 1**
- Young DR, Haskell WL, Taylor CB, Fortmann SP. *American Journal of Epidemiology* 1996;**144**(3):264-74. **464 - 1**
- Zapka J, Costanza M, Harris D, Hosmer D, Stoddard A, Barth R, Gaw V. *Preventive Medicine* 1993;**22**:34-53. **243 - 1**
- Zapka JG, Harris DR, Hosmer D, Costanza ME, Mas E, Barth R. *Health Services Research* 1993;**28**(2):223-35. **809 - 1**
- Zapka JG, Stoddard AM, McCusker J. *AIDS Education and Prevention* 1993;**5**(4):352-66. **720 - 1**



# Appendix 7

## Decision details by study design (grouped by health area)

**TABLE 25** Description of the decision, the interventions, and the measures recorded in studies: RCTa

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
23 - I	Adherence to self-care behaviour in adults undergoing radiotherapy	None	Cancer	Control – routine care for radiation therapy patients Group 1 – routine plus video	Age, sex, level of education, marital status, ethnicity, knowledge, usefulness of information
39 - I	Uptake of cervical cancer screening in adult well women	None	Cancer	Control – no additional information Group 1 – general practitioner (GP) reminder letter Group 2 – letter headed by clinic Group 3 – additional information leaflet posted	Age, level of education, social class, marital status, actual risk status, medical history, health behaviours-general knowledge, perception of risk, intention
41 - I	Uptake of mammography in adult well women	None	Cancer	Control – letter not signed by GP Group 1 – letter signed by GP posted to women 'refusers'	Age, social class, health behaviours-general
52 - I	Uptake of mammography in well adult women	None	Cancer	Control – no additional information Group 1 – physician reminder telephone calls Group 2 – medical assistant telephone call Group 3 – reminder letter	Age, social class, marital status, actual risk status, health insurance
85 - I	Adherence to breast self-examination (BSE) in adult, well women	None	Cancer	Group 1 – information only Group 2 – information BSE and prompt on contraceptive pill packet	Age, level of education, ethnicity, health behaviours-general, self-efficacy
108 - I	Uptake of faecal occult blood (FOB) screening in well adults	None	Cancer	Control – usual screening pad: manipulation of stools Group 1 – experimental testing stools: pad dangled in toilet after bowel movement	Age, sex, health insurance
122 - I	Colorectal screening in well adults	None	Cancer	Group 1 – haemoccult kit: manipulation stools and information on diet Group 2 – early detector pad: used like toilet paper. Information about diet Group 3 – coloscreen: tissue paper placed in toilet after use. Information on diet	Age, sex
316 - I	Uptake of colorectal screening in well adults (workplace)	None	Cancer	Control – letter: availability of FOB testing at work Group 1 – risk appraisal form assessing personal risk plus letter invite colorectal screening at work	Age, sex, level of education, marital status, ethnicity, actual risk status, family history, health behaviours-general, knowledge, attitudes, perception of risk, intention
509 - I	Uptake of mammography in well adult women	None	Cancer	Group 1 – one-to-one counselling, leaflet plus free mammography voucher Group 2 – same information as Group 1, no voucher	Age, level of education, social class, marital status, actual risk status, family history, medical history, health behaviours-general, health insurance, living arrangements, knowledge, reasons, health professional measures, efficacy
101 - I	Uptake of mammography in well women	Framing information	Cancer	Control – routine letter invitation Group 1 – letter plus section on worrisome facts about breast cancer (anxiety condition) Group 2 – letter plus encouraging (coping) things about breast cancer	Age, family history, health behaviours-general

*continued*

**TABLE 25 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTa

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
117 - 1	Uptake of colorectal screening in elderly well adults	Health promotion	Cancer	Control – routine slide plus tape presentation Group 1 – elderly information giver gave talk before slide show (close identity participants) Group 2 – adapted slide show, more accessible, more procedural information about screening Group 3 – elderly educator and modified slide show	Age, sex, level of education, social class, marital status, ethnicity
197 - 1	Uptake of colorectal cancer screening in well adults	Social cognition model	Cancer	Group 1 – verbal plus written information about colorectal cancer and screening Group 2 – Group 1 plus free screening kit	Sex, level of education, ethnicity, health behaviours-general, perception of risk
275 - 1	Uptake of mammography in well adult women	Social cognition model	Cancer	Control – routine letter invitation Group 1 – letter plus more explanation risk breast cancer Group 2 – risk invitation plus self-evaluation risk questionnaire Group 3 – risk invitation, risk questionnaire plus personalised risk letter	Age, family history, medical history, reproductive history, health behaviours-general
53 - 1	Adherence to medication in adults with hypertension	None	Medicine	Control – routine medical care Group 1 – telephone link, regular care plus weekly calls using computerised voice and touch tone answers for feedback	Age, sex, level of education, social class, marital status, actual risk status
50 - 1	Adherence to health lifestyle, cessation of unhealthy lifestyle in well adults	None	Medicine	Group 1 – illustrated leaflet about cardiovascular disease (CVD) risk-reducing factors Group 2 – group counselling (1–2 hours), goal achievement, weight loss programmes	Age, sex, level of education, social class, actual risk status, physiological/psychological assessment
64 - 1	Cessation of smoking, alcohol and uptake of exercise in medicine well adults	None	Medicine	Control – routine CVD risk examination plus leaflet Group 1 – paid GPs, routine examination plus additional counselling, follow-up visit plus patient reimbursement	Age, sex, social class, marital status, ethnicity, actual risk status, health behaviours-general, health insurance
118 - 1	Cessation of smoking in well adults with asthmatic children	None	Medicine	Control – routine smoking risk information plus quit smoking in home advice Group 1 – child's physician gave information on passive smoking plus leaflet plus ways reducing child exposure smoke	Age, level of education, marital status, ethnicity, actual risk status
216 - 2	Adherence to cessation of smoking/fatty food programme in well adults	None	Medicine	Group 1 – invitation letter, leaflet on smoking and weight programmes, cost US\$5 flat fee Group 1 – letter invitation, leaflet and refundable deposit US\$60	Age, sex, level of education, actual risk status, physiological/psychological assessment
229 - 1	Cessation of smoking in well adults	None	Medicine	Group 1 – verbal information on smoking and nicotine gum plus free gum Group 2 – verbal information on smoking and gum plus gum cost US\$6 Group 3 – information on smoking and gum plus gum cost US\$20	Age, sex, level of education, social class, actual risk status
350 - 1	Cessation of smoking in well patients	None	Medicine	Group 1 – two appointments to discuss quitting and progress, offered gum plus leaflet Group 2 – same Group 1 plus offered additional visit	Age, sex, level of education, marital status, actual risk status, health behaviours-general, usefulness information, intention, social reinforcement/norms, motivation
483 - 1	Adherence to medication in elderly adults	None	Medicine	Control – pharmacists informed study but training deferred Group 1 – pharmacists trained in workshops in identifying problems with adherence plus leaflets for patients	Age, sex, level of education, social class, marital status, actual risk status, attitudes, usefulness of information

continued

**TABLE 25 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTa

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
648 - I	Cessation of smoking in well adults (worksites)	None	Medicine	Group 1 – minimal counselling: leaflet plus told to wear nicotine patch (22 mg or 44 mg strength) Group 2 – individual counselling: leaflet plus motivational message by physician plus quit letter Group 3 – group counselling: leaflet plus attend clinic for counselling and skill acquisition	Age, sex, level of education, ethnicity, actual risk status, medical history, physiological/psychological assessment,
793 - I	Cessation of in well adults	None	Medicine	Group 1 – nicotine patch plus usual leaflet Group 2 – nicotine patch plus additional information Group 3 – placebo patch plus leaflet Group 4 – placebo patch plus additional information	Age, sex, actual risk status, perception of well-being
781 - I	Adherence to self-management in adult with asthma	None	Medicine	Control – routine clinic care Group 1 – verbal information self-management, asthma, therapies, feedback on inhaler use	Age, sex, actual risk status, health behaviours-general, physiological/psychological assessment
755 - I	Changes to CVD risk factors in well adults	None	Medicine	Control 1 – another company delayed intervention Control 2 – delayed intervention within company Group 1 – health information Group 2 – health information plus cholesterol score feedback Group 3 – health information plus cholesterol risk figure Group 4 – health information plus cholesterol level plus risk score	Age, sex, social class, actual risk status, health behaviours-general, physiological/psychological assessment
67 - I	Intention to have cardio-pulmonary resuscitation (CPR) in well adults	None	Medicine	Control – no additional information Group 1 – leaflet CPR, framed in terms of senility Group 2 – leaflet CPR, framed in terms of Alzheimers disease	Age, sex, level of education, social class, marital status, religion, ethnicity, living arrangements knowledge
196 - I	Cessation of smoking, adherence to diet, use of services in well, pregnant women	None	Medicine, obstetrics and gynaecology (midwifery)	Control – routine leaflet Group 1 – leaflet plus weekly phone call from interviewer	Age, marital status, ethnicity, reproductive history anxiety – state/trait, depression, other affect, social support/stress
714 - I	Utilisation of care in parents of children recently attended emergency	None	Medicine, surgery (A&E)	Control – assessment only Group 1 – health professional follow-up phone call to parents to answer any questions	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, health insurance, perceived satisfaction
200 - I	Adherence to exercise, cessation of poor diet in adult well premenopausal women	None	Medicine, obstetrics and gynaecology (midwifery)	Control – no additional information Group 1 – group programmes by dietician: meal plans, exercise information, goal setting, newsletter	Age, level of education, social class, marital status, ethnicity, reproductive history, health behaviours-general, physiological/psychological assessment, health professional measures, worry, attitudes, social support/communication, usefulness of information, coping
724 - I	Cessation of smoking in pregnant women	None	Medicine, obstetrics and gynaecology (midwifery)	Control – assessment only Group 1 – routine information plus explanation on leaflet use plus follow-up session plus note prompt plus buddy	Age, level of education, ethnicity, reproductive history
581 - I	Cessation of smoking in adults undergoing minor surgery	None	Medicine, surgery	Control – blank tape played during surgery Group 1 – suggestion to give up smoking tape played during surgery	Age, sex, actual risk status, medical history, motivation, awareness of intervention

continued

**TABLE 25 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTa

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
542 - I	Participation in clinical trial in ambulatory adult patients	Framing information	Medicine	Group 1 – treatment A twice as effective Group 2 – treatment A half as effective	Age, level of education, social class, ethnicity, medical history reasons
120 - I	Cessation of alcohol in school children	Health promotion	Medicine	Control – control group Group 1 – modified 'alcohol misuse prevention study' information: more role-play against social pressure, interactive sessions with feedback	Sex, knowledge, attitudes, efficacy
301 - I	Compliance with medication in well and epileptic adults	Communication	Medicine	Group 1 – information medication using standard 'medical practice' form Group 2 – modified information written for lay person plus risk and side-effect information	Age, sex, level of education, marital status, actual risk status, anxiety – state, knowledge, perception of risk
325 - I	Uptake of exercise in adults with CVD	Shared consultation	Medicine	Group 1 – leaflet and verbal information on exercise, checking pulse Group 2 – Group 1 plus behavioural strategies: goal setting, reinforcement (phone) and diary	Age, sex, level of education, social class, marital status, actual risk status, health behaviours-general, physiological/psychological assessment, knowledge, social reinforcement/norms
202 - I	Uptake of appointment in adult substance users	None	STDs (HIV)	Control – no additional contact Group 1 – telephone call emphasising importance of attendance	Age, sex, social class, ethnicity, actual risk status, medical history, perceived satisfaction
393 - I	Cessation of/adherence to risky/safe sex in adolescents with STD	None	STDs (HIV)	Control – individual risk assessment HIV, leaflet and counselling on condom use/HIV prevention Group 1 – routine information plus physician asking about knowledge and skills to modify behaviour	Age, sex, ethnicity, actual risk status, medical history, health behaviours-general
741 - I	Cessation of risky sex/uptake of safe sex in adults with STD	None	STDs (HIV)	Control – counselling only on HIV risk factors plus blood test for STD Group 1 – HIV counselling plus STD blood plus HIV test	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, medical history, worry, perception of well-being, knowledge
107 - I	Use of condoms/safer sex in school children	Social cognition model	STDs (HIV)	Control – no additional information Group 1 – verbal information: sexual development, STD, contraception, skill acquisition in groups and communication skills	Age, sex, social class, actual risk status, health behaviours-general, knowledge, attitudes, self-efficacy
585 - I	Adherence to safe sex in well adolescents	Social cognition model, health promotion	STDs (HIV)	Group 1 – video information: HIV risks and behaviour and discussion groups Group 2 – information structured by protection-motivation theory, skill acquisition	Age, sex, actual risk status, knowledge
799 - I	Changes HIV risky behaviour in well adult women	Social cognition model	STDs (HIV)	Control – delayed information Group 1 – social skills acquisition, culturally appropriate information on HIV Group 2 – HIV information only	Age, level of education, social class, actual risk status, reproductive history, health behaviours-general, living arrangements knowledge, perception of control, self-efficacy, social reinforcement/norms
299 - I	Uptake of immunisation in well adults	None	Primary care	Control – routine Group 1 – physician reminder – printed on 'bill' after consultation Group 2 – telephone call made by nurse Group 3 – letter signed by physician	Age, sex, actual risk status, reproductive history
378 - I	Uptake of immunisation in well adults	None	Primary care	Control – routine care Group 1 – reminder postcard to visit physician	Age, sex, social class
638 - I	Utilisation of services in adult male patients primary care	None	Primary care	Control – usual care Group 1 – re-appointment times made and telephone contact made at three time points by physician	Age, marital status, actual risk status, medical history, health behaviours-general, perception of well-being, perceived satisfaction

continued



**TABLE 25 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTa

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
387 - I	Increased participation in consultation in adult general practice patients	Shared consultation	Primary care	Control – health eating leaflet, note examples high fat foods Group I – leaflet encouraging patients to be more active in consultation, note concerns for doctor, ask questions diagnosis and treatment (piloted)	Age, sex, social class, health professional measures perception of control, perceived satisfaction, self-efficacy, usefulness of information
333 - I	Adherence to immunogens in elderly well adults	Health promotion	Primary care	Control – assessment only Group I – verbal information about medication, appointments, illness	Age, sex, level of education, social class, marital status, actual risk status, medical history, health behaviours-general, physiological/psychological assessment, depression
425 - I	Uptake of prenatal services in pregnant well women	None	Obstetrics and gynaecology (midwifery)	Control – usual 13 visits with health professional Group I – reduced to seven visits with health professional	Age, level of education, social class, marital status, ethnicity, reproductive history, health behaviours-general, physiological/psychological assessment, health professional measures, worry, attitudes, social support communication, usefulness of information, coping
778 - I	Uptake of prenatal screening in well pregnant women	None	Obstetrics and gynaecology (midwifery), genetics	Control – routine care Group I – additional individual session plus leaflet and verbal information serum screening, ultrasound, carrier testing for cystic fibrosis (CF) Group 2 – women attended group sessions plus leaflets	Age, social class, ethnicity, reproductive history, anxiety – state, depression, worry
678 - I	Uptake of antenatal care in well pregnant women	None	Obstetrics and gynaecology (midwifery)	Control – usual antenatal care: attend clinics by different professionals Group I – team-care: known team of midwives provided continuity care throughout pregnancy	Age, social class, marital status, reproductive history, physiological/psychological assessment, autonomy in decision making, perception of informed consent, perceived satisfaction, usefulness of information
606 - I	Uptake of appointments in adults who attempted suicide	None	Mental health	Control – usual care: patients contact clinic Group I – nurse visit to patients not attending follow-up appointment	Age, sex, level of education, social class, marital status, actual risk status, medical history, anxiety – state, depression
427 - I	Adherence to medication in psychotic adults	Health promotion	Mental health	Control – supportive (listening) counselling Group I – counselling guided by 'compliance' therapy: beliefs, coping, attitudes to drug use	Age, sex, ethnicity, actual risk status, medical history, attitudes, disability scale

**TABLE 26** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
352 - 1	Uptake of mammography in well women	None	Cancer	Control – routine reminder postcard Group 1 – routine reminder postcard plus voucher on attendance (US\$2)	Age
352 - 2	Uptake of mammography in well women	None	Cancer	Control – routine reminder postcard Group 1 – telephone reminder	Age
352 - 3	Uptake of mammography in well adults	None	Cancer	Control – clinic-based reminder letter Group 1 – signed letter physician reminder	Age
353 - 1	Uptake of mammography in well women	None	Cancer	Control – no additional contact Group 1 – volunteers 'taught' about purpose mammography and to contact five acquaintances to encourage their attendance	Age, level of education, social class, marital status, ethnicity, health behaviours-general
420 - 1	Uptake of colorectal screening in well adults	None	Cancer	Control: letter invitation, FOB test plus leaflet plus reminder letter: half receive gain frame, half loss frame information Group 1 – routine care plus reminder phone call: gain/loss framed information Group 2 – routine plus information leaflet (readability score) about efficacy screening plus phone reminder: gain/loss framing Group 3 – routine plus additional leaflet plus reminder phone with additional information on use: gain/loss frame information	Age, sex, reasons
482 - 1	Uptake of colorectal screening in well adults	None	Cancer	Control – routine care: letter invitation plus testing kit Group 1 – routine care plus reminder phone call Group 2 – routine care plus leaflet plus reminder call Group 3 – routine care plus leaflet plus 'how to use test' call plus reminder call	Age, sex
653 - 1	Uptake of mammography in well adult women	None	Cancer	Control – routine letter invitation Group 1 – letter invitation plus information on taking part study for 'additional scientific investigation'	Age, marital status, ethnicity, living arrangements
675 - 1	Uptake of colorectal screening in well adults	None	Cancer	Group 1 – perform test over 3 days without diet restriction Group 2 – perform test over 6 days without diet restriction Group 3 – perform test over 3 days with diet restriction Group 4 – perform test over 6 days with diet restriction	Age, sex
683 - 1	Uptake of colorectal cancer screening in well adults	None	Cancer	Group 1 – signed GP letter plus diet restrictions plus FOB test Group 2 – signed GP letter plus faecal kit Group 3 – signed letter, leaflet plus FOB testing kit Group 4 – signed letter plus instructions to phone for free kit Group 5 – letter signed by 'professor' and FOB testing kit	Age, sex, marital status
491 - 1	Uptake of BSE in well women (work)	None	Cancer	Control – group session on BSE, video plus leaflet plus practice movement Group 1 – control plus feedback on BSE on model Group 2 – control plus Group 2 plus individual session plus intensive feedback	Age, level of education, marital status, ethnicity, health behaviours-general, physiological/psychological assessment
533 - 1	Uptake of BSE in well women	None	Cancer	Control – verbal, video and leaflet information on BSE and 'over-clothing' technique Group 1 – control plus individual supervision on model with lumps Group 2 – Group 1 plus individual counselling	

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
566 - I	Returning colorectal screening cards in well adults	None	Cancer	Control – hemocult cards by post, no stamped addressed envelope Group 1 – hemocult cards sent with stamped addressed envelopes	Age, sex, marital status, ethnicity, health insurance
705 - I	Adherence to safe-sun behaviours in well adolescents	None	Cancer	Control – questionnaire only Group 1 – read through leaflet Group 2 – read leaflet plus video Group 3 – read leaflet plus designed poster Group 4 – leaflet plus discussion one week later	Age, sex, actual risk status, knowledge, attitudes
728 - I	Uptake of cervical smear in well women	None	Cancer	Control – usual care Group 1 – leaflet plus fact sheet by post (available in several Asian languages) Group 2 – personal visit plus fact sheet plus video Group 3 – personal visit plus fact sheet plus leaflet	Age, social class, marital status, religion, ethnicity, actual risk status, usefulness of information
630 - I	Uptake of breast screening well women non-attendees	None	Cancer	Group 1 – home visits by nurse, ascertained reasons non-attendance and information on screening Group 2 – home visit nurse, ascertained reasons non-attendance Group 3 – signed reminder letter GP	Age, perception of control, self-esteem
285 - I	Uptake of colorectal cancer screening in well adults	None	Cancer	Group 1 – letter to all households plus leaflet information colorectal cancer screening, media campaign plus refund for buying test Group 2 – Group 1 plus given free test at surgery Group 3 – Group 1 plus test sent by post to non-attendees	Age, sex
727 - I	Uptake of mammography in well women	None	Cancer	Group 1 – free screening plus media information plus open invitation letter to non attendees Group 2 – Group 1 plus fixed appointment letter	Age
543 - I	Uptake of cervical smear in well women	None	Cancer	Control – routine breast screening appointment Group 1 – timed cervical screening appointment as well as breast screening	Age, ethnicity, health behaviours-general
587 - I	Uptake of in prostate screening in well male adults	None	Cancer	Group 1 – verbal information availability screening Group 2 – description of procedure screening	Age, level of education, social class, ethnicity, family history, health insurance
713 - I	Uptake of cervical smears in well women with screen-positive smears	None	Cancer	Control – routine care Group 1 – information leaflet	Age, level of education, marital status, worry, knowledge
48 - I	Uptake of cervical screening in well women	None	Cancer	Control – comparison women in hospital ward Group 1 – offered smear in hospital Group 2 – offered leaflet when discharged	Age, level of education, marital status, ethnicity, reproductive history, health professional measures
488 - I	Uptake of hypothetical medication trial in adults with cancer	None	Cancer	Group 1 – audio-tape plus verbal information cancer, trial and treatment provided Group 2 – computer screen touch control, programme same information as Group 1	Age, sex, level of education, knowledge, perceived satisfaction, usefulness of information
86 - I	Uptake of breast and cervical screening in well adult women	Health promotion	Cancer	Control – information about community living Group 1 – information about the importance of screening, BSE, nutrition and attending health centres	Age, level of education, social class, marital status, ethnicity, attitudes, health behaviours-general, service utilisation of, health insurance, knowledge, reasons, self-efficacy
114 - I	Uptake of mammography in adult well women	Social cognition model	Cancer	Control – no additional information Group 1 – belief intervention: 'stages of change/HBM' assessment, tailored belief intervention plus leaflet plus professional contact Group 2 – information only: facts, timing and procedure mammography Group 3 – Group 1 plus 2: facts on procedure plus targeted beliefs	Age, level of education, ethnicity, health behaviours-general, knowledge, attitudes, perception of risk, reasons, operationalised model, motivation

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
232 - I	Uptake of mammography in well women	Social cognition model	Cancer	Control – leaflet by post: breast cancer plus thank you for questionnaire completion Group 1 – extra leaflet: importance screening, procedure and attendance encouragement plus thank you note	Age, education, social class, marital status, ethnicity, health behaviours, insurance, knowledge, attitudes, perception of risk, reasons, intent, efficacy
438 - I	Uptake of mammography in well adult women (workplace)	Framing information	Cancer	Group 1 – video: positive frame mammography screening (saves lives) Group 2 – video: negative frame mammography screening (not attend is life threatening)	Age, level of education, social class, marital status, religion, ethnicity, family history, health behaviours-general, knowledge, attitudes, perception of risk, self-efficacy, intention, efficacy, other affect
784 - I	Uptake of mammography in well women	Social cognition model	Cancer	Control – women not interviewed until after screening Group 1 – women interviewed before screening Group 2 – postal questionnaire sent before screening	Age, level of education, social class, marital status, religion, ethnicity, family history, medical history, health behaviours-general, physiological/psychological assessment, living arrangements, reproductive history, perception of well-being, attitudes, perception of risk, perception of susceptibility, intention, efficacy
700 - I	Participation in consultation in adults with cancer	Shared consultation	Cancer	Group 1 – informed cancer services are available Group 2 – information encouraged writing questions down before consultation	Age, sex, actual risk status, medical history, living arrangements, anxiety – state, knowledge, preference consultation style, perceived satisfaction, disability scale, coping
789 - I	Uptake of mammography in well women	Attribution theory	Cancer	Group 1 – neutral information video about breast screening Group 2 – video plus internal responsibility frame to get mammography Group 3 – video plus external (physician) responsibility frame to order mammography	Age, level of education, social class, marital status, religion, ethnicity, actual risk status, family history, medical history, perception of well-being, relief/reassurance, regret/responsibility, knowledge, attitudes, usefulness of information
207 - I	Cessation of smoking/adherence to diet in well children	Social learning theory	Cancer, medicine	Control – questionnaire only Group 1 – verbal information: tobacco prevention plus culturally appropriate peer volunteer in school Group 2 – information to improve diet Group 3 – information aimed at diet and smoking	Age, sex, knowledge, attitudes
288 - I	Uptake of radon testing in well adults	Social cognition model	Cancer, other	Group 1 – telephoned adults plus leaflet plus advised to test home Group 2 – telephoned adults plus leaflet plus encourage house testing plus told high risk area and house likely to have radon	Age, sex, level of education, reproductive history, health behaviours-general, perception of risk, perception of susceptibility, reasons
690 - I	Uptake of BSE	Social cognition model	Cancer	Control – data collection only Group 1 – belief targeted verbal plus leaflets (HBM): breast cancer and BSE Group 2 – routine information plus skills on BSE Group 3 – Groups 1 plus 2	Age, level of education, ethnicity, attitudes, perception of risk, perception of susceptibility, reasons, operationalised model, motivation
458 - I	Cessation of skin cancer risks in well adolescents	Social cognition model	Cancer	Control – assessment only Group 1 – videotape risks skin cancer, self-assessment form, feedback and ways change behaviour	Age, sex, level of education, actual risk status, health behaviours-general, knowledge, attitudes, perception of susceptibility, intention, social reinforcement/norms
307 - I	Uptake of BSE in older well women	Social cognition model	Cancer	Control – basic one-to-one BSE instruction Group 1 – peer BSE 'buddy' selected Group 2 – women brought partner of choice for BSE buddy	Age, level of education, marital status, attitudes, self-efficacy
140 - I	Cessation of smoking in adults with cancer	Social cognition model	Cancer, medicine	Control – routine care Group 1 – leaflet plus quit smoking sessions plus trained health professional	Age, sex, level of education, marital status, actual risk status, medical history, physiological/psychological assessment, knowledge, attitudes, operationalised model

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
757 - 1	Uptake of mammography in well women	Framing information	Cancer	Control – no assessment Control 2 – assessment only Group 1 – leaflet mammography negatively framed plus invitation letter Group 2 – leaflet positively framed plus letter	Age, level of education, ethnicity, actual risk status, medical history
434 - 1	Uptake of trial cancer treatment in adults with cancer	Framing information	Cancer	Group 1 – scenario: surgical removal intestine followed by chemotherapy, neutrally framed Group 2 – scenario: surgery followed by chemotherapy positively framed Group 3 – scenario: surgery followed by chemotherapy negatively framed	Age, sex, level of education, actual risk status, medical history, attitudes, autonomy in decision making
816 - 1	Participation in clinical trials in adults with cancer	Shared consultation	Cancer	Control – routine informed consent procedures Group 1 – control plus follow-up phone calls by research nurses (information giving)	Age, sex, level of education, marital status, actual risk status, medical history, anxiety – state, perceived satisfaction, memory, awareness intervention
667 - 1	Participation in decision making in women with breast cancer	Shared consultation	Cancer	Group 1 – leaflet breast cancer treatments and others experience Group 2 – computer interaction, pictures and text	Age, level of education, ethnicity, actual risk status, knowledge, autonomy in decision making
671 - 1	Cancellation surgery in adults with CVD	None	Medicine	Group 1 – information: procedure, side-effects and complications Group 2 – more information risks and complications	Age, sex, level of education, actual risk status, medical history, anxiety – state, disability scale, usefulness of information
1 - 1	Uptake of exercise in well adults with risk factors for CVD	None	Medicine	Control – usual care: GP suggests changes weight, exercise, and smoking Group 1 – control plus attendance at local group session led by a health professional	Age, sex
10 - 1	Use of health care by medicine male patients	None	Medicine	Control – internist recommends specialist to patients when attend primary care Group 1 – evaluation patient's needs, develop personalised care plan, nurse coordinated care from clinic.	Age, level of education, social class, marital status, ethnicity, actual risk status, depression, perception of well-being, disability scale, social support/communication
28 - 1	Use of health services in adult patients with Parkinson's disease	None	Medicine, mental health	Control – usual care Group 1 – propath programme: provide in-depth advice on coping with Parkinson's disease, use leaflet and video.	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, perception of well-being, other affect, perceived satisfaction, disability scale
63 - 1	Uptake of exercise in adult back pain patients	None	Medicine	Control – routine care Group 1 – control plus leaflet information on back pain and coping techniques Group 2 – Group 1 information delivered by nurse plus telephone call for additional information	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, health behaviours-general, depression, perception of well-being, knowledge, perceived satisfaction
47 - 1	Uptake of immunisations and CVD risk factor reduction in well adults	None	Medicine, primary care	Group 1 – health screening by local hospital Group 2 – health screening by physician	Age, sex, level of education, social class, marital status, medical history, health insurance, living arrangements
55 - 1	Cessation of smoking in adult well children	None	Medicine	Control – routine information giving video Group 1 – culturally sensitive video	Age, sex, level of education, social class, knowledge, usefulness of information
55 - 2	Cessation of smoking in adult well children	None	Medicine	Control – discussion group about smoking Group 1 – showing previously developed videos	Age, sex, level of education, social class, ethnicity, self-efficacy, usefulness of information
57 - 1	Adherence to medication in adults with hypertension	None	Medicine	Group 1 – medication once per day Group 2 – medication twice a day	Age, sex, physiological/psychological assessment
88 - 1	Adherence to medication in adult medicine physically ill sample	None	Medicine	Control – medication only Group 1 – medication plus information about side-effects, name, adherence to regimen	Age, medical history, knowledge

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
102 - I	Cessation of smoking in well adults	None	Medicine	Control – smoking status assessed Group 1 – US\$10 incentive plus win 'ex-smoker of the month' money	Age, sex, level of education, social class, actual risk status, intention
105 - I	Adherence to medication in medicine adults	None	Medicine	Control – usual medication regimen Group 1 – same packaging but twice daily pill taking Group 2 – modified information and twice daily dosing	Age, sex, ethnicity, actual risk status, perception of well-being, perception of risk, disability scale
110 - I	Cessation of smoking/adherence to healthy lifestyle in medicine well population	None	Medicine	Control – informed that a programme running and health claims monitored Group 1 – underwent risk appraisal, personalised risk feedback, letters, leaflets Group 2 – underwent risk appraisal only	Age, sex, level of education, actual risk status, health behaviours-general, physiological/psychological assessment, stress
147 - I	Cessation of fatty diet in well adults	None	Medicine	Control – recorded and weighed food Group 1 – group counselling by nutritionist, feedback on fat intake, asked to make changes	Age, sex, actual risk status, attitudes, reasons
155 - I	Consent to participate in well adults	None	Medicine	Group 1 – patient consent to observation only Group 2 – consent to observation plus videotape	Age, sex, level of education, other affect, perceived satisfaction
183 - I	Cessation of alcohol use in well adult women	None	Medicine	Control – newsletters of 'dummy' interests Group 1 – newsletter included articles on alcohol use and abuse	Age, level of education, social class, marital status, ethnicity, actual risk status, knowledge, attitudes, reasons
216 - I	Adherence to cessation of smoking and poor diet in well adults	None	Medicine	Group 1 – letter invitation, leaflet smoking and weight loss programme, incentive (US\$) option and/or flat fee payment to programme Group 2 – invitation letter, programme leaflets sent if returned further information card	Age, sex, level of education, actual risk status, physiological/psychological assessment
217 - I	Cessation of smoking in well adolescents	None	Medicine	Control – no intervention Group 1 – group sessions: smoking, skills to resist smoking, feedback Group 2 – media intervention Group 3 – health information plus Group 1 Group 4 – Group 1 plus Group 2 plus Group 3	Sex, level of education, ethnicity, actual risk status, knowledge, self-efficacy, intention, coping
249 - I	Cessation of smoking in well adults	None	Medicine	Control – routine physician advice Group 1 – routine plus two-part counselling with trained 'smoking' counsellor	Age, sex, level of education, social class, actual risk status, medical history, health behaviours-general, perception of well-being, self-efficacy, intention, motivation, stress
259 - I	Cessation of alcohol use in well adult women	None	Medicine	Control – no additional contact Group 1 – leaflet information about alcohol level screen positive, cessation of, follow-up letter Group 2 – reasons screen positive, asked about drinking habits, monthly consultations	Age, sex, level of education, social class, actual risk status, health behaviours-general, physiological/psychological assessment
279 - I	Cessation of smoking/adherence to healthy lifestyle in well children	None	Medicine	Control – screening only Group 1 – verbal and leaflet information on diet, exercise regimens and smoking ; obese children additional counselling	Actual risk status, physiological/psychological assessment
356 - I	Cessation of salt in diet – adults with hypertension	None	Medicine	Control – no intervention Group 1 – nurses trained information on: addition salt, diet and hypertension, take blood pressure	Age, level of education, marital status
574 - I	Cessation of smoking in well adults	None	Medicine	Group 1 – immediate cessation of counselling Group 2 – tailored cessation of plus placebo gum Group 3 – tapered quitting plus active gum	Age, sex, actual risk status, depression, perception of well-being, other affect
652 - I	Attendance of follow-up cholesterol screening appointment in well adults (mobile unit in car-park)	None	Medicine	Control – routine testing plus leaflet Group 1 – letter plus notification receive US\$17 food coupon on attendance Group 2 – reminder letter to attend	Age, sex, level of education, social class, marital status, actual risk status, medical history, health behaviours-general

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
679 - I	Uptake of exercise in well children	None	Medicine	Control – routine health instruction and assessment Group 1 – trained teacher, leaflets on: diet, smoking and exercise plus extra exercise classes	Age, sex, social class, ethnicity, health behaviours-general, physiological/psychological assessment, living arrangements, knowledge
489 - I	Uptake of cessation of lifestyle change in adults with hyperlipidaemia	None	Medicine	Control – content unclear Group 1 – patient-designed intervention to reduce CVD risk factors delivered by health professionals Group 2 – professional-designed programme carried out by same health professionals, no further information given	Age, sex, level of education, marital status, actual risk status, family history, medical history, health behaviours-general, perception of risk, reasons, self-efficacy, operationalised model
772 - I	Cessation of drinking in adult problem drinkers	None	Medicine	Control – waiting list control Group 1 – directive style consultation: drinking defined as a problem, advice total giving up Group 2 – patient-centred: reflective listening, not labelled alcoholics, not confront problem	Age, sex, level of education, marital status, actual risk status, family history, attitudes
807 - I	Adherence to self-management in adults with medicine	None	Medicine	Control – training programme only Group 1 – training programme plus support groups	
447 - I	Adherence to medication in adults with hypertension	None	Medicine	Control – health education package Group 1 – letter before appointment, follow-up on non-attendance plus card with clinic details	Age, sex, social class, marital status, medical history, health behaviours-general, physiological/psychological assessment
783 - I	Cessation of smoking in well adults	None	Medicine	Control – enhanced usual care: leaflet and access self-help materials Group 1 – leaflet plus self-quitting guidelines: coping techniques plus manual plus no smoking prompts plus feedback cards to return Group 2 – Group 1 plus letter encouraging adoption smoker buddy Group 3 – Group 2 plus additional telephone contact with counsellor	Age, sex, level of education, marital status, ethnicity, actual risk status, medical history, health behaviours-general, attitudes, reasons, self-efficacy, intention
336 - I	Cessation of/adherence to lifestyle CVD factors in well adolescents	None	Medicine	Control – assessment only Group 1 – assessment plus screening Group 2 – know your body information: lifestyle behaviours and changes in diet, exercise, smoking Group 3 – assessment plus screening plus Group 2	Age, sex, ethnicity, physiological/psychological assessment, knowledge, attitudes, social reinforcement/norms
530 - I	Compliance with medication in adults with asthma	None	Medicine	Control – assessment only Group 1 – four asthma management sessions: lectures, leaflet and feedback skills	Age, sex, level of education, marital status, ethnicity, physiological/psychological assessment knowledge
564 - I	Cessation of drinking in adults with drinking habit (workplace)	None	Medicine	Group 1 – hospitalisation plus attendance AA on release Group 2 – attended AA only Group 3 – offered choice of AA, hospitalisation, alternative counselling or no treatment	Age, sex, social class, ethnicity, actual risk status, health behaviours-general, law infringements
702 - I	Adherence to medication in medicine patients	None	Medicine	Control – routine care Group 1 – teaching session plus pill review Group 2 – Group 1 plus follow-up phone call	Age, actual risk status
213 - I	Cessation of smoking in elderly adults	None	Medicine	Control – delayed intervention group Group 1 – physician trained on quitting plus leaflet for older adults (readability) plus letter	Age, sex, actual risk status
327 - I	Adherence to self-care in adults with asthma	None	Medicine	Control – routine verbal plus leaflet information: house mites and asthma Group 1 – control plus computer programme feedback given on responses (piloted)	Sex, level of education, social class, marital status, actual risk status

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
328 - I	Adherence to self-care in adults with medicine	None	Medicine	Control – delayed intervention Group 1 – verbal information by health professional: diet, exercise, self-monitoring, adherence to medication plus skills plus group meetings	Age, sex, actual risk status, medical history, physiological/psychological assessment, self-efficacy
612 - I	Adherence to minor surgical therapy in adults with sleep apnoea	None	Medicine	Group 1 – oral 'snore-guard' used when sleeping Group 2 – use nasal continuous positive airway pressure treatment	Age, sex, actual risk status, physiological/psychological assessment, perception of well-being, attitudes, perceived satisfaction
629 - I	Uptake of advance directives in elderly adults with acute illness	None	Medicine	Control – routine care Group 1 – counsellor information on health proxy, noted in charts	Age, sex, ethnicity, actual risk status, health insurance, physiological/psychological assessment, perception of well-being
268 - I	Cessation of CVD risk factors in men at increased risk CVD and families	None	Medicine	Control – screened but not informed at risk Group 1 – physician letter about family risk by lifestyle plus physician/ dietician counselling plus newsletter prompt	Actual risk status, health behaviours-general, physiological/psychological assessment
395 - I	Uptake of screening/ immunisation in well adults	None	Medicine, cancer, primary care	Control – routine care Group 1 – computer/note prompt for physician Group 2 – computer/note prompt plus patient reminder letter Group 3 – patient reminder letter	Age, sex, ethnicity, health behaviours-general, health insurance
396 - I	Self-management/diet in adults with medicine	None	Medicine	Control – delayed information Group 1 – use of computer and diet package to check diet	Age, sex, social class, actual risk status, medical history, physiological/psychological assessment knowledge, perceived satisfaction
398 - I	Diet/exercise adherence to in adults with medicine	None	Medicine	Group 1 – behaviour therapy plus low calorie diet (1200 kcal per day) Group 2 – behaviour therapy plus very low calorie diet (500 kcal on alternate days)	Age, sex, level of education, actual risk status, medical history, physiological/psychological assessment depression
402 - I	Adherence to treatment in adult alcoholics	None	Medicine	Group 1 – all patients offered same behavioural programme Group 2 – patients randomised to either psychiatric or behavioural therapy	
439 - I	Uptake of CVD screening/ utilisation of service in well elderly adults	None	Medicine	Control – routine Group 1 – staff training, counselling about CVD risks and prevention, Medicaid voucher, free screening	Age, sex, level of education, social class, marital status, actual risk status, health insurance, health behaviours-general, living arrangements anxiety – state, depression, perception of well-being
527 - I	Adherence to medication in adults with chronic obstructive pulmonary disease	None	Medicine, cancer	Control – yearly assessment plus inhaler Group 1 – intensive smoking cessation of, health professional contact and feedback every 4 months plus inhaler with medication or placebo	Age, sex, level of education, ethnicity, actual risk status, medical history
168 - I	Adherence to medication in adults with hypercholesterolaemia	None	Medicine	Control – routine drug allocation Group 1 – drug allocation plus five phone calls	Age, ethnicity, actual risk status, medical history, physiological/psychological assessment
265 - I	Cessation of CVD risks in children with family history CVD	None	Medicine	Control – re-assessed at one year only (and matched) Group 1 – home visits, letter and phone calls: information shopping, cooking, signing diet contracts Group 2 – Group 1 but delayed start intervention	Age, sex, actual risk status, knowledge, health behaviours-general, physiological/psychological assessment
365 - I	Dietary change in well men	None	Medicine	Control – assessment only Group 1 – information from nutritionist – dietary changes	Age, level of education, social class, actual risk status, family history, medical history, health behaviours-general
372 - I	Utilisation of services/ lifestyle change in adults screened positive for cholesterol	None	Medicine	Control – usual care Group 1 – patient level: personalised risk letter, reminder for doctor, fridge magnet Group 2 – doctor received patient pack: letter for patient, result screening test Group 3 – Group 1 plus Group 2	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, family history, physiological/psychological assessment, health insurance

continued



**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
263 - I	Changes in diet in well women at risk for breast cancer	None	Medicine, cancer	Control – assessment only Group 1 – group sessions – weekly sessions with nutritionist	Age, level of education, social class, actual risk status, family history, physiological/psychological assessment
693 - I	Adherence to medication in elderly patients	None	Medicine	Group 1 – leaflet Group 2 – leaflet plus researcher explaining information Group 3 – Group 2 plus medication prompt plus additional home visits Group 4 – Group 3 plus additional information about medication	Age, sex, level of education, physiological/psychological assessment, knowledge
137 - I	Adherence to medication in breast-feeding women requiring antibiotics	None	Medicine, obstetrics and gynaecology (midwifery)	Group 1 – information about safety antibiotics while breast feeding Group 2 – Group 1 plus information about diarrhoea in babies as side-effect	Age, sex
139 - I	Adherence to self-management in elderly adults with medicine	None	Medicine	Control – routine care Group 1 – monthly phone calls reminding clinic appointments, encouragement healthy lifestyle and drug adherence	Age, sex, ethnicity, actual risk status, physiological/psychological assessment perceived satisfaction
143 - I	Cessation of smoking in well adults	None	Medicine	Group 1 – television-based information and leaflet Group 2 – Group 1 plus monthly newsletters plus support line numbers	Age, sex, level of education, social class, ethnicity, actual risk status, self-efficacy, awareness intervention
668 - I	Cessation of smoking	Social cognition model	Medicine	Group 1 – routine smoking leaflet by post Group 2 – 'appropriate' leaflet to stages of change of smoker Group 3 – 'appropriate' leaflet plus report of progress Group 4 – 'appropriate' leaflet plus feedback progress plus phone call	Age, sex, social class, marital status, ethnicity, actual risk status, reasons
332 - I	Uptake of living will in well elderly adults	Framing information (prospect theory)	Medicine	Group 1 – positively framed information: PVS, stroke, dementia and living wills (vignettes readability) Group 2 – Group 1 information, negatively framed Group 3 – Group 1 information, in style of medical guidelines	Age, sex, level of education, religion, living arrangements
20 - I	Use of services in adults with sickle cell	Health promotion	Medicine, sickle cell	Control – disease education information Group 2 – disease information plus training in cognitive coping skills for pain	Age, sex, level of education, social class, marital status, actual risk status, disability scale, coping
27 - I	Adherence to self-care behaviour in adult patients with arthritis	Shared consultation	Medicine, primary care	Control – patients only contacted for assessment Group 1 – symptom monitoring (placebo): asked about symptoms, no mention arthritis Group 3 – counselling group: skills talking with doctor plus clarifying information about disease and medication plus reality therapy focusing on behaviour change	Age, sex, level of education, ethnicity, actual risk status, other affect, disability scale, social support/communication
104 - I	Self-maintenance in adults with medicine	Health promotion	Medicine	Control – routine clinic care Group 1 – compliance: focus on medication behaviours Group 2 – behavioural: focus on behavioural strategies to change behaviour Group 3 – Group 2 plus instruction: focus on behaviour strategies plus write contract	Age, sex, social class, marital status, actual risk status, physiological/psychological assessment
220 - I	Adherence to exercise and healthy diet in well children	Social cognition model	Medicine	Control – school-based health education only Group 1 – Child and Adolescent Trial for Cardiovascular Health (CATCH) programme: specific CVD risk factor information plus feedback over 3 years plus no smoking policy plus alter food available in school	Age, sex, ethnicity, actual risk status, knowledge, attitudes, self-efficacy, social support/communication, intention, social reinforcement/norms

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
233 - I	Adherence to exercise and healthy diet in well adults	Health promotion	Medicine	Control – assessment only: no feedback or information Group 1 – assessment plus face-to-face information plus feedback on adherence to exercise and diet plus phone call	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, medical history, physiological/psychological assessment
241 - I	Adherence to exercise in well adults	Social learning theory	Medicine	Control – leaflet plus video plus lectures: exercise, diet, smoking, use gym Group 1 – tailored self-efficacy theory information: log exercise achieved plus feedback from instructor plus social modelling plus further explanation exercises plus buddy groups	Age, sex, actual risk status, health behaviours-general, physiological/psychological assessment, self-efficacy
276 - I	Cessation of smoking in well adolescents	Health promotion	Medicine	Control – no additional information Group 1 – information given by older students plus skills to resist smoking plus no smoking contract	Sex, actual risk status, knowledge, attitudes, social reinforcement/norms
295 - I	Adherence to diet in adults with hypertension	Health promotion	Medicine	Control – no additional information Group 1 – task-centred instruction (Glaser): less emphasis on knowledge disease more on ways of coping/functioning Group 2 – task centred plus goal setting plus self-monitoring (use contract)	Age, sex, level of education, actual risk status, physiological/psychological assessment, knowledge
305 - I	Adherence to screening/healthy lifestyle in adults with hypertension (workplace)	Lay model illness	Medicine	Control – routine information hypertension: definition, link with disease, importance health checks, reduction lifestyle behaviours Group 1 – action plan message: control plus leaflet plus illustrations on reducing risk factors Group 2 – wellness thinking: routine plus information on experience of hypertension (no symptoms) plus emphasis on health, feeling good, visual information on enjoying exercise	Age, sex, ethnicity, actual risk status, health behaviours-general, physiological/psychological assessment, knowledge, attitudes, self-efficacy, coping
321 - I	Adherence to exercise in well women	Social learning theory	Medicine	Group 1 – exercise instruction plus verbal reinforcement Group 2 – exercise instruction plus verbal reinforcement plus positive cognitive statements	Age, physiological/psychological assessment, anxiety – state/trait, depression, perception of control, motivation, personality type
157 - I	Cessation of smoking in well women with children (effects passive smoking)	Health promotion	Medicine	Control – data collection only Group 1 – social learning theory verbal plus leaflet (piloted): skills to maintain smoke-free environment plus memory prompt plus feedback plus nurse home visits	Age, level of education, ethnicity, actual risk status, medical history, reproductive history, health behaviours-general
376 - I	Attendance at smoking cessation programme in well adults	Social cognition model	Medicine	Control – physician advise to quit plus health counsellor session Group 1 – control plus quitting leaflets Group 2 – control plus video cessation of group from former quitter plus told sessions free plus additional leaflets Group 3 – control plus video plus choice of self-quitting leaflets or group counselling	Age, sex, ethnicity, actual risk status, perception of well-being, self-efficacy, operationalised model, motivation
469 - I	Medication compliance in adults with hypertension	Health promotion	Medicine	Control 1 – new patients: no additional information Group 1 – new patients: medication plus leaflet: importance adhering and lifestyle change plus monthly newsletter Control 2 – existing patients: no additional information Group 2 – existing patients plus Group 1 information	Age, sex, actual risk status
502 - I	Self-management in children with medicine	Framing information	Medicine	Control – small group instructions: lecture plus video plus feedback Group 1 – video plus lecture framed using anchored instructions and context patient's life	Age, sex, actual risk status, knowledge, reasons

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
503 - 1	Uptake of/adherence to lifestyle in adults with medicine	Social cognition model	Medicine	Control – no details Group 1 – nurse counselling: modifying eating/exercise plus monitor calorie intake plus social support plus identify barriers	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, health behaviours-general, physiological/psychological assessment, perception of well-being, knowledge, social support/communication
707 - 1	Cessation of/adherence to lifestyle behaviours in adults following infarction	Social learning theory	Medicine	Control – usual care verbal information: smoking, nutrition and exercise Group 1 – self-efficacy based information: feedback, skill acquisition, additional contact. HCP.	Age, sex, level of education, social class, ethnicity, actual risk status, physiological/psychological assessment
158 - 1	Cessation of smoking in well adults	Social cognition model, attrib. theory	Medicine	Control – generic health letter Group 1 – personalised, theoretically driven leaflet plus signed letter	Age, sex, actual risk status
158 - 2	Cessation of smoking in well adults	Social cognition model, attrib. theory	Medicine	Control – questionnaires only Group 1 – tailored letter and personal details	Age, sex, level of education, actual risk status
210 - 1	Uptake of/cessation of healthy diet in well adults (workplace)	Social learning theory	Medicine	Group 1 – standard healthy eating leaflets Group 2 – information tailored to results preliminary questionnaire	Age, sex, level of education, actual risk status, attitudes, physiological/psychological assessment, usefulness of information, intention
211 - 1	Cessation of smoking in well adults (church based)	Social cognition model	Medicine	Group 1 – 'fair' to provide health feedback self-help leaflet Group 2 – fair, sermons, additional volunteers, audio-tapes, script-guided information	Age, sex, level of education, actual risk status, operationalised model
644 - 1	Cessation of smoking/CVD risk factor reduction in well adults (workplace)	Social cognition model	Medicine	Group 1 – risk assessment and feedback Group 2 – Group 1 plus leaflet and video information: risk factors Group 3 – Group 2 plus behavioural counselling: target setting based on stages change model Group 4 – Group 3 plus financial incentives	Age, sex, level of education, social class, marital status, physiological/psychological assessment
269 - 1	Adherence to diet in elderly well adults	Health promotion	Medicine, cancer	Control – routine letter to double fibre rate Group 1 – leaflet plus free fibre supplement plus group counselling plus newsletters plus recipe competitions Group 2 – letter plus fibre supplement free for 3 months	Age, sex
291 - 1	Adherence to self-management in parents of children with asthma	Social learning theory	Medicine	Control – routine medical care Group 1 – family counselling: self-management, feedback and skill acquisition	Age, sex, ethnicity, worry
541 - 1	Adherence to/cessation of healthy diet in women with breast cancer	Social learning theory	Medicine, cancer	Control – routine nutritional information Group 1 – fat content information common foods plus food diary plus counselled to reduce fat intake	Age, actual risk status, medical history, health behaviours-general, physiological/psychological assessment
255 - 2	Cessation of smoking in well adolescents	Health promotion	Medicine	Control Group 1 – Minnesota smoking prevention campaign (social influences model) Group 2 – shorter version Group 1 Group 3 – smoke free generation programme	Age, sex, actual risk status, social reinforcement/norms
145 - 1	Cessation of smoking in well adolescents	Social cognition model, health promotion	Medicine	Control – assessment smoking behaviour only Group 1 – group lessons: smoking, skill acquisition, changing social norms over 3 years	Age, sex, actual risk status, other affect, perception of risk, social reinforcement/norms, stress, personality type

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
264 - I	Cessation of/adherence to diet and exercise in well adults	Social learning theory	Medicine	Control – exercise intervention programme Group 1 – information about healthy diet and food preparing skills Group 2 – information about healthy diet Group 3 – information weight management and exercise	Age, sex, level of education, actual risk status, health behaviours-general, physiological/psychological assessment
659 - I	Change in diet in well men	Expected utility theory	Medicine	Control – assessment only Group 1 – women attended small group sessions weekly	Age, education, social class, risk status, medical history, health behaviours, perception of well-being, knowledge, attitudes, social reinforcement/norms
622 - I	Adherence to treatment in adults with acute and chronic illness	Social cognition model	Medicine, surgery (A&E)	Control – routine care Group 1 – routine care plus phone call information based on HBM Group 2 – routine care plus HBM clinic intervention Group 3 – routine plus HBM clinic intervention plus phone call	Age, sex, level of education, marital status, ethnicity, actual risk status, medical history, knowledge, perception of risk, perception of susceptibility, reasons, operationalised model
814 - I	Uptake of exercise in well adult women	Social learning theory	Medicine	Control – exercise classes only Group 1 – information based on relapse prevention plus exercise class Group 2 – Group 1 plus incentive	Age, physiological/psychological assessment
490 - I	Uptake of rehabilitation exercise in adults with hip replacements	Stress – coping model	Medicine, surgery	Control – routine 'pre surgery' information Group 1 – routine 'pre-surgery' information plus further information on hip replacement	Age, sex, anxiety – state, perceived satisfaction, waiting time/consultation, usefulness of information, stress
22 - I	Uptake of condoms in well adults	None	STDs (HIV)	Control – no health information Group 1 – house campaigns plus community plus calendars/T-shirts plus posters plus leaflets: HIV, condom use and prevention information	Age, sex, level of education, anxiety – state, knowledge
732 - I	Cessation of risky risk in well adults	None	STDs (HIV)	Control – comparison, delayed intervention community Group 1 – STD information plus leaflets plus medication plus free condoms	Sex, actual risk status, medical history
736 - I	Cessation of risky HIV behaviours/uptake of safer practices in adult drug users	None	STDs (HIV)	Control – waiting list control Group 1 – group education session on HIV Group 2 – group counselling session plus opportunity to be tested	Age, sex, level of education, social class, marital status, actual risk status, medical history, health behaviours-general
341 - I	Cessation of drug use in adult substance users	None	STDs (HIV)	Control – interview plus waiting list Group 1 – information HIV (video, feedback) plus waiting list for test counselling Group 2 – HIV information plus counselling about having test plus test	Age, sex, level of education, marital status, ethnicity, actual risk status, health behaviours-general, knowledge, attitudes, perception of risk
156 - I	Cessation of drug use in adult drug users	None	STDs (HIV)	Control – reward for 'drug free' behaviour delayed 3 months Group 1 – high reward: 4 stars for a prize for drug free behaviour Group 2 – low reward: 8 stars for a prize for drug free behaviour	Age, sex, marital status, ethnicity, actual risk status, physiological/psychological assessment
212 - I	Cessation of drug use in well adolescents	None	STDs (HIV)	Control – school curriculum: might be similar to intervention group Group 1 – Drug Abuse Resistance Education (DARE) project: skill acquisition, drug use lesson-based information	Sex, ethnicity, actual risk status, health behaviours-general, attitudes, social reinforcement/norms
639 - I	Cessation of/adherence to HIV behaviours in well mothers	None	STDs (HIV)	Group 1 – leaflets and opportunity to ask questions Group 2 – video-tape information Group 3 – nurse educator	Age, level of education, ethnicity, reproductive history, worry, knowledge, attitudes
289 - I	Adherence to safe sex/uptake of HIV testing in well young adults	None	STDs (HIV)	Control – assessment plus free condoms Group 1 – anticipatory guidance: lecture plus video: efficacy condoms, individual counselling plus free condoms Group 2 – Group 1 plus individual counselling by peer plus free condoms	Age, sex, social class, actual risk status

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
465 - I	Cessation of/adherence to risky/safe sex in well women	None	STDs (HIV)	Group 1 – peer educator information, video, leaflet Group 2 – health professional trained to provide brief information	Age, ethnicity, knowledge, attitudes
733 - I	Cessation of drug use/ attendance post-natal care in women drug-abusers	None	STDs (HIV), obstetrics and gynaecology (midwifery)	Control – no additional visits Group 1 – health professional visited client for 18 months: support plus information plus problem solving	Age, level of education, marital status, actual risk status, medical history, health behaviours-general, other affect, attitudes, reproductive history, stress
4 - I	Cessation of cocaine/ adherence to therapy sessions in adult cocaine users	Social cognition model, health promotion	STDs (HIV)	Control – 90-min counselling session twice a week Group 1 – 120-min session, five times a week	Sex, level of education, social class, ethnicity, actual risk status, family history, health behaviours-general, personality
24 - I	Intention to use condoms in adult well males	Health promotion	STDs (HIV)	Control – given non-alcoholic drink Group 1 – given alcoholic drink	Age, level of education, marital status, ethnicity, health behaviours, knowledge, attitudes, perception of susceptibility, reasons, efficacy
223 - I	Cessation of drug use in female adolescent users	Health promotion	STDs (HIV), medicine	Control – verbal information: prevalence drug use, teenage development and sexual behaviour Group 1 – group plus individual sessions tailored to drug use, skills acquisition, problem solving	Age, level of education, social class, marital status, ethnicity, actual risk status, reproductive history
226 - I	Cessation of drug use in well adolescents	Health promotion	STDs (HIV), medicine	Control – information: health and social consequences drug use Group 1 – information: consequences drugs plus resistance skills Group 2 – information aimed at beliefs: drug use, norms within school Group 3 – resistance skills plus targeted information	Sex, social class, ethnicity, actual risk status
247 - I	Cessation of drug use in well adolescents	Health promotion	STDs (HIV), medicine	Group 1 – social skills: resist drug taking Group 2 – affect management: focus decision making skills plus stress management Group 3 – social skills plus affect management	Sex, ethnicity, actual risk status, health behaviours-general
277 - I	Cessation of drug use in well adolescents	Health promotion	STDs (HIV), medicine	Control – no additional information Group 1 – health belief model information by teachers: smoking, marijuana and alcohol use, resistance skills Group 2 – Group 1 information plus peer volunteer	Sex, level of education, ethnicity, actual risk status, knowledge, attitudes, perception of risk, self-efficacy, intention, social reinforcement/norms
424 - I	Cessation of/adherence to risky/safe sex in children (school)	Social cognition model	STDs (HIV)	Control – delayed programme Group 1 – video plus classroom information: HIV and skill acquisition plus competitions plus feedback plus training teacher Group 2 – Group 1 plus parents encouraged to attend and help child with homework	Sex, ethnicity, actual risk status, knowledge, attitudes, health behaviours-general, reasons, intention, social reinforcement/norms
478 - I	Cessation of/adherence to risky/safe sex in well adolescents (school)	Social learning theory	STDs (HIV)	Control – routine curriculum Group 1 – information based on social learning theory: contraception plus role-play plus skill acquisition	Sex, education, religion, ethnicity, family history, knowledge, attitudes, self-efficacy, social support/communication, intention, social reinforcement/norms
600 - I	Participation in research/ cessation of substance abuse in well adolescents	Social cognition model	STDs (HIV), medicine	Group 1 – letter plus leaflet plus questionnaire completion plus option to take part in counselling (TRS) Group 2 – Group 1 information plus no choice about counselling	
685 - I	Uptake of condoms in well female adolescents	Social cognition model	STDs (HIV)	Control – family planning professional information: condom use plus demonstration Group 1 – control plus encouraged client to handle condom Group 2 – Group 1 plus feedback plus elicitation clients condom use self-efficacy	Age, actual risk status, reproductive history, knowledge, attitudes, self-efficacy

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
742 - 1	Cessation of/uptake of safer sex in pregnant women	Social learning theory	STDs (HIV)	Control – no additional information Group 1 – general health promotion video plus role play plus peer volunteer Group 2 – Group 1 delivery information but specifically about AIDS	Age, level of education, social class, ethnicity, actual risk status, reproductive history, health behaviours-general, knowledge, perception of risk, social support/communication, intention
492 - 1	Cessation of/adherence to risky sex in well adolescents	Social cognition model, social learning theory	STDs (HIV)	Control – delayed intervention Group 1 – information: HIV and condoms plus overcoming barriers plus skill acquisition	Age, sex, ethnicity, health behaviours-general, knowledge, attitudes, perception of risk, self-efficacy, intention
766 - 1	Cessation of/adherence to risky/safe sex in well adolescents	Health promotion, social learning theory	STDs (HIV)	Group 1 – Group discussion information: HIV and AIDS Group 2 – information plus skills acquisition: decision making and communication plus support and exposure to peers with HIV	Age, sex, actual risk status, health behaviours-general, knowledge, attitudes, reasons, self-efficacy, social support/communication, efficacy
776 - 1	Change of sex behaviour in 'high risk' women	Social cognition model	STDs (HIV)	Control – 'placebo' information: nutrition, child and family care Group 1 – HIV-AIDS information plus skill acquisition plus support	Actual risk status, health behaviours-general, knowledge, perception of risk, social support/communication
165 - 1	Cessation of risky drug use in drug users	Social learning theory	STDs (HIV)	Control – no group sessions Group 1 – Group session information AIDS Group 2 – Group 1 plus additional sessions on skill acquisition	Age, sex, level of education, social class, ethnicity, actual risk status, health insurance, living arrangements, law infringements
632 - 1	Cessation of/adherence to HIV risk behaviours in adult drug users	Social cognition model, social learning theory	STDs (HIV)	Group 1 – leaflets Group 2 – leaflets plus group discussion information: HIV and prevention, skill acquisition plus homework	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, anxiety – state, knowledge, attitudes, perception of susceptibility, self-efficacy, social support/communication, sexual orientation, efficacy
632 - 2	Cessation of/adherence to for drug using adults	Social cognition model, social learning theory	STDs (HIV)	Group 1 – leaflets Group 2 – group discussion plus leaflet information (social learning theory): HIV risks and prevention plus homework	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, anxiety – state, knowledge, attitudes, perception of susceptibility, self-efficacy, social support/communication, alienation
635 - 1	Cessation of risky sex/ drug use in black, male adult drug users	Social cognition model, social learning theory	STDs (HIV)	Group 1 – group information session viewing video plus leaflets Group 2 – psychologist led group session plus leaflet plus skill acquisition plus feedback	Age, education, physiological/ psychological assessment, anxiety – state, knowledge, perception of susceptibility, self-efficacy, social support/communication, efficacy
636 - 1	Cessation of risky sex/ drugs in adult drug users	Social cognition model	STDs (HIV)	Group 1 – HIV information in first week Group 2 – HIV information in second week Group 3 – enhanced theory-based group information plus additional counselling	Age, sex, education, ethnicity, actual risk status, living arrangements, law infringements
525 - 1	Cessation of risky drug use in adult substance users	Social cognition model, health promotion	STDs (HIV)	Control – assessment of risk Group 1 – stages change assessment to tailor cognitive-behaviour intervention	Age, sex, education, social class, marital status, actual risk status, health behaviours-general, living arrangements, reasons, operationalised model
660 - 1	Change in HIV risk factors amongst well adolescents	Social cognition model, social learning theory	STDs (HIV)	Control – assessment only Group 1 – group information guided by theory plus skill acquisition	Age, sex, level of education, ethnicity, knowledge, attitudes, self-efficacy, social reinforcement/norms

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
454 - 1	Changes in HIV risks in well women	Stress – coping model	STDs (HIV)	Control – information: culturally sensitive HIV prevention plus free condoms plus free test plus free bleach Group 1 – control plus more information specific to the woman's problems plus skill acquisition plus leaflet	Age, level of education, social class, marital status, ethnicity, actual risk status, anxiety – trait, depression, worry, knowledge, attitudes, perception of risk, reproductive history, stress, coping
706 - 1	Utilisation of primary care in mothers of children with childhood illnesses	None	Primary care, paediatrics	Control – no information or questionnaire Group 1 – questionnaire plus leaflet (readability plus piloting)	Age, level of education, marital status, reproductive history
6 - 1	Adherence to medicine in physically ill children	None	Primary care	Control – 'official' drug Group 1 – generic or alternative to official drug	Age, sex
6 - 2	Adherence to medication in physically ill children	None	Primary care	Control – official drug Group 1 – generic product, not official	Age, sex
6 - 3	Adherence to medication in physically ill children	None	Primary care	Control – official drug Group 1 – generic, 'non-official' group	Age, sex
32 - 1	Compliance with healthier diet in adult obese women	None	Primary care, medicine	Group 1 – regular food/time: usual food depends on weeks of treatment Group 2 – regular food/weight: usual food depending on reaching target weight Group 3 – stimulus/time: new diet depends on weeks of treatment Group 4 – stimulus/weight: new diet depends on reaching target weight	Age, level of education, actual risk status, medical history, attitudes, perception of control, disability scale, social support/communication
51 - 1	Adherence to medication in children with pneumonia	None	Primary care	Group 1 – medication given a syrup and cup to take medication Group 2 – medication in the form of a syrup and a spoon Group 3 – medication in tablet form Group 4 – medication given as a powdered sachet	Age, sex, level of education, actual risk status, living arrangements
174 - 1	Uptake of exercise and diet in obese women	None	Primary care, medicine	Control – cancer screening plus information plus option to attend delayed intervention Group 1 – group plus one-to-one counselling: weight issues plus ways to change diet/ increase exercise plus feedback plus problem solving plus buddy support system	Age, level of education, social class, marital status, ethnicity, physiological/ psychological assessment, knowledge, self-efficacy
302 - 1	Utilisation of service in well adult parents	None	Primary care	Control – kept diary children's illnesses Group 1 – diary children's illness plus leaflet on management common illness (piloted)	Age, level of education, social class, family history, reproductive history, reasons
589 - 1	Uptake of health screening in well adults	None	Primary care	Group 1 – signed letter invitation with fixed appointment Group 2 – prompt on GPs notes to encourage screening	Health behaviours-general, physiological/psychological assessment, worry, attitudes, usefulness of information
572 - 1	Uptake of immunisation well children	None	Primary care	Control – non-attendees sent reminder letter Group 1 – letter plus leaflet plus phone call	
747 - 1	Uptake of osteoporosis screening in well women	None	Primary care	Group 1 – fixed time letter plus leaflet Group 2 – fixed appointment but needed to confirm plus leaflet Group 3 – letter requiring women to make appointment plus leaflet	Social class
769 - 1	Use of primary care service in patients attending A&E	None	Primary care, medicine	Group 1 – patient in A&E seen by GP Group 2 – A&E patient seen by senior house officer Group 3 – A&E patient seen by registrar	Perceived satisfaction
520 - 1	Utilisation of doctor in parents of ill children	None	Primary care, paediatrics	Control – usual care Group 1 – leaflet on common childhood illnesses and symptoms	Age, level of education, social class, medical history, living arrangements, worry, perception of well-being, perception risk, perception of susceptibility,

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
234 - I	Uptake of influenza vaccine in elderly well patients	None	Primary care	Control – newsletter on vaccine plus results of health status Group 1 – newsletter on vaccine plus free immunisation at hospital Group 2 – newsletter on vaccine plus free immunisation by primary care physician	Age, sex, level of education, marital status, medical history, health insurance
337 - I	Utilisation of physician in adults with coughs	None	Primary care	Control – assessment only Group 1 – leaflet: causes coughs, self-care remedies, when contact physician plus explained by nurse/physician	Age, sex, actual risk status, health insurance
796 - I	Utilisation of primary care in adults with coughs	None	Primary care	Control – no additional information after contact GP Group 1 – doctor explained leaflet in consultation	Age, sex, actual risk status, health insurance
414 - I	Adherence to therapy/diet in obese adults	None	Primary care, medicine	Group 1 – exercise classes plus leaflet weight loss plus group discussion diet plus minimum support: weekly letter and feedback weight loss Group 2 – Group 1 plus 2 plus booster sessions: feedback and skill acquisition plus phone contact	Age, sex, health behaviours-general, physiological/psychological assessment, depression
553 - I	Adherence to exercise in adults with arthritis	None	Primary care	Control – usual treatment: relaxation Group 1 – aerobic walking: exercise to increase heart rate Group 2 – water aerobics: same as Group 1 but in water	Age, sex, ethnicity, actual risk status, physiological/psychological assessment, anxiety – state, depression, perception of well-being, disability scale, social support/communication
408 - I	Uptake of immunisation in well children	None	Primary care	Control – routine information Group 1 – telephone reminder recorded message	Age, sex, ethnicity, actual risk status
19 - I	Uptake of hypothetical treatments in male geriatric patients	Framing information	Primary care	Group 1 – survival data of therapies presented as point estimates Group 2 – survival data presented in the form of a curve	Age, sex, level of education, medical history
167 - I	Adherence to well-child clinic in well parents	Social cognition model	Primary care, paediatrics	Control – no additional information Group 1 – letter reminder informed by HBM Group 2 – postcard reminder prompt	Age, level of education, social class
357 - I	Adherence to medication in women with children acute otitis media	Health promotion	Primary care	Control – routine care: verbal information plus explanation syringe use in Spanish Group 1 – written leaflet (readability score) plus leaflet with pictures plus verbal information	Age, ethnicity, reproductive history
426 - I	Uptake of health screening in well adults	Social cognition model	Primary care, medicine	Control – letter invite to make an appointment Group 1 – letter invite with fixed appointment	Attitudes, perception of risk, perception of susceptibility, reasons, intention, efficacy
506 - I	Participation in consultation in physically ill patients	Communication	Primary care	Control – placebo leaflet: description clinics available Group 1 – leaflet encouraging question asking (readability)	Age, sex, level of education, social class, health insurance, perceived satisfaction, awareness intervention
633 - I	Uptake of flu vaccine in well elderly	Social cognition model	Primary care	Control – nurse visit plus information on safety Group 1 – nurse visit plus vaccine information	Age, sex, actual risk status, perception of well-being
369 - I	Adherence to medication in adults with osteoarthritis	Stress – coping model	Primary care	Control – routine care Group 1 – monthly phone call Group 2 – information within clinic Group 3 – information in clinic plus phone call	Age, sex, level of education, social class, ethnicity, perception of well-being, perceived satisfaction, disability scale, social support/communication
620 - I	Uptake of health check in well adult males	Social cognition model	Primary care, medicine	Group 1 – invitation letter Group 2 – invitation within surgery	Age, worry, attitudes, perception of control, perception of susceptibility, reasons, operationalised model, intention, efficacy

continued



**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
135 - I	Decision to breast feeding in pregnant well women	None	Obstetrics and gynaecology (midwifery)	Control – usual care groups: breast feeding vs. formula feeding plus leaflets on breast pump plus T-shirts plus posters Group 1 – feedback groups: fears, misperceptions and positioning advice plus breast pump and wipes plus free football ticket for husband plus peer counsellor	Knowledge, attitudes, social support/communication, intention
698 - I	Adherence to hormone replacement therapy (HRT) in well women	None	Obstetrics and gynaecology (midwifery)	Control – no HRT Group 1 – HRT that induces cyclical withdrawal menorrhoea Group 2 – HRT induces complete amenorrhoea	Age, medical history, reproductive history, physiological/psychological assessment, perception of well-being, reasons
634 - I	Uptake of osteoporosis screening in well women	None	Obstetrics and gynaecology (midwifery)	Group 1 – open letter: contact clinic to make appointment Group 2 – letter with fixed appointment time	Social class
768 - I	Maintenance of breast feeding in well women	Expected utility theory	Obstetrics and gynaecology (midwifery)	Group 1 – commercial pack containing formula only Group 2 – pack contained breast pump, pads and cream (no formula)	Age, level of education, social class, marital status, religion, ethnicity, actual risk status, attitudes, intention
616 - I	Uptake of breast feeding in well women	Health promotion	Obstetrics and gynaecology (midwifery)	Control – leaflets only Group 1 – culturally sensitive video-tape plus three group discussions	Age, level of education, social class, marital status, ethnicity, health behaviours-general, knowledge, attitudes, reproductive history, intention
403 - I	Uptake of breast feeding in well women	Health promotion	Obstetrics and gynaecology (midwifery)	Control – routine information Group 1 – women phoned by nurses Group 2 – women visited by nurse	Age, level of education, social class, medical history, reproductive history, attitudes, stress
562 - I	Adherence to teeth hygiene in well children	None	Dentistry	Control – no additional information Group 1 – school lecture to parents and children, small group discussion and feedback skills for children Group 2 – lecture information only	Knowledge, reasons
576 - I	Adherence to oral hygiene in well adults	Health promotion	Dentistry	Control – routine dental care Control 2 – routine plus patient phone in results of 'plaque test' Group 1 – additional consultation plus feedback from health professional Group 2 – Group 1 plus checklist of tasks	
758 - I	Uptake of oral hygiene in adults with gingivitis	Social cognition model	Dentistry	Control – routine dental hygiene instructions Group 1 – video-tape information plus feedback session with hygienist	Age, sex, social class, actual risk status, physiological/psychological assessment, self-efficacy, attitudes, perception of risk, intention, social reinforcement/norms, motivation
371 - I	Adherence to dental care in well adolescents	Social cognition model	Dentistry	Control – information about teeth Group 1 – verbal information: teeth, prevention, health and social factors	Knowledge, intention
457 - I	Adherence to dental hygiene in well adults	Social learning theory	Dentistry	Control – toothbrush only Group 1 – electric toothbrush and gel toothpaste Group 2 – shower-based oral hygiene system: toothbrush, paste and mouth rinse	Attitudes, perceived satisfaction
304 - I	Adherence to medication in children screened positive for tuberculosis (TB) bacteria	None	Infectious disease	Control – no additional information Group 1 – telephone call every 3 months to encourage drug adherence Group 2 – nurse visited child's home every 3 months Group 3 – child saw doctor every 3 months	

continued

**TABLE 26 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTb

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
308 - I	Adherence to anti-malaria procedures in well children	None	Infectious disease	Control – assessment only Group I – small group teaching about malaria	Age, actual risk status, knowledge
96 - I	Adherence to medication in adult psychotic patients	None	Mental health	Group I – counselling about schizophrenia with family: one family per clinician Group 2 – counselling about schizophrenia: six families per counsellor	Age, sex, level of education, social class, marital status, ethnicity, medical history, utilisation of services
479 - I	Adherence to medication in adults with schizophrenia	None	Mental health	Control – routine medication Group I – medication plus family counselling: group meetings plus visit by health professional	Age, sex, actual risk status, medical history, physiological/psychological assessment, attitudes, disability scale plus supervision
511 - I	Adherence to exercise/health care/utilisation of in adults with Parkinson's disease	None	Mental health	Control – delayed onset trial Group I – signed physician letter plus leaflet: exercise, diets and side-effects plus bi-monthly feedback by physician	Age, actual risk status, medical history, health behaviours-general, self-efficacy, stress
494 - I	Utilisation of services in adult carers of patients with Alzheimer's disease	None	Mental health	Control – non-active counselling: carers told where to obtain information Group I – counselling on getting support plus obtained information plus counselling with carer plus family plus directive means to encourage attendance	Age, sex, level of education, social class, religion, ethnicity, actual risk status, depression, perception of well-being, perception of risk, disability scale, social support/communication
554 - I	Utilisation of services in adults with mental health disabilities	None	Mental health	Control – usual services of Medicaid beneficiaries Group I – Medicaid beneficiaries enrolled in pre-paid services and advice on selecting services	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, living arrangements anxiety – state, depression, perception of well-being, social support/communication
794 - I	Adherence to medication in adults with depression	Health promotion	Mental health	Control – routine care Group I – trained physicians plus patient watched video plus question form for participation in consultation	Age, sex, level of education, social class, marital status, actual risk status, anxiety – state, depression, other affect, perceived satisfaction
493 - I	Decision confidence/skill in carers of people with Alzheimer's disease	Expected utility theory	Mental health	Control – assessment only Group I – trained to use computer decision programme: information, decision tree, elicitation utilities and best outcome plus computer installation	Age, sex, depression, other affect, perception of control, usefulness of information, alienation
176 - I	Choice of hypothetical medication in adult well patients	Framing (prospect theory)	Other	Group I – choice of two medications for hypothetical disease framed risk figures using either relative or absolute figures.	Age, sex, level of education, actual risk status, medical history, reasons
150 - I	Adherence to wound care in adults with emergency lacerations	None	Surgery (A&E)	Control – leaflet on wound care (readability formula applied) Group I – leaflet plus cartoon illustration	Age, sex, level of education, ethnicity, waiting time/consultation, perceived satisfaction

**TABLE 27** Description of the decision, the interventions, and the measures recorded in studies: RCTs

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
37 - I	Uptake of cervical cancer screening in adult well women	None	Cancer	Control – no additional information Group 1 – video information (culturally sensitive) displayed in waiting area before physician appointments	Age, ethnicity, health insurance
154 - I	Uptake of mammography in well women with family history breast cancer	None	Cancer	Control – letter invitation plus questionnaire Group 1 – written plus audio-tape information: screening, breast cancer plus feedback BSE plus decision making kit (reasons for having mammography: information aid not decision aid)	Age, level of education, marital status, actual risk status, family history, health behaviours-general, attitudes, perception of risk, efficacy
296 - I	Adherence to colorectal screening in well adults	None	Cancer	Control – no additional information Group 1 – physician trained plus verbal information plus screening leaflet plus free kit Group 2 – Group 1 plus reminder letter	Age, sex, actual risk status, health professional measures
377 - I	Uptake of cervical screening in well women	None	Cancer	Group 1 – immediate offer smear by physician, if refused made appointment to re-attend Group 2 – physician offered smear immediately plus discussion about refusal plus encouragement to have test immediately	Age, marital status, actual risk status, medical history, attitudes
412 - I	Uptake of colorectal screening in well adults	None	Cancer	Control – return hemoccult cards in person Group 1 – return in addressed envelope Group 2 – return in stamped addressed envelope	Age, sex, ethnicity, health behaviours-general, health insurance
695 - I	Uptake of mammography in well Asian women	None	Cancer	Control – routine invitations only Group 1 – volunteer attend women's home plus verbal information breast cancer	Age, ethnicity, knowledge
712 - I	Uptake of mammography in well women	None	Cancer	Group 1 – reminder card and appointment date Group 2 – Group 1 plus additional reminder letter Group 3 – reminder letter Group 4 – verbal recommendation to attend for follow-up	Age, reasons, usefulness of information
310 - I	Cessation of smoking in well adults (workplace)	None	Cancer, medicine	Control – placebo video: seat-belt use Group 1 – video: man with lung cancer explains risks Group 2 – Group 1 video plus increase self-efficacy message Group 3 – Group 1 video with 'gruesome' scene taken out Group 4 – information about smoker who had both legs amputated from smoking	Age, sex, social class, perception of risk, intention
752 - I	Colorectal cancer screening/health checks in well adults	None	Cancer, medicine, primary care	Group 1 – hemoccult test posted Group 2 – Group 1 plus letter invite health check Group 3 – invite health check plus explanation hemoccult test Group 4 – invite health check	Age, sex
691 - I	Choice irradiation therapy in women with breast cancer	Shared consultation	Cancer	Control – routine consultation Group 1 – clinician given checklist prompt Group 2 – consultation plus checklist plus women used a decision board	Age, level of education, marital status, actual risk status, knowledge, attitudes autonomy in decision making, preference consultation style, reasons, usefulness of information
248 - I	Attendance for follow-up appointment in adult women with screen-positive cervical smear	Social cognition model	Cancer	Control – routine letter invitation Group 1 – letter plus leaflet explaining cervical smear results	Age, level of education, social class, marital status, ethnicity, medical history, reproductive history, health behaviours-general
354 - I	Uptake of mammography in well women	Social cognition model, health promotion	Cancer	Control – letter about survey plus incentive to return questionnaire Group 1 – letter plus video plus group discussions plus leaflets (piloted and informed by health belief model) plus times mammography	Age, level of education, marital status, ethnicity, actual risk status, family history, health behaviours-general knowledge, attitudes, perception of risk, reasons

continued

**TABLE 27 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTc

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
380 - I	Adherence to colposcopy in women with screen-positive smears	Communication	Cancer	Control – routine phone call to reschedule appointment Group 1 – counselling confronting barriers to attendance	Age, level of education, marital status, ethnicity, reasons
484 - I	Uptake of cervical smears in well women	Social cognition model	Cancer	Group 1 – signed letter plus brief explanation smear plus leaflet Group 2 – free bus tickets mailed plus letter Group 3 – video in waiting room on cervical screening	Age, level of education, marital status, ethnicity, actual risk status, health insurance
507 - I	Uptake of mammography in well adult women	Social cognition model	Cancer	Control – routine leaflet plus letter Group 1 – tailored leaflet (social cognition model): positive framing previous screening experiences Group 2 – Group 1 plus opinion of GP and peer member in glossier leaflet	Age, level of education, marital status, attitudes, perceived satisfaction, self-efficacy, intention, social reinforcement/norms
508 - I	Uptake of cervical screening in adult women with dysplasia/atypia	Expected utility theory	Cancer	Control – routine consultation Group 1 – piloted leaflets explaining about dysplasia/atypia	Age, level of education, marital status, ethnicity, actual risk status, reproductive history, health behaviours-general, health insurance
303 - I	Adherence to medication in elderly patients	None	Medicine	Control – general health information given by nurse in groups Group 1 – control plus leaflet plus personalised prompt of current medication	Age, sex, marital status, actual risk status, knowledge, self-efficacy
320 - I	Cessation of/adherence to healthy lifestyle and attendance clinic in adults with hypertension	None	Medicine	Control – routine clinic visit Group 1 – letter invitation for patient plus relatives plus written and verbal information, hypertension, medication, diet plus feedback	Age, sex, actual risk status, health behaviours-general, physiological/psychological assessment
381 - I	Attendance referral appointment in adults with raised cholesterol (workplace)	None	Medicine	Control – leaflet plus screening plus feedback on raised cholesterol level Group 1 – screening leaflet plus feedback screen positive plus letter of cholesterol figure plus leaflet explaining link between cholesterol and CVD	Age, sex, actual risk status, family history, medical history, health behaviours-general, reasons, intention
471 - I	Cessation of alcohol use in well adults	None	Medicine	Group 1 – self-help leaflet: steps to sensible drinking plus routine therapy session Group 2 – brief summary information in leaflet during routine therapy sessions Group 3 – additional counselling as required	Age, sex, level of education, social class, marital status, actual risk status, physiological/psychological assessment
473 - I	Cessation of smoking in well adults	None	Medicine	Control – no additional information Group 1 – quit for good illustrated leaflets: reasons for quitting, minimising weight increase, people to contact, coping skills Group 2 – quit and win illustrated leaflets: identify problems, structured quitting	Age, sex, level of education, social class, marital status, actual risk status, perception of risk, self-efficacy, usefulness of information, intention, social reinforcement/norms, awareness intervention, coping
475 - I	Uptake of self-care in elderly adults with medicine	None	Medicine	Control – no additional information Group 1 – patients contacted by phone for 4 weeks after discharge, additional information given as needed	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, health behaviours-general, physiological/psychological assessment, knowledge
485 - I	Cessation of alcohol use in well adults	None	Medicine	Group 1 – inpatient rehabilitation Group 2 – advised attend alcoholics anonymous Group 3 – advice from counsellors plus offered either AA or hospital	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, health behaviours-general, depression, perceived satisfaction, stress
763 - I	Adherence to medication in elderly discharged hospital patients	None	Medicine	Control – routine care: medication administered by nurses Group 1 – pharmacist discussed medication patients plus nurse supervised patient self-medication plus memory prompt	Age, sex, marital status, actual risk status, waiting time/consultation, knowledge

continued

**TABLE 27 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTc

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
459 - I	Cessation of smoking in women with children	None	Medicine	Group 1 – leaflet for young mothers on quitting (readability) Group 2 – general giving up smoking leaflet Group 3 – illustrated and coloured leaflets	Age, level of education, ethnicity, actual risk status, reasons, usefulness of information
754 - I	Changes diet in children with high cholesterol	None	Medicine	Control – screened negative controls Control 2 – screened positive control group Group 1 – leaflet information (social cognitive theory): diet and cholesterol plus audio-taped feedback Group 2 – counselling with dietician plus specific dietary instructions plus leaflets plus phone contact	Age, sex, ethnicity, knowledge
708 - I	Changes CVD risk factors in well adults	None	Medicine	Control – health information plus cholesterol test (no feedback) Group 1 – health information plus reading plus feedback of cholesterol	Age, sex, social class
711 - I	Adherence to treatment in adult substance users	None	Medicine	Group 1 – routine care Group 2 – introduction of a follow-up telephone call for patients	Age, sex, social class, marital status
170 - I	Accident prevention in well parents	None	Medicine, other	Control – no additional information Group 1 – counselling by nurse plus leaflets: cessation of smoking, effects passive smoking and prevention home injuries	Level of education, social class
343 - I	Uptake of advance directives in geriatric patients	None	Medicine	Control – notes used for analysis Group 1 – information advance directives, copies forms, reminder card to discuss with physician	Age, sex, social class, marital status, ethnicity, actual risk status, medical history, health insurance perception of well-being, attitudes, reasons
56 - I	Attendance for CVD risk screening in adult well males	Social cognition model	Medicine	Group 1 – invitation to attend free screening for CVD risk factors Group 2 – invitation to attend for screening but charged US\$40	Age, social class, marital status, actual risk status, health behaviours, physiological/psychological assessment
472 - I	Involvement in consultation in adults with diabetes	Shared consultation	Medicine	Control – routine consultation Group 1 – health professional consultation plus decision tree plus active role in treatment information plus skill acquisition plus audio and leaflet home package plus prompt for next consultations	Age, sex, level of education, social class, actual risk status, medical history, physiological/psychological assessment, depression, worry, need cognition, autonomy in decision making, preference consultation style, perceived satisfaction, alienation
72 - I	Cessation of smoking well adults via school-based programme	Social cognition model	Medicine	Control – screening session only Group 1 – screening plus leaflet plus group counselling plus feedback	Age, sex, level of education, marital status, ethnicity, actual risk status, health behaviours-general, health insurance, knowledge, operationalised model
242 - I	Cessation of smoking in well male adults	Social cognition model	Medicine	Control – usual care Group 1 – self-help leaflet tailored to stages of change on smoking	Age, actual risk status, medical history, operationalised model, usefulness information
254 - I	Cessation of smoking in well adults	Social cognition model	Medicine	Control – 'stop smoking' statement plus leaflet Group 1 – self-quit advice statement plus decision to quit video plus quitting aids plus calendar prompt plus leaflet Group 2 – group recruitment: advice statement plus attendance group video plus offer free group plus leaflet Group 3 – Group 1 plus 2	Age, sex, level of education, social class, ethnicity, actual risk status, physiological/psychological assessment, perception of well-being
375 - I	Adherence to exercise in adults with CVD	Social learning theory	Medicine	Control – routine group information on medication, diet and exercise Group 1 – additional control information by health professional plus an exercise contract	Age, sex, actual risk status, knowledge, disability scale

continued

**TABLE 27 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTc

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
792 - I	Cessation of smoking in adolescents	Social cognition model	Medicine	Control – usual lesson Group 1 – media plus school-based information based on social cognition models	Age, level of education, social class, ethnicity, health behaviours-general, attitudes, reasons, social reinforcement/norms
640 - I	Cessation of smoking in well adolescents	Social learning theory	Medicine	Control – no additional information Group 1 – peer volunteer group information: tobacco use plus T-shirts plus skill acquisition plus video plus feedback plus competitions plus letter plus phone booster messages	Age, sex, ethnicity, actual risk status, social reinforcement/norms
287 - I	Cessation of smoking in adult hospital patients	Social cognition model	Medicine	Control – assessment only Group 1 – stage of change assessed and targeted video plus verbal plus leaflet information plus free gum plus telephone contact	Age, sex, level of education, actual risk status, medical history
169 - I	Adherence to medication in elderly patients	Communication	Medicine	Control – routine verbal plus leaflet information Group 1 – leaflet plus contact pharmacist to discuss medication	Age, sex, level of education, actual risk status, medical history, living arrangements, knowledge, disability scale
405 - I	Cessation of smoking in well adolescents	Health promotion	Medicine	Control – assessment only Group 1 – peer volunteer group information on cessation of smoking plus skill acquisition	Sex, knowledge, attitudes
442 - I	Changes in CVD risk factors in well adults (work)	Social cognition model	Medicine	Control – delayed intervention Group 1 – assessed stage of change plus target information: CVD risks plus skill acquisition plus changes workplace	Sex, social class, ethnicity, health behaviours-general
360 - I	Cessation of alcohol, smoking and drug use in well children	Health promotion	Medicine, STDs (HIV)	Control – waiting list control Group 1 – weekly lessons: alcohol, tobacco and marijuana plus skill refusal acquisition plus key-ring prompts plus stickers	Age, sex, level of education, ethnicity, actual risk status, knowledge, attitudes, self-efficacy, social reinforcement/norms
115 - I	Adherence to healthy diet in pregnant women	Social cognition model	Medicine, obstetrics and gynaecology (midwifery)	Group 1 – self-assessment 'quiz' plus leaflet plus shopping list pad plus stickers Group 2 – letter from GP plus leaflet with recipes and stickers	Age, social class, marital status, reproductive history, health behaviours-general, knowledge, attitudes, reasons, intention, social reinforcement/norms
329 - I	Cessation of smoking in pregnant women	Social cognition model	Medicine, obstetrics and gynaecology (midwifery)	Control – routine no smoking information by physician Group 1 – health professional explained no smoking leaflet plus feedback Group 2 – video informed by HBM (piloted) plus leaflet plus opportunity for feedback	Age, level of education, marital status, ethnicity
80 - I	Adherence to healthy lifestyles, cessation of unhealthy lifestyle in coronary artery by-pass graft patients	Health promotion	Medicine, surgery	Control – routine care and assessments Group 1 – routine plus behavioural education programme: goal setting, feedback, additional contact, group exercise sessions	Age, sex, level of education, social class, marital status, physiological/psychological assessment, anxiety – state, depression, perception of well-being
649 - I	Uptake of safer sex in well adults	None	STDs (HIV)	Control – questionnaires only Group 1 – comic of 'real-life' stories plus leaflet plus group discussions plus feedback plus skill acquisition	Age, sex, actual risk status, knowledge, attitudes, intention
684 - I	Uptake of safe sex in adult drug abusers	None	STDs (HIV)	Group 1 – information about HIV Group 2 – enhanced information: emphasis on skill acquisition	Age, level of education, marital status, ethnicity, actual risk status, health behaviours-general, living arrangements, attitudes, perception of susceptibility, self-efficacy, social support/communication, social reinforcement/norms
720 - I	Cessation of HIV risk in adult substance users	None	STDs (HIV)	Control – video and group routine information about HIV Group 1 – control plus additional sessions plus feedback plus skills	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, reproductive history, health behaviours-general, social support/communication, social reinforcement/norms, living arrangements

continued

**TABLE 27 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTc

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
407 - I	Cessation of risky HIV behaviours in adult substance users	None	STDs (HIV)	Control – usual counselling Group 1 – HIV skill information: use condoms, bleaching and risk reduction	Age, sex, level of education, actual risk status
181 - I	Adherence to oral contraception in female adolescents	Communication	STDs (HIV)	Control – routine contraception information Group 1 – additional nurse information: examine barriers to adherence, ways of coping	Age, marital status, actual risk status, attitudes, efficacy
762 - I	Cessation of/adherence to sexual and drug behaviour in adult well women	Social cognition model, framing information	STDs (HIV)	Control – usual public health video (white males in video) Group 1 – video: black women gave information Group 2 – video: black women with more culturally specific message (piloted)	Age, level of education, social class, marital status, actual risk status, health behaviours-general, anxiety – state, knowledge, attitudes, perception of risk, usefulness of information
528 - I	Cessation of risky HIV behaviours in adult drug users	Social cognition model	STDs (HIV)	Group 1 – verbal plus leaflet plus video information (HBM): HIV test, condom and safe drug use Group 2 – Group 1 plus additional follow-up visits plus specialist information	Age, sex, level of education, social class, ethnicity, actual risk status, health behaviours-general, living arrangements
646 - I	Cessation of risky HIV behaviours in Latino adult female drug users	Health promotion, stress – coping model	STDs (HIV)	Control – routine video information: HIV and risk reduction Group 1 – culturally sensitive video plus peer volunteer plus leaflet plus discussion: coping and skill acquisition plus leaflet plus free condoms and bleach	Age, level of education, social class, marital status, religion, actual risk status, health behaviours-general, depression, worry, perception of well-being, knowledge, attitudes, stress, coping, self-esteem
696 - I	Uptake of condoms in adults with STDs	Social cognition model	STDs (HIV)	Control – routine clinic information Group 1 – watched video based on theory reasoned action Group 2 – video plus group discussion	Sex, ethnicity, knowledge, attitudes, perception of risk, self-efficacy
817 - I	Cessation of risky drug use in adult substance users	Social cognition model	STDs (HIV)	Group 1 – information based on HBM, TRA Group 2 – enhanced information (relapse prevention): additional contact plus skill building	Sex, level of education, ethnicity, actual risk status, health behaviours-general, knowledge, attitudes, perception of risk, perception of susceptibility, self-efficacy
749 - I	Adherence to/cessation of risky sex behaviour in well male adolescents	Social cognition model	STDs (HIV)	Control – career planning and opportunities information Group 1 – information HIV, skill acquisition, feedback	Age, level of education, actual risk status, health behaviours-general, knowledge, attitudes, intention, personality type
777 - I	Uptake of face washing in children treated for trachoma	None	Primary care, surgery	Control – treatment trachoma only Group 1 – treatment trachoma plus face-washing information	Age, sex, actual risk status
172 - I	Attendance for appointment in well adults (primary care)	None	Primary care	Control – no additional invitation Group 1 – postcard reminder	Satisfaction, health insurance, health professional measures, awareness of intervention
363 - I	Uptake of immunisation rates in well women with children	None	Primary care	Control – no additional information Group 1 – discussion with nurse and GP plus leaflet (readability) plus reminder letter	Age, ethnicity, reproductive history
470 - I	Uptake of primary care health services in well adults	None	Primary care	Control – routine 'new patient' treatment Group 1 – new patients interviewed by physician and nurse	
512 - I	Uptake of immunisation in well children	None	Primary care	Control – no additional information Group 1 – pre-recorded general reminder plus recall message Group 2 – pre-recorded specific recall plus reminder message	Age, sex, ethnicity, actual risk status, living arrangements
98 - I	Adherence to self-management in adult patients with arthritis	Health promotion	Primary care	Control – saw physiotherapist only Group 1 – physiotherapist plus group sessions, feedback, information on arthritis, coping and problem solving, relaxation	Age, sex, actual risk status, medical history, physiological/psychological assessment, anxiety – state, depression, knowledge, self-efficacy, disability scale
601 - I	Request information/change lifestyle in well adults	Shared consultation	Primary care, medicine	Control – consent form only Group 1 – prompt card asking questions/ brief information CVD	Age, sex

continued

**TABLE 27 contd** Description of the decision, the interventions, and the measures recorded in studies: RCTc

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
601 - 2	Request information/change lifestyle in well adults	Shared consultation	Primary care, medicine	Control – consent only Group 1 – nurse prepared patients for consultation using script to discuss health behaviours	Age, sex
83 - 1	Utilisation of (prenatal) services amongst pregnant women	None	Obstetrics and gynaecology (midwifery)	Control – routine visits from health professionals plus informed study Group 1 – reduced visits: consultation focused on 'prenatal milestones' plus leaflets	Age, medical history, reproductive history, physiological/psychological assessment, health insurance, knowledge, usefulness of information
106 - 1	Attendance postpartum appointment in adolescents	None	Obstetrics and gynaecology (midwifery)	Control – routine care Group 1 – routine care plus teddy bear	Age, marital status, ethnicity, reproductive history, health behaviours-general, health insurance, alienation
788 - 1	Utilisation of postnatal care in new mothers	None	Obstetrics and gynaecology (midwifery)	Control – routine nursery care Group 1 – health professional information: caring new-born plus feedback plus 24 hour phone access plus appropriate utilisation of services	Age, level of education, ethnicity, medical history, reproductive history, anxiety – state, depression, knowledge, social support/communication
481 - 1	Attendance for appointment in adults with TB	None	Infectious disease	Control – no reminder message Group 1 – pre-recorded telephone message with appointment time plus clinic number Group 2 – pre-recorded message plus time appointment plus emphasis government scheme (authority) Group 3 – recorded message plus appointment time plus emphasis importance attending (importance) Group 4 – recorded message plus appointment time plus both authority and importance	Age, sex, ethnicity, attitudes
411 - 1	Adherence to treatment in adults with TB	None	Infectious disease	Control – routine information given by clinic staff Group 1 – additional information given by another professional	Age, sex, level of education, social class, marital status, religion, ethnicity, health insurance, living arrangements, knowledge, attitudes
743 - 1	Uptake of CF carrier screening in pregnant/well adults	None	Genetics	Group 1 – leaflet plus counselling plus offer of stepwise screening Group 2 – leaflet plus counselling plus couple testing only	Age, sex, social class, marital status, family history, anxiety – state, perception of well-being, knowledge, attitudes, perception of risk, reproductive history, social support/communication, intention
790 - 1	Uptake of carrier testing (CF) in well adults	None	Genetics	Group 1 – letter in year 1 Group 2 – letter plus leaflet Group 3 – leaflet at clinic Group 4 – approached researcher plus leaflet plus explanation plus test Group 5 – invite at clinic plus leaflet plus explanation plus test appointment Group 6 – letter year 2	Age, sex, anxiety – state, knowledge
293 - 1	Uptake of infant car seat in well mothers	None	Other, obstetrics and gynaecology (midwifery)	Control – routine care only Group 1 – car seat plus information video Group 2 – car seat plus video plus instructions	Age, level of education, social class, ethnicity
349 - 1	Attendance for sexual-difficulties clinic in well adults	None	Other	Control – sent appointment times Group 1 – questionnaire plus information sheet plus stamped addressed envelope plus appointment	Age, sex, living arrangements, health professional measures
58 - 1	Having donor card in well adults	Social cognition model	Other	Control – no specific information Group 1 – leaflet plus inclusion donor cards by post Group 2 – trained health professionals to participate in meetings and promote donor cards plus media Group 3 – Group 1 plus 2	Age, sex, social class, religion, knowledge, attitudes
324 - 1	Adherence to safety advice in well mothers	Social learning theory	Other	Control – questionnaire only Group 1 – illustrated leaflet (readability)	Age, level of education, ethnicity, reproductive history, knowledge, attitudes, self-efficacy, social desirability
717 - 1	Utilisation of health care in adults with mental health problems	Communication	Mental health – general	Control – appointment letter only Group 1 – letter plus information about services	Anxiety – state, preference consultation style, perceived satisfaction



**TABLE 28** Description of the decision, interventions and measures recorded in concurrent studies

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
44 - 1	Uptake of mammography in adult well women	None	Cancer	Control – community comparison Group 1 – community intervention: posted leaflets plus calendar prompts plus trained health professionals	Age, level of education, social class, family history, health behaviours-general, health insurance, knowledge, perception of risk, awareness of intervention
61 - 1	Uptake of cervical cancer screening in well adult women	None	Cancer	Control – no additional information Group 1 – letter invitation sent to eligible women	Age
203 - 1	Uptake of mammography in adult well women	None	Cancer	Control – comparative community Group 1 – increasing awareness in community (variety methods) plus health professionals (by letter) Group 2 – increase community awareness plus women leaflet posted	Age
243 - 1	Uptake of mammography in adult well women	None	Cancer	Control – no additional programme Group 1 – trained physicians plus leaflet plus community groups plus memory prompts	Age, level of education, social class, actual risk status, family history, health behaviours-general, health insurance, knowledge, attitudes, reasons, social support/communication
246 - 1	BSE in well adult women	None	Cancer	Control – pre- and post-test measurements only Control 2 – annual survey's only Group 1 – trained nurse information at women's community groups plus video plus feedback plus calendar reminder plus skill acquisition plus media Group 2 – Group 1 plus second maintenance reminder programme	Age, level of education, social class, self-efficacy, social support/communication, awareness of intervention
385 - 1	Cessation of tobacco (not smoking) use in well children	None	Cancer	Control – completed measures Group 1 – school-based intervention plus acquisition skills (resisting) plus drug information Group 2 – Group 1 plus parental education plus community leader 'enforcers'	Age, sex, level of education, actual risk status, reproductive history, health behaviours-general, depression, attitudes, perceived satisfaction, intention, social reinforcement/norms
686 - 1	Uptake of mammography in well adult women	None	Cancer	Control – comparison community Group 1 – physician information breast cancer screening plus media plus community campaigns plus cheaper mammograms	Age, level of education, social class, ethnicity, health insurance, health professional measures, knowledge, attitudes, intention, awareness intervention
171 - 1	Uptake of general cancer prevention in well adult women	None	Cancer	Control – no additional information Group 1 – cancer-related consultations made priority plus group discussions: self-examination and prevention plus leaflet plus newsletter	Age, level of education, ethnicity, knowledge, awareness of intervention
205 - 1	Uptake of mammography in well women	None	Cancer	Group 1 – media promotion organised by hospitals Group 2 – community: elected representatives from business and social groups plus show videos plus leaflet distribution	Age, awareness of intervention
205 - 2	Uptake of mammography screening in well women	None	Cancer	Group 1 – community: elected organisers from business and social groups plus show video plus leaflet distribution Group 2 – family practitioner recommend attendance plus notes prompt plus leaflets plus posters in surgery	Age, awareness of intervention
397 - 1	Uptake of cervical/breast screening in well women	None	Cancer	Control – no additional information Group 1 – opportunistically nurse invitation: information plus free screening plus appointment for next day	Age, ethnicity, actual risk status, knowledge, awareness of intervention
546 - 1	Uptake of cervical screening in well women	None	Cancer	Control – routine invitation by national programme Group 1 – control plus letter invitation from GP	Age, medical history, living arrangements

*continued*

**TABLE 28 contd** Description of the decision, interventions and measures recorded in concurrent studies

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
448 - I	Uptake of colorectal cancer screening in elderly well adults	None	Cancer	Control – routine slide show offer plus free hemoccult kit plus diet change Group 1 – control plus elderly framed information plus memory prompts Group 2 – Group 1 plus demonstration hemoccult test using peanut butter	Age, sex, level of education, social class, marital status, ethnicity, family history, medical history, physiological/psychological assessment, perception of well-being, disability scale
418 - I	Uptake of screening tests in well adults	None	Cancer, primary care, medicine	Control – north clinic: no information Group 1 – south clinic: no information Group 2 – north clinic: leaflet on screening and health preventive behaviours (readability)	Age, sex, level of education
386 - I	Uptake of BSE in well women	Health promotion	Cancer	Control – verbal information on placebo topic Group 1 – American Cancer Society (ACS) leaflet and verbal information: BSE, cancer plus model demonstration Group 2 – ACS information plus additional feedback session	Age, level of education, actual risk status, family history, medical history, reproductive history, health behaviours-general, knowledge, efficacy
760 - I	Uptake of mammography in well adult women	Social cognition model	Cancer	Control – assessment only Group 1 – group information (HBM): breast cancer and screening Group 2 – Group 1 information plus arguments for and against plus role model plus planning action plus letter	Age, level of education, social class, marital status, ethnicity, health behaviours-general, knowledge, attitudes, perception of risk, perception of susceptibility, reasons, operationalised model, intention
608 - I	Uptake of mammography in well adults women	Social cognition model	Cancer	Control – usual offer mammography with mobile vans Group 1 – routine offer plus media campaign	Age, level of education, social class, ethnicity, living arrangements, worry, knowledge, attitudes, perception of risk, perception of susceptibility, awareness of intervention
404 - I	Uptake of mammography in well women	Health promotion	Cancer	Control – routine breast screening Group 1 – leaflets plus free screening plus faster feedback reports plus reminder letters plus training staff plus staff feedback	Age, level of education, social class, marital status, ethnicity, family history, health professional measures
446 - I	Uptake of mammography in well women	Social cognition model	Cancer	Control – no information Group 1 – information leaflet plus GP reminder letter plus counselling by phone	Age, level of education, social class, marital status, ethnicity, health insurance
136 - I	Adherence to sun protection in well children	Health promotion	Cancer	Control – assessment only Group 1 – group information plus posters plus free sunscreen Group 2 – Group 1 plus skill acquisition plus leaflets plus continued information from teachers	Age, sex, social class, actual risk status, knowledge, attitudes
237 - I	Adherence to/cessation of to healthy lifestyle in well adults	Social cognition model	Cancer, medicine	Control – assessment only Group 1 – trained group leaders plus leaflet plus verbal group information plus media plus letter plus activities	Age, sex, level of education, social class, marital status, actual risk, health behaviour
180 - I	Uptake of safer sun in well adults	Social cognition model	Cancer	Control – routine information through schools and media, information on wearing protective clothing and sunscreen Group 1 – more intensive programme using more media, message based on theory of identification/attitude confronting, more shady spaces, reduction in sunscreen prices	Age, sex, knowledge, attitudes, perception of risk, social reinforcement/norms, motivation, efficacy
45 - I	Uptake of cholesterol screening in well adults	None	Medicine	Control – no additional information Group 1 – package intervention: media plus setting up network in community plus workplace plus trained health professionals	Age, sex, ethnicity, medical history, health behaviours-general knowledge, perception of risk
82 - I	Health care utilisation in asthma patients	None	Medicine	Control – no exposure to information Group 1 – teaching session on asthma: knowledge and management	Age, sex, actual risk status, medical history, physiological/psychological assessment, knowledge, quality of life

continued

**TABLE 28 contd** Description of the decision, interventions and measures recorded in concurrent studies

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
218 - I	Cessation of smoking in adult smokers	None	Medicine	Control – comparison community Group 1 – trained volunteers plus media plus quit-line plus leaflet plus group sessions	Sex, age, level of education, health behaviour, self-efficacy, social norm, awareness of intervention
222 - I	Cessation of/adherence to healthy/risky lifestyle in well adolescents	None	Medicine	Control – matched no intervention schools Group 1 – teachers screened/feedback	Age, sex, level of education, social class, ethnicity, marital status, medical history, physiological/psychological assessment, health behaviours, attitude
235 - I	Cessation of smoking in well adults	None	Medicine	Group 1 – no restrictions on smoking Group 2 – employees agree smoking contract within workplace Group 3 – smoking ban imposed	Age, sex, level of education, marital status
245 - I	Cessation of smoking in CVD adult patients	None	Medicine	Group 1 – motivational counselling: health and well-being, anticipation quitting problems Group 2 – behavioural counselling: quit date set, abrupt quitting, compliance contract, self-reward and relaxation Group 3 – behavioural counselling plus nicotine fading: gradual quitting	Age, sex, level of education, social class, marital status, actual risk status, health behaviours-general, anxiety – trait, depression, social support/communication, coping
256 - I	Cessation of smoking in well adults	None	Medicine	Control – physician advice only Group 1 – advice plus cessation counselling by physicians (behaviour oriented model) Group 2 – Group 1 plus prescription nicotine gum	Age, sex, marital status, ethnicity, actual risk status, medical history, health behaviours-general, self-efficacy, social support/communication, intention
258 - I	Cessation of/adherence to lifestyle in well adults and children	None	Medicine	Control – Group 1 – intervention 1980–1986 involved school plus community plus media information	Age, sex, level of education, actual risk status, physiological/psychological assessment
260 - I	Cessation of smoking in well adult males	None	Medicine	Control – usual smoking cessation information by physicians Group 1 – intervention by research staff: dosage reduction and coping 'risky' situations	
280 - I	Cessation of smoking in well adults	None	Medicine	Control – television based smoking cessation programme Group 1 – North Karelia intervention plus control plus best 'non smoker' competition Group 2 – control plus competition in Turku community	Sex, level of education, actual risk status, awareness of intervention
283 - I	Cessation of smoking	None	Medicine	Group 1 – group meetings – behavioural strategies to reduce smoking, buddy system, stress management, relaxation, leaflet, gum Group 2 – all information Group 1 plus competition for best 'no smoker'	Age, sex, level of education, social class, actual risk status, stress
284 - I	Cessation of smoking in well adults	None	Medicine	Control – health screen only for smokers and non smokers Group 1 – health screen plus programmes on healthy lifestyle plus attend in own time programmes about diet, smoking, stress, etc. (buddy systems, ways coping smoking cessation, etc.), smoking areas plus no smoking signs	Age, sex, level of education, social class, actual risk status, medical history, perception of well-being
315 - I	Adherence to medication in elderly adults	None	Medicine	Control – medication prompt card, feedback on taking medication (pills from a bottle) Group 1 – medication card, feedback on taking medication and pills in foil packet – dates marked so know if taken pill	Age, sex, actual risk status
390 - I	Cessation of/adherence to healthy lifestyle in well adults (workplace)	None	Medicine	Control – health education plus exercise classes (none attendance) Group 1 – workplace introduced a well publicised fitness room Group 2 – counselling to high risk factor employees plus encouraged to take up exercise Group 3 – counselling plus organised exercise sessions plus buddy system plus fitness facility	Actual risk status, health behaviours-general, physiological/psychological assessment

continued

**TABLE 28 contd** Description of the decision, interventions and measures recorded in concurrent studies

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
504 - I	Uptake of services in parents with asthmatic children	None	Medicine	Control – waiting list control Group 1 – trained GPs plus parent leaflet (piloted) plus consultation content prompt plus feedback	Age, sex, actual risk status, medical history, perception of risk
657 - I	Cessation of smoking in well adults	None	Medicine	Control – smokers exposed to the 'kick the habit' TV programme Group 1 – control plus asked for self-quit leaflet	Sex, level of education, social class, ethnicity, actual risk status, medical history
731 - I	Cessation of smoking in well adolescents	None	Medicine	Control – no planned intervention Group 1 – family smoking education (no details) Group 2 – adolescent focused smoking intervention Group 3 – Group 1 plus 2	Age, sex, social class, family history, health behaviours-general, knowledge, attitudes, perception of control, reproductive history, self-esteem
740 - I	Cessation of smoking in well adults	None	Medicine	Control – cholesterol screening result letter, contact physician Group 1 – control plus media campaign: change diet, smoking cessation, exercise uptake. Group 2 – Group 1 plus managing risk leaflet plus public meetings plus health professional contact	Age, sex, actual risk status, physiological/ psychological assessment, knowledge
779 - I	Cessation of smoking in well adults	None	Medicine	Control – comparison village (monk smoking norm) Group 1 – monk discourages smoking in ceremonies and asks all new monks to give up smoking	Age, sex, level of education, social class, actual risk status
206 - I	Cessation of CVD risk factors in well adults	None	Medicine	Control – national population sample Group 1 – health centre/community/social groups plus leaflets plus meetings plus poster information: CVD change, diet, changes in canteens and shops	Age, sex, physiological/ psychological assessment
215 - I	Cessation of fatty diet in well adults	None	Medicine	Control – comparison community Group 1 – health screenings plus school information plus prompts and healthy changes in grocery stores Group 2 – media information plus changes grocery stores Group 3 – information diet change at community and health centre levels	
409 - I	Change diet in well adolescents	None	Medicine	Control – routine school nutrition information Group 1 – trained teachers: nutrition for life programme	Knowledge, attitudes, health professional measures
312 - I	Change diet in well adults	None	Medicine	Control – comparison community Group 1 – community plus media plus health centres plus workplace plus schools information: reduction CVD risk factors	Age, sex, social class
273 - I	Cessation of smoking in well adolescents	None	Medicine	Control – no additional information Group 1 – media and school-based information plus leaflets plus peer volunteers	Age, sex, ethnicity, living arrangements
394 - I	Uptake of eye checks in adults with medicine	None	Medicine, surgery	Control – interviewed only Group 1 – examined eyes at home plus leaflet plus result letter plus phone call reminder	Age, sex, level of education, social class, marital status, ethnicity, medical history, health behaviours-general, physiological/ psychological assessment, knowledge
194 - I	Intention to drive safely in school children	None	Medicine, other	Control – Group 1 – information plus skills teaching about driving safety	Sex, level of education, knowledge, attitudes
89 - I	Adherence to exercise in adult well population	Social cognition model	Medicine	Control – no information about control Group 1 – pre-intervention group (same workplace as post-intervention group) Group 2 – post intervention group information: feedback risk factors, screening, and CVD risks plus encouraged to attend GP	Age, sex, attitudes, health behaviours-general, physiological/psychological assessment, self-efficacy, intention, social reinforcement/norms
184 - I	Cessation of smoking in well adolescents	Social cognition model	Medicine	Group 1 – school intervention plus video plus peers volunteer information plus skill acquisition Group 2 – media intervention aimed at adolescents plus competition	Age, sex, social class, ethnicity, actual risk status, health behaviours-general, knowledge, social support/communication, awareness of intervention

continued

**TABLE 28 contd** Description of the decision, interventions and measures recorded in concurrent studies

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
253 - 1	Cessation of/adherence to healthy diet in well parents of school children	Social learning theory	Medicine	Control – assessment only Group 1 – trained teachers plus verbal plus leaflet information: change salt, fibre and diet plus parent information plus feedback Group 2 – letter information plus newsletter plus classes plus fridge magnet prompt Group 3 – Group 1 plus 2	Level of education, social class, living arrangements, ethnicity, knowledge, intention, attitude, self-efficacy, social support
345 - 1	Cessation of smoking in well adults	Social cognition model	Medicine	Control – routine clinic advice on smoking cessation of Group 1 – trained physician assessed and targeted information to patient's stages change smoking cessation	Age, sex, ethnicity, actual risk status, health behaviours-general, health professional measures, reasons, operationalised model, awareness of intervention
416 - 1	Adherence to exercise/diet in overweight adults	Social cognition model	Medicine	Control – assessments only Group 1 – self-directed change group information: exercise and diet plus skill acquisition plus feedback problem solving plus support Group 2 – Group 1 plus individual counselling with health professional Group 3 – Group 1 plus individual counselling with peer	Age, sex, level of education, actual risk status, physiological/psychological assessment, health behaviours-general, anxiety – state/trait, perception of control, self-efficacy, social support/communication, motivation, awareness of intervention, stress, coping
440 - 1	Cessation of smoking and CVD risk factors in well adults	Social learning theory	Medicine	Control – no additional information Group 1 – social learning theory based community group leader information: risk factors and skill acquisition plus support groups	Age, sex, level of education, social class, ethnicity, physiological/psychological assessment
209 - 1	Uptake of exercise in well adults	Social cognition model	Medicine	Control – routine doctor information on hepatitis B Group 1 – counselling by doctor (stages of change) on exercise uptake	Age, sex, social class, marital status, ethnicity, actual risk status
631 - 1	Adherence to/cessation of diet in well adults	Social learning theory	Medicine	Control – no media information Group 1 – information informed by SLT: media plus leaflet plus group meetings plus letters plus school-based information plus memory aids	Age, sex, level of education, physiological/psychological assessment knowledge
464 - 1	Uptake of exercise in well adults	Social learning theory	Medicine	Control – assessment only Group 1 – Stanford five city project 6 year dissemination information: verbal plus leaflets plus media plus posters plus leaders community groups	Age, sex, ethnicity, actual risk status, physiological/psychological assessment, knowledge, attitudes, self-efficacy
658 - 1	Cessation of/uptake of lifestyle change in well adults	Social cognition model, social learning theory	Medicine	Control – comparison community Group 1 – school-based, community-based, grocery shop changing, healthy restaurant menus, walking clubs, videos, newspapers	Age, sex, level of education, social class, ethnicity, actual risk status, medical history, health behaviours-general, physiological/psychological assessment, awareness of intervention
726 - 1	Cessation of smoking/adherence to exercise in well adults	Social learning theory	Medicine	Control – assessment only Group 1 – group leaders plus institutions trained about CVD risk and prevention plus introduction screening plus media plus school plus health centres plus communities	Age, sex, actual risk status, medical history, physiological/ psychological assessment, knowledge, attitudes, awareness of intervention
255 - 1	Cessation of smoking in well adolescents	Health promotion	Medicine	Control – comparison community Group 1 – trained teachers school-based intervention	Age, sex, social reinforcement/norms
313 - 1	Adherence to self-management in adults with medicine	Social cognition model	Medicine	Control – assessment only Group 1 – health professional provided information on self-care Group 2 – peer provided information about self-care	Age, sex, actual risk status, anxiety – state, knowledge, attitudes, perception of risk, perception of control, operationalised model, intention, social reinforcement/norms
367 - 2	Cessation of smoking in well adults	Social learning theory	Medicine	Group 1 – quit and win programme Group 2 – attendees of heart check programme	Age, sex, level of education, marital status, actual risk status

continued

**TABLE 28 contd** Description of the decision, interventions and measures recorded in concurrent studies

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
370 - I	Cessation of alcohol use in well adolescents	Social learning theory	Medicine	Control – school-based assessment information only Group 1 – group lessons: alcohol use plus skill acquisition plus booster session	Actual risk status, knowledge, perception of control, social reinforcement/norms
238 - I	Cessation of smoking in well adolescent	Social learning theory	Medicine	Control – comparison community Group 1 – leaflet plus verbal information: risks smoking and skill acquisition plus trained teachers plus ban smoking	
261 - I	Change CVD risk factors in well adolescents	Social learning theory	Medicine	Control – assessment only Group 1 – video plus group information (social learning theory) plus posters plus feedback	Actual risk status, physiological/psychological assessment knowledge
456 - I	Change CVD risks in well elderly adults	Social learning theory	Medicine	Control – assessment only Group 1 – peer-led group information sessions on diet, exercises and stressors	Age, sex, level of education, marital status, medical history, knowledge, self-efficacy
709 - I	Change CVD risk behaviours in well adults	Social learning theory	Medicine	Control – assessment only Group 1 – community leaders diffuse information plus media plus exercise classes	Age, sex, level of education, marital status, self-efficacy, social reinforcement/norms, awareness of intervention
311 - I	Cessation of substance use in well adolescents	Health promotion	Medicine, STDs (HIV)	Control – no additional information Group 1 – information framed by social influences behaviour plus skill acquisition plus media Group 2 – information framed to control affect plus stress management plus decision making	Sex, ethnicity
125 - I	Cessation of drug use, adherence to safe sex in adolescent drug users	None	STDs (HIV)	Control – waiting list control Group 1 – Group format health information HIV plus application problem solving therapy	Age, social class, ethnicity, actual risk status, health behaviours-general, sexual orientation, law infringements
161 - I	Cessation of/adherence to safer sex in adults with HIV	None	STDs (HIV)	Control – counselling for HIV test only (clinic 1) Group 1 – counselling HIV test plus information by trained counsellors on risk reduction (clinic 2)	Age, sex, level of education, marital status, actual risk status, health behaviours-general attitudes
177 - I	Cessation of drug use and risky sex in drug users	None	STDs (HIV)	Group 1 – illustrated, leaflet plus verbal information presented by sex worker on HIV risks and prevention plus free condoms plus transport to health centre Group 2 – Group 1 plus additional contact sex worker and provide feedback on changes	Age, sex, level of education, social class, ethnicity, actual risk status, health behaviours-general, knowledge, perception of risk
392 - I	Uptake of safe sex in well, married women	None	STDs (HIV)	Control – routine government family planning project Group 1 – information: family planning plus child and parental immunisations plus re-hydration therapy for diarrhoea	Age, level of education, actual risk status, reproductive history, health behaviours-general, perceived satisfaction
791 - I	Adherence to/cessation of risky sex in adult well female sex workers	None	STDs (HIV)	Control – routine care Group 1 – video plus leaflet plus group discussion: HIV risk factors tailored sex workers plus free condoms plus posters plus feedback	Age, level of education, ethnicity, actual risk status, medical history, reproductive history, knowledge, self-efficacy
555 - I	Cessation of/adherence to risky sex in well adolescents	None	STDs (HIV)	Control – no additional HIV/STD information Group 1 – HIV/STD information given to classes by nurse plus skill acquisition plus free condoms	Age, sex, actual risk status, living arrangements, knowledge, attitudes
665 - I	Change in sexual practice in well gay adult males	None	STDs (HIV)	Control – routine leaflets plus posters Group 1 – identify group leaders and train about HIV risk factor, prevention and skill acquisition plus place in clubs to disseminate information	Age, ethnicity, knowledge, social reinforcement/norms
579 - I	Cessation of risky sex in adult well women (sex workers)	None	STDs (HIV)	Control – delayed intervention group Group 1 – group plus leaflet plus video information: STD and condoms	Age, level of education, ethnicity, knowledge
748 - I	Utilisation of services in adults with HIV	None	STDs (HIV)	Group 1 – physician actively involved in patient's health care Group 2 – physician unaware patient HIV Group 3 – patients not registered GP Group 4 – mixed group of motivated patients and GPs	Age, sex, social class, ethnicity, actual risk status

continued

**TABLE 28 contd** Description of the decision, interventions and measures recorded in concurrent studies

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
455 - I	Adherence to contraception in well adolescent girls	None	STDs (HIV)	Control – assessment only Group I – trained staff on adolescent behaviours plus introduction consultation protocols	Age, religion, ethnicity, knowledge, perceived satisfaction
71 - I	Cessation of cocaine use, attendance sessions in adult, pregnant cocaine users	None	STDs (HIV), obstetrics and gynaecology (midwifery)	Control – routine collection urine samples plus counselling plus coping Group I – control plus additional management counselling	Level of education, social class, ethnicity, actual risk status, reproductive history, attitudes
486 - I	Reduction pregnancies in well adolescents	Conflict theory (Janis and Mann)	STDs (HIV), obstetrics and gynaecology (midwifery)	Control – baseline measures control group Group I – counselling (Janis and Mann): decision-making skills plus prenatal care plus adoption plus future pregnancies	Age, level of education, social class, marital status, ethnicity, actual risk status, reproductive history, self-efficacy
17 - I	Cessation of risky sex, adherence to condoms in school children	Social cognition model	STDs (HIV)	Control – question and answers on AIDS Group I – presentation by person with AIDS Group 2 – role-play (feedback) plus skill acquisition	Knowledge, attitudes
18 - I	Cessation of steroids, male school children	Health promotion	STDs (HIV)	Control – delayed intervention Group I – Adolescents Training and Learning to Avoid Steroids (ATLAS) classroom information: diet and alternatives to steroids plus skill acquisition plus weight room plus incentives	Level of education, social class, ethnicity, knowledge, attitudes, self-efficacy
240 - I	Uptake of condoms/cessation of risky sex in well adolescents	Social cognition model	STDs (HIV)	Control – no additional HIV information Group I – trained teachers skill-based information plus feedback plus video plus skills acquisition	Age, sex, ethnicity, actual risk status, health behaviours-general, knowledge, attitudes, perception of risk, intention
251 - I	Cessation of drug use in well adolescents	Health promotion	STDs (HIV), medicine	Control – no additional information Group I – school and community based information plus acquisition skills plus norms non-drug use plus media plus peers volunteers	Age, level of education, social class, ethnicity, actual risk status, health behaviours-general, living arrangements
391 - I	Uptake of safe sex in adult well males	Health promotion	STDs (HIV)	Control – comparative community Group I – male motivation project plus leaflets plus radio drama plus group discussion	
515 - I	Cessation of risky sex in well adolescents	Social cognition model	STDs (HIV)	Control – delayed intervention Group I – trained teachers group information (theory planned behaviour) plus feedback plus skill acquisition plus parental involvement plus T-shirts	Age, sex, health behaviours-general, living arrangements, knowledge, attitudes, social support/communication, social reinforcement/norms, awareness of intervention
759 - I	Adherence to safe sex in well adolescents	Social learning theory	STDs (HIV), medicine	Control – routine sex education – matched schools Group I – verbal group information on puberty, contraception, reproduction and relationships and empowerment skills	Sex, level of education, social class, actual risk status, health behaviours-general, living arrangements, knowledge, attitudes, usefulness of information, social reinforcement/norms
628 - I	Cessation of/adherence to risky sex	Health promotion	STDs (HIV)	Control – usual information for runaways Group I – additional counselling plus video: HIV risk and prevention plus feedback plus skills	Age, sex, ethnicity, actual risk status, sexual orientation
780 - I	Cessation of/adherence to risky/safe sex in well adults	Social learning theory	STDs (HIV)	Control – comparison community Group I – comics plus newsletters plus free condoms plus house call information on HIV risk factor cessation	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, reproductive history, knowledge, self-efficacy
463 - I	Cessation of risky sex in gay well males	Health promotion	STDs (HIV)	Control – delayed onset intervention Group I – leaflets plus posters in bars plus university and community areas	Age, level of education, marital status, ethnicity, actual risk status, sexual orientation, self-efficacy, awareness of intervention
663 - I	Adherence to/cessation of risky/safe sex in well adolescents	Health promotion	STDs (HIV)	Control – assessment only Group I – trained teachers group lessons plus leaflet plus audio-tape plus skill acquisition	Age, sex, level of education, social class, religion, health behaviours-general, knowledge, attitudes, reasons

continued

**TABLE 28 contd** Description of the decision, interventions and measures recorded in concurrent studies

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
373 - I	Change sexual activity in well adolescents	Social cognition model	STDs (HIV)	Control – routine sex information Group 1 – verbal plus leaflet plus group information (HBM/SLT) plus skill acquisition plus feedback	Age, sex, ethnicity, reproductive history, knowledge, attitudes, perception of susceptibility, reasons, self-efficacy, operationalised model
453 - I	Cessation of drug use in well adolescents	Social learning theory	STDs (HIV), medicine	Control – assessment only Group 1 – peer volunteer verbal information plus skill acquisition Group 2 – Group 1 plus booster sessions	Age, sex, ethnicity, actual risk status, living arrangements, health behaviours-general, knowledge, attitudes, social reinforcement/norms
822 - I	Cessation of risky sex in well adolescents	Social cognition model, social learning theory	STDs (HIV)	Group 1 – brief verbal information about safe sex (school clinic) Group 2 – peer volunteer verbal information: HIV and prevention pregnancy (school clinic) Group 3 – pregnancy prevention information plus vouchers for contraception (school clinic) Group 4 – target high-risk groups plus provision contraception plus prompt repeat prescriptions (school clinic) Group 5 – pregnancy prevention information plus contraception prescribed plus prompt for repeat prescription (school clinic) Group 6 – all sex-related information available plus school dispensed contraception	Age, sex, reasons, awareness intervention
823 - I	Change CVD/HIV risk factors in well adolescents	Health promotion	STDs (HIV), medicine	Control – assessment only Group 1 – school based module on teenage health plus skill acquisition	Age, sex, health behaviours-general knowledge, attitudes
813 - I	Change in CVD risk behaviours in elderly adults	None	Primary care, medicine	Control – assessment only Group 1 – letters plus leaflets plus newsletter plus feedback CVD risk from physician	Age, sex, level of education, actual risk status, health behaviours-general, physiological/psychological assessment
146 - I	Uptake of influenza vaccination in well elderly adults with a history pneumonia	None	Primary care	Control – cases documented before intervention Group 1 – cases documented after intervention: trained physicians plus reimbursed for vaccine plus patient letter plus media plus leaflet	Age, sex, actual risk status, health behaviours-general
257 - I	Adherence to health checks in elderly well adults	None	Primary care, medicine	Group 1 – letter invitation Group 2 – letter plus phone call Group 3 – letter plus phone call plus fixed appointment time Group 4 – Group 3 plus elicited reasons non-attendance plus encouraged attendance	Age, sex, level of education, medical history, physiological/psychological assessment, health insurance
775 - I	Uptake of immunisation in parents of well children	None	Primary care	Group 1 – children registered with general practice Group 2 – children registered with health clinics	Age, living arrangements
331 - I	Uptake of immunisation in parents of children attended A&E	None	Primary care	Control – children attended A&E not immunised Group 1 – children attended A&E immunised	Age, sex, actual risk status, physiological/psychological assessment, health insurance
95 - I	Adherence to self-management in adults with arthritis	Social learning theory	Primary care	Control – no additional information Group 1 – group counselling information session for over 6 weeks	Age, sex, level of education, actual risk status, physiological/psychological assessment, self-efficacy, disability scale
3 - I	Uptake of breast feeding in adult well women	None	Obstetrics and gynaecology (midwifery)	Control – no additional information Group 1 – trained health workers gave verbal information breast feeding plus posters plus leaflets at routine post-natal care visits	Age, level of education, marital status, religion, ethnicity, reproductive history, knowledge, perception of risk
529 - I	Uptake of xylitol chewing gum in adolescents	None	Dentistry	Control – no information Group 1 – xylitol and oral hygiene information by dental assistance plus free chewing gum	Sex, knowledge
618 - I	Uptake of dental services in well children and young adults	Marketing model	Dentistry	Control – no change in dentistry practices Group 1 – reimbursement costs for children/adults under 26 years	Age, sex, level of education, health insurance

continued



**TABLE 28 contd** Description of the decision, interventions and measures recorded in concurrent studies

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
430 - I	Reproductive decision in Tay-Sachs carriers	None	Genetics	Group 1 – genetic counselling information Tay-Sachs and reproductive options	Marital status, ethnicity, attitudes, perception of risk, affect, intentions, autonomy decision making
516 - I	Uptake of carrier testing in well adults	Social cognition model	Genetics	Group 1 – letter invitation plus if interested counsellor appointment plus leaflet Group 2 – approached patients in health centre plus leaflet plus counselled and tested	Age, sex, level of education, marital status, ethnicity, reproductive history, knowledge, attitudes, perception of risk, need cognition/certainty, need cognitive closure/ambiguity, intention
323 - I	Adherence to medication and attendance clinic in symptomatic adults	None	Infectious disease	Control – no additional information Group 1 – social network group, researchers attend community group meetings: information plus offer free health centre treatment Group 2 – media group: posters plus town criers give same information	Age, sex, social class
586 - I	Adherence to treatment in adolescents with TB	None	Infectious disease	Group 1 – directly observed preventive therapy taken within school Group 2 – hospital based medication (not observed)	Age, sex, ethnicity
452 - I	Adherence to treatment in patients with TB	None	Infectious disease	Routine – routine patient care for TB Group 1 – staff training and supervision on TB	Age, sex, actual risk status, medical history, living arrangements
124 - I	Adherence to medication in adults with schizophrenia	None	Mental health	Control – remained on oral medication Group 1 – patients changed from oral to injecting medication	Sex, ethnicity, actual risk status
825 - I	Adherence to medication in adults with schizophrenia	None	Mental health	Control – no information about control group Group 1 – only told group meetings	Actual risk status, medical history, attitudes, reasons
335 - I	Adherence to medication in parents of children with diarrhoea	None	Paediatrics	Control – interviewed only Group 1 – posters plus demonstration how to mix salt-sugar solution plus feedback	Age, level of education, social class, religion, reproductive history, knowledge
570 - I	Adherence to safety in well adolescents	None	Other	Control – assessment only Group 1 – peer presented information: accidents, disabilities and risk taking Group 2 – Group 1 information given by health professional	Age, sex, level of education, actual risk status, medical history, knowledge, perception of risk, perception of control, reasons, self-efficacy, intention, awareness of intervention
773 - I	Uptake of bicycle safety in well children plus parents	None	Other	Control – letter to parents and children plus teen magazine plus voucher for bike shop Group 1 – control plus high profile in school: posters, stickers, stunt bike rider demonstration	Age, sex, social class, family history, health behaviours-general, reasons, knowledge, attitudes, intention, social reinforcement/norms
40 - I	Reporting symptoms in adult women with bladder complaints	None	Surgery	Control – paper-pen diary of symptom recording Group 1 – computer symptom reporting	Age, ethnicity, actual risk status, depression, other affect
75 - I	Use of health care in adult well patients	None	Surgery	Group 1 – medical assistance and no health insurance Group 2 – private insurance	Age, sex, social class, marital status, health insurance attitudes

**TABLE 29** Description of decision, intervention and measures recorded in studies: before/after different samples

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
34 - I	Uptake of mammography in adult well women	None	Cancer	Group I – prompt on patient's notes	
100 - I	Uptake of mammography in well women	None	Cancer	Group I – letter of invitation	Age, actual risk status, health insurance
192 - I	Uptake of cervical screening in adult well women	None	Cancer		Age
383 - I	Uptake of mammography in well based adult women	None	Cancer	Group I – reduced-fee mammogram plus media plus presentations in workplaces/health centres	Age, ethnicity, social class, level of education, medical history, knowledge, awareness of intervention
421 - I	Uptake of mammography/BSE in well women	None	Cancer	Group I – health professional encouragement plus monthly clinics plus calendar prompts plus self-referral contact number	Age, level of education, social class, attitude
575 - I	Uptake of cervical screening in well women	None	Cancer	Group I – introduction mobile screening unit plus information	Age, marital status, level of education, social class, knowledge
401 - I	Uptake of mammography in well women	None	Cancer	Group I – trained staff plus reminder system implemented plus prompt sheet in reviewed notes plus noted women's breast cancer history	Age
765 - I	Adherence to sun protective behaviours in well adults	Social learning theory	Cancer	Group I – poster plus leaflet: skin cancer and protection plus feedback plus peer volunteers plus raffle plus free sunscreen	
495 - I	Uptake of living will in recently hospitalised adults	None	Medicine	Group I – leaflet on patient self-determination act and information on living will	Age, sex, level of education, social class, ethnicity, knowledge, attitudes
340 - I	Cessation of smoking in well adults	None	Medicine	Group I – community intervention: media plus posters plus leaflets plus volunteers plus free cessation classes plus prize-draw	Age, sex, level of education, ethnicity, knowledge, awareness of intervention
423 - I	Cessation of smoking in drug dependent adults	None	Medicine, STDs (HIV)	Group I – change in hospital policy to no-smoking zone plus patients sign no-smoking contract	
550 - I	Cessation of drug use in elderly substance using men	None	Medicine, STDs (HIV)	Group I – introduction no-smoking ban in hospital	Age, level of education, social class, marital status, actual risk status, medical history, health behaviours-general
514 - I	Cessation of smoking in pregnant well women	None	Medicine, obstetrics and gynaecology (midwifery)	Group I – media campaign plus trained professionals plus quitline: emphasis quitting not cutting down	Age, social class, level of education, reproductive history, knowledge, attitudes, awareness of intervention
461 - I	Cessation of drink-driving in well adults	None	Medicine, other	Group I – media campaigns increase seat belt use, maintain speed limits, reduce drunk driving, fines	Awareness of intervention
462 - I	Cessation of/adherence to CVD risk factors in well adults	Social cognition model, social learning theory	Medicine	Group I – information based social learning theory: group leaders plus increase exercise clubs plus health fairs plus health checks plus school competitions	Age, sex, level of education, ethnicity
670 - I	Uptake of HIV screening in well adults	None	STDs (HIV)	Group I – all media campaigns over 4-year period	Sexual orientation
680 - I	Uptake of safe sex/cessation of risky sex in adults with STDs	None	STDs (HIV)	Group I – media campaigns advocating condom use	Sex, ethnicity, sexual orientation, medical history, social class

continued

**TABLE 29 contd** Description of decision, intervention and measures recorded in studies: before/after different samples

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
662 - I	Adherence to safer sex in adults with STDs	None	STDs (HIV)	Group I – announcement 'magic johnson' (basketball player) as HIV positive	Age, sex, level of education, marital status, ethnicity, actual risk status, health behaviours-general, reasons, living arrangements knowledge, attitudes, perception of risk, perceived satisfaction
142 - I	Change HIV risks in adult substance users	None	STDs (HIV)	Group I – introduction of sale syringes in pharmacies within France	Age, sex, level of education, social class, marital status, ethnicity, actual risk status, medical history, reproductive history, health behaviours-general, living arrangements, knowledge
619 - I	Change HIV behaviours in well adult men	None	STDs (HIV)	Group I – media plus posters plus group information and feedback sessions	Age, social class, reasons, living arrangements, sexual orientation
647 - I	Cessation of risky sex in well gay men	Health promotion	STDs (HIV)	Group I – 'trend-setter' peer trained to give information in clubs, identified by eye-catching badge to start conversations	Age, sex, ethnicity, knowledge, social norms
193 - I	Uptake of immunisation in mothers of new-borns	None	Primary care, paediatrics	Control – noted attendance at clinics Group I – leaflets about childhood immunisations	Age, level of education, social class, ethnicity, reproductive history, health behaviours-general, memory, reasons
767 - I	Uptake of immunisation in parents of well children	None	Primary care	Group I – region-based intervention: feedback non attendees to health professional plus trained health professionals	Deprivation indices
413 - I	Uptake of immunisation in well adults (high-risk influenza)	None	Primary care	Group I – introduction of immunisation status in notes	Reasons
623 - I	Uptake of breast feeding in well women	None	Obstetrics and gynaecology (midwifery)	Group I – introduction new staff position for baby feeding and changes on ward	Perceived satisfaction
26 - I	Breast feeding in adult well women	Marketing model	Obstetrics and gynaecology (midwifery)	Group I – trained health professionals plus leaflets and video in clinic increased profile breast screening plus prompts in notes	Age, social class, religion, reproductive history, health insurance, perception of risk, reasons
97 - I	Adherence to dental hygiene in well adults	None	Dentistry	Group I – large public campaign media plus posters plus leaflets	Knowledge
737 - I	Attendance for dental check ups in well adults	None	Dentistry	Group I – verbal information on importance periodontal care plus made fixed appointments	
563 - I	Utilisation of services and uptake of dental treatment in well children	None	Dentistry	Group I – changed examination/treatment from attending fixed clinic to attending mobile unit	Age, ethnicity
239 - I	Adherence to sleeping position baby in adolescent mothers	None	Paediatrics	Group I – home visit midwife information plus feedback plus telephone contact	Age, marital status, level of education, social class, health behaviour, physiological/psychological assessment, reasons, knowledge
294 - I	Utilisation of services plus testing for fragile X in well adults	None	Genetics	Group I – dissemination fragile X video and leaflet information by agencies	Usefulness information
674 - I	Attendance for surgery in women undergoing hysterectomies	None	Surgery	Group I – introduction nurse information at pre-operative assessment clinic	Satisfaction, anxiety
149 - I	Uptake of bicycle helmet in school children	None	Other	Group I – media plus posters plus leaflets plus community events plus verbal information plus feedback plus reduction price helmets plus changing roads for bike paths	Age, sex

**TABLE 30** Description of decision, intervention and measures recorded in studies: before/after same sample

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
99 - I	Uptake of mammography in adult well women	None	Cancer	Group counselling with feedback and ways to overcome barriers to screening (devised a board game)	Age, knowledge, attitudes
266 - I	Adherence to safe sun in adults with skin cancer	None	Cancer	Verbal plus leaflet information on sun protection following removal of non-melanoma skin cancer plus feedback own behaviour	Age, sex, ethnicity, medical history, actual risk
746 - I	Uptake of breast and cervical screening in well adult women	None	Cancer	Trained volunteers plus community workshops plus posters plus media plus video	Age, sex, ethnicity, knowledge, reasons
815 - I	Uptake of testicular self-examination in well adolescents	None	Cancer	Leaflet plus lecture from health professional on testicular self-examination	Knowledge, attitudes
795 - I	Uptake of cervical screening in well women	None	Cancer	GP contract plus computerised systems	Age
809 - I	Uptake of mammography in well women	None	Cancer	One-to-one counselling plus trained health professional plus posters plus leaflets plus media plus prompt on notes plus reminders	Age, health insurance
133 - I	Adherence to/cessation of healthy lifestyle in school children	None	Cancer, medicine	Schools funded projects, trained teachers information: heart disease, cancer nutrition and exercise programmes	Actual risk status, health behaviours-general, physiological/psychological assessment, knowledge, attitudes
466 - I	Uptake of screening/immunisation in well adults	None	Cancer, primary care, medicine	Group I – prompt in notes for GP: screenings and immunisations	Age, sex, ethnicity, health insurance
46 - I	Uptake of mammography in adult well women (family history breast cancer)	Self-regulation theory	Cancer	Group I – targeted women family history breast cancer: diet, BSE, and mammography	Knowledge, attitudes, perception of risk
126 - I	Uptake of mammography and BSE in well women	Expected utility theory	Cancer	Group I – leaflets about mammography and BSE	Age, level of education, ethnicity, attitudes, perception of susceptibility, efficacy
198 - I	Adherence to safe sun in adults with skin cancer	Social cognition model	Cancer	Group I – trained volunteer plus leaflet information skin care	Age, sex, level of education, social class, health behaviour, knowledge, attitudes, self-esteem, perception of control, perception of risk
42 - I	Utilisation of services and adherence to medication	None	Medicine	Patient chart with personalised medication details, symptoms and course of action	Actual risk status, health behaviours-general
74 - I	Adherence to medication adult patients with medicine	None	Medicine	Self-management educational control programme	Age, sex, physiological/psychological assessment knowledge
79 - I	Adherence to lifestyle in school children	None	Medicine	Video information: diet, exercise, smoking cessation plus additional exercise activities	Age, sex, level of education, ethnicity, actual risk status, family history, health behaviours-general, physiological/psychological assessment, attitudes, perception of risk, stress
111 - I	Completion advance directives in patients with haemodialysis	None	Medicine	Leaflet on advance directives plus opportunity to talk with nurse	Age, sex, level of education, marital status, religion, ethnicity, actual risk status, medical history, knowledge, attitudes, reasons
244 - I	Uptake of exercise in well adults	None	Medicine	Trained physicians plus leaflet plus verbal information exercise plus follow-up phone call plus note prompt	Age, sex, level of education, social class, marital status, living arrangement, health insurance, social norms, self-efficacy

continued

**TABLE 30 contd** Description of decision, intervention and measures recorded in studies: before/after same sample

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
270 - I	Smoking cessation of in well adults	None	Medicine	Group I – verbal information plus group sessions plus self-monitoring plus skill acquisition	Age, sex, ethnicity, level of education, social class, affect, reasons, perceived satisfaction
355 - I	Uptake of exercise in well men (worksite)	None	Medicine	Group I – free exercise classes plus group activities plus feedback plus prize for weight loss	Age, level of education, social class, ethnicity, marital status, health behaviour, physiological/ psychological assessment
384 - I	Uptake of screening, cessation of/adherence to healthy lifestyle in well adults (workplace)	None	Medicine	Group I – free screening plus health professional feedback on blood pressure, cholesterol levels and lifestyles changes	Age, sex, ethnicity, social class, family history, medical history, physiological/ psychological assessment
521 - I	Self-care in elderly adults with medicine	None	Medicine	Group I – group counselling plus self care skill acquisition for medicine	Physiological/psychological assessment, knowledge, affect, coping
676 - I	Self-management in adults with asthma	None	Medicine	Group I – self-management 'credit card' to record peak flow rates and other self-management behaviours	Age, sex, ethnicity, actual risk status, physiological/ psychological assessment
522 - I	Adherence to self-management in adults with asthma	None	Medicine	Group I – asthma action plan: checklist for patients and GPs to assess together plus media	Age, sex, physiological/psychological assessment
544 - I	Adherence to self-management in adults with asthma	None	Medicine	Group I – self-monitoring skills plus information asthma plus summary management plan	Age, sex, social class, actual risk status, medical history, knowledge
262 - I	Smoking cessation of in well adults (workplace)	None	Medicine	Group I – smoking cessation information plus buddy system plus gum	Age, sex, social class, ethnicity, actual risk status, physiological/ psychological assessment
449 - I	Adherence to medication in children with asthma	None	Medicine	Group I – leaflet plus diary card plus feedback on medication within daily routine	Age, sex, medical history
798 - I	Change in CVD behaviours in well adult males	None	Medicine	Group I – information about CVD risks and screening	Age, actual risk status, health behaviours-general, physiological/ psychological assessment
821 - I	Uptake of exercise/cessation of smoking in adults with CVD	None	Medicine	Group I – information CVD risk and prevention plus exercise programme	Age, sex, actual risk status, medical history, physiological/psychological assessment
621 - I	Uptake of weaning food in children of well adults	None	Medicine, obstetrics and gynaecology (midwifery)	Group I – piloted and culturally appropriate information on making weaning food, groups, skill acquisition	Age, level of education, social class, actual risk status, reproductive history, reasons, knowledge, attitudes, awareness of intervention
348 - I	Uptake of/cessation of lifestyle and osteoporosis screening in women	None	Medicine, primary care	Group I – leaflet plus opportunity group information plus free screening	Age, level of education, social class, family history, health behaviours, health insurance knowledge
162 - I	Adherence to healthy diet in elderly, physically ill adults	None	Medicine, primary care	Group I – nurse education programme plus memory prompts plus media plus change menu in residential home plus social reinforcement plus incentive game	Age, sex
84 - I	Cessation of drink-driving in adolescents	None	Medicine, other	Nurse information on car crashes injuries plus interactive sessions on prevention	
594 - I	Adherence to medication in adults with hypertension	Framing information, shared consultation	Medicine	Group I – interactive video, content depends on disease type: mild hypertension, alter risk through lifestyle change; BPH, encouraging systematic decision making of all alternatives	Age, sex, level of education, actual risk status, medical history, physiological/ psychological assessment, worry, attitudes, perception of risk, need cognitive closure, autonomy in decision making, usefulness of information

continued

**TABLE 30 contd** Description of decision, intervention and measures recorded in studies: before/after same sample

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
179 - I	Cessation of smoking in adults with asthmatic children	Health promotion	Medicine	Group I – counselling sessions advising patients risk smoking plus coping strategies	Actual risk status
367 - I	Cessation of smoking in well adults	Social learning theory	Medicine	Group I – up in smoke – no-smoking groups, lottery incentive, recruited through media	Age, sex
91 - I	Use of condoms, cessation of risky sex in adults with HIV	None	STDs (HIV)	HIV patients counselled about HIV and high-risk behaviours	Age, sex, ethnicity, knowledge
38 - I	Uptake of condoms, cessation of risky sex in school children	None	STDs (HIV)	Introduction health centres at schools plus routine visits scheduled plus information about safer sex	Age, social class, medical history, reproductive history, health behaviours-general
49 - I	Cessation of risky sex, uptake of condoms in school children	None	STDs (HIV)	AIDS information given by medical students plus interactive sessions plus video	Age, sex, knowledge, attitudes, perception of risk
123 - I	Adherence to safer sex, cessation of drug use in adolescent gay males	None	STDs (HIV)	Peer volunteer plus video plus group information: on safe sex and drug use	Age, level of education, social class, marital status, ethnicity, actual risk status, health behaviours-general knowledge, attitudes
159 - I	Cessation of/adherence to safer sex in well adult men	None	STDs (HIV)	Group I – workshops discussing risky sex plus skills acquisition	Age, ethnicity, level of education, social class, marital status, sexual orientation, religion, health behaviours, actual risk attitudes, knowledge, self-efficacy, intentions
480 - I	Cessation of/adherence to safe sex in well adolescent women	None	STDs (HIV)	Group I – peer volunteer individual counselling plus leaflet on HIV risks	Age, ethnicity, health insurance, level of education, social class, knowledge
651 - I	Adherence to safe sex/ HIV testing in adult well plus seropositive women	None	STDs (HIV)	Group I – video plus Group discussions led by physician plus free condoms plus spermicides	Age, marital status, actual risk status, medical history
655 - I	Uptake of contraception in well women	None	STDs (HIV)	Group I – contraception video plus free contraception	Age, marital status, actual risk status, reproductive history, reasons
688 - I	Condom use in seropositive adults (work)	None	STDs (HIV)	Group I – counselling couples to use condoms plus free condoms plus calendar to mark off when used condoms	Age, sex, marital status, medical history, reasons
729 - I	Cessation of risky needle behaviour in drug using adults	None	STDs (HIV)	Group I – verbal information on HIV, drug and sex behaviour plus clean needles and cleaning kit	Age, sex, ethnicity, actual risk knowledge
770 - I	Change HIV behaviours in drug using adults	None	STDs (HIV)	Group I – trained outreach workers plus additional visits to drug users plus free bleach and condoms plus leaflets plus free transport to centre	Age, sex, level of education, social class, actual risk
610 - I	Cessation of/adherence to risky sex in well adults	None	STDs (HIV)	Group I – use of puppets show: seriousness, transmission, avoidance of HIV (shows video)	Age, sex, ethnicity, worry, knowledge, perception of risk, usefulness of information
467 - I	Adherence to/cessation of HIV risk factors in well adolescents	None	STDs (HIV)	Group I – video plus leaflet plus group discussion about HIV	Age, sex, social class, knowledge, attitudes, perception of risk
723 - I	Use condoms in female sex workers with STD	None	STDs (HIV)	Group I – free condoms plus group discussions plus design poster	Age, level of education, marital status, knowledge, living arrangements, efficacy
545 - I	Cessation of risky sex in adults with HIV	None	STDs (HIV)	Group I – HIV counselling and testing plus scheduling follow-up visit (no details information)	Age, sex, ethnicity, actual risk status
650 - I	Cessation of risky HIV behaviours in pregnant women	None	STDs (HIV), obstetrics and gynaecology (midwifery)	Group I – group session HIV/AIDS transmission and testing	Age, marital status, ethnicity, knowledge, attitudes, intention

continued

**TABLE 30 contd** Description of decision, intervention and measures recorded in studies: before/after same sample

Study number	Summary decision and area of health	Theory	Health area	Comparison groups	Variables referred to
537 - I	Attendance clinic in adults with HIV	None	STDs (HIV), primary care	Group I – separate waiting area for women and children plus increase female staff plus free transport plus onsite colposcopy	Age, sex, ethnicity, actual risk status
35 - I	Uptake of condoms, cessation of risky sex in school children	None	STDs (HIV)	Group I – video plus leaflets plus large and small group information plus peer volunteer plus skills training	Age, sex, ethnicity, actual risk status, health behaviours-general, reproductive history, attitudes, social reinforcement/norms
364 - I	Adherence to safer sex in adult well women	Social learning theory	STDs (HIV)	Group I – small group plus video social cognition theory driven information (piloted): skill acquisition plus exercises plus HIV and condom use	Age, level of education, religion, knowledge, attitudes, self-efficacy, perception of norms
526 - I	Adherence to/cessation of risky sex in gay adolescent males	Social cognition model, social learning theory	STDs (HIV)	Group I – video plus small group information plus art plus skill acquisition plus confronting barriers information plus access to services	Age, actual risk, health behaviour, knowledge, perception of risk, self efficacy, perception of control, social support, perception of norms, stress, coping, anxiety, depression
654 - I	Cessation of/adherence to HIV behaviours in gay well adult males	New model testing	STDs (HIV)	Group I – Stop AIDS project: group information HIV and testing plus skill acquisition plus feedback plus 'contract' to change behaviour	Age, ethnicity, knowledge, attitudes, self-efficacy
450 - I	Change HIV risks in women with substance using partners	Social cognition model, social learning theory	STDs (HIV)	Group I – HIV information plus skill acquisition plus free bleach plus condoms plus incentives	Level of education, social class, ethnicity, actual risk status, reproductive history, perception of control
573 - I	Uptake of immunisation in well adults	None	Primary care	Group I – lecture plus group discussions plus leaflet	Knowledge, attitudes
342 - I	Uptake of immunisation (influenza) in well adults	None	Primary care	Group I – information by physician at day health centre plus free vaccine	Social class, attitudes, reasons
132 - I	Uptake of breast feeding in well women	None	Obstetrics and gynaecology (midwifery)	Trained health professionals on breast feeding and skills plus encourage 'rooming in' in hospital plus encourage support groups	Reasons
163 - I	Adherence to handwashing in pregnant women with children	None	Obstetrics and gynaecology (midwifery)	Group I – information on transmission of cytomegalovirus and affects fetus plus provision latex gloves plus liquid soap	Age
361 - I	Adherence to behavioural therapy programme in mentally ill adults	None	Mental health	Group I – discussion of patient targets within group plus problem solving	Age, sex, marital status, level of education, actual risk





## Appendix 8

### Intervention details by study design (grouped by health area in same order as appendix 7)

**TABLE 31** Description of the quality of the study and summary results: RCTa

Study number	Intervention level	Sample size	Summary of results
23 - 1	Patient	? / 69 / ? Convenience sample	Intervention associated with increase in amount of self-care behaviours
39 - 1	Patient	878 / 659 / ? Total sample available	Signed letter by GP more likely to increase self-report smear attendance than letter sent by clinic or information alone
41 - 1	Patient	? / 465 / 450 Total sample available	Signed letter by GP increased uptake of refusers from 10% to 21%
52 - 1	Patient	151 / 151 / 151 Total sample available	Uptake of mammography greater with telephone call reminder irrespective of identity caller. Medical assistant group most cost-effective
85 - 1	Patient	907 / 722 / 722 Total sample available	Authors claim prompt on contraceptive packet increased performance of BSE (57% vs. 49% at 6 months). Both interventions increased adherence to BSE after intervention. Note: large attrition and only a random sample of women's results reported
108 - 1	Patient	? / 283 / 283 Convenience sample	Uptake of FOB testing not associated with manipulation of stools or simplification of screening technique
122 - 1	Patient	? / 1842 / 1842 Not adequately described	Uptake of colorectal cancer screening not associated with type of screening method
316 - 1	Patient	1455 / 278 / 212 Systematic sample	No significant increase in attendance for colorectal screening following the letter-based information intervention (4% vs. 9%)
509 - 1	Patient	153 / 119 / 105 Total sample available	Voucher intervention increased attendance for mammography (44% vs. 10%)
101 - 1	Patient	150 / 150 / 150 Total sample available	Intervention not associated with uptake of mammography. Note: positively framed message associated with greater intention to attend than the routine letter
117 - 1	Other	228 / 171 / 171 Total sample available	Elderly educator and modified information was associated with increased adherence to screening
197 - 1	Patient	100 / 81 / 81 Systematic sample	Supplying screening kit plus information increased uptake of colorectal screening from 0% to 51%. HBM operationalised
275 - 1	Patient	2076 / 2076 / 1476 Total sample available	Interventions not associated with differential uptake rates (49%, 41%, 42%, 45%)
53 - 1	Patient	? / 267 / ? Total sample available	Adherence to self-report medication improved from 12% to 18% with use of the computerised telephone ( $p = 0.03$ )
50 - 1	Patient	184 / 184 / ? Total sample available	Counselling group reported changes in exercise uptake (16% vs. 50%), diet (19% vs. 33%) but no changes for smoking or alcohol intake. Biophysical markers not associated with changes
64 - 1	Patient	5282 / 4195 / 1098 Total sample available	Trend towards significance for smoking cessation but no significant difference on alcohol use and exercise uptake
118 - 1	Other	92 / 72 / 72 Systematic sample	Intervention associated with increase in number of quit attempts and a trend towards significance for smoking outside the home
216 - 2	Not recorded	? / ? / ? Not adequately described	Flat fee intervention had higher uptake than refundable deposit intervention ( $p < 0.0001$ )

continued

**TABLE 31 contd** Description of the quality of the study and summary results: RCTa

Study number	Intervention level	Sample size	Summary of results
229 - I	Patient	386 / 106 / 106 Total sample available	No differences in number people quitting smoking. Note, increasing cost of nicotine gum associated with decreasing attempts at obtaining gum and quit attempts
350 - I	Patient	? / 647 / 647 Convenience sample	Both physician-based information intervention groups associated with the same rate of cessation (approximately 11%)
483 - I	Other	762 / 410 / 352 Convenience sample	No association between training pharmacists and patient adherence
648 - I	Patient	? / 504 / 398 Convenience sample	No differential effects groups on smoking cessation (56% vs. 66% vs. 68%). Caution: two work-site samples differed at baseline
793 - I	Patient	1772 / 1686 / 1686 Not adequately described	Additional information not associated with smoking cessation. Nicotine patch associated with increase in cessation (11% to 19%)
781 - I	Patient	? / 122 / 115 Systematic sample	Self-management associated with reduction in use of antibiotics, ambulatory care and days taken off work ( $p < 0.05$ )
755 - I	Not recorded	1600 / 1381 / 1053 Systematic sample	Analysis performed control 2 and Group 4: Group 4 associated changes in alcohol and fat consumption, increase fruit and vegetables ( $p < 0.01$ ) but no change in smoking or exercise
67 - I	Patient	1581 / 661 / ? Systematic sample	Information intervention not associated with changes in preferences for CPR. Analysis suggested knowledge Alzheimer's disease, age, income and ethnicity were associated with reduced rates of resuscitation
196 - I	Patient	221 / 131 / 122 Total sample available	Telephone support associated with adherence to more regular meals and use of community resources but no change in smoking cessation
714 - I	Patient	190 / 190 / 184 Total sample available	Intervention associated with fewer missed appointments and increased adherence to treatment ( $p < 0.01$ )
200 - I	Patient	? / 535 / 520 Total sample available	Intervention associated with increase in physical activity and reductions in CVD risk factors
724 - I	Patient	1171 / 994 / 814 Total sample available	Intervention associated with increase in smoking cessation (9% vs. 14%; $p < 0.01$ ). Cost calculation provided
581 - I	Patient	? / 363 / 326 Convenience sample	Information intervention not associated with changes in smoking cessation (8% vs. 10%)
542 - I	Patient	? / 100 / 100 Convenience sample	Positive frame associated with increased consent to study (67% vs. 42%; $p < 0.01$ ). Interaction with reasons for choice
120 - I	Other	3989 / 3704 / 2082 Total sample available	Participants in control group less likely to respond. No association between the intervention and self-report of alcohol use. Note: little description of control group so difficult to interpret findings
301 - I	Patient	46 / 41 / 28 Total sample available	Increased information on side-effects not associated with changes in adherence to medication. Caution: small sample size
325 - I	Patient	74 / 54 / 54 Convenience sample	No reliable differential effects between intervention groups on increased uptake of exercise. Both interventions increased uptake over time ( $p < 0.001$ )
202 - I	Patient	80 / 80 / 80 Total sample available	Intervention not associated with increased attendance (68% vs. 53%). However, those failing to attend first appointment more likely to respond to a reminder phone call (84% vs. 52%)
393 - I	Patient	97 / 90 / 83 Total sample available	Intervention not associated with changes in risky or safe sex
741 - I	Patient	500 / 256 / 186 Total sample available	Attendance at STD clinic associated with reduction in sexual partners, HIV testing intervention and associated increased use condoms ( $p = 0.05$ ). Purpose of study to assess impact of offering HIV testing
107 - I	Other	? / 1213 / 808 Systematic sample	Intentions to engage in safer sex greater in intervention group ( $p < 0.0001$ ). Operationalising social cognitive framework
585 - I	Other	383 / 138 / 98 Convenience sample	Authors claim intervention associated with increase in condom use
799 - I	Patient	? / 128 / 100 Convenience sample	Social skills and culturally appropriate information associated with increase in safer sex ( $p < 0.05$ )
299 - I	Other	8069 / 5242 / ? Total sample available	All reminders associated with increase in attendance for tetanus immunisation ( $p < 0.0001$ ). Letter reminder produced greater attendance than physician or nurse reminder

continued

**TABLE 31 contd** Description of the quality of the study and summary results: RCTa

Study number	Intervention level	Sample size	Summary of results
378 - I	Patient	1068 / 1068 / 1068 Total sample available	Postcard reminder increased immunisation rates from 9% to 25%
638 - I	Patient	? / 497 / 434 Not adequately described	Telephone intervention associated with less utilisation of primary and hospital care, shorter stays in hospital, less medication. Cost calculation provided
387 - I	Patient	163 / 120 / 120 Total sample available	Intervention associated with greater question-asking by patient. Results presented with socio-economic status interaction
333 - I	Patient	5132 / 254 / 254 Systematic sample	Intervention associated with increase in immunisation and need for physician, decrease in use of medication ( $p < 0.01$ )
425 - I	Patient	3252 / 2794 / ? Total sample available	Intervention associated with fewer scans, day admissions and decreased satisfaction with service
778 - I	Patient	3368 / 1691 / 1691 Total sample available	Individual information increased attendance for serum screening. General information reduced attendance for CF carrier testing. Note: low attendance at classes (52%)
678 - I	Patient	854 / 814 / 814 Total sample available	Team care intervention associated with increased attendance to ante-natal clinic and perception of more informed choice for patient decision making
606 - I	Patient	516 / 516 / 516 Total sample available	Nurse intervention associated with greater attendance at follow-up (51% vs. 40%) and reduced number of attempted suicides (11% vs. 17%)
427 - I	Patient	68 / 47 / ? Total sample available	Adherence to medication more likely to be achieved in medication group

**TABLE 32** Description of the quality of the study and summary results: RCTb

Study number	Intervention level	Sample size	Summary of results
352 - 1	Patient	187 / 187 / 187 Total sample available	Small incentive not associated with increased uptake of mammography
352 - 2	Patient	184 / 184 / 184 Total sample available	No differential effect of telephone and postcard reminder with uptake of mammography
352 - 3	Patient	63 / 63 / 63 Total sample available	Signed physician letter associated with increased attendance for mammography and breast screening clinic ( $p < 0.05$ )
353 - 1	Other	? / 769 / 594 Convenience sample	Phone call reminders by acquaintances associated with increased uptake of mammography (49% vs. 34%; $p < 0.001$ )
420 - 1	Patient	? / 2201 / 2201 Systematic sample	Intervention associated with increased uptake of colorectal screening (Group 3 significantly greater than other groups: 48% vs. 27%, 37%, 37%). Gain/loss framing of messages not associated with uptake of colorectal screening
482 - 1	Patient	? / 2201 / 1565 Systematic sample	Interventions not associated with increase adherence to colorectal screening
653 - 1	Patient	? / 3726 / 3726 Systematic sample	Informing women about research project not associated with uptake of breast cancer screening (53% vs. 53%)
675 - 1	Patient	153 / 153 / 153 Total sample available	Completion colorectal screening not associated with length of time to take samples (3 or 6 days). Adherence associated with dietary restrictions (28% vs. 53%)
683 - 1	Patient	? / 966 / 966 Total sample available	Differential effects in uptake rate (Group 1: 48%; Group 2: 55%; Group 3: 46%; Group 4: 31%; Group 5: 23%). Cost-benefit analysis carried out
491 - 1	Patient	? / 614 / 439 Convenience sample	Interventions not associated with differential effect self-report BSE. Increase in self-report over time. Inappropriate statistics and erroneous exclusion of subjects
533 - 1	Patient	? / 614 / 479 Convenience sample	Differential effects associated with intervention, individual feedback on own breast associated with better BSE technique ( $p < 0.01$ )
566 - 1	Patient	? / 325 / 325 Convenience sample	Stamped addressed envelope increased return rate from 61% to 74%. Interaction reported with whether insured or not
705 - 1	Other	? / 543 / 262 Total sample available	Interventions not associated with changes in safe-sun behaviour
728 - 1	Patient	1143 / 834 / 834 Systematic sample	Differential effects in increasing uptake of cervical smear (5%; 11%; 30%; 26%). Personal visit associated with increased attendance rather than video
630 - 1	Patient	799 / 799 / 782 Total sample available	No differential effect interventions on women's attendance for breast screening (11% vs. 8% vs. 13%)
285 - 1	Not recorded	? / ? / ? Not adequately described	Poor explanation study, results reported only for refund vs. free test. Free test increased screening (42% vs. 56%)
727 - 1	Patient	437771 / 2266 / 2266 Systematic sample	Fixed appointment associated with increase in attendance (27% vs. 40%). Second intervention study not reported as unclear explanation
543 - 1	Patient	2131 / 1912 / 1912 Total sample available	Intervention associated increase cervical smear attendance (28% vs. 13%). Higher in those attending breast screening (42% vs. 20%)
587 - 1	Patient	669 / 205 / 205 Total sample available	Increased information associated with decreased interest in prostate screening ( $p < 0.05$ )
713 - 1	Patient	? / 125 / 108 Systematic sample	Intervention associated with increased attendance at follow-up clinic (75% vs. 46%). Note: little information on attrition
48 - 1	Other	995 / 553 / ? Total sample available	Women already in hospital had greater uptake of cervical smears (72%). No difference in uptake of smears between information only and control group (24% vs 20%)
488 - 1	Patient	169 / 102 / 100 Total sample available	Interactive computer screen associated with increased intention to take part in clinical trial compared with audio-taped information. Concerned with informed patient decision making
86 - 1	Other	? / ? / ? Not adequately described	Intervention associated with increase in cervical and mammography uptake

continued

**TABLE 32 contd** Description of the quality of the study and summary results: RCTb

Study number	Intervention level	Sample size	Summary of results
114 - I	Patient	? / 1104 / 405 Systematic sample	Patients belief interventions were twice as effective as routine information
232 - I	Patient	? / 802 / 626 Systematic sample	Intensive education intervention not associated with a differential effect uptake of mammography. Operationalised HBM
438 - I	Patient	? / 181 / 133 Convenience sample	Intervention associated with a trend towards significance (gain 52%, loss 66%; $p = 0.07$ ). Operationalised prospect theory
784 - I	Patient	3291 / 3291 / 3290 Total sample available	Authors claim attendance increased to 70% if woman interviewed (42% overall). Actual uptake by group unclear as analysis to assess predictors of uptake
700 - I	Patient	? / 142 / 92 Not adequately described	Question-based intervention increased question asking (35% vs. 16%). Assessed preference of consultation style
789 - I	Patient	? / 250 / 197 Convenience sample	Information framed using internal responsibility associated with increased uptake of mammography (55% vs. 66% vs. 57%; $p < 0.01$ )
207 - I	Other	? / ? / 117 Convenience sample	Smoking and smoking/diet group associated with reduction in smoking; diet and smoking/diet intervention associated with changes in diet
288 - I	Patient	1927 / 647 / 390 Convenience sample	No differential effect of interventions on number kits ordered or intention to test. Assessed changes in perceptions of risk
690 - I	Patient	2822 / 1404 / 511 Systematic sample	Belief and information group associated with greater increase in BSE than other groups ( $p < 0.05$ )
458 - I	Other	1703 / 1703 / 1270 Total sample available	Intervention associated with trend towards intention to change behaviour ( $p < 0.01$ )
307 - I	Patient	875 / 175 / 157 Convenience sample	No differential effect in interventions. BSE improved over time
140 - I	Patient	? / 186 / 114 Convenience sample	Intervention not associated with changes in smoking cessation
757 - I	Patient	535 / 535 / 446 Total sample available	No differential effects from framing information but additional information associated with increased uptake of mammography (66% vs. 53%; $p < 0.01$ )
434 - I	Patient	264 / 96 / 90 Convenience sample	Framing information not associated with willingness to enter trial
816 - I	Patient	218 / 180 / 180 Total sample available	Intervention not associated with increase or decrease in participation (13% control vs. 24% intervention). Analysis looked at predictors of attendance
667 - I	Patient	64 / 60 / 60 Total sample available	Intervention not associated with changes in participation consultation or final decision
671 - I	Patient	? / 100 / 81 Not adequately described	Additional information not associated with cancellation of procedure. Note: small sample size and non-validated measures
1 - I	Patient	? / 681 / 626 Not adequately described	No change in exercise behaviour, though reduction in anxiety
10 - I	Patient	200 / 160 / 140 Not adequately described	No association between care plan group and hospital attendance. Main results between intervention, patient satisfaction and mortality
28 - I	Patient	50 / 46 / 46 Not adequately described	Intervention not associated with change in use of healthcare services
63 - I	Patient	430 / 391 / 294 Total sample available	Nurse intervention associated with increased uptake of exercise. Exercise uptake not associated with routine and routine plus information group
47 - I	Patient	? / 2441 / ? Convenience sample	Physician group more likely to eat more healthily and be immunised than hospital-based group. No association with smoking behaviour
55 - I	Other	? / 267 / 267 Not adequately described	Intervention not associated with intention to stop smoking

continued

**TABLE 32 contd** Description of the quality of the study and summary results: RCTb

Study number	Intervention level	Sample size	Summary of results
55 - 2	Other	? / 450 / 220 Not adequately described	Intervention not associated with intention to stop smoking
57 - 1	Not recorded	427 / 320 / 213 Total sample available	Adherence decreased if medication taken twice a day (95% vs. 82%)
88 - 1	Patient	71 / 60 / 60 Total sample available	Intervention not associated with changes in adherence. Presented results of demographic associations with adherence
102 - 1	Other	1425 / 1097 / 770 Convenience sample	Intervention not associated with cessation of smoking. Interaction between level education and treatment condition
105 - 1	Patient	74 / 36 / 31 Total sample available	Modified information associated with increase in adherence to medication (79%, 83%, 93%). Small sample size
110 - 1	Other	4712 / 4712 / 1452 Systematic sample	Intervention associated with decrease in self-report alcohol use, increase in healthy eating but no change in smoking or exercise. Caution in interpreting the results as inappropriate use of statistics and potential confounders between randomised groups
147 - 1	Patient	? / 70 / 61 Convenience sample	Intervention group associated with reduction in fatty food intake
155 - 1	Patient	? / 128 / 128 Systematic sample	Patients less likely to sign consent form if given information about videotaping consultation
183 - 1	Patient	? / 453 / 290 Not adequately described	No association with information group and cessation of alcohol
216 - 1	Patient	6000 / 3257 / 140 Systematic sample	One-stage intervention had a higher uptake rate than the two-stage approach 3% vs. 2% ( $p < 0.05$ )
217 - 1	Other	7351 / 2577 / ? Total sample available	No changes in smoking behaviour either before/after intervention or between varying amounts and types of information groups
249 - 1	Patient	? / 179 / 155 Not adequately described	No association between counselling and cessation of smoking
259 - 1	Patient	? / 338 / 336 Convenience sample	Control group drinking twice as much as either intervention group ( $p < 0.001$ ). No difference between intervention groups
279 - 1	Other	1005 / 1005 / 766 Total sample available	Intervention associated with reduced uptake of smoking (17% vs. 26%). No baseline measure of smoking in intervention group
356 - 1	Other	2002 / 1001 / 1001 Systematic sample	Blood pressure and diet monitoring by nurse not associated with change in salt intake/diet change
574 - 1	Patient	? / 44 / 22 Not adequately described	No differences in cessation between groups (68% vs. 71% vs. 60%). High cessation rates indicate that sample not representative
652 - 1	Patient	? / 1975 / 1948 Convenience sample	Authors suggest intervention associated with increase in attendance (46%, 61%, 58%)
679 - 1	Other	? / 1274 / 1274 Systematic sample	Intervention not associated with changes in exercise behaviour
489 - 1	Patient	? / 36 / 36 Convenience sample	Interventions not associated with any changes to lifestyle behaviours
772 - 1	Patient	? / 44 / 42 Convenience sample	No differential effect of patient-centred and directive style with cessation of drinking. Counselling better than no counselling ( $p < 0.02$ )
807 - 1	Patient	? / 204 / ? Convenience sample	Intervention associated with increase in self-management of diabetes
447 - 1	Patient	? / 224 / 209 Not adequately described	Intervention associated with increased clinic attendance ( $p < 0.001$ ). Note: interaction analysis performed between 'new' hypertensives and known infrequent attendees
783 - 1	Patient	2021 / 2021 / 1877 Convenience sample	Telephone support counselling increased cessation rates (16% vs. 15% vs. 14% vs. 23%)

continued

**TABLE 32 contd** Description of the quality of the study and summary results: RCTb

Study number	Intervention level	Sample size	Summary of results
336 - I	Other	2753 / 2202 / 1652 Total sample available	Interventions not associated with changes in diet and smoking. Curriculum plus screening associated with increase in aerobic exercise ( $p < 0.05$ )
530 - I	Patient	192 / 116 / 116 Convenience sample	Intervention associated with increase adherence at follow-up ( $p < 0.05$ )
564 - I	Not recorded	371 / 227 / 227 Total sample available	No differences in treatment adherence by group (14%, 12%, 10%). Differential effects of groups on other drinking outcomes: hospitalisation > choice > AA
702 - I	Patient	70 / 70 / 70 Total sample available	Information plus phone call associated with greater increase in adherence behaviour ( $p < 0.01$ )
213 - I	Other	? / ? / 659 Convenience sample	Intervention associated with increased rates of cessation (18% vs. 9%)
327 - I	Patient	? / 52 / 52 Convenience sample	No association between computer group and increase in house-mite reduction activities. Both groups increased house-mite reduction activities over time
328 - I	Patient	? / 102 / 102 Convenience sample	Intervention associated with increase in healthy diet and self-monitoring of glucose. No association between exercise and medication adherence
612 - I	Patient	? / 27 / 25 Convenience sample	Type of device not associated with adherence therapy
629 - I	Patient	200 / 190 / 190 Total sample available	Intervention associated with increase in completion healthcare proxy (48% vs. 6%, $p < 0.001$ )
268 - I	Patient	? / 5354 / 3122 Total sample available	Presentation results by men, wives and children. Intervention associated with changes in diet ( $p < 0.01$ ). No change in smoking cessation
395 - I	Other	7397 / 7397 / 7397 Total sample available	Differential effects by group. Authors claim greatest increase associated with physician and patient reminder group ( $p < 0.01$ )
396 - I	Patient	? / 105 / 105 Not adequately described	Computer group associated changes in fat and calorie intake ( $p < 0.05$ )
398 - I	Patient	? / 93 / 75 Convenience sample	Results not reported for exercise/diet adherence by group. Analysis reporting differences between black/white participants. Exercise uptake but not diet intake improved over time
402 - I	Patient	? / 211 / 211 Not adequately described	Randomisation of patients resulted in lower uptake treatment (71% vs. 56%; $p < 0.05$ )
439 - I	Patient	5281 / 4195 / 3097 Total sample available	No data provided for uptake of screening in control group. Intervention a predictor for better health at 2-year follow-up
527 - I	Patient	? / 241 / 231 Convenience sample	Feedback intervention associated with increased adherence with inhaler ( $p < 0.01$ )
168 - I	Patient	122 / 102 / 102 Total sample available	Telephone intervention not associated with adherence to medication (34% vs. 29%)
265 - I	Patient	? / 229 / 169 Total sample available	Interventions not associated with exercise or smoking behaviours. Control group increased intake of cholesterol, early intervention associated with greater decrease in cholesterol intake than late group ( $p < 0.05$ )
365 - I	Patient	321 / 253 / 231 Convenience sample	Intervention aimed at women associated with change in diet in men ( $p < 0.05$ ). Note: excluded men no longer living in marital home
372 - I	Patient	? / 207 / 173 Convenience sample	Intervention not associated with consultation attendance or changes to lifestyle
263 - I	Patient	? / 303 / 264 Convenience sample	Intervention associated with reduction of fat in diet (17% vs. 36%)
693 - I	Patient	? / 52 / 42 Total sample available	Authors claim that medication schedule associated with increased compliance but sample size small
137 - I	Patient	? / 181 / 138 Convenience sample	Intervention not associated with adherence to medication

continued

**TABLE 32 contd** Description of the quality of the study and summary results: RCTb

Study number	Intervention level	Sample size	Summary of results
139 - 1	Patient	307 / 275 / 275 Total sample available	Analysis performed by risk group: intervention, obese patients more likely to make appointment with dietician; intervention, smoking group more likely to report quitting (though not confirmed by cotinine level)
143 - 1	Patient	? / 1745 / 1745 Convenience sample	No differential effects of intervention on smoking cessation (16% vs. 17%)
668 - 1	Patient	? / 870 / 602 Convenience sample	Differential effects of intervention over time. Personalised feedback plus stage of change leaflet associated with greater cessation. Applied transtheoretical model
332 - 1	Patient	? / 201 / 201 Convenience sample	Differential effects for way medical interventions described: 31% cases positively described, 12% negatively described and 19% medical guideline accepted treatment
20 - 1	Patient	85 / 64 / 43 Convenience sample	Intervention not associated with change in use of healthcare service
27 - 1	Patient	500 / 329 / 303 Convenience sample	Association between groups and use of services by disease: osteoporosis, counselling group fewer clinic visits than other groups; rheumatoid arthritis no association by groups
104 - 1	Patient	? / 156 / 156 Convenience sample	Different associations by different behaviours: behavioural strategy group more likely to monitor glucose level; attention group lower adherence to smoking cessation and medication but better diet and weight behaviours
220 - 1	Other	8565 / 7795 / ? Systematic sample	Adherence to healthy diet associated with intervention ( $p < 0.001$ )
233 - 1	Patient	? / 1800 / 1381 Not adequately described	Significant group by time interactions for adherence exercise and cessation fat and caffeine intake
241 - 1	Patient	? / 125 / 114 Convenience sample	Intervention associated with increased duration and frequency of exercise. Path analysis assessed relationship self-efficacy and exercise adherence
276 - 1	Other	1013 / 570 / 570 Total sample available	Intervention not associated with changes in smoking cessation
295 - 1	Patient	? / 66 / 56 Not adequately described	Interventions not associated with self-reported changes in diet at 3 months
305 - 1	Patient	? / 293 / 177 Convenience sample	Both theory-based groups associated with cessation of fatty diet and adherence to exercise. No changes for smoking, use of health professional, and adherence to medication
321 - 1	Patient	? / 26 / 22 Convenience sample	Intervention not associated with differences in adherence (no figures provided)
157 - 1	Other	2332 / 933 / 583 Convenience sample	Information group not associated with infant smoke exposure reduction. Both control and intervention groups increased smoking over time (53% vs. 46%)
376 - 1	Patient	1387 / 1378 / 1378 Total sample available	Intervention associated with increased attendance at smoking cessation programme (after GP advice, 1%; after recruitment 9%)
469 - 1	Patient	? / 453 / 453 Not adequately described	Intervention associated with increased adherence to medication in both new and existing patients
502 - 1	Patient	? / 84 / 58 Not adequately described	Intervention not associated with changes in self-management of illness
503 - 1	Patient	? / 66 / 48 Not adequately described	Intervention associated with changes in lifestyle behaviours ( $p = 0.04$ ). Behaviours not noted and percentages not stated. Regression analyses performed looking at knowledge
707 - 1	Patient	? / 585 / 585 Not adequately described	No difference in adherence to therapies (11% vs. 9%). Biochemical assessment smoking, diet and exercise suggest changes by group
158 - 1	Patient	72 / 51 / 51 Total sample available	Intervention associated with increased cessation in moderate smokers (31% vs. 7%)
158 - 2	Patient	410 / 296 / 197 Convenience sample	Intervention associated with smoking cessation in moderate smokers ( $p < 0.05$ adjusted for non-responders)
210 - 1	Patient	686 / 507 / 347 Not adequately described	Tailored intervention associated with self-reported changes in diet ( $p < 0.01$ )

continued



**TABLE 32 contd** Description of the quality of the study and summary results: RCTb

Study number	Intervention level	Sample size	Summary of results
211 - 1	Other	? / 340 / 292 Convenience sample	No differential effects of interventions on smoking cessation (27% vs. 22%). Operationalised stages of change model
644 - 1	Other	488 / 431 / 364 Not adequately described	Analysis reported by group and collapsed groups: behavioural counselling associated with higher cigarette cessation rate than information-feedback groups (18% vs. 3%, $p < 0.01$ )
269 - 1	Patient	? / 345 / 139 Convenience sample	Differential effect of intervention on fibre consumption (29% vs. 88% vs. 66%, $p < 0.01$ ). Cost-benefit analysis
291 - 1	Patient	? / 290 / 225 Convenience sample	Education intervention associated with increase in self-management for parents and children
541 - 1	Patient	524 / 290 / 196 Total sample available	Intervention associated with reduction in fat intake ( $p < 0.001$ )
255 - 2	Other	8992 / 8992 / 7180 Systematic sample	No differential effects of interventions on smoking behaviour
145 - 1	Other	? / 5589 / 3884 Convenience sample	Intervention not associated with cessation of smoking. Analysis assessed rebelliousness, norms and smoking status
264 - 1	Patient	? / 208 / 169 Not adequately described	Intervention associated with increased exercise uptake for control and Group 3, a decrease in fat consumption for Group 2
659 - 1	Patient	? / 550 / 304 Not adequately described	Women's intervention associated with reduction in men's consumption of fat ( $p < 0.01$ ) but no increase in eating healthy foods
622 - 1	Patient	? / ? / 842 Not adequately described	No differential effects of intervention groups, both increased adherence over control group ( $p < 0.01$ ). Operationalised HBM
814 - 1	Patient	? / 136 / 120 Convenience sample	Intervention associated with greater adherence to exercise programme ( $p < 0.01$ )
490 - 1	Patient	133 / 88 / 80 Total sample available	Intervention leaflet associated with increase in rehabilitation exercises ( $p < 0.001$ ). Main analysis assessed gender differences and anxiety
22 - 1	Other	? / 2271 / ? Systematic sample	Results presented by gender. Males and females in intervention group showed increase in intention to use condoms
732 - 1	Other	14749 / 12537 / 8845 Systematic sample	Intervention not associated with changes in risky sex behaviour. Purpose of study to look at prevalence of HIV
736 - 1	Patient	? / 313 / 218 Convenience sample	Authors claim intervention not associated with change in behaviours but sample group highly selective and attrition
341 - 1	Patient	313 / 313 / 218 Total sample available	Intervention not associated with changes in drug use or sexual behaviour. Sub-group analysis between used vs. not used alcohol
156 - 1	Patient	? / 99 / 61 Not adequately described	High reward associated with increase in attendance, decrease 'drug' behaviours. No difference in low reward intervention over time
212 - 1	Other	? / ? / ? Not adequately described	Intervention not associated with any changes in cigarette, alcohol or marijuana over 5-year period
639 - 1	Patient	243 / 217 / 217 Total sample available	Video group associated with greater intention to reduce risk behaviour immediately after information. No effect at 2 months
289 - 1	Patient	? / 107 / 77 Convenience sample	Acquisition of condoms associated with intervention (38% vs. 77% vs. 75%). No differential effects between intervention groups
465 - 1	Patient	? / 214 / 97 Convenience sample	Results on behaviour not clearly reported, no reported change in behaviour
733 - 1	Patient	? / 70 / 60 Convenience sample	Intervention associated with trend towards drug cessation and attendance for health appointments ( $p = 0.06, 0.07$ )
4 - 1	Patient	184 / 184 / 184 Convenience sample	Contact therapy not associated with greater decrease in cocaine use. Authors looked at predictors of cocaine use

continued

**TABLE 32 contd** Description of the quality of the study and summary results: RCTb

Study number	Intervention level	Sample size	Summary of results
24 - 1	Patient	? / 20 / ? Convenience sample	Alcohol was associated with reduced discussion of condoms. No change in intention to use condoms
223 - 1	Patient	? / ? / 296 Convenience sample	Social skills intervention no more effective than routine information in increasing cessation of drug use
226 - 1	Other	3011 / 2416 / 2416 Total sample available	Groups confronting beliefs and normative behaviours associated with reduced drug use. Control and resistance training alone not useful in reducing drug use behaviour
247 - 1	Other	5070 / 3549 / ? Total sample available	Authors claim psycho-social interventions associated with changes in alcohol, smoking and marijuana use. Analysis performed by gender
277 - 1	Other	? / ? / ? Total sample available	No association of theory-based interventions and changes in alcohol, smoking and marijuana use, 2 years after intervention. Analysis performed on process measures
424 - 1	Other	? / ? / ? Convenience sample	Authors claim intervention associated with increased safer sex practices. Note: analysis only on sexually active sub-group
478 - 1	Other	? / 1033 / 758 Total sample available	Intervention associated with greater contraceptive discussions with parents. In female, first-time sexual encounters intervention associated with safer sex practices
600 - 1	Other	387 / 220 / 208 Total sample available	Directive intervention associated with lower recruitment rates (65% vs. 52%). No association between intervention and attendance at counselling sessions. No demographic results presented although reported
685 - 1	Patient	? / 291 / 291 Not adequately described	Theory-based information associated with increase in attainment of free condoms ( $p < 0.01$ ). Assessed relationship experience and attitude formation (Fazio)
742 - 1	Patient	? / ? / 206 Not adequately described	Analysis suggests AIDS information group more likely to use condoms
492 - 1	Patient	? / 89 / 81 Convenience sample	Intervention not associated with changes in risky behaviour
766 - 1	Patient	246 / 246 / 225 Total sample available	Intervention associated with increased use of condoms over time ( $p < 0.01$ ), reduction in the initiation of sexual activity (31% vs. 12%)
776 - 1	Patient	? / 187 / 93 Convenience sample	Intervention associated with increased use of condoms ( $p < 0.001$ )
165 - 1	Patient	? / 1770 / 994 Convenience sample	Intervention not associated with cessation of risky drug behaviours although a time effect
632 - 1	Patient	55 / 50 / 47 Convenience sample	Analysis of safe sex/drug use not reported. Intervention associated with skill acquisition for condom and needle sterilisation use
632 - 2	Patient	? / 98 / 60 Not adequately described	Intervention not associated with behaviour change
635 - 1	Patient	176 / 152 / 106 Total sample available	Risky sex/drug behaviour decreased over time ( $p < 0.001$ ). No reliable differential effects by intervention but for sex/ drugs more in information group increased risk behaviour (11% vs. 4%)
636 - 1	Patient	? / 497 / 407 Not adequately described	Enhanced intervention associated with cessation of cocaine use and risky drug use
525 - 1	Patient	? / 200 / 88 Convenience sample	No differential effect between groups on drug use, risky sex and study attrition. Over time, reduction in risky drug use but not sex
660 - 1	Other	1315 / 1201 / 867 Systematic sample	Intervention associated with adherence to safer sex ( $p < 0.01$ )
454 - 1	Patient	? / 916 / 858 Convenience sample	Decreased risky behaviours for both groups over time, traditional group associated with greater cessation of behaviours than intensive group
706 - 1	Patient	465 / 419 / 419 Total sample available	Intervention associated with mixed findings: reduction in home visits for coughs, colds, fevers but increase for diarrhoea and vomiting
6 - 1	Patient	10 / 10 / ? Convenience sample	Adherence to medication not associated with taste of medication

continued

**TABLE 32 contd** Description of the quality of the study and summary results: RCTb

Study number	Intervention level	Sample size	Summary of results
6 - 2	Not recorded	15 / 12 / ? Not adequately described	Adherence to medication not associated with taste of medication
6 - 3	Patient	10 / 10 / ? Convenience sample	Adherence to medication not associated with taste of medication
32 - 1	Patient	201 / 194 / 174 Total sample available	Intervention associated with differences in attendance and adherence. Stimulus group had lower attendance than regular class; weight-dependence group had better adherence than time-dependent group
51 - 1	Patient	? / 400 / 385 Not adequately described	Medication provided with precise unit of measurement in liquid form associated with increase in adherence to medication
174 - 1	Patient	? / 44 / 39 Convenience sample	Intervention associated with adherence to exercise programme but not with diet
302 - 1	Other	? / 100 / 98 Not adequately described	Leaflet associated with fewer home visits and physician appointments and more self-treatments of children's symptoms ( $p > 0.01$ )
589 - 1	Patient	? / 325 / 159 Systematic sample	No difference between two methods of invitation (72% vs. 66%)
572 - 1	Patient	239 / 239 / 103 Total sample available	Intervention associated with increased uptake of immunisation ( $p < 0.001$ )
747 - 1	Patient	? / 1200 / 1200 Systematic sample	Differential effect of interventions on uptake (75% vs. 69% vs. 54%)
769 - 1	Not recorded	4641 / 855 / 567 Systematic sample	No differential effects of interventions on attendance at general practice. Article reports costs of each alternative
520 - 1	Patient	3913 / 3404 / 3404 Total sample available	Leaflet associated with lower utilisation of services. Interaction with perception of health risk
234 - 1	Patient	? / 3884 / 1989 Not adequately described	Free vaccination increased uptake of flu immunisation more than the control group. Uptake of immunisation increased over time. Note: less than a 40% response rate
337 - 1	Other	2624 / 2624 / 2624 Total sample available	Attendance increased in intervention group, particularly for 'alarming' coughs
796 - 1	Other	548 / 548 / 548 Total sample available	Intervention associated with adherence to cough care treatment (30% vs. 56%; $p < 0.01$ )
414 - 1	Patient	? / 43 / 32 Convenience sample	No association of intervention and therapy adherence, diet or exercise changes
553 - 1	Patient	? / 120 / 86 Not adequately described	No association between groups and changes in exercise behaviour. Predictors of behaviour analysis only reported
408 - 1	Patient	229 / 229 / 222 Total sample available	Telephone reminder not associated with increase in immunisation
19 - 1	Patient	? / 236 / 236 Not adequately described	Choice of treatments associated with presentation data: fewer patients chose immediate survival when data presented as a curve rather than point estimates
167 - 1	Patient	? / 288 / 288 Not adequately described	Interventions associated with increased attendance at well-baby clinic ( $p < 0.05$ ). No differential effects interventions (68%, 75%, 74%)
357 - 1	Patient	? / 66 / 62 Not adequately described	Illustrated leaflet not associated with increased adherence to medication ( $p = 0.07$ )
426 - 1	Patient	299 / 299 / 299 Total sample available	Fixed appointment increased uptake from 27%– 59%. Operationalised HBM
506 - 1	Patient	141 / 67 / 67 Total sample available	Intervention not associated with increase in patient participation consultation
633 - 1	Patient	422 / 422 / 359 Total sample available	Intervention not associated with uptake of vaccination

continued

**TABLE 32 contd** Description of the quality of the study and summary results: RCTb

Study number	Intervention level	Sample size	Summary of results
369 - I	Patient	587 / 439 / 439 Total sample available	Intervention not associated with adherence to medication
620 - I	Patient	299 / 254 / 121 Not adequately described	No differential effect in uptake of health check by group. Fewer people in opportunistic group approached, resulting in differences in uptake (79% vs. 33%; $p < 0.01$ )
135 - I	Patient	68 / 68 / 55 Convenience sample	Authors suggest intervention group associated with increase in exclusively breast feeding for 3 months ( $p < 0.02$ ). (published again <i>Journal of The American Dietetic Association</i> 1995;95:323-8.)
698 - I	Patient	300 / 210 / 210 Not adequately described	Adherence trial associated with effectiveness of HRT in controlling menopausal symptoms (47% vs. 49% vs. 73%)
634 - I	Patient	? / 800 / 800 Not adequately described	Fixed appointment letter associated with increased attendance (75% vs. 54%)
768 - I	Patient	270 / 216 / 87 Total sample available	Breast pump rather than formula associated with increased breast feeding only if women perceived bottle to make night-time feed easier ( $p < 0.05$ )
616 - I	Patient	? / 194 / 175 Convenience sample	Intervention associated with increased breast-feeding (38% vs. 70%; $p < 0.001$ )
403 - I	Patient	? / ? / 180 Total sample available	No association between interventions and uptake or duration of breast feeding
562 - I	Patient	232 / 232 / 186 Total sample available	Feedback and information group associated with increase in dental hygiene (76% vs. 67% vs. 90%; $p < 0.01$ )
576 - I	Patient	? / 20 / 20 Not adequately described	Interventions associated with increase in dental hygiene (additional feedback plus checklist)
758 - I	Patient	700 / 167 / 91 Convenience sample	Intervention not associated with changes in dental hygiene behaviour. Operationalised the theory of reasoned action
371 - I	Other	? / 119 / 98 Not adequately described	Intervention not associated with changes in self-reported dental behaviour, but changes in plaque level
457 - I	Patient	? / 84 / 84 Not adequately described	Group 2 associated increase in dental hygiene activities (36% vs. 41% vs. 100%; $p < 0.01$ )
304 - I	Patient	392 / 318 / ? Convenience sample	Differential attendance rates for final visit (65% vs. 94% vs. 95% vs. 78%). Despite inappropriate statistics, nurse home-based interventions associated with increased attendance ( $p < 0.05$ )
308 - I	Other	? / 200 / 179 Systematic sample	Intervention group more likely to clear mosquito breeding sites (82% vs. 25%)
96 - I	Patient	? / 191 / 172 Not adequately described	Intervention not associated with adherence to medication
479 - I	Patient	? / 77 / 77 Not adequately described	Family intervention associated with increased adherence to treatment. Results indicate patient's clinical state improved
511 - I	Patient	400 / 322 / 290 Total sample available	Intervention associated with exercise uptake ( $p < 0.001$ ) and trend towards decreased physician visits ( $p = 0.06$ )
494 - I	Not recorded	? / 206 / 199 Convenience sample	Intervention associated with fewer referrals to nursing homes ( $p < 0.05$ ). Main analysis between social support, health and behaviour
554 - I	Patient	? / 816 / 783 Systematic sample	Pre-paid plans not associated with Medicaid beneficiaries utilisation services
794 - I	Patient	281 / 217 / 177 Total sample available	Authors claim intervention associated with increased adherence to medication (76% vs. 50%; $p < 0.01$ )
493 - I	Patient	? / 102 / 91 Convenience sample	Computer associated with increased confidence in decision making but not decision skill. Computer used for information and not as a decision aid
176 - I	Patient	511 / 492 / 470 Total sample available	57% patients chose medication phrased in relative-risk terms, 15% when phrased in absolute-risk terms. Good experimental design but no control over order effects
150 - I	Patient	400 / 234 / 205 Total sample available	Use of cartoons plus leaflet increased adherence to wound care

**TABLE 33** Description of the quality of the study and summary results: RCTc

Study number	Intervention level	Sample size	Summary of results
37 - I	Patient	? / 1744 / ? Total sample available	Video intervention associated with increased uptake of rate cervical smear
154 - I	Patient	511 / 369 / 369 Total sample available	Intervention associated with increased mammography screening. Note: women 'dropped out' perceived themselves to be less susceptible to breast cancer and perceived mammography to be less effective than those who took part
296 - I	Patient	527 / 395 / 179 Not adequately described	Free test by physician increased adherence to screening ( $p < 0.05$ ). Addition of reminder letter not associated with increase in adherence
377 - I	Patient	621 / 204 / 202 Total sample available	No differential effect in uptake of either intervention (55% vs. 67%)
412 - I	Patient	? / 146 / 146 Not adequately described	Stamped addressed envelope associated with higher return rate for hemocult cards (37% vs. 57% vs. 71%; $p < 0.01$ ). Calculated cost intervention
695 - I	Not recorded	527 / 498 / 147 Systematic sample	Uptake of mammography not associated intervention
712 - I	Patient	220 / 220 / 178 Total sample available	Differential effects of interventions by experimental group (73% vs. 72% vs. 44% vs. 36%; $p < 0.01$ )
310 - I	Other	? / 1618 / 603 Convenience sample	No differential effects associated with any interventions on smoking cessation. Analysis carried out for number of quit attempts and differences in participants/non-participants
752 - I	Other	1588 / 1588 / 1588 Total sample available	Sending hemocult by post associated with greater uptake (26% vs. 32% vs. 21%; $P < 0.01$ ). Hemocult screening not associated with invitation to attend health check
691 - I	Other	89 / 82 / 82 Total sample available	No differential effect in uptake irradiation therapy by group (95%: 96%: 93%). Differences with decision-making process: consultation less directive and more involved in decision process
248 - I	Patient	170 / 161 / 161 Total sample available	Leaflet not associated with increased attendance at appointment ( $P = 0.097$ ). No evidence leaflet readable
354 - I	Other	752 / 616 / 412 Systematic sample	Theory-based intervention associated with increased uptake of mammography (45% vs. 12%; $p < 0.001$ ). Operationalised HBM
380 - I	Patient	90 / 90 / 90 Total sample available	Telephone counselling confronting barriers to attendance increased attendance at clinic (67% vs. 43%, $p < 0.05$ )
484 - I	Other	2044 / 2044 / 2044 Total sample available	Free transportation associated with highest attendance at clinic, letter only with lowest. Logistic regression analysis, so no percentages
507 - I	Patient	3035 / 3035 / 963 Total sample available	Interventions not associated with increased uptake rates (89%, 90%, 90%). Note all uptake rates very high
508 - I	Other	541 / 541 / 541 Total sample available	Intervention associated with increased cervical screening. Logistic regression presented, interaction with type of cervical abnormality
303 - I	Other	268 / 219 / 219 Total sample available	Information intervention not associated increased with adherence. Interactions showed least compliant patients improved following intervention (55% vs. 32%)
320 - I	Patient	379 / 287 / 287 Total sample available	Information intervention associated with decreased attendance at clinic $p < 0.05$ , no change in diet, smoking and alcohol behaviour
381 - I	Patient	? / 313 / 238 Convenience sample	Additional information and letter prompt not associated with increased referral to physician (25% vs. 24%) or self-reported changes in diet, exercise or smoking
471 - I	Patient	? / 139 / 96 Convenience sample	No results reported evaluating impact interventions alone. Results analysed by gender group interaction
473 - I	Patient	? / 921 / 570 Convenience sample	Interventions not associated with any changes in smoking cessation
475 - I	Patient	? / 36 / 27 Convenience sample	Intervention associated with increase in checking blood glucose, diet, exercise, service utilisation
485 - I	Patient	371 / 227 / 200 Total sample available	Hospitalisation associated with greatest cessation of drinking after 2 years (37% vs. 17% vs. 16%)
763 - I	Other	? / 88 / 79 Not adequately described	Intervention associated with increase adherence (95% vs. 83%; $p = 0.02$ )

continued

**TABLE 33 contd** Description of the quality of the study and summary results: RCTc

Study number	Intervention level	Sample size	Summary of results
459 - I	Other	? / 873 / 630 Convenience sample	No differential effects of cessation smoking by group (10% vs. 9% vs. 10%)
754 - I	Patient	924 / 342 / 305 Convenience sample	Both intervention groups associated with changes in diet ( $p < 0.05$ ). Assessed knowledge
708 - I	Other	? / 580 / 510 Not adequately described	Intervention not associated with changes in smoking
711 - I	Other	? / 309 / 309 Not adequately described	Follow-up phone call decreased drop-out rate for 6 months only after treatment ( $p < 0.01$ )
170 - I	Patient	1142 / 1015 / 747 Not adequately described	Information intervention associated with small change in preventive behaviour in better educated parents. Likely some group contamination and no evidence leaflet readable
343 - I	Other	309 / 149 / 137 Total sample available	Information intervention not associated with increase in advance directives. Note: questionnaire only offered to intervention group, no self-report data for controls
56 - I	Other	2452 / 1272 / 1260 Systematic sample	Attendance twice as high when free service provision (66% vs. 37%)
472 - I	Other	67 / 61 / 61 Total sample available	Intervention patients asked more questions at discharge (7.8 vs. 3.1; $p < 0.01$ ). An association between perception of physical and psychological functioning and intervention
72 - I	Other	? / 446 / 218 Convenience sample	Results for both groups at all time points not reported. No differences in smoking cessation by group across all time points. Operationalised stages of change model
242 - I	Patient	482 / 375 / 268 Convenience sample	Stages of change intervention associated with increased smoking cessation and more quitting attempts. Note: only two groups reported although three randomised
254 - I	Patient	3161 / 2707 / 2707 Total sample available	Prompts plus skill acquisition group increased cessation smoking twice as much as advice only or group session (from 6% to 12%)
375 - I	Patient	? / 28 / 28 Convenience sample	Intervention not associated with greater attendance at exercise classes. Purpose of study to assess increase in knowledge
792 - I	Other	? / 5458 / 2540 Total sample available	Theory-based intervention associated with increase in smoking cessation ( $p < 0.01$ )
640 - I	Other	3655 / 3655 / 2688 Total sample available	Intervention not associated with changes in tobacco use
287 - I	Patient	1119 / 1119 / 996 Total sample available	Intervention associated with increased smoking cessation (11% vs. 16%; $p < 0.009$ )
169 - I	Patient	1383 / 719 / 706 Total sample available	Intervention associated with increased adherence ( $p < 0.001$ )
405 - I	Other	? / 146 / 146 Not adequately described	Intervention not associated with changes in tobacco use
442 - I	Other	? / 2791 / 1222 Convenience sample	Intervention not associated with changes in CVD risk factors
360 - I	Other	? / 511 / 434 Not adequately described	Intervention not associated with changes in children's intention to change drug use behaviour. Reports changes in process measures
115 - I	Patient	328 / 328 / 225 Total sample available	Study operationalised the theory of planned behaviour and results analysed accordingly. Interventions not associated with differential adherence to diet. No control group
329 - I	Patient	193 / 193 / 109 Total sample available	No differential effect on cessation of smoking (42% vs. 38% vs. 48% quit)
80 - I	Patient	131 / 91 / 86 Total sample available	No association between intervention and changes in self-report smoking cessation (too few smokers), dietary change or uptake of exercise. Both groups changed diet and uptake of exercise
649 - I	Other	337 / 337 / 240 Total sample available	Intervention not associated with changes in safer sex practices

continued

**TABLE 33 contd** Description of the quality of the study and summary results: RCTc

Study number	Intervention level	Sample size	Summary of results
684 - 1	Patient	? / 407 / 301 Convenience sample	No differential effects of interventions on condom use
720 - 1	Other	? / 385 / 243 Total sample available	Analysis assessed relationship between social network and behaviour. Unclear relationship between intervention and behaviour
407 - 1	Other	? / 2144 / 1866 Convenience sample	Intervention associated with decreased in risky drug use ( $p < 0.05$ )
181 - 1	Patient	60 / 51 / 39 Total sample available	Authors suggest intervention associated with greater adherence ( $p < 0.05$ ). Unclear sample size in final analysis, large attrition
762 - 1	Patient	? / 106 / 76 Convenience sample	Interventions associated with increased acquisition of condoms and initiation of HIV conversations ( $p < 0.01$ ). Interventions not associated with self-reported behaviour change safer sex
528 - 1	Patient	907 / 693 / 381 Convenience sample	Authors claim intensive information associated with safer drug practices
646 - 1	Other	? / 233 / 213 Convenience sample	Intervention not associated with cessation of drug/ sex behaviours
696 - 1	Other	3348 / 3226 / 1665 Total sample available	Interventions associated with differential effect on uptake of free condoms (21% vs. 27% vs. 41%)
817 - 1	Other	567 / 411 / 339 Total sample available	No differential effect of interventions and risky behaviour. Both interventions associated with decreased risky behaviours
749 - 1	Patient	? / 157 / 157 Convenience sample	HIV risk intervention associated with appropriate change in sex behaviours ( $p < 0.01$ )
777 - 1	Other	? / 1417 / 1417 Total sample available	Intervention associated with increase in facewashing ( $p < 0.05$ )
172 - 1	Other	? / 1303 / 1303 Total sample available	Postcard reminders not associated with increased attendance. Predictors of non-attendance analysis carried out
363 - 1	Patient	? / 238 / 238 Not adequately described	Intervention not associated with change in immunisation uptake (31% vs. 33%)
470 - 1	Patient	207 / 177 / 157 Total sample available	Intervention associated with fewer problems but more health supervision surgery visits. No change in late and emergency visits
512 - 1	Other	8002 / 8002 / 6623 Total sample available	Odds ratios reported: intervention associated with increased immunisation of children; specific message more impact than general; invitation call greater impact than reminder
98 - 1	Patient	140 / 75 / 57 Convenience sample	Intervention associated with changes in self-management of arthritis but not exercise
601 - 1	Patient	? / 129 / 129 Convenience sample	Intervention increased request for information from 10% to 27%. No change in self-report of behaviour
601 - 2	Patient	? / 163 / 163 Convenience sample	Nurse intervention associated with increased question asking 8% to 54%
83 - 1	Patient	1566 / 549 / 401 Total sample available	Intervention group associated with fewer prenatal visits and ultrasounds. No differences in uptake serum screening or perinatal outcomes
106 - 1	Patient	? / 240 / 240 Not adequately described	Provision of toy associated with greater attendance post-partum. It is likely that the 'toy' group had more contact with health professional
788 - 1	Other	251 / 251 / 226 Total sample available	Additional information not associated with postpartum healthcare utilisation
481 - 1	Other	2008 / 2008 / 2008 Total sample available	Interventions not associated with differential effect attendance. Before/after analysis suggest reminder call increased attendance from 53% to 62%
411 - 1	Patient	? / 40 / 40 Not adequately described	Intervention associated with greater adherence TB medication (64% vs. 11%; $p < 0.001$ )

continued

**TABLE 33 contd** Description of the quality of the study and summary results: RCTc

Study number	Intervention level	Sample size	Summary of results
743 - I	Patient	? / 2002 / 1908 Not adequately described	Uptake of CF carrier screening not associated with type of screening technique. Purpose of study to assess impact testing
790 - I	Patient	5529 / 5529 / 5529 Not adequately described	Differential effects of interventions on uptake CF carrier testing (12% vs. 9% vs. 17% vs. 70% vs. 25% vs. 4%)
293 - I	Other	210 / 93 / 93 Total sample available	No differential effects between groups on use of car seats. Note: majority of sample using car seats
349 - I	Patient	202 / 93 / 61 Total sample available	Questionnaire plus leaflet reduced number of missed initial appointments and increased attendance for therapy at 6 months ( $p < 0.05$ )
58 - I	Other	5600 / ? / ? Systematic sample	No baseline figures, statistical tests or indication sample size. Twice as likely to sign donor card if 'intensive package' than information only or control group (5%, 5%, 12%, 13%)
324 - I	Patient	? / 149 / 98 Convenience sample	Illustrated leaflet associated with increased intention to check heat of water ( $p < 0.0001$ )
717 - I	Patient	74 / 74 / 74 Total sample available	Intervention associated with increased attendance (57% vs. 82%; $p < 0.05$ )



**TABLE 34** Description of the quality of the study and summary results: concurrent

Study number	Intervention level	Sample size	Summary of results
44 - 1	Other	1750 / ? / ? Systematic sample	No significant difference between sampled women's uptake of breast screening. Difficult to interpret results as an additional intervention conducted within the communities at the same time
61 - 1	Other	? / ? / ? Total sample available	Results from medical records by age. Difficult to assess significance association intervention with uptake rates (71% vs. 65%)
203 - 1	Other	1937 / 1697 / 1697 Total sample available	Intervention not associated with differential effects on uptake by group. Interaction between age and uptake of screening
243 - 1	Other	? / ? / ? Systematic sample	Community and physician intervention associated with increased attendance for mammography
246 - 1	Other	? / 1405 / 875 Systematic sample	Both intervention groups associated with increase in BSE
385 - 1	Other	? / 1200 / ? Not adequately described	No significant differences in tobacco use between intervention groups (18% vs. 20% vs. 15%)
686 - 1	Other	? / 970 / 970 Systematic sample	Intervention associated with increased mammography (from 35% to 55% vs. 30% to 40%)
171 - 1	Other	? / 1500 / 966 Systematic sample	Information intervention not associated with changes in smoking, sun-tanning or breast examination behaviours
205 - 1	Other	1346 / 1346 / 559 Total sample available	Community-based intervention associated with greater uptake than media intervention (50% vs. 32%)
205 - 2	Other	1013 / 1013 / 1013 Total sample available	GP intervention associated with greater attendance at breast cancer screening than community intervention (64% vs. 49%)
397 - 1	Other	? / ? / ? Convenience sample	Results not reported for uptake cervical and mammography screening by group. Note: awareness programme increased in both groups so contamination groups
546 - 1	Other	1616 / 1332 / 1332 Total sample available	Personalised invitation associated with greater attendance at cervical screening than national rate (55% vs. 41%)
448 - 1	Other	225 / 135 / 135 Total sample available	Skills group associated with increased uptake of colorectal cancer (65% vs. 41% vs. 94%); $p < 0.05$
418 - 1	Other	? / 294 / 294 Not adequately described	Intervention not associated with change in uptake of sigmoidoscopy examinations, immunisations, cholesterol screening or installation of smoke detectors
386 - 1	Other	? / 189 / 179 Not adequately described	Uptake and maintenance of BSE improved in both intervention groups ( $p < 0.001$ ). No differential effect between intervention groups
760 - 1	Other	? / 615 / 295 Convenience sample	Differential effects of interventions on attendance for mammography (10% vs. 30% vs. 34%; $p < 0.01$ ). Operationalised HBM
608 - 1	Other	1194 / 628 / 628 Convenience sample	Media-based intervention associated with increase in mammography uptake (15% vs. 24%; $p < 0.05$ )
404 - 1	Other	? / 1857 / 1857 Total sample available	Women within the intervention community more likely to report having a mammogram (30% vs. 19% increase; $p < 0.05$ ). Analysis carried out to assess predictors attendance
446 - 1	Other	? / 2724 / 2724 Systematic sample	Intervention associated with changes in mammography uptake (10% vs. 27%; $p < 0.01$ )
136 - 1	Other	648 / 648 / 612 Systematic sample	Group 2 associated with increased sun protection behaviours over time ( $p < 0.01$ )
237 - 1	Other	2000 / 1506 / 1231 Systematic sample	Intervention associated with a decreased fat intake over time ( $p < 0.01$ ). No change in smoking habits
180 - 1	Other	? / ? / ? Not adequately described	Self-report hat wearing and sunscreen increased, no change in covering up after intervention. Less change in men's behaviour
45 - 1	Other	? / ? / ? Systematic sample	Self-report attending cholesterol checks higher in white intervention than control population. No increase amongst black respondents. Sample randomly selected by phone

continued

**TABLE 34 contd** Description of the quality of the study and summary results: concurrent

Study number	Intervention level	Sample size	Summary of results
82 - I	Patient	? / 84 / 84 Convenience sample	Patients within intervention group fewer emergency visits and days off work than matched controls. No differences between number hospitalisations. Note: data suggest non-parametric analysis more appropriate
218 - I	Other	2812 / 1506 / 1093 Total sample available	Intervention not associated with reliable differences in smoking cessation
222 - I	Other	286 / 209 / 125 Total sample available	Screening teachers associated with uptake of exercise but no change in smoking behaviour
235 - I	Other	1052 / 966 / 931 Not adequately described	Prevalence of smoking not associated with type of work policy; daily rate cigarette consumption associated with type of work policy
245 - I	Patient	? / 62 / ? Convenience sample	Intervention groups not associated with smoking cessation at 6 months follow-up. Caution: hospital had a no-smoking policy
256 - I	Patient	1946 / 1286 / 997 Total sample available	Cessation rates: advice only (13%), counselling only (24%), counselling and gum (30%). Logistic regression analysis modelling demographic variables and cessation
258 - I	Other	3558 / 708 / 708 Convenience sample	Authors suggest intervention associated with changes in CVD risk factors. Caution: low response rate over large period of time
260 - I	Patient	? / 8184 / 8184 Not adequately described	Special care intervention reported greater change in smoking cessation than usual care (43% vs. 14%). Little information about intervention or process measures of change
280 - I	Other	16089 / 3382 / 2972 Convenience sample	Participation in survey greater in North Karelia (0.6% vs. 0.3%, 0.3%). Reported change cessation smoking higher in North Karelia (25% vs. 20%, 19%)
283 - I	Other	5445 / 56 / 56 Convenience sample	Competition intervention associated with greater interest in programme (2% vs. 0.6%). Maintenance abstinence greater in competition intervention ( $p < 0.05$ )
284 - I	Other	? / 2147 / 1220 Convenience sample	Cessation of smoking greater within intervention group (32% vs. 17%; $p < 0.05$ )
315 - I	Other	96 / 84 / 55 Total sample available	Packaging pills in foil holders with 'date to take' increased compliance up to 3 months after discharge (49% vs. 23%)
390 - I	Other	? / ? / 1880 Convenience sample	Authors claim counselling and exercise compatible with daily life is effective in increasing exercise uptake. Note: results not clearly reported and unclear
504 - I	Other	? / 46 / 46 Total sample available	Intervention associated with fewer GP contacts and emergency visits ( $p < 0.01$ ) but no change in referrals to specialists
657 - I	Patient	? / 4487 / 2767 Systematic sample	Intervention group three times more likely to quit smoking (6% vs. 2%)
731 - I	Other	? / 5078 / 4538 Systematic sample	Interventions not associated with changes in smoking behaviour
740 - I	Other	? / ? / ? Not adequately described	Authors claim that smoking cessation greater within intervention community but analysis not carried out to illustrate this
779 - I	Other	1087 / 1036 / 1036 Total sample available	Intervention village greater long-term smoking cessation rate ( $p = 0.01$ ) but no difference in short-term cessation rates
206 - I	Other	? / ? / ? Not adequately described	Intervention associated with increased smoking cessation in men only (8% vs. 3%). Results presented for physiological estimates CVD risk
215 - I	Other	? / 1500 / ? Systematic sample	Authors claim Group 1 associated diet change. However, small sub-section of sample. Group 2 and 3 not associated with changes
409 - I	Other	? / ? / 1863 Convenience sample	No differential effect of information on children's report of diet. Note: differences between trained/not trained teachers, e.g. funding schools; inappropriate statistics
312 - I	Other	? / ? / ? Systematic sample	Intervention associated with reduction in fatty diet ( $p < 0.05$ )
273 - I	Other	? / 4270 / 2605 Systematic sample	Intervention not associated with cessation of smoking, though there was a time effect ( $p < 0.01$ )

continued

**TABLE 34 contd** Description of the quality of the study and summary results: concurrent

Study number	Intervention level	Sample size	Summary of results
394 - 1	Patient	? / 1024 / 860 Convenience sample	Intervention not associated with change in ophthalmic specialist use
194 - 1	Other	203 / 184 / ? Total sample available	Authors present changes in intention to behave more safely for: seat-belt use as a driver and passenger, speeding and drink-driving. Intervention associated with increased intention of passenger seat-belt use
89 - 1	Patient	738 / 227 / 227 Convenience sample	Authors suggest post-intervention group had greater intention to change exercise behaviour than pre-intervention group ( $p < 0.05$ ). Operationalise the theory planned behaviour
184 - 1	Other	276 / 232 / 146 Total sample available	Interventions not associated with differential effect of smoking cessation but reduction in overall pre-/post-intervention
253 - 1	Other	? / 465 / 346 Convenience sample	Parent information associated with greater increase of intention and actual food buying than child/control groups. Operationalised SLT
345 - 1	Other	252 / 252 / 187 Total sample available	Intervention not associated with increased smoking cessation rates. Operationalised stages of change theory
416 - 1	Patient	? / 48 / 48 Convenience sample	Interventions not associated with changes in self-report behaviour
440 - 1	Other	? / 2500 / ? Systematic sample	Behavioural-based intervention assessed mainly by physiological changes. There was no association with smoking cessation
209 - 1	Other	407 / 255 / 212 Convenience sample	Stages of change intervention associated with increased exercise uptake ( $p < 0.01$ )
631 - 1	Other	? / ? / 6814 Systematic sample	Intervention not associated with reliable changes in self-report of diet change at 5 years
464 - 1	Other	? / 907 / 907 Systematic sample	Intervention not associated with exercise adherence but effects reported for subsidiary analysis
658 - 1	Other	? / 2136 / 1674 Systematic sample	Intervention increased attendance at lifestyle change activities
726 - 1	Other	? / ? / ? Systematic sample	Little association between intervention and self-reports of smoking or exercise behaviour
255 - 1	Other	? / ? / ? Systematic sample	Intervention not associated with changes in tobacco use
313 - 1	Other	625 / 558 / 516 Total sample available	Both interventions associated with increased medication adherence and self-monitoring ( $p < 0.05$ ). Neither intervention associated with changes in diet and exercise
367 - 2	Patient	? / 341 / 119 Convenience sample	Differential attrition (56% vs. 48%) and smoking cessation rates (14% vs. 57%). Difficult to draw any conclusions as poor design
370 - 1	Other	? / 5356 / 3833 Total sample available	Intervention associated with decreased self-report of alcohol use ( $p < 0.01$ )
238 - 1	Other	? / ? / ? Systematic sample	Intervention not associated with changes in smoking behaviour
261 - 1	Other	171 / 171 / 157 Total sample available	Intervention associated with favourable changes in smoking, diet and dental health ( $p < 0.01$ )
456 - 1	Other	405 / 155 / 55 Not adequately described	Intervention associated with changes in diet ( $p < 0.05$ ) but not smoking or uptake exercise
709 - 1	Other	? / ? / ? Systematic sample	Intervention not associated with changes in diet, smoking, exercise. Interaction between social support and intervention
311 - 1	Other	? / ? / 2863 Not adequately described	Authors claim social intervention associated with lower onset substance use than control, whereas the affective intervention group was associated with increased substance use
125 - 1	Patient	? / 117 / 77 Convenience sample	Intervention associated with intention to practice safer sex. No association with drug or alcohol use

continued

**TABLE 34 contd** Description of the quality of the study and summary results: concurrent

Study number	Intervention level	Sample size	Summary of results
161 - I	Other	? / 600 / 600 Convenience sample	Intervention associated with increased condom use and reduced number of sexual partners ( $p < 0.001$ )
177 - I	Patient	? / 2624 / 1616 Convenience sample	Results not reported for comparison interventions but as before/ after design, predictors of behaviour change. Cessation of risky needle use and adherence to safer sex in males after intervention
392 - I	Other	? / 7946 / 7946 Systematic sample	Uptake of contraception use in intervention area over time greater than control community (57% vs. 27%)
791 - I	Patient	? / 541 / 524 Convenience sample	Intervention associated with increased use of condoms and raising issue of condom use with clients ( $p < 0.01$ )
555 - I	Other	? / 1398 / 1398 Convenience sample	Authors claim intervention not associated with use of condoms but increase in intention to use condoms
665 - I	Other	? / 1267 / 1267 Convenience sample	Authors claim reduction in self-report of risky sex practices within intervention city only
579 - I	Other	1226 / 442 / 442 Convenience sample	Information intervention not associated with changes in condom use
748 - I	Patient	? / 209 / 20 Convenience sample	Group 1 associated with reduced length of hospital stay and fewer outpatient visits ( $p < 0.01$ ) but increased GP visits for Groups 1 and 4
455 - I	Other	? / 1261 / 1256 Not adequately described	Control group associated with increased utilisation of care ( $p < 0.01$ ), intervention group less likely to change method of contraception (85% vs. 92%, $p < 0.01$ )
71 - I	Not recorded	29 / 22 / 22 Convenience sample	Those with lower addiction (baseline group 1) were more likely to attend antenatal classes. Those with a higher addiction (intensive Group 2) expressed more intention to quit. Design of study unable to evaluate intervention
486 - I	Patient	? / 248 / 113 Not adequately described	Intervention associated with greater number of children adopted (43% vs. 0%), less sexually active (57% vs. 75%) at 24 months. Low response rate and samples significantly different at beginning of study
17 - I	Other	738 / 734 / 520 Total sample available	Authors suggest intention to use condoms and practice safer sex increased after role play. Carried out attitude-behaviour analysis
18 - I	Other	173 / 120 / 80 Systematic sample	Intervention associated with less intention to use steroids. Operationalised model
240 - I	Other	? / 2844 / 1477 Not adequately described	Theory-based (skills) intervention associated with increased use of condoms, reduction in number of partners and behavioural intentions. Operationalised theory of reasoned action
251 - I	Other	? / 5378 / 3875 Not adequately described	Authors claim programme associated with increased smoking cessation in intervention school
391 - I	Patient	? / 763 / 763 Systematic sample	Authors suggest intervention associated with increased condom use (42% vs. 30%). Unclear description of design either before/after or concurrent
515 - I	Other	? / 2026 / 1785 Systematic sample	Intervention associated with greater intention to practice safer sex. Operationalisation theory planned behaviour
759 - I	Other	? / 6573 / 6573 Not adequately described	Intervention not associated with changes in behaviour, though control schools increased sexual activity and intervention group remained the same. Analysis looked at predictors of sexual activity
628 - I	Other	? / 197 / 145 Convenience sample	Authors claim intervention associated with increased consistent condom use ( $p < 0.01$ )
780 - I	Other	? / 2239 / 2239 Systematic sample	Intervention not associated with changes in risky sex
463 - I	Other	? / 298 / 191 Convenience sample	Intervention community decreased number of risky sex behaviours, not changed in comparison community
663 - I	Other	880 / 845 / 804 Systematic sample	Intervention not associated with intention to change behaviour ( $p = 0.07$ )
373 - I	Other	? / 1444 / 888 Convenience sample	Intervention not associated with sexual activity. Analysis operationalised HBM

continued

**TABLE 34 contd** Description of the quality of the study and summary results: concurrent

Study number	Intervention level	Sample size	Summary of results
453 - I	Other	? / 377 / 161 Convenience sample	Both intervention groups associated with changes in alcohol use but not cigarette or marijuana use ( $p < 0.05$ )
822 - I	Other	? / ? / ? Convenience sample	Differential effects between groups and in comparison with intervention groups for safer sex
823 - I	Other	? / ? / 4806 Not adequately described	Intervention associated with changes in smoking and drug use ( $p < 0.05$ )
813 - I	Other	5686 / 3779 / 1786 Total sample available	Intervention associated with changes in seat belt use and diet ( $p < 0.01$ ). No changes for smoking, exercise or alcohol
146 - I	Other	2978 / 2978 / 2978 Total sample available	Authors suggest intervention associated with increased uptake of influenza vaccination (40% to 56%). Main analysis was predictors uptake
257 - I	Other	? / ? / 3884 Total sample available	More aggressive recruitment associated with greater attendance for health check (14%, 21%, 32%, 37%)
775 - I	Other	3616 / 3616 / 3616 Total sample available	Children more likely to be immunised in general practice than in health centre (80% vs. 68%). Analysis assessed differences between rural and inner city locations
331 - I	Patient	? / 80 / 80 Not adequately described	Adherence to routine health/immunisation screenings was not associated with immunisation in A&E
95 - I	Patient	? / 175 / ? Not adequately described	No improvement in self-management behaviours by intervention group. Main outcomes were knowledge and satisfaction
3 - I	Other	? / 256 / 206 Not adequately described	Intervention associated with an increase in breast feeding
529 - I	Other	? / 2610 / 2508 Systematic sample	Intervention associated with increased chewing of xylitol ( $p < 0.05$ )
618 - I	Other	? / ? / ? Convenience sample	Intervention not associated with increased use of reimbursed dental service
430 - I	Patient	? / ? / 230 Not adequately described	Differences in reproductive choices found between counselled/not counselled. Note: not a controlled sample
516 - I	Patient	? / 1607 / 1607 Total sample available	Health centre intervention higher uptake than letter invitation (24% vs. 4%). Assessed process variables
323 - I	Other	9150 / 1265 / 551 Total sample available	Both information interventions associated with higher recruitment to study and adherence at 6 months, no differential effect between intervention groups
586 - I	Patient	161 / 127 / 127 Convenience sample	Observation of medication taking associated with greater adherence (88% vs. 50%)
452 - I	Other	1300 / 1300 / 1300 Total sample available	Intervention associated with increased completion treatment (65% vs. 78%; $p < 0.05$ )
124 - I	Patient	? / 93 / 93 Convenience sample	Whether drug was taken orally or injected had no impact on medication adherence post-discharge
825 - I	Not recorded	? / ? / ? Not adequately described	Authors claim intervention associated with increased adherence
335 - I	Other	? / ? / 334 Convenience sample	Intervention women more likely to assemble rehydration solution accurately
570 - I	Other	? / 106 / 106 Not adequately described	Intervention not associated with any changes in risk taking behaviours
773 - I	Other	1620 / 1168 / 1043 Total sample available	No association between intensive intervention and increased adherence to helmet use
40 - I	Not recorded	? / 72 / ? Convenience sample	More symptoms noted when using computer also preferred method
75 - I	Patient	? / 108 / 108 Convenience sample	Insured patients more likely to keep appointments, less likely to make appointments and willing to pay more for insurance

**TABLE 35** Description of the quality of the study and summary results: before/after different samples

Study number	Intervention level	Sample size	Summary of results
34 - I	Other	? / ? / ? Not adequately described	Mammography uptake improved over 5 years
100 - I	Patient	? / ? / ? Convenience sample	Uptake rate for mammography increased from 47% to 72% a year
192 - I	Other	410 / 316 / ? Total sample available	Cervical screening increased from 51% to 78%
383 - I	Other	? / 516 / 516 Systematic sample	Intervention not associated with uptake of mammography (47% vs. 41%: aware intervention vs. not aware)
421 - I	Other	? / ? / ? Not adequately described	Actual results not reported but authors claim intervention associated with increased BSE
575 - I	Patient	? / 2972 / 2972 Systematic sample	Intervention associated with increased smear uptake (20% to 58%)
401 - I	Other	985 / 656 / 656 Total sample available	Trained health professionals and provision prompts increased referral and attendance in women for mammography (55% to 88%)
765 - I	Patient	618 / 109 / 109 Convenience sample	Authors claim intervention associated increase sun protective behaviours (6.5% to 27%)
495 - I	Patient	? / 533 / 533 Systematic sample	Intervention not associated with increase in making living wills. Analysis looked at predictors of living will completion
340 - I	Other	? / 7667 / 7667 Systematic sample	Authors claim smoking decreased over a 4-year period. Actual percentages not reported and likely self-selecting sample bias
423 - I	Patient	912 / 706 / 706 Total sample available	Authors claim media interventions associated with increase testing for HIV but no increase in number of positives
550 - I	Other	344 / 314 / 314 Total sample available	No-smoking policy not associated changes marijuana and alcohol use. Associated with cessation of cigarette use ( $p = 0.02$ )
514 - I	Not recorded	? / 1232 / 1232 Systematic sample	Intervention not associated cessation smoking
461 - I	Other	? / 15188 / 15188 Systematic sample	Self report increase seat belt use (17% more) and reduction drunk driving (40% less) greater in intervention city
462 - I	Other	? / 2016 / 2016 Systematic sample	Intervention community associated increase physical activity and cholesterol checks ( $p < 0.05$ ), no change smoking cessation or fruit consumption
670 - I	Patient	? / ? / ? Convenience sample	Analysis suggests greatest change in testing after television campaigns
680 - I	Patient	? / ? / ? Convenience sample	Authors claim interventions associated with decreased risky sexual behaviours. Interactions with ethnicity and sexual orientation
662 - I	Patient	? / 276 / 266 Convenience sample	Announcement associated with greater intention to test for HIV ( $p < 0.01$ )
142 - I	Other	? / ? / ? Convenience sample	Authors claim changes in risky sex and safe drug taking practices
619 - I	Other	? / 1515 / 1515 Convenience sample	Surveys demonstrated no change in behaviour over the two time points. Behaviour change probably made before first survey
647 - I	Other	? / 1469 / 1469 Not adequately described	Intervention associated reduction high risk behaviours in all three cities
193 - I	Patient	? / ? / ? Systematic sample	Intervention not associated with an increase in childhood immunisations
767 - I	Other	? / ? / ? Total sample available	Intervention associated increase immunisation (53% to 78%). Analysis assessed uptake by deprivation indices

continued

**TABLE 35 contd** Description of the quality of the study and summary results: before/after different samples

Study number	Intervention level	Sample size	Summary of results
413 - I	Other	? / 299 / 299 Total sample available	Intervention associated increase vaccination from 50% to 73%
623 - I	Other	? / 652 / 524 Not adequately described	Intervention not associated changes in breast feeding behaviour
26 - I	Other	167 / 167 / 167 Total sample available	Intervention associated increase breast feeding in hospital, not maintained at home
97 - I	Other	? / ? / ? Convenience sample	More people reported brushing teeth and buying toothbrushes /fluorinated toothpaste. No reduction in those that do not attend a health professional
737 - I	Patient	? / 604 / 604 Not adequately described	Adherence to treatment increased between two time points
563 - I	Patient	? / 928 / 928 Not adequately described	Use of mobile clinic increased completion of examination forms and adherence to treatment (from 9% to 91%)
239 - I	Other	6168 / 5925 / 5385 Total sample available	Risky child sleep position decreased from 30% to 5%
294 - I	Other	? / ? / ? Convenience sample	Authors claim 523 people requested fragile X information, no increase uptake diagnostic tests. Note: no baseline, post intervention or population figures given to support authors claims
674 - I	Patient	? / 1339 / 1339 Total sample available	Introduction clinic associated with fewer non-attendees (0.3% vs. 4.3%)
149 - I	Other	? / 783 / 783 Convenience sample	Authors claim helmet use increased

**TABLE 36** Description of the quality of the study and summary results: before/after same sample

Study number	Intervention level	Sample size	Summary of results
99 - I	Patient	1087 / 727 / 526 Convenience sample	Intention to attend for mammography increased (60% vs. 51%; $p < 0.01$ ). Analysis performed by intention to have vs. not to have
266 - I	Patient	1592 / 1592 / 1022 Total sample available	Increased sun protective behaviours were observed over 6 years. Unable to attribute to clinic intervention alone
746 - I	Patient	1628 / 1000 / 100 Systematic sample	Intervention associated with increased uptake of cervical and mammography screening. However, less than target rates
815 - I	Patient	? / 127 / 127 Not adequately described	Authors claim increased uptake of testicular cancer screening over three years ( $p < 0.01$ )
795 - I	Other	9650 / 9650 / 9650 Total sample available	Cervical smear uptake increased from 78% to 85% (80% target rate)
809 - I	Other	? / 268 / 268 Systematic sample	Intervention associated with increased mammography uptake. Results presented by age group
133 - I	Other	? / 1600 / ? Systematic sample	Intervention not associated with changes in health behaviours over time
466 - I	Other	? / ? / ? Total sample available	Prompt on notes associated with increased attendance for screening and immunisations
46 - I	Patient	? / 10 / ? Not adequately described	Study designed to compare a psycho-educational programme and control but no measure of behaviour. Before/after results associated with increased BSE and mammography
126 - I	Patient	3686 / 1585 / 948 Total sample available	Authors suggest intervention associated with uptake of mammography. Results analysed using age and income interactions
198 - I	Patient	320 / 200 / 178 Total sample available	Intervention associated with increased safer sun practices ( $p < 0.05$ )
42 - I	Not recorded	? / 110 / 102 Not adequately described	Intervention associated with decreased GP visits, increased use of prevention medication, decreased use of other medication
74 - I	Patient	? / 15 / 15 Not adequately described	Actual results not reported, authors claim participants more likely to test blood and adjust insulin level after programme ( $p = 0.01$ )
79 - I	Other	? / 70 / 54 Convenience sample	Claimed to be comparative design but only results of before/after measures intervention group reported. Intervention associated with changes in diet adherence but not exercise
111 - I	Patient	57 / 43 / 31 Convenience sample	Receipt of information on living wills increased patients completion of wills and discussions with family members and health professionals
244 - I	Patient	? / 396 / 257 Convenience sample	Intervention associated with increased exercise uptake
270 - I	Patient	111 / 89 / 89 Total sample available	Intervention associated with 25% smoking cessation rate. Analysis assessed predictors of cessation
355 - I	Patient	? / 94 / 38 Convenience sample	Intervention associated with increased exercise uptake. Analysis assessed attendees vs. non-attendees
384 - I	Patient	? / 2244 / 338 Convenience sample	Intervention associated with changes in diet and exercise uptake but not smoking
521 - I	Patient	208 / 186 / 148 Total sample available	Intervention associated with increased self-management behaviours ( $p < 0.01$ )
676 - I	Patient	223 / 38 / 30 Convenience sample	Intervention associated with fewer emergency visits, improved use of inhalers and appropriate self-administration of steroids
522 - I	Patient	2500 / 588 / 300 Convenience sample	Intervention not associated with changes in medication use or utilisation of health services
544 - I	Not recorded	142 / 142 / 132 Total sample available	Authors report adherence to medication increased, reduction in visits by health professionals ( $p < 0.001$ )

continued



**TABLE 36 contd** Description of the quality of the study and summary results: before/after same sample

Study number	Intervention level	Sample size	Summary of results
262 - I	Patient	2215 / 1113 / 1097 Convenience sample	Intervention associated with 13% cessation rate. Only 5% of smokers who did not take part in intervention quit
449 - I	Patient	? / 78 / 53 Convenience sample	Intervention associated with increased adherence from 73% to 83% ( $p < 0.01$ )
798 - I	Patient	1006 / 526 / 520 Total sample available	Self-reported changes at follow-up: exercise (11%), smoking (15%), alcohol (24%), diet (less fat 46%)
821 - I	Patient	? / 68 / 45 Not adequately described	Authors claim intervention associated with implementation of exercise programme at home and smoking cessation
621 - I	Other	? / 937 / 596 Convenience sample	Participants more likely to adopt new diet (16% vs. 1%). Poor design but looked at predictors of uptake
348 - I	Patient	2324 / 771 / 512 Total sample available	Authors claim intervention associated with changes in osteoporosis prevention activities. Analysis assessed differences in attendees/non-attendees
162 - I	Patient	160 / 3 / 3 Not adequately described	Authors claim intervention associated with increased healthy diet choice
84 - I	Patient	? / ? / 274 Convenience sample	Intervention associated with changes in self-report of car safety
594 - I	Patient	? / 83 / 64 Convenience sample	Interactive video associated with uptake of medication from 0% to 25% of patients. Study assessed several decision-making process measures
179 - I	Patient	? / 7 / 5 Convenience sample	Sample selection of before/after design too small to draw any conclusions
367 - I	Patient	? / 112 / 78 Convenience sample	15% sample quit after the programme
91 - I	Patient	? / 20 / ? Convenience sample	Intervention associated with increased self-report of safe sex behaviours, cessation of risky sex
38 - I	Not recorded	? / 143 / 56 Convenience sample	Intervention associated with increased use of condoms and decreased risky sex encounters
49 - I	Not recorded	1180 / 1161 / ? Total sample available	Intervention associated with intentions to engage in safer sex, in one of the two schools. No association in the other school
123 - I	Patient	? / 239 / 139 Convenience sample	Intervention associated with increased self-report use of condoms and decreased anal intercourse
159 - I	Patient	202 / 193 / 147 Convenience sample	Intervention not associated with changes in risky sexual behaviours. Assessed predictors of behaviour
480 - I	Patient	414 / 414 / 241 Total sample available	Intervention associated with reduction in sexual activity (33% to 25%)
651 - I	Patient	3702 / 1458 / 1385 Systematic sample	Use of condoms increased from 7% to 22% after intervention
655 - I	Patient	? / 586 / 502 Convenience sample	Intervention associated with increased uptake of hormonal contraception
688 - I	Patient	? / 195 / 149 Convenience sample	Intervention associated with increased condom use (62% to 82%, $p = 0.01$ )
729 - I	Patient	? / 659 / 402 Convenience sample	Authors claim intervention associated with some change in drug use behaviours
770 - I	Patient	? / 1113 / 983 Convenience sample	Authors claim a mixed association between intervention and drug use behaviours
610 - I	Patient	? / ? / 208 Convenience sample	Authors claim intervention associated with increased intention to use condom (44% vs. 65%) and reduce number of partners (43% vs. 53%)

continued

**TABLE 36 contd** Description of the quality of the study and summary results: before/after same sample

Study number	Intervention level	Sample size	Summary of results
467 - I	Other	? / 2378 / 1688 Total sample available	Intervention associated with greater intention to engage in safe sex
723 - I	Patient	? / 292 / 134 Not adequately described	Intervention associated with increased use condom (64% to 70%)
545 - I	Patient	? / 755 / 61 Not adequately described	Authors claim association between counselling and reduction risky sex
650 - I	Patient	814 / ? / 614 Total sample available	Clinic information not associated with changes in risk behaviours
537 - I	Patient	? / 700 / 700 Total sample available	Intervention not associated with increased attendance at clinic. No differences in attendance between sexes
35 - I	Other	97 / 77 / 31 Total sample available	Small sample size and self-selected. Self-report of condom use increased and discussions of safer sex increased after the intervention
364 - I	Patient	? / 109 / 109 Convenience sample	Intervention associated with increased intention to use condoms. Operationalised social cognitive theory
526 - I	Patient	141 / 138 / 119 Total sample available	Intervention associated with changes in risky sex behaviours. Analysis assessed predictors behaviour change, operationalised five models
654 - I	Patient	? / 327 / 327 Convenience sample	Authors claim intervention associated with intention to change sex practices ( $p < 0.01$ )
450 - I	Patient	? / 93 / 69 Convenience sample	Authors claim intervention associated with risk-reducing behaviours
573 - I	Patient	? / 1915 / 1145 Total sample available	Intervention associated with 20% increased uptake of Hepatitis B vaccination ( $p < 0.01$ )
342 - I	Patient	? / 195 / 195 Total sample available	Uptake of immunisation increased from 8% to 46%
132 - I	Patient	893 / 893 / 893 Total sample available	Breast feeding increased from 18% to 52%
163 - I	Patient	? / 11 / 11 Convenience sample	Authors claim intervention associated with increased hygiene activities
361 - I	Patient	? / 11 / 11 Not adequately described	Authors claim intervention associated with increased appropriate behaviour (library visit)

# Health Technology Assessment panel membership

This report was identified as a priority by the Screening Panel.

## Acute Sector Panel

### Current members

<b>Chair:</b> <b>Professor Francis H Creed,</b> University of Manchester	Dr Katherine Darton, M.I.N.D. Mr John Dunning, Papworth Hospital, Cambridge	Ms Grace Gibbs, West Middlesex University Hospital NHS Trust	Dr Duncan Keeley, General Practitioner, Thame
Professor Clifford Bailey, University of Leeds	Mr Jonathan Earnshaw, Gloucester Royal Hospital	Dr Neville Goodman, Southmead Hospital Services Trust, Bristol	Dr Rajan Madhok, East Riding Health Authority
Ms Tracy Bury, Chartered Society of Physiotherapy	Mr Leonard Fenwick, Freeman Group of Hospitals, Newcastle-upon-Tyne	Professor Mark P Haggard, MRC	Dr John Pounsford, Frenchay Hospital, Bristol
Professor Collette Clifford, University of Birmingham	Professor David Field, Leicester Royal Infirmary	Professor Robert Hawkins, University of Manchester	Dr Mark Sculpher, University of York
			Dr Iqbal Sram, NHS Executive, North West Region

### Past members

Professor John Farndon, University of Bristol*	Professor Cam Donaldson, University of Aberdeen	Mrs Wilma MacPherson, St Thomas's & Guy's Hospitals, London	Professor Michael Sheppard, Queen Elizabeth Hospital, Birmingham
Professor Senga Bond, University of Newcastle- upon-Tyne	Professor Richard Ellis, St James's University Hospital, Leeds	Dr Chris McCall, General Practitioner, Dorset	Professor Gordon Stirrat, St Michael's Hospital, Bristol
Professor Ian Cameron, Southeast Thames Regional Health Authority	Mr Ian Hammond, Bedford & Shires Health & Care NHS Trust	Professor Alan McGregor, St Thomas's Hospital, London	Dr William Tarnow-Mordi, University of Dundee
Ms Lynne Clemence, Mid-Kent Health Care Trust	Professor Adrian Harris, Churchill Hospital, Oxford	Professor Jon Nicholl, University of Sheffield	Professor Kenneth Taylor, Hammersmith Hospital, London
	Dr Gwyneth Lewis, Department of Health	Professor John Norman, University of Southampton	

## Diagnostics and Imaging Panel

### Current members

<b>Chair:</b> <b>Professor Mike Smith,</b> University of Leeds	Dr Barry Cookson, Public Health Laboratory Service, Colindale	Mrs Maggie Fitchett, Association of Cytogeneticists, Oxford	Professor Chris Price, London Hospital Medical School
Dr Philip J Ayres, Leeds Teaching Hospitals NHS Trust	Professor David C Cumberland, University of Sheffield	Dr Peter Howlett, Portsmouth Hospitals NHS Trust	Dr William Rosenberg, University of Southampton
Dr Paul Collinson, Mayday University Hospital, Thornton Heath	Professor Adrian Dixon, University of Cambridge	Professor Alistair McGuire, City University, London	Dr Gillian Vivian, Royal Cornwall Hospitals Trust
	Mr Steve Ebdon-Jackson, Department of Health	Dr Andrew Moore, Editor, <i>Bandolier</i>	Dr Greg Warner, General Practitioner, Hampshire
		Dr Peter Moore, Science Writer, Ashtead	

### Past members

Professor Michael Maisey, Guy's & St Thomas's Hospitals, London*	Professor MA Ferguson-Smith, University of Cambridge	Professor Donald Jeffries, St Bartholomew's Hospital, London	Professor John Stuart, University of Birmingham
Professor Andrew Adam, Guy's, King's & St Thomas's School of Medicine & Dentistry, London	Dr Mansel Hacney, University of Manchester	Dr Ian Reynolds, Nottingham Health Authority	Dr Ala Szczepura, University of Warwick
Dr Pat Cooke, RDRD, Trent Regional Health Authority	Professor Sean Hilton, St George's Hospital Medical School, London	Professor Colin Roberts, University of Wales College of Medicine	Mr Stephen Thornton, Cambridge & Huntingdon Health Commission
Ms Julia Davison, St Bartholomew's Hospital, London	Mr John Hutton, MEDTAP International Inc., London	Miss Annette Sergeant, Chase Farm Hospital, Enfield	Dr Jo Walsworth-Bell, South Staffordshire Health Authority

\* Previous Chair  
continued

continued

## Methodology Panel

### Current members

<b>Chair:</b> <b>Professor Martin Buxton</b> , Brunel University	Professor Ann Bowling, University College London Medical School	Professor Jeremy Grimshaw, University of Aberdeen	Dr Nick Payne, University of Sheffield
Professor Doug Altman, Institute of Health Sciences, Oxford	Dr Mike Clarke, University of Oxford	Dr Stephen Harrison, University of Leeds	Professor Margaret Pearson, NHS Executive North West
Dr David Armstrong, Guy's, King's & St Thomas's School of Medicine & Dentistry, London	Professor Michael Drummond, University of York	Mr John Henderson, Department of Health	Professor David Sackett, Centre for Evidence Based Medicine, Oxford
Professor Nick Black, London School of Hygiene & Tropical Medicine	Dr Vikki Entwistle, University of Aberdeen	Professor Richard Lilford, Regional Director, R&D, West Midlands	Dr PAG Sandercock, University of Edinburgh
	Professor Ewan Ferlie, Imperial College, London	Professor Theresa Marteau, Guy's, King's & St Thomas's School of Medicine & Dentistry, London	Dr David Spiegelhalter, Institute of Public Health, Cambridge
	Professor Ray Fitzpatrick, University of Oxford	Dr Henry McQuay, University of Oxford	Professor Joy Townsend, University of Hertfordshire

### Past members

Professor Anthony Culyer, University of York*	Professor George Davey-Smith, University of Bristol	Mr Nick Mays, King's Fund, London	Professor Charles Warlow, Western General Hospital, Edinburgh
Professor Michael Baum, Royal Marsden Hospital	Professor Stephen Frankel, University of Bristol	Professor Ian Russell, University of York	
Dr Rory Collins, University of Oxford	Mr Philip Hewitson, Leeds FHSA	Dr Maurice Slevin, St Bartholomew's Hospital, London	

## Pharmaceutical Panel

### Current members

<b>Chair:</b> <b>Professor Tom Walley</b> , University of Liverpool	Professor Rod Griffiths, NHS Executive West Midlands	Dr Andrew Mortimore, Southampton & SW Hants Health Authority	Dr Frances Rotblat, Medicines Control Agency
Dr Felicity Gabbay, Transcrip Ltd	Mrs Jeanette Howe, Department of Health	Mr Nigel Offen, Essex Rivers Healthcare, Colchester	Dr Eamonn Sheridan, St James's University Hospital, Leeds
Mr Peter Golightly, Leicester Royal Infirmary	Professor Trevor Jones, ABPI, London	Mrs Marianne Rigge, The College of Health, London	Mrs Katrina Simister, Liverpool Health Authority
Dr Alastair Gray, Health Economics Research Unit, University of Oxford	Ms Sally Knight, Lister Hospital, Stevenage	Mr Simon Robbins, Camden & Islington Health Authority, London	Dr Ross Taylor, University of Aberdeen

### Past members

Professor Michael Rawlins, University of Newcastle- upon-Tyne*	Ms Christine Clark, Hope Hospital, Salford	Dr Tim Elliott, Department of Health	Dr John Posnett, University of York
Dr Colin Bradley, University of Birmingham	Mrs Julie Dent, Ealing, Hammersmith & Hounslow Health Authority, London	Dr Desmond Fitzgerald, Mere, Bucklow Hill, Cheshire	Dr Tim van Zwanenberg, Northern Regional Health Authority
Professor Alasdair Breckenridge, RDRD, Northwest Regional Health Authority	Mr Barrie Dowdeswell, Royal Victoria Infirmary, Newcastle-upon-Tyne	Professor Keith Gull, University of Manchester	Dr Kent Woods, RDRD, Trent RO, Sheffield
		Dr Keith Jones, Medicines Control Agency	

## Population Screening Panel

### Current members

<b>Chair:</b> <b>Professor Sir John Grimley Evans,</b> Radcliffe Infirmary, Oxford	Professor Howard Cuckle, University of Leeds	Professor Dian Donnai, St Mary's Hospital, Manchester	Professor Alexander Markham, St James's University Hospital, Leeds
Ms Stella Burnside, Altnagelvin Hospitals Trust, Londonderry	Dr Carol Dezateux, Institute of Child Health, London	Dr Tom Fahey, University of Bristol	Dr Ann McPherson, General Practitioner, Oxford
Mr John Cairns, University of Aberdeen	Dr Anne Dixon Brown, NHS Executive, Anglia & Oxford	Mrs Gillian Fletcher, National Childbirth Trust	Dr Susan Moss, Institute of Cancer Research
		Dr JA Muir Gray, Institute of Health Sciences, Oxford	Dr Sarah Stewart-Brown, University of Oxford

### Past members

Dr Sheila Adam, Department of Health*	Dr Anne Ludbrook, University of Aberdeen	Professor Catherine Peckham, Institute of Child Health, London	Professor Nick Wald, University of London
Professor George Freeman, Charing Cross & Westminster Medical School, London	Professor Theresa Marteau, Guy's, King's & St Thomas's School of Medicine & Dentistry, London	Dr Connie Smith, Parkside NHS Trust, London	Professor Ciaran Woodman, Centre for Cancer Epidemiology, Manchester
Dr Mike Gill, Brent & Harrow Health Authority		Ms Polly Toynbee, Journalist	

## Primary and Community Care Panel

### Current members

<b>Chair:</b> <b>Dr John Tripp,</b> Royal Devon & Exeter Healthcare NHS Trust	Ms Judith Brodie, Age Concern, London	Mr Andrew Farmer, Institute of Health Sciences, Oxford	Dr Chris McCall, General Practitioner, Dorset
Mr Kevin Barton, East London & City Health Authority	Mr Shaun Brogan, Daventry & South Northants Primary Care Alliance	Professor Richard Hobbs, University of Birmingham	Dr Robert Peveler, University of Southampton
Professor John Bond, University of Newcastle- upon-Tyne	Mr Joe Corkill, National Association for Patient Participation	Professor Allen Hutchinson, University of Sheffield	Professor Jennie Popay, University of Salford
Dr John Brazier, University of Sheffield	Dr Nicky Cullum, University of York	Dr Phillip Leech, Department of Health	Ms Hilary Scott, Tower Hamlets Healthcare NHS Trust, London
	Professor Pam Enderby, University of Sheffield	Dr Aidan Macfarlane, Oxfordshire Health Authority	Dr Ken Stein, North & East Devon Health Authority

### Past members

Professor Angela Coulter, King's Fund, London*	Professor Andrew Haines, RDRD, North Thames Regional Health Authority	Mr Lionel Joyce, Chief Executive, Newcastle City Health NHS Trust	Professor Dianne Newham, King's College London
Professor Martin Roland, University of Manchester*	Dr Nicholas Hicks, Oxfordshire Health Authority	Professor Martin Knapp, London School of Economics & Political Science	Professor Gillian Parker, University of Leicester
Dr Simon Allison, University of Nottingham	Mr Edward Jones, Rochdale FHSA	Professor Karen Luker, University of Liverpool	Dr Mary Renfrew, University of Oxford
Professor Shah Ebrahim, Royal Free Hospital, London	Professor Roger Jones, Guy's, King's & St Thomas's School of Medicine & Dentistry, London	Dr Fiona Moss, Thames Postgraduate Medical & Dental Education	

## National Coordinating Centre for Health Technology Assessment, Advisory Group

### Current members

---

**Chair:**

**Professor John Gabbay,**  
Wessex Institute  
for Health Research  
& Development

Ms Lynn Kerridge,  
Wessex Institute for Health  
Research & Development

Professor James Raftery,  
Health Economics Unit,  
University of Birmingham

Professor Andrew  
Stevens,  
Department of Public  
Health & Epidemiology,  
University of Birmingham

Dr Ruairidh Milne,  
Wessex Institute for Health  
Research & Development

Professor Ian Russell,  
Department of Health Sciences  
& Clinical Evaluation,  
University of York

Professor Mike  
Drummond,  
Centre for Health Economics,  
University of York

Ms Kay Pattison,  
Research & Development  
Directorate, NHS Executive

Dr Ken Stein,  
North & East Devon  
Health Authority

### Past member

---

Dr Paul Roderick,  
Wessex Institute for Health  
Research & Development

# HTA Commissioning Board

## Current members

---

### Chair:

**Professor Charles Florey**,  
Department of Epidemiology &  
Public Health, Ninewells  
Hospital & Medical School,  
University of Dundee

**Professor Doug Altman**,  
Director of ICRF/NHS Centre  
for Statistics in Medicine,  
Oxford

**Professor John Bond**,  
Professor of Health Services  
Research, University of  
Newcastle-upon-Tyne

**Mr Peter Bower**,  
Independent Health Advisor,  
Newcastle-upon-Tyne

**Ms Christine Clark**,  
Honorary Research Pharmacist,  
Hope Hospital, Salford

**Professor Shah Ebrahim**,  
Professor of Epidemiology  
of Ageing, University of Bristol

**Professor Martin Eccles**,  
Professor of  
Clinical Effectiveness,  
University of Newcastle-  
upon-Tyne

**Dr Mike Gill**,  
Director of Public Health &  
Health Policy, Brent & Harrow  
Health Authority

**Dr Alastair Gray**,  
Director, Health Economics  
Research Centre,  
University of Oxford

**Professor Mark Haggard**,  
MRC Institute of  
Hearing Research

**Dr Jenny Hewison**,  
Senior Lecturer,  
Department of Psychology,  
University of Leeds

**Professor Sir Miles Irving**  
(Programme Director),  
Professor of Surgery,  
University of Manchester,  
Hope Hospital, Salford

**Professor Alison Kitson**,  
Director, Royal College of  
Nursing Institute

**Dr Donna Lamping**,  
Senior Lecturer, Department of  
Public Health, London School  
of Hygiene & Tropical Medicine

**Professor Alan Maynard**,  
Professor of Economics,  
University of York

**Professor Jon Nicholl**,  
Director, Medical Care  
Research Unit,  
University of Sheffield

**Professor Gillian Parker**,  
Nuffield Professor of  
Community Care,  
University of Leicester

**Dr Tim Peters**,  
Reader in Medical Statistics,  
Department of Social Medicine,  
University of Bristol

**Professor Martin Severs**,  
Professor in Elderly  
Health Care,  
Portsmouth University

**Dr Sarah Stewart-Brown**,  
Director, Institute of  
Health Sciences,  
University of Oxford

**Professor Ala Szczepura**,  
Director, Centre for  
Health Services Studies,  
University of Warwick

**Dr Gillian Vivian**,  
Consultant, Royal Cornwall  
Hospitals Trust

**Professor Graham Watt**,  
Department of General Practice,  
Woodside Health Centre,  
Glasgow

**Professor Kent Woods**,  
Regional Director of R&D  
NHS Executive Trent

**Dr Jeremy Wyatt**,  
Senior Fellow, Health &  
Public Policy, School of Public  
Policy, University College,  
London

## Past members

---

**Professor Ian Russell**,  
Department of Health  
Sciences & Clinical Evaluation,  
University of York\*

**Professor David Cohen**,  
Professor of Health Economics,  
University of Glamorgan

**Mr Barrie Dowdeswell**,  
Chief Executive,  
Royal Victoria Infirmary,  
Newcastle-upon-Tyne

**Dr Michael Horlington**,  
Head of Corporate Licensing,  
Smith & Nephew Group  
Research Centre

**Professor Martin Knapp**,  
Director, Personal Social  
Services Research Unit,  
London School of Economics  
& Political Science

**Professor Theresa Marteau**,  
Director, Psychology & Genetics  
Research Group, Guy's, King's  
& St Thomas's School of  
Medicine & Dentistry,  
London

**Professor Sally McIntyre**,  
MRC Medical Sociology Unit,  
Glasgow

**Professor David Sackett**,  
Centre for Evidence Based  
Medicine, Oxford

**Dr David Spiegelhalter**,  
MRC Biostatistics Unit,  
Institute of Public Health,  
Cambridge

**Professor David Williams**,  
Department of  
Clinical Engineering,  
University of Liverpool

**Dr Mark Williams**,  
Public Health Physician,  
Bristol

\* Previous Chair

---

Copies of this report can be obtained from:

The National Coordinating Centre for Health Technology Assessment,  
Mailpoint 728, Boldrewood,  
University of Southampton,  
Southampton, SO16 7PX, UK.  
Fax: +44 (0) 1703 595 639 Email: [hta@soton.ac.uk](mailto:hta@soton.ac.uk)  
<http://www.soton.ac.uk/~hta>

ISSN 1366-5278