

Infrared-Induced Rotamerization of Oxalic Acid Monomer in Argon Matrix

Ermelinda M. S. Maçôas and Rui Fausto*

Department of Chemistry—CQC, University of Coimbra, P-3049 Coimbra, Portugal

Mika Pettersson, Leonid Khriachtchev, and Markku Räsänen

Laboratory of Physical Chemistry, University of Helsinki, P.O. Box 55 (A.I. Virtasen aukio 1), FIN-00014 Helsinki, Finland

Received: February 17, 2000; In Final Form: April 27, 2000

Infrared-induced conformational isomerization of oxalic acid monomer isolated in an argon matrix at 7.5 K was studied by infrared spectroscopy. For the first time, three conformational states of this molecule were identified experimentally and their vibrational spectra are assigned. In good agreement with density functional theory predictions, all the observed conformers exhibit a trans O=C–C=O axis, differing in the relative conformation of their O–C–O–H axes. In the most stable conformer (belonging to the C_{2h} symmetry point group), two intramolecular OH···O= hydrogen bonds are present. The second (C_s) most stable conformer shows a single OH···O= bond, and the third one (C_{2h}) does not exhibit any intramolecular hydrogen bond. Using narrowband tunable irradiation in the near-infrared region it was possible to promote very efficiently conformer interconversions, which was followed spectroscopically.

Introduction

Oxalic acid (ethanedioic acid; $C_2H_2O_4$) is the smallest member in the series of aliphatic dicarboxylic acids. At room temperature, it exists in two crystalline varieties, the stable orthorhombic α -phase and the metastable monoclinic β -phase.^{1,2} In both crystalline phases, where extensive intermolecular hydrogen bonding occurs, individual molecules were found to be exclusively in a conformation where the O–C–C–O dihedral angle is 180° , that is, the two carbonyl groups are trans to each other, and the two H–O–C–C dihedral angles assume a trans configuration (conformer **tTt**[†], in Figure 1).^{1,2} However, in both gaseous phase and in inert gas matrices at low temperatures, monomeric oxalic acid has been found to exist predominantly in a different conformation, where the two H–O–C–C dihedrals adopt the cis configuration and are involved in intramolecular hydrogen bonding (form **cTe**).^{3–5}

High-level ab initio and density functional theory (DFT) calculations on monomeric oxalic acid have also been reported previously.^{6–8} In general, the highest-level theoretical calculations predict the existence of at least five conformers of oxalic acid. In good agreement with the available experimental results,^{3–5} the calculations systematically predict conformer **cTe** as corresponding to the most stable form. In an earlier matrix isolation infrared (IR) spectroscopic study, some evidence could be found pointing to the existence of more than one conformer of oxalic acid trapped in the matrices.⁴ However, only the second most stable conformer predicted by the calculations (form **cTt**, in Figure 1) could be unequivocally identified experimentally.³ A recent Stark-modulated free jet microwave spectroscopic study⁸ has confirmed the structure of this conformer. The other




Figure 1. B3LYP/6-31G**⁷-calculated conformers of monomeric oxalic acid.⁷ The difference in energy (kJ mol^{-1}), including zero-point vibrational energy-corrected relative conformational energies to the most stable form,⁷ are given in parentheses. Form **cGc** was found to correspond to an energy maximum both at ha HF/6-31G**² and MP2/6-311++G**⁸ (see text).

theoretically predicted conformers of oxalic acid have not been observed until now, though at least the third- and fourth-lowest-energy conformers (forms **tTt** and **tCt**) were calculated to have relatively accessible excess energies to the conformational ground state.^{7,8} The B3LYP/6-31G**⁷ and MP2/6-31G**³ calculations provided identical results predicting the existence of six conformers, all but the highest-energy form (**cGc** in Figure 1) being planar. However, a more recent study carried out at the MP2/6-311++G** level of theory, where energy profiles associated with internal rotation around the C–C and C–O bonds were calculated, indicates that **cGc** does not correspond

* Author to whom all correspondence should be addressed.

[†] The nomenclature here adopted is the same used in the previous experimental paper on oxalic acid;³ an uppercase letter designates the O–C–C–O dihedral angle (φ ; C, cis; T, trans; G, gauche), and the first and last lowercase letters designate the HOCC (τ_1) and CCOH (τ_2) dihedral angles, respectively.

to a true energy minimum, whereas the **tTt**, **tCt**, and **cCt** forms are slightly distorted from planarity.⁸

Identification of conformer **cTt** was achieved by observation of IR- and ultraviolet (UV)-induced conformational processes for oxalic acid isolated in noble gas matrices at low temperature, followed by IR spectroscopy.³ A Xe arc lamp with a narrowband filter (max 270 nm, full width at half-maximum 30 nm, corresponding to the $\pi^* \leftarrow n$ transition⁹) was used for UV irradiation, which was found to induce **cTc** \rightarrow **cTt** isomerization, accompanied by partial decomposition of oxalic acid monomers. IR irradiation was carried out using the globar of the spectrometer either without filter or with a low-pass filter (transmitting ca. 80% between 2000 and 600 cm^{-1}). Irradiation above 2000 cm^{-1} was found to be active at initiating the reverse process (**cTt** \rightarrow **cTc** isomerization).³

It has repeatedly been reported for other carboxylic acid monomers that selective excitation of the OH stretching vibrations are of great utility to study conformational isomerization processes in this type of compound.^{10–12} Furthermore, contrary to UV irradiation, IR irradiation does not usually lead to decomposition of the matrix-isolated molecules. Because the IR irradiation carried out in the previous study on oxalic acid was not selective, we decided to submit this compound to a new investigation, where selective narrowband and tunable IR irradiation was used, to better characterize the observed isomerization processes and, in particular, to search for other conformational species.

Experimental and Computational Details

The matrix samples were prepared by continuously flushing a Teflon U-tube containing oxalic acid (Aldrich, 99+%), heated to 100 °C, with high-purity argon (>99.995%). The gaseous mixture was deposited onto a cooled CsI window in a close-cycle helium cryostat (APD, DE202A) at 7.5 K. The IR spectra (4000–450 cm^{-1}) were measured with a Nicolet SX-60 Fourier transform IR spectrometer equipped with a liquid-nitrogen-cooled mercury–cadmium–telluride detector and a KBr beam splitter, with spectral resolutions of 0.25 or 1.0 cm^{-1} . Typically 200 interferograms were coadded. In our samples, monomeric species were almost exclusively formed, which was achieved by choosing an appropriate temperature of the substance container and optimizing the matrix gas flow rate.

To promote conformational interconversions, tuneable pulsed IR irradiation was used, as provided by the idler beam of an optical parametric oscillator (OPO Sunlite, Continuum). The pulse duration of the IR irradiation was ca. 5 ns with a line width of $\sim 0.1 \text{ cm}^{-1}$. The absolute accuracy of the IR radiation was established with an accuracy better than 1 cm^{-1} by a Burleigh WA-4500 wavemeter used to control the OPO signal radiation wavelength.

The DFT calculations were performed with the Becke's three-parameter hybrid method using the Lee–Yang–Parr's correlation functional^{13,14} and the 6-31G**¹⁵ basis set. All the calculations were made with the GAUSSIAN 92/DFT package of programs,¹⁶ installed on a DEC ALPHA 7000 computer, at the Center of Informatics of the University of Coimbra.

The force constants (symmetry internal coordinates) to be used in the normal coordinate analysis were obtained from the DFT Cartesian harmonic force constants using the program TRANSFORMER.¹⁷ This program was also used to prepare the input data for the normal coordinate analysis programs used in this study (BUILD-G, and VIBRAT¹⁸). The calculated frequencies were then scaled by using a single scale factor (0.9614⁸) to obtain a better agreement with the experimental frequencies.

Although very simple, this scaling procedure preserves the potential energy distributions (PEDs) as they emerge from the DFT calculations, thus having an important advantage over the more elaborate force-field scaling procedures, which use more than one scale factor.

Results and Discussion

Conformer Interconversion. The high-level DFT calculations by Higgins et al.⁷ indicate that the energies of the second- to the fifth-lowest-energy forms of oxalic acid shall not exceed the energy of the most stable conformer by more than 25 kJ mol^{-1} , whereas the highest-energy form (conformer **cGc**) has a predicted relative energy of 63.5 kJ mol^{-1} . Thus, the relative populations of the different conformers of oxalic acid at the temperature used to prepare the matrix (373 K) are **cTc:cTt:tTt:tCt:cCt:cGc** = 95.15:3.64:0.71:0.44:0.06: $< 10^{-5}$ assuming a Boltzmann distribution. The MP2 results of Godfrey et al.⁸ do not differ qualitatively from the DFT data, although the relative energy differences between the conformers are somewhat smaller. In addition, as mentioned above, the MP2 study indicated that **cGc** does not correspond to a minimum in the potential energy surface. The potential energy profiles associated with the internal rotation around the C–C bond for trans/trans or trans/cis arrangements of the carboxylic moieties calculated at the MP2 level of theory have revealed the following features:⁸ (a) Without taking into consideration zero-point vibrational energy corrections, the **tTt**, **tCt**, and **cCt** planar conformations should correspond to local maxima between two equivalent slightly distorted structures (the planar structures were calculated to be 0.26, 1.02, and 0.07 kJ mol^{-1} above the corresponding nonplanar minima, respectively). In the last case, however, the zero-point vibrational energy level of the true minimum energy conformation is 0.12 kJ mol^{-1} above the energy of the planar structure. Thus, the most probable structure falls at the symmetric planar position (**cCt**). In addition, when zero-point vibrational energy is considered, the **tTt** planar structure becomes only 0.10 kJ mol^{-1} higher in energy than the corresponding distorted forms, and then the planar structure must also be the most probable geometry at temperatures higher than 13 K. (b) The calculated energy barrier separating the distorted **tCt** forms from **tTt** is only ca. 1.6 kJ mol^{-1} . Thus, at the temperature used in our study to prepare the sample (373 K), the available thermal energy can easily be used to overcome this barrier and, consequently, in the present context **tCt** is best regarded as vibrationally excited **tTt** rather than a separate species.

Taking into consideration the theoretical predictions, it can be expected that, in the gaseous phase at 373 K, four conformers of monomeric oxalic acid exist (**cTc**, **cTt**, **tTt**, and **cCt**), although the highest-energy form (**cCt**) should have a population smaller than 0.1%. Assuming that gas-phase populations are trapped in the matrices, the presence of forms **cTc**, **cTt**, and **tTt** should in principle be detected in the vibrational spectra of matrix-isolated oxalic acid monomer.

The IR spectrum of a freshly prepared matrix of oxalic acid is dominated by intense bands due to the most stable **cTc** conformer. The assignments of these bands were discussed in detail elsewhere.^{3,4}

As expected, apart from the bands due to the conformational ground state (**cTc**), some minor bands ascribable to other conformers are also present in the spectrum. As mentioned in the *Introduction*, by irradiating the matrix in the OH stretching fundamental region, Nieminen et al. were able to observe conformational isomerization, the reaction product correspond-

TABLE 1: Definition of Internal Symmetry Coordinates Used in Normal-Coordinate Analysis

coordinate	cTc/tTt		cTt		definition ^a	
	approx. desc.	symmetry (C_{2h})	approx. desc.	symmetry (C_s)	cTc/tTt	cTt
S_1	ν OH s.	A_g	ν OH'	A'	$\nu O_5H_7 + \nu O_6H_8$	νO_6H_8
S_2	ν OH	B_u	ν OH	A'	$\nu O_5H_7 - \nu O_6H_8$	νO_5H_7
S_3	ν C=O s.	A_g	ν C=O	A'	$\nu C_1=O_3 + \nu C_2=O_4$	$\nu C_2=O_4$
S_4	ν C=O	B_u	ν C=O'	A'	$\nu C_1=O_3 - \nu C_2=O_4$	$\nu C_1=O_3$
S_5	ν C-O s.	A_g	ν C-O s.	A'	$\nu C_1-O_5 + \nu C_2-O_6$	
S_6	ν C-O	B_u	ν C-O	A'	$\nu C_1-O_5 - \nu C_2-O_6$	
S_7	ν C-C	A_g	ν C-C	A'	νC_1-C_2	
S_8	δ OCO s.	A_g	δ OCO s.	A'	$2\delta O_5C_1O_3 - \delta C_2C_1=O_3 - \delta C_2C_1-O_5 + 2\delta O_6C_2O_4 - \delta C_1C_2=O_4 - \delta C_1C_2-O_6$	
S_9	δ OCO	B_u	δ OCO	A'	$2\delta O_5C_1O_3 - \delta C_2C_1=O_3 - \delta C_2C_1-O_5 - 2\delta O_6C_2O_4 + \delta C_1C_2=O_4 + \delta C_1C_2-O_6$	
S_{10}	δ CC=O s.	A_g	δ CC=O s.	A'	$\delta C_2C_1=O_3 - \delta C_2C_1-O_5 + \delta C_1C_2=O_4 - \delta C_1C_2-O_6$	
S_{11}	δ CC=O	B_u	δ CC=O	A'	$\delta C_2C_1=O_3 - \delta C_2C_1-O_5 - \delta C_1C_2=O_4 + \delta C_1C_2-O_6$	
S_{12}	δ COH s.	A_g	δ COH s.	A'	$\delta C_1O_5H_7 + \delta C_2O_6H_8$	
S_{13}	δ COH	B_u	δ COH	A'	$\delta C_1O_5H_7 - \delta C_2O_6H_8$	
S_{14}	γ C=O	A_u	γ C=O	A''	$\gamma C_2-C_1(=O_3)-O_5 + \gamma C_1-C_2(=O_4)-O_6$	
S_{15}	γ C=O s.	B_g	γ C=O s.	A''	$\gamma C_2-C_1(=O_3)-O_5 - \gamma C_1-C_2(=O_4)-O_6$	
S_{16}	τ C-C	A_u	τ C-C	A''	$\tau O_3=C_1-C_2=O_4$	
S_{17}	τ C-O	A_u	τ C-O	A''	$\tau O_3=C_1-O_5-H_7 + \tau O_4=C_2-O_6-H_8$	
S_{18}	τ C-O s.	B_g	τ C-O s.	A''	$\tau O_3=C_1-O_5-H_7 - \tau O_4=C_2-O_6-H_8$	

^a For atom numbering see Figure 1. ν , stretching; δ , bending; γ , rocking; τ , torsion.

ing to the lowest-energy conformer, whereas this conformer behaved as the reactive species when the sample was irradiated in the UV region.³ The IR-reactive species was identified as corresponding to form **cTt** and a tentative assignment of the bands due to this conformer was also proposed in this previous study. However, the relatively low efficiency of the photoisomerization process did not allow the unequivocal assignment of all the observed features. In addition, a possible contribution from other conformers to the observed spectra could not be discarded.

In the present study, IR irradiation in the 6800–6700 cm^{-1} region was undertaken. In this spectral region, the OH stretching combination bands ($S_1 + S_2$; see Table 1 for definition of symmetry coordinates) or first overtones of the OH stretching vibrations of appropriate symmetry are expected to occur. Because the most stable conformer belongs to the C_{2h} symmetry point group, the first overtone of its OH stretching modes does not absorb in the IR radiation. Nevertheless, the $S_1 + S_2$ combination band belongs to the B_u symmetry species and then must correspond to the active vibration of the **cTc** conformer found at $\approx 6712 \text{ cm}^{-1}$, which we have used to promote photochemical isomerization of this conformer to other forms. After 5 min of irradiation at this wave number, a noticeable reorganization of band intensities was observed. Figures 2 and 3 show difference spectra (irradiated sample–nonirradiated sample) in the C=O and OH stretching regions obtained at different stages of irradiation. In Figure 3 the experimental features are compared with the DFT-calculated spectra.

In the C=O stretching region, irradiation leads to an increase of the triplet around 1830 cm^{-1} , and the doublets near 1780 and 1760 cm^{-1} , whereas a decrease was observed in the case of the features at ca. 1810 cm^{-1} (see Figures 2 and 3). Comparison of the observed spectrum with the calculated spectra for the three most stable conformers (**cTc**, **cTt**, and **tTt**) clearly indicates that the irradiation leads to conversion of conformer **cTc** to both forms **cTt** and **tTt**. Thus, the irradiation is effective in promoting internal rotation about the C–O bond.

It is known that when the carboxylic group assumes the cis arrangement (C–C–O–H dihedral angle equals 0°) the carbonyl stretching vibration occurs at a higher frequency than when it takes the trans geometry. This trend is essentially due to the fact that the $[O=C-O-H \leftrightarrow O^--C=O^+-H]$ mesom-




Figure 2. Observed spectrum of a nonirradiated sample of oxalic acid isolated in an Ar matrix (3650–3400 cm^{-1} and 1900–1700 cm^{-1} regions; intensities are shown in arbitrary units) and difference spectra obtained at different stages of irradiation at 6712 cm^{-1} . It is clear from this figure that during the first stages of irradiation conformer **cTt** is the main product, whereas conformer **tTt** starts to become the dominant product at longer times of irradiation.

erism is more important in the cis than in the trans conformation of this group.^{19,20} Hence, the double-bond character of the carbonyl bond decreases and, consequently, the force constant associated with the C=O stretching vibration reduces. On the other hand, intramolecular hydrogen bonding involving the carbonyl oxygen atom as acceptor also leads to a red shift in the frequency of the carbonyl stretching mode.

In the **cTt** conformer, the cis carboxylic group is not involved in hydrogen bonding and its carbonyl stretching mode absorbs near 1830 cm^{-1} , whereas the hydrogen-bonded trans carboxylic group gives rise to the doublet occurring at a considerably lower frequency (ca. 1760 cm^{-1}). In conformer **cTc**, both carbonyl groups adopt the cis arrangement but they also participate in intramolecular hydrogen bonding. Thus, in this conformer, the effects of *carboxylic group conformation* and *intramolecular hydrogen bonding* compensate partially each other. Consequently, the IR active carbonyl stretching mode (B_u symmetry) is expected to occur at a frequency between those of the modes




Figure 3. Calculated and observed spectra (3650–3400 cm^{-1} and 1900–1700 cm^{-1} regions) of oxalic acid isolated in an Ar matrix. Calculated spectra simulate the absolute intensities using Gaussian functions with an arbitrary full width at half-height of 1 cm^{-1} for the three most stable conformers of oxalic acid. Experimental data correspond to difference spectra (6712 cm^{-1} irradiated sample – nonirradiated sample).

due to the **cTt** conformer, giving rise to the doublet observed near 1810 cm^{-1} (see Figures 2 and 3). Finally, the doublet at ca. 1780 cm^{-1} is ascribable to the IR active carbonyl stretching mode of **tTt** where the two carboxyl groups are trans and nonintramolecularly hydrogen bonded. The splittings of the bands are probably due to matrix field effects that are rather commonly observed in spectra of matrix-isolated species, especially if the vibrational modes involve large changes in the molecular dipolar moment.^{3,21}

In the OH stretching region, the irradiation at 6712 cm^{-1} leads to an increase of two triplets at 3565–3555 and 3550–3545 cm^{-1} that can be ascribed respectively to the IR active OH stretching modes of the non-hydrogen-bonded hydroxylic groups of **tTt** and **cTt**. A doublet at 3485–3480 cm^{-1} is assigned to the hydrogen-bonded hydroxylic group of the **cTt** conformer. The OH stretching fundamental mode of the **cTc** conformer corresponds to the downward multiplet at 3460–3450 cm^{-1} .

The next irradiation at 6755 cm^{-1} excites the first overtone of the OH stretching mode of the **cTt** conformer (intramolecularly bonded OH group). This irradiation at 6755 cm^{-1} followed the first pumping at 6712 cm^{-1} . The observed difference spectrum in the C=O and OH stretching regions together with the calculated spectra are presented in Figure 4. In these spectra, the bands belonging to conformer **cTt** appear as negative peaks, whereas those belonging to the **cTc** and **tTt** conformers point upward. Thus, irradiation at this frequency leads to conversion of **cTt** to both **cTc** and **tTt**. Once again, the irradiation is shown to be effective in promoting internal rotation about the C–O bond. Similarly to the irradiation at 6712 cm^{-1} , no evidence of rotamerization about the C–C bond was observed.

The energy differences between the three observed conformers can be estimated on the basis of the experimental IR band intensities recorded immediately after deposition, taking into account the theoretically predicted intensities⁷ and assuming that no conformational isomerization takes place during the deposition process. An energy difference of ca. 10 kJ mol^{-1} between **cTt** and **cTc** was obtained by this approach, whereas conformer **tTt** was estimated to be higher in energy than the conformational ground state by ca. 16 kJ mol^{-1} , corresponding to relative populations at the deposition temperature of **cTc**: 95.9%, **cTt**: 3.6%, and **tTt**: 0.5%. These results were obtained from the intensities of the O–H or C=O stretching modes by weighting the experimental integrated intensities by the corresponding




Figure 4. Calculated and observed spectra (3650–3400 cm^{-1} and 1900–1700 cm^{-1} regions) of oxalic acid isolated in an Ar matrix. Calculated spectra simulate the absolute intensities using Gaussian functions with an arbitrary full width at half-height of 1 cm^{-1} for the three most stable conformers of oxalic acid. Experimental data correspond to difference spectra (6755 cm^{-1} irradiated sample – 6712 cm^{-1} irradiated sample).

calculated IR absorption cross-sections. It is worth mentioning that the DFT calculated values are 10.11 and 15.17 kJ mol^{-1} , respectively, corresponding to relative populations at 373 K of **cTc**: 95.15, **cTt**: 3.64, and **tTt**: 0.71,⁷ thus being in very good agreement with the observed data.

After irradiation at 6755 cm^{-1} during 10 min, ca. 46% of the population of **cTt** was converted into **cTc** (27%) and **tTt** (19%) while taking into account the calculated IR absorption cross-sections. This result is in qualitative agreement with the relative order of magnitude of the calculated energy barriers associated with the **cTt**→**cTc** and **cTt**→**tTt** isomerization processes, the first being 2.9 kJ mol^{-1} lower than the last.⁷ On the other hand, **cTc**→**cTt** isomerization was found to be considerably more efficient than **cTc**→**tTt** when irradiation at 6712 cm^{-1} was undertaken, as can be expected because **cTc**→**tTt** involves the breaking of two hydrogen bonds instead of one hydrogen bond as in **cTc**→**cTt**.

Selective irradiation of the OH stretching vibrations have been found to be of great utility to study conformational isomerization processes in carboxylic compounds.^{10–12} Nevertheless, it should be emphasized that the elementary mechanism of IR-induced rotamerization is far from being completely understood. For instance, direct coupling of the excited overtone levels with the torsional manifold is a very high-order process, thus quite improbable. The role of the solid host in intramolecular energy transfer is not clear either.

As mentioned earlier, observation of other conformers (**tCt** and **cCt**) in the nonirradiated matrix was not expected, taking into consideration their predicted relative populations. Indeed, the estimated **cCt** population in the gas at the deposition temperature (373 K) is lower than 0.1%.⁷ For **tCt**, though instantaneous cooling of the 373 K gas would lead to a population of ca. 0.5% as estimated on the basis of the DFT calculations,⁷ the population in the matrix can be expected to be much smaller, because fast conversion of this form to **tTt** during sample deposition may easily occur, considering the very low energy barrier separating these two forms (ca. 1.6 kJ mol^{-18}).

Another relevant observation obtained in the irradiation experiments is the fact that no evidence was found of any photochemically induced isomer with a cis O=C–C=O axis. In fact, taking into consideration the theoretically calculated energy barriers associated with conformational interconversion,

TABLE 2: Observed and Scaled^a B3LYP/6-31G Frequencies of Three Most Stable Oxalic Acid Conformers Isolated in an Argon Matrix**

cTc decreasing bands with 6712 cm ⁻¹ excitation			tTt increasing bands with 6755 cm ⁻¹ excitation			cTt decreasing bands with 6755 cm ⁻¹ excitation		
ν_{obs}^b (cm ⁻¹)	$\nu_{\text{calc.}}$ (cm ⁻¹)	PED (%) ^c	ν_{obs}^b (cm ⁻¹)	$\nu_{\text{calc.}}$ (cm ⁻¹)	PED (%) ^c	ν_{obs}^b (cm ⁻¹)	$\nu_{\text{calc.}}$ (cm ⁻¹)	PED (%) ^c
			3565	3613	ν OH (100)	3549	3603	ν OH (100)
			3560			3548		
3461	3500	ν OH (100)	3556			3546		
3459						3484	3523	ν OH' (100)
3457						3479		
3453								
1812	1807	ν C=O (83), ν C-O (10)				1834	1825	ν C=O (81)
1810						1829		
						1825		
			1781	1781	ν C=O (87)	1762	1748	ν C=O' (78)
			1780			1760		
						1416	1398	ν C-O s. (35), δ COH (22), ν C-C (18), δ OCO s. (15)
						1414		
1321	1301 ^d	δ COH (64), ν C-O (27)	non obs.	1295	δ COH (56), ν C-O (26), δ OCO (16)	1319	1303 ^e	δ COH s. (62), ν C-O (23)
1316						1313		
1315						1213		
1267						1210		
1260						1208		
1259						1203		
1173	1180	ν C-O (62), δ COH (17), ν C=O (13)				1161	1171	δ COH (49), ν C-O s. (29)
1172						1159		
						1157		
			1112	1112	ν C-O (59), δ COH (34),	1137	1142	ν C-O (58), δ COH (22)
			1110			1134		
			1108					
						799	777	ν C-C (39), δ OCO s. (31), ν C-O s. (18)
						796	772	γ C=O (78), τ C-O s. (11), τ C-O (11)
669	696	τ C-O (100)	641	655	τ C-O (84), γ C=O (16)	650	660	τ C-O (89)
664			637			633	637	τ C-O s. (84), γ C=O s. (12)
657	644	δ OCO (84)	636			621	629	δ OCO (83), δ COH s. (10)
656			631	612	δ OCO (81), δ COH (14)	616		
459	435	γ C=O (66), τ C-C (24), τ C-O (10)				612		
458						558	526	δ CC=O s. (81)
							411	γ C=O s. (78), τ C-C (17)
				397	γ C=O (96)		409	δ OCO s. (62), ν C-C (29)
264 ^f	257	δ CC=O (92)		253	γ CC=O (100)		258	δ CC=O (94)
	126	τ C-C (84), τ C-O (16)		37	τ C-C (96)		95	τ C-C (82), τ C-O (17)
E^g (kJ mol ⁻¹)		-993176.4916			-993161.4254			-993166.446
ΔE^h					15.1			10.1

^a Scale factor 0.9614. ^b Minor bands not listed, attributed to nonfundamental modes, were observed at: 1855, 1853, 1483, 1481, 1290, 1286, 1284, 1153, 1129, 1127, 1103, 1100, 1098, 996, 994, 990 (cTc); 1187 (tTt), and 1459, 1449, 1062, 1060, 1057, 824 (cTt). ^c Only contributions higher than 10% are listed. ^d Fermi resonance of δ COH with [τ C-O (B_g) + γ C=O (A_u)]; ^e Fermi resonance of δ COH with 2 δ OCO. ^f Taken from ref 5. ^g B3LYP/6-31G** energy. ^h B3LYP/6-31G** energy compared with the most stable conformer. ν , stretching; δ , bending; τ , torsion; γ , rocking.

one can expect both **cCt** and **tCt** to be found. Conformer **cCt** might be obtained from **cTt** because the calculated **cTt**→**cCt** rotational barrier is estimated to be 4.62 kJ mol⁻¹ lower than that associated with the observed **cTt**→**tTt** process.⁷ Conversion into **tCt** form is possible via more complex pathways involving either **cCt** or **tTt** as intermediates. Nevertheless, despite the energy transferred to the molecules during the irradiation being ca. 80 kJ mol⁻¹, that is, above the calculated energy barriers (the highest calculated energy barrier for these processes is <75 kJ mol⁻¹⁷), the internal rotation around the C-C bond does not take place. A possible explanation for this finding requires consideration of volume restrictions to the internal rotation imposed by the matrix. Internal rotation about the C-C bond involves the motion of heavy atoms within the noble gas cage, which requires significant reorganization of the matrix. Consequently, the effective energy barriers for C-C internal rotation in the matrix-isolated molecules may be considerably higher than for the molecule in a vacuum, thus preventing the reaction to occur. Note that the observed interconversion processes involve only motion of hydrogen atoms and do not require any major matrix rearrangement.

Assignment of IR Spectra of cTc, tTt, and cTt Conformers. Because the IR-irradiation experiments enabled the clear discrimination of the bands of the three observed conformers (see Figures 2–4), the complete assignment of the vibrational signature of these conformers could be undertaken. In Table 2 we present the experimental and calculated wave numbers and the potential energy distribution for the conformational ground state (**cTc**) and conformers **cTt** and **tTt**, respectively. The symmetry coordinates used to perform the normal coordinate analysis are shown in Table 1.

The assignments here made for the lowest-energy conformer follow closely those reported in ref 3. The two observed complex bands at ca. 1260–1270 cm⁻¹ and 1315–1320 cm⁻¹ are ascribed to a Fermi resonance doublet (split because of matrix site effects) resulting from interaction between the fundamental δ COH (B_u) vibration and the τ C-O s. + γ C=O ($B_g \times A_u = B_u$) combination band.

In the case of conformer **cTt**, some of the assignments of ref 3 are now revised. In particular, the C-O s. stretching and the δ CC=O s. modes (which were not assigned in ref 3) are now ascribed respectively to the doublet appearing at 1415/1416

cm^{-1} and to the band at 558 cm^{-1} , whereas the two δCOH in-plane bending vibrations are now assigned to the doublet at $1161/1157\text{ cm}^{-1}$ (δCOH ; see Table 2) and to the Fermi resonance doublet (split because of matrix site effects) whose component bands appear at $1319/1313\text{ cm}^{-1}$ and $1213-1203\text{ cm}^{-1}$ (δCOH s. interacting with the first overtone of $\delta\text{O}=\text{C}-\text{O}$).

The assignments made for the **tTt** conformer observed now for the first time are straightforward. As for the other observed forms, the calculated and experimental data for this form are in excellent agreement. It is worth mentioning that we did not observe clearly a band ascribable to the δCOH mode of this conformer because this vibration is predicted to occur near the very intense and structured band due to the most stable conformer at $1267-1259\text{ cm}^{-1}$ (calculated frequency for δCOH in **tTt**: 1295 cm^{-1} ; see Table 2).

It is also interesting to point out that the difference of about 70 cm^{-1} between the observed frequencies associated with the free ($3549-3546\text{ cm}^{-1}$) and hydrogen-bonded ($3484-3479\text{ cm}^{-1}$) O-H group in conformer **cTt** indicates that in oxalic acid the intramolecular hydrogen bond is weaker than in glycolic acid (80 cm^{-1})²² or in water dimer (106 cm^{-1})²³

Conclusion

Using narrowband tunable irradiation in the $6800-6700\text{ cm}^{-1}$ region, it was possible to unequivocally identify the three lowest-energy conformers of oxalic acid monomer and perform a complete vibrational assignment of their IR spectra in an argon matrix at 7.5 K . In consonance with the previous studies, the most stable conformer was found to be the **cTc** conformer, which is characterized by two $\text{OH}\cdots\text{O}=\text{C}$ intramolecular hydrogen bonds, and the second most stable conformer (**cTt**) exhibits a single intramolecular hydrogen bond. The third most stable conformer was observed experimentally for the first time and found to be the **tTt** conformer, with the two carboxyl groups being in a trans configuration and having no intramolecular hydrogen bonds. All observed conformers exhibit a trans structure around the central C-C bond.

The selective IR pumping promotes conformational inter-conversion between the three observed conformers (which implies internal rotation around a C-O bond). On the other hand, C-C internal rotation leading to conformers exhibiting a cis $\text{O}=\text{C}-\text{C}=\text{O}$ axis was not observed, which can be explained taking into consideration volume restrictions imposed by the rigidity of the matrix that increases the effective barrier for the

rotation around this bond. The same behavior has been observed in other dicarboxylic acids that are actually being studied in our laboratories (e.g., malonic acid, fumaric acid).

Acknowledgment. This work has been held within the PRAXIS XXI (QUI/2/2.1/412/94) research program that is also partially funded by FEDER. The financial support of Academy of Finland is also acknowledged.

References and Notes

- (1) Derissen, J. L.; Smit, P. H. *Acta Crystallogr. B* **1974**, *B30*, 2240-2242.
- (2) Maçôas, E. M. S. *Caracterização Estrutural e Espectroscópica de Ácidos Carboxílicos*; Department of Chemistry, University of Coimbra, Portugal, Internal Report, 1997 (and references therein).
- (3) Nieminen, J.; Räsänen, M.; Murto, J. *J. Phys. Chem.* **1992**, *96*, 5303-5308.
- (4) Stace, B. C.; Oralratmanee, B. C. *J. Mol. Struct.* **1973**, *18*, 339.
- (5) Redington, L. R.; Redington, T. E. *J. Mol. Struct.* **1978**, *48*, 165.
- (6) Bock, C. W.; Redington, R. L. *J. Chem. Phys.* **1986**, *85*, 5391.
- (7) Higgins, J.; Zhou, X.; Liu, R.; Huang T. T.-S. *J. Phys. Chem. A* **1997**, *101*, 2702-2708.
- (8) Godfrey, P. D.; Mirabella, M. J.; Brown R. D. *J. Phys. Chem. A* **2000**, *104*, 258.
- (9) Bock, R. A. *Can. J. Chem.* **1984**, *62*, 1414.
- (10) Kulbida, A.; Nosov, A. *J. Mol. Struct.* **1992**, *265*, 17.
- (11) Kulbida, A.; Fausto, R. *J. Chem. Soc., Faraday Trans.* **1993**, *89*, 4257.
- (12) Pettersson, M.; Lundell, J.; Khriachtchev, L.; Räsänen, M. *J. Am. Chem. Soc.* **1997**, *119*, 11715.
- (13) Becke, A. D. *J. Chem. Phys.* **1993**, *98*, 5648.
- (14) (a) Lee, C.; Yang, W.; Parr, R. G. *Phys. Rev.* **1988**, *B37*, 785. (b) Miehlich, B.; Savin, A.; Stoll, H. Preuss, H. *Chem. Phys. Lett.* **1989**, *157*, 200.
- (15) Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Gill, P. M. W.; Johnson, B. G.; Wong, M. W.; Foresman, J. B.; Robb, M. A.; Head-Gordon, M.; Replogle, E. S.; Gomperts, R.; Andres, J. L.; Raghavachari, Binkley, J. S.; Gonzalez, C.; Martin R. L.; Fox, D. J.; Defrees, D. J.; Baker, J.; Stewart, J. J. P.; Pople, J. A. GAUSSIAN92/DFT (rev. G.2), Gaussian Inc., Pittsburgh, PA, 1993.
- (16) Hehre, W. J.; Ditchfield, R.; Pople J. A. *J. Chem. Phys.* **1972**, *56*, 2257.
- (17) Fausto R. TRANSFORMER (version 2.0), Chemistry Department, University of Coimbra, Portugal, 1997.
- (18) Faria, M. D. G.; Fausto R. BUILD-G (version 2.0) and VIBRAT (version 2.0). Chemistry Department, University of Coimbra, Portugal, 1997 (these programs incorporate several routines from programs GMAT and FPERT, Fuher, H.; Kartha, V. B.; Krueger, P. J.; Mantsch, H. H. *N. R. C. Bull.* **1976**, *15*, 1.
- (19) Teixeira-Dias, J. J. C.; Fausto, R. *THEOCHEM* **1993**, *282*, 123.
- (20) Fausto, R. *THEOCHEM* **1994**, *315*, 123.
- (21) Hallamasek, D.; Babka, E.; Knözinger, E. *J. Mol. Struct.* **1997**, *408/409*, 125.
- (22) Redington, R. L.; Liang, C. K. *J. Mol. Spectrosc.* **1984**, *104*, 25.
- (23) Fröchtenicht, B.; Kaloudis, M.; Koch, M.; Huisken, F. *J. Chem. Phys.* **1996**, *105*, 6128.