

# Inhibition of biofilm formation and swarming of *Bacillus subtilis* by (5*Z*)-4-bromo-5-(bromomethylene)-3-butyl-2(5*H*)-furanone

D. Ren<sup>1</sup>, J.J. Sims<sup>2</sup> and T.K. Wood<sup>1</sup>

<sup>1</sup>Departments of Chemical Engineering and Molecular & Cellular Biology, University of Connecticut, Storrs, and <sup>2</sup>Department of Plant Pathology, University of California, Riverside, USA

2001/307: received 26 October 2001 and accepted 7 January 2002

D. REN, J.J. SIMS AND T.K. WOOD. 2002.

**Aims:** (5*Z*)-4-Bromo-5-(bromomethylene)-3-butyl-2(5*H*)-furanone (furanone) of the marine alga *Delisea pulchra* was synthesized, and its inhibition of swarming motility and biofilm formation of *Bacillus subtilis* was investigated.

**Methods and Results:** Furanone was found to inhibit both the growth of *B. subtilis* and its swarming motility in a concentration-dependent way. In addition, as shown by confocal scanning laser microscopy, furanone inhibited the biofilm formation of *B. subtilis*. At 40 µg ml<sup>-1</sup>, furanone decreased the biofilm thickness by 25%, decreased the number of water channels, and reduced the percentage of live cells by 63%.

**Conclusions, Significance and Impact of the Study:** Natural furanone has potential for controlling the multicellular behaviour of Gram-positive bacteria.

## INTRODUCTION

The incidence of Gram-positive bacterial infections has increased significantly in recent years. For example, the number of Gram-positive infections per 1000 hospital admissions increased sixfold in the UK from 1985 to 1996, while the number of Gram-negative infections increased fourfold in the same period (Crowe *et al.* 1998). In addition, Gram-positive bacteria were responsible for more than 60% of the nosocomial bloodstream infections in the United States, while Gram-negative bacteria were responsible for only 27% of such infections (Edmond *et al.* 1999). Since antibiotics are not equally efficient for Gram-negative and Gram-positive bacteria, the appearance of antibiotic-resistant Gram-positive bacteria poses a serious problem (Edmond *et al.* 1999). Furthermore, Gram-positive biofilms cause significant corrosion problems, such as that of *Desulfosporosinus orientis* which corrodes mild steel seriously (Kumar *et al.* 1999). Therefore, it would be advantageous to have additional Gram-positive antimicrobials.

Swarming and biofilm formation are strongly related to disease (Allison *et al.* 1992; Elvers and Lappin-Scott 2000).

Although the mechanisms of these two important multicellular activities have been investigated (Allison and Hughes 1991; Davies *et al.* 1998; Kolter and Losick 1998), techniques for efficient control are still insufficient.

Swarming exists for both Gram-positive and Gram-negative bacteria, and many species have been shown to exhibit this motility, including *Proteus mirabilis* and *Bacillus subtilis* (Allison and Hughes 1991). Swarming has three stages: (i) recognition of appropriate environmental signals with the formation of elongated and hyperflagellated swarming cells; (ii) co-ordinated movement of the swarming cells; (iii) consolidation to vegetative cells and initiation of a new round of swarming (Allison and Hughes 1991). It occurs at specific surface viscosities (e.g. 0.7–0.8% agar plates for *Serratia marcescens*) and nutrient conditions (Allison and Hughes 1991; Fujikawa 1994). In the laboratory, swarming can take place on semi-solid agar plates with a speed as high as 16 µm s<sup>-1</sup> (Allison and Hughes 1991). *Bacillus subtilis* has been shown to exhibit swarming motility that is affected by conditions such as nutrient status (Fujikawa 1994), surface wetness (Mendelson and Salhi 1996), and the existence of chemicals such as surfactants and detergents (Thampuran and Surendran 1996).

Biofilm formation is a form of multicellular behaviour that is ubiquitous in aquatic environments, and *B. subtilis* has been shown to form biofilms on metal surfaces (Jayaraman

Correspondence to: Dr T.K. Wood, Departments of Chemical Engineering and Molecular & Cellular Biology, University of Connecticut, 191 Auditorium Road, Storrs, CT 06269–3222, USA (e-mail: twood@enr.uconn.edu).

*et al.* 1997). By attachment to submerged surfaces and production of microbial products such as polysaccharides (Elvers and Lappin-Scott 2000), bacteria are able to form biofilms with complex, three-dimensional structures. The top or base of the biofilm has the highest cell density, and water channels (lateral variation) exist for the transport of nutrients and waste (Lawrence *et al.* 1991; Kolter and Losick 1998; Elvers and Lappin-Scott 2000).

It has been found that cell communication, based on density-dependent chemical inducers termed quorum sensing, is important for bacterial multicellular behaviour such as swarming (Eberl *et al.* 1996b) and biofilm formation (Davies *et al.* 1998). Cell communication in quorum sensing is based on small molecules, called autoinducers (AI), such as acylated homoserine lactones for Gram-negative bacteria and peptide signalling molecules for Gram-positive bacteria (Bassler 1999). Quorum sensing is a complex system in which one strain may encode multiple autoinducers and may have interspecies signalling (sense the population of other bacteria). For example, *Vibrio harveyi* has two AI systems, AI-1 (hydroxybutanoyl-L-homoserine lactone) (Cao *et al.* 1989) for species-specific quorum sensing, and AI-2 (furanone-related compound furanosyl borate diester, derived from 4,5-dihydroxy-2,3-pentanedione) (Schauder *et al.* 2001; Chen *et al.* 2002) for non-specific quorum sensing (Bassler *et al.* 1997; Surette and Bassler 1998; Freeman and Bassler 1999). Whilst the AI-1 system is exclusive to Gram-negative bacteria, the AI-2 system is found in both Gram-positive (including *B. subtilis* (Surette *et al.* 1999)) and Gram-negative bacteria (Surette *et al.* 1999).

Recently, several natural furanones produced by the seaweed *Delisea pulchra* have been found to reduce the growth rate of Gram-positive bacteria, including *Staphylococcus aureus* and *Staph. epidermidis*, but to have no effect on the growth of mammalian cells (Kjelleberg 1998). It has also been shown that such furanones can inhibit several activities of Gram-negative bacteria without affecting the growth rate. These include the swarming of *Proteus mirabilis* (Gram *et al.* 1996), the biofilm formation of *Escherichia coli* (Ren *et al.* 2001) and siderophore synthesis of *Pseudomonas putida* F1 (Ren *et al.* unpublished). It is likely that these furanones can inhibit the multicellular behaviour of Gram-negative bacteria by suppression of the autoinduction circuit (Givskov *et al.* 1996), and furanone has been shown to inhibit quorum sensing based on either the AI-1 or AI-2 system (Ren *et al.* 2001).

In this paper, the antimicrobial properties of furanone were investigated by examining its impact on the growth rate, swarming motility and biofilm formation of *B. subtilis*. This is one of the first reports of the antibiotic effect of furanone on *B. subtilis* and its effect on the swarming and biofilm formation of a Gram-positive bacterium.

## MATERIALS AND METHODS


### Bacterial strains and culture media

*Bacillus subtilis* BE1500 (Nagarajan *et al.* 1992) with plasmid pBE92 (henceforth *B. subtilis*) was obtained from E.I. du Pont de Nemours Inc. (Wilmington, DE, USA) and was grown at 37°C in LB medium supplemented with kanamycin at 50 µg ml<sup>-1</sup>. pBE92 is a derivative of pBE60 that encodes kanamycin resistance (Nagarajan *et al.* 1992) and was used to avoid contamination while culturing the *Bacillus* strain.

LB medium supplemented with 0.4% agar (20 ml) was used to study swarming in 100 × 15 mm Petri dishes which were dried for 2 h at 30°C. The furanone was dissolved in dichloromethane at various concentrations (119–2380 µg ml<sup>-1</sup>) and added to the top of the agar plate in 40 µl (40 µl dichloromethane was added to all plates, including those with no furanone). To determine whether the furanone would form a gradient in the agar plate by diffusion, or remain at the agar surface, tests were conducted by dissolving hydrophobic ink in dichloromethane and adding it to the top of agar plates. It was found that the dichloromethane did not diffuse into the agar. Hence, the concentration of furanone was based on the area of this layer but not the volume. After adding the furanone, swarming plates were dried for another 3 h at 30°C. *Bacillus subtilis* cells were inoculated with toothpicks at the centre of the furanone-containing area from 2-day-old streaks on LB medium supplemented with 50 µg ml<sup>-1</sup> kanamycin. The swarming colonies were grown at 37°C. At least seven plates were used for each furanone concentration.

### Furanone synthesis

4-Bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone (Fig. 1) was synthesized as reported previously (Beechan and Sims 1979), with the exception that the furanone was purified using column chromatography with hexane and ethyl acetate at a ratio of 100:1 (2.5 × 60 cm column, Spectrum Chromatography, Houston, TX, USA). The structure was verified with <sup>1</sup>H-NMR (Bruker DRX-400 MHz, Billerica, MA, USA; 6.24 single peak, vinylidene; 2.39 triple peaks, coupling constant J 7.2 Hz, allylic methylene; 0.93 triple peaks, coupling constant J 7.2 Hz, terminal methyl), mass spectroscopy (Series II 5890, Hewlett Packard, San Fernando, CA, USA; molecular ion peaks M<sup>+</sup>: 308, 310, 312) and infrared spectroscopy (Nicolet-Magna-IR 560, Madison, WI, USA; reciprocal absorbing wavelength: 2958, 1793, 1610, 1276, 1108, 1030 cm<sup>-1</sup>) by comparing with literature values (Beechan and Sims 1979; Manny *et al.* 1997). Routinely, the furanone product was verified by comparing with standard furanone synthesized by the method of Beechan and Sims (1979) using thin layer chromatography (hexane and ethyl acetate at a ratio of 20:1, R<sub>f</sub> = 0.8; Silica Gel 60, F-254).



**Fig. 1** Structure of (5Z)-4-bromo-5-(bromomethylene)-3-butyl-2(5H)-furanone

### Growth rate

The growth rates in liquid cultures to which furanone had been added were measured in duplicate in 250 ml shake flasks with LB medium (25 ml) supplemented with 50  $\mu\text{g ml}^{-1}$  kanamycin. The furanone was dissolved in 95% ethanol (14.9  $\text{mg ml}^{-1}$ ) and added to each flask to give a final concentration of 10, 20 and 40  $\mu\text{g ml}^{-1}$ . All of the flasks (including those with no furanone) were supplemented with the same amount of ethanol (67.1  $\mu\text{l}$  of 95% ethanol to 25 ml medium) to eliminate the effects of the solvent.

To determine the effect of furanone on colony growth of *B. subtilis*, furanone was added to the top of the LB agar (1.5%) plates which support growth but not swarming. Furanone was added by dissolving in dichloromethane at different concentrations in a volume of 40  $\mu\text{l}$ . *Bacillus subtilis* was inoculated as in the swarming experiments. The same concentrations of furanone as those in the swarming experiments were tested.

### Biofilm formation

The impact of furanone on biofilm formation was determined with SAE1018 steel coupons (25.5 mm diameter and 1.2 mm thick, polished with P240 polishing paper, 3M Imperial Wet-or-Dry, St Paul, MN, USA). A single coupon was placed in a 100  $\times$  15 mm Petri dish with 20 ml LB supplemented with 50  $\mu\text{g ml}^{-1}$  kanamycin. The furanone was dissolved in 95% ethanol (14.9  $\text{mg ml}^{-1}$ ) and added to the culture at the start of the experiments with the final concentration of 20 or 40  $\mu\text{g ml}^{-1}$  (the same amount of 95% ethanol, 0.6%, was added to all samples, including those without furanone). Coupons were incubated at 37°C without shaking for 3 days, then analysed by confocal microscopy. Metal coupons with attached biofilms were removed gently from Petri dishes and immersed immediately in 0.85% NaCl solution to remove the bulk of planktonic cells. The biofilm was stained with the Live/Dead Baclit bacteria viability assay kit (L-7007, Molecular Probes, Inc., Eugene, OR, USA) at a concentration of 1.125  $\mu\text{l ml}^{-1}$  (for both staining components, A and

B, of this kit). The staining process was performed in the dark at room temperature for 30 min with 4 ml staining solution for each sample.

After staining, the coupons were protected with a cover glass and observed with confocal scanning laser microscopy (MRC 600, BIO-RAD, Hercules, CA, USA) using the dual channel mode (K1/K2 filter block combination); the sample was excited at 488 nm and 568 nm and detected from 585–infinite nm and 506–538 nm. The biofilm thickness was measured by focusing through the biofilm from the solid–air interface to the metal surface, and averaging five or six positions for each coupon (17 coupons were tested). The images were processed with the National Institutes of Health Image 1.6 and Adobe Photoshop 5.5 (Adobe, San Jose, CA, USA). Z-section images (single image of the sectioning plane parallel to the metal surface) and vertical section images (a collection of images orthogonal to the metal surface at one position) were obtained.

## RESULTS

### Effect of furanone on growth

Furanone strongly affected the growth of *B. subtilis* on LB agar plates (Table 1) in a concentration-dependent manner; 52  $\mu\text{g cm}^{-2}$  furanone completely inhibited growth in 48 h. Inhibition of growth was also seen in suspension cultures (Table 2). At 0 and 10  $\mu\text{g ml}^{-1}$  furanone, the cultures grew immediately after inoculation, although 10  $\mu\text{g ml}^{-1}$  furanone caused a 28% decrease in the specific growth rate. At a concentration of 20  $\mu\text{g ml}^{-1}$  furanone, the lag phase was extended significantly (about 13 h), and the specific growth rate was decreased by 36% compared with that of the control. At a concentration of 40  $\mu\text{g ml}^{-1}$ , furanone inhibited growth completely (no growth obtained in 24 h).

### Effect of furanone on swarming motility

The swarming motility of *B. subtilis* was highly reproducible, and the morphology of the swarming colonies was consistent with that reported previously (Allison and

**Table 1** Effect of furanone on colony growth of *Bacillus subtilis*

Time (h)	Colony diameter (mm)			
	Concentration of furanone ( $\mu\text{g/cm}^2$ )			
	0	13	26	52
0	NG	NG	NG	NG
24	8.3 $\pm$ 0.3	2.0 $\pm$ 1.0	NG	NG
48	15.2 $\pm$ 0.3	9.3 $\pm$ 1.2	8.5 $\pm$ 0.5	0.5 $\pm$ 0.5
72	24.3 $\pm$ 1.2	15.7 $\pm$ 1.5	13.0 $\pm$ 0	9.0 $\pm$ 0


$\pm$ : Standard deviation; NG: no growth.

**Table 2** Effect of furanone on the growth rate of planktonic (suspension) cells and on the biofilm formation of *Bacillus subtilis*

Furanone concentration ( $\mu\text{g ml}^{-1}$ )	Growth rate		Biofilm formation	
	$\mu$ ( $\text{h}^{-1}$ )	Lag period (h)	Thickness of biofilm ( $\mu\text{m}$ )	Percentage of live cells in the biofilm
0	$1.71 \pm 0.09$	0	$14 \pm 1$	82.7
10	$1.23 \pm 0.03$	0	ND	ND
20	$1.09 \pm 0.01$	13	$13 \pm 2$	77.2
40	0	ND	$10 \pm 1$	30.9

±: Standard deviation; ND: not determined.

Hughes 1991). Fourteen hours after inoculation, colonies without furanone began to swarm at a rate of  $0.3 \mu\text{m s}^{-1}$  (average speed of active and quiescent phases of growth), which is similar to previous reports (around  $0.41 \mu\text{m s}^{-1}$ ) (Mendelson and Salhi 1996). However, the presence of furanone reduced the swarming speed in a concentration-dependent manner (Fig. 2), with 99.9% swarming inhibited at  $52 \mu\text{g cm}^{-2}$ . The swarming speed of the colonies with furanone was  $0.24 \mu\text{m s}^{-1}$  ( $13 \mu\text{g cm}^{-2}$  furanone),  $0.12 \mu\text{m s}^{-1}$  ( $26 \mu\text{g cm}^{-2}$  furanone) and  $0 \mu\text{m s}^{-1}$  ( $52 \mu\text{g cm}^{-2}$  furanone) (Fig. 2). Furanone at  $52 \mu\text{g cm}^{-2}$  inhibited swarming completely, but growth was still present as evidenced by the slow increase in colony size 26 h after inoculation (Fig. 2). The shape of the colonies with furanone was very similar to that of the colonies without furanone (Fig. 3a). In addition, a high concentration of furanone was also tested;  $250 \mu\text{g cm}^{-2}$  furanone inhibited not only swarming but also growth, as no growth appeared after 3 days (the inoculated colonies did not expand). Streaking the inoculated cells on a LB plate indicated that the cells were dead. This is consistent with the growth rate experiment in that furanone inhibited the growth of *B. subtilis* with a concentration as low as



**Fig. 2** Colony swarming of *Bacillus subtilis* in the presence of furanone (increasing colony diameter with time). (●) Control; (○)  $13 \mu\text{g cm}^{-2}$  (62.3% inhibited); (■)  $26 \mu\text{g cm}^{-2}$  (95.1% inhibited); (□)  $52 \mu\text{g cm}^{-2}$  (99.9% inhibited)

$13 \mu\text{g cm}^{-2}$ . However, when the *B. subtilis* colony that began swarming on the side of the plate without furanone entered the furanone-containing area, its swarming was not affected (Fig. 3b,c). This suggested that the swarming colony is more resistant to furanone once swarming is initiated.


It should be noted that the furanone was only added to a small region of the top of the agar plate ( $40 \mu\text{l}$  dichloromethane formed a  $1.7 \text{ cm}^2$  round layer), so the swarming speed can only be calculated with the initial part of the curves when the colonies were young (Fig. 2). Once the swarming colonies matured (both within and without the furanone-containing zone), the swarming speed increased almost to the speed of a swarming colony in the absence of furanone. This further indicates that mature swarming colonies are more resistant to furanone.

### Effect of furanone on biofilm formation


Using confocal scanning laser microscopy, it was observed that 20 and  $40 \mu\text{g ml}^{-1}$  furanone caused a 5.6% and 25% decrease in the biofilm thickness, respectively (Table 2). In addition,  $20 \mu\text{g ml}^{-1}$  furanone did not significantly affect the ratio of live to dead cells in the biofilm (reduced by 6.7%), while  $40 \mu\text{g ml}^{-1}$  decreased the percentage of live cells by 63% compared with the control (no furanone, Table 2). Vertical sections of the biofilm indicated no change in the biofilm structure with 0 and  $20 \mu\text{g ml}^{-1}$  furanone, and the water channels could be seen clearly (dark area between the cell clusters). At  $40 \mu\text{g ml}^{-1}$ , the water channels began to collapse (Fig. 4). Hence, furanone affects *B. subtilis* biofilm formation significantly. Interestingly,  $40 \mu\text{g ml}^{-1}$  furanone completely inhibited the growth of *B. subtilis* in shake flasks (no growth in 24 h), while it did not completely prevent biofilm formation. However,  $60 \mu\text{g ml}^{-1}$  furanone inhibited biofilm formation completely (data not shown).

### DISCUSSION

Besides swarming, swimming is an important mechanism for bacterial surface translocation (Allison and Hughes 1991). At



**Fig. 3** Effect of furanone on the swarming motility of *Bacillus subtilis* (the circle indicates the area in which furanone was added). (a)  $52 \mu\text{g cm}^{-2}$  furanone, 40 h after inoculation; (b)  $250 \mu\text{g cm}^{-2}$  furanone, 40 h after inoculation; (c)  $250 \mu\text{g cm}^{-2}$  furanone, 70 h after inoculation



**Fig. 4** Effect of furanone on the *Bacillus subtilis* biofilm (representative confocal images). Bar =  $10 \mu\text{m}$ . A1 (Z section, i.e. planar view at top of the biofilm at water interface) and A2 (vertical section, i.e. biofilm viewed on edge by compiling Z sections), no furanone; B1 (Z section) and B2 (vertical section),  $20 \mu\text{g ml}^{-1}$  furanone; C1 (Z section) and C2 (vertical section),  $40 \mu\text{g ml}^{-1}$  furanone. Live cells were stained green and dead cells were stained red using the BacLit viability kit. The void spaces between cell clusters are water-filled channels for nutrient and waste transfer (Lawrence *et al.* 1991; Kolter and Losick 1998)

low agar concentrations, cells swim through water-filled channels and form colonies with morphologies similar to those of chemotactic rings (Eberl *et al.* 1996a). Swimming is the result of rotation of the bacterial flagella rotor and is not a multicellular behaviour (Moat and Foster 1995). After

investigating its behaviour on LB plates of different agar concentrations, *B. subtilis* was found to be capable of swimming at agar concentrations lower than 0.3%, and of swarming on 0.4% agar plates. Hence, swimming behaviour was eliminated by controlling the agar concentration. The

non-fractal morphology of the swarming colonies obtained here agrees well with that reported previously (Allison and Hughes 1991). Since the furanone was dissolved in dichloromethane before adding to the swarming plates, experiments were conducted to study the effect of dichloromethane on swarming, and no effect was found (data not shown). From the results of this study, it can be seen that furanone strongly inhibits the swarming of *B. subtilis*.

*Bacillus subtilis* is predicted to produce, via the LuxS synthase, a quorum sensing signal, AI-2 (Surette *et al.* 1999), and this AI-2 probably has a structure related to that of furanone (Chen *et al.* 2002). Since the brominated furanone used in this study has been shown to inhibit quorum sensing based on AI-2 (Ren *et al.* 2001), it probably inhibits swarming and biofilm formation of *B. subtilis* not only by suppression of growth but also by interfering with quorum sensing. However, it has not been possible to discriminate between these two possible effects. Given that the furanone has been shown to inhibit the growth of Gram-positive bacteria at a concentration that is non-toxic to mammalian cells (Kjelleberg 1998), it has potential as an antimicrobial agent. This study therefore provides preliminary information for the potential use of furanone to inhibit the multicellular behaviour of Gram-positive bacteria. The increased resistance to furanone exhibited by *B. subtilis* swarming colonies and biofilm suggests that it may be advantageous to combine furanone with other antibiotics for better performance. In addition, modifying the structure of natural furanone may make it more efficient.

## ACKNOWLEDGEMENTS

The authors thank Prof. David Knecht for assistance with confocal scanning laser microscopy. The assistance of Mr Xing Dong (NMR analysis) and Mr Haihu Qin (IR analysis) is also appreciated.

## REFERENCES

- Allison, C., Coleman, N., Jones, P.L. and Hughes, C. (1992) Ability of *Proteus mirabilis* to invade human urothelial cells is coupled to motility and swarming differentiation. *Infection and Immunity* **60**, 4740–4746.
- Allison, C. and Hughes, C. (1991) Bacterial swarming: an example of prokaryotic differentiation and multicellular behaviour. *Science Progress Edinburgh* **75**, 403–422.
- Bassler, B.L. (1999) How bacteria talk to each other: regulation of gene expression by quorum sensing. *Current Opinions in Microbiology* **2**, 582–587.
- Bassler, B.L., Greenberg, E.P. and Stevens, A.M. (1997) Cross-species induction of luminescence in the quorum-sensing bacterium *Vibrio harveyi*. *Journal of Bacteriology* **179**, 4043–4045.
- Beechan, C.M. and Sims, J.J. (1979) The first synthesis of fimbrolides, a novel class of halogenated lactones naturally occurring in the red seaweed *Delisea fimbriata* (Bonnemaisoniaceae). *Tetrahedron Letters* **19**, 1649–1652.
- Cao, J. and Meighen, E.A. (1989) Purification and structural identification of an autoinducer for the luminescence system of *Vibrio harveyi*. *Journal of Biological Chemistry* **264**, 21670–21676.
- Chen, X., Schauder, S., Potier, N. and Hughson, F.M. (2002) Structural identification of a bacterial quorum-sensing signal containing boron. *Nature* **415**, 545–549.
- Crowe, M., Ispahani, P., Humphreys, H., Kelley, T. and Winter, R. (1998) Bacteraemia in the adult intensive care unit of a teaching hospital in Nottingham, UK, 1985–1996. *European Journal of Clinical Microbiological Infection and Disease* **17**, 377–384.
- Davies, D.G., Parsek, M.R., Pearson, J.P., Iglewski, B.H., Costerton, J.W. and Greenberg, E.P. (1998) The involvement of cell-to-cell signals in the development of a bacterial biofilm. *Science* **280**, 295–298.
- Eberl, L., Christiansen, G., Molin, S. and Givskov, M. (1996a) Differentiation of *Serratia liquefaciens* into swarm cells is controlled by the expression of the *flhD* master operon. *Journal of Bacteriology* **178**, 554–559.
- Eberl, L., Winson, M.K., Sternberg, C. *et al.* (1996b) Involvement of N-acyl-L-homoserine lactone autoinducers in controlling the multicellular behaviour of *Serratia liquefaciens*. *Molecular Microbiology* **20**, 127–136.
- Edmond, M.B., Wallace, S.E., McClish, D.K., Pfaller, M.A., Jones, R.N. and Wenzel, R.P. (1999) Nosocomial bloodstream infections in United States hospitals: a three-year analysis. *Clinical Infectious Diseases* **29**, 239–244.
- Evers, K.T. and Lappin-Scott, H.M. (2000) *Encyclopedia of Microbiology*. ed. Lederberg, J. San Diego, California: Academic Press.
- Freeman, J.A. and Bassler, B.L. (1999) A genetic analysis of the function of LuxO, a two-component response regulator involved in quorum sensing in *Vibrio harveyi*. *Molecular Microbiology* **31**, 665–677.
- Fujikawa, H. (1994) Diversity of the growth patterns of *Bacillus subtilis* colonies on agar plates. *FEMS Microbiology Ecology* **13**, 159–168.
- Givskov, M., de Nys, R., Manefield, M. *et al.* (1996) Eukaryotic interference with homoserine lactone-mediated prokaryotic signaling. *Journal of Bacteriology* **178**, 6618–6622.
- Gram, L., de Nys, R., Maximilien, R., Givskov, M., Steinberg, P. and Kjelleberg, S. (1996) Inhibitory effects of secondary metabolites from the red alga *Delisea pulchra* on swarming motility of *Proteus mirabilis*. *Applied and Environmental Microbiology* **62**, 4284–4287.
- Jayaraman, A., Cheng, E.T., Earthman, J.C. and Wood, T.K. (1997) Importance of biofilm formation for corrosion inhibition of SAE 1018 steel by pure-culture aerobic biofilm. *Journal of Industrial Microbiology and Biotechnology* **18**, 396–401.
- Kjelleberg, S. (1998) Inhibition of Gram positive bacteria. International patent WO 99/53915.
- Kolter, R. and Losick, R. (1998) One for all and all for one. *Science* **280**, 226–227.
- Kumar, A.V.R., Singh, R. and Nigam, R.K. (1999) Mossbauer spectroscopy of corrosion products of mild steel due to microbologically influenced corrosion. *Journal of Radioanalytical and Nuclear Chemistry* **242**, 131–137.
- Lawrence, J.R., Korber, D.R., Hoyle, B.D., Costerton, J.W. and Caldwell, D.E. (1991) Optical sectioning of microbial biofilms. *Journal of Bacteriology* **173**, 6558–6567.

- Manny, A.J., Kjelleberg, S., Kumar, N., de Nys, R., Read, R.W. and Steinberg, P. (1997) Reinvestigation of the sulfuric acid-catalysed cyclisation of brominated 2-alkyllevulinic acids to 3-alkyl-5-methylene-2 (5H)-furanones. *Tetrahedron* **53**, 15813–15826.
- Mendelson, N.H. and Salhi, B. (1996) Patterns of reporter gene expression in the phase diagram of *Bacillus subtilis* colony forms. *Journal of Bacteriology* **178**, 1980–1989.
- Moat, A.G. and Foster, J.W. (1995) *Microbial Physiology*. New York: Wiley-Liss.
- Nagarajan, V., Albertson, H., Chen, M. and Ribbe, J. (1992) Modular expression and secretion vectors for *Bacillus subtilis*. *Gene* **114**, 121–126.
- Ren, D., Sims, J.J. and Wood, T.K. (2001) Inhibition of biofilm formation and swarming of *Escherichia coli* by (5Z)-4-Bromo-5-(Bromomethylene)-3-Butyl-2(5H)-Furanone. *Environmental Microbiology* **3**, 731–736.
- Schauder, S., Shokat, K., Surette, M.G. and Bassler, B.L. (2001) The LuxS-family of bacterial autoinducers: biosynthesis of a novel quorum-sensing signal molecule. *Molecular Microbiology* **41**, 463–476.
- Surette, M.G. and Bassler, B.L. (1998) Quorum sensing in *Escherichia coli* and *Salmonella typhimurium*. *Proceedings of the National Academy of Sciences USA* **95**, 7046–7050.
- Surette, M.G., Miller, M.B. and Bassler, B.L. (1999) Quorum sensing in *Escherichia coli*, *Salmonella typhimurium*, and *Vibrio harveyi*: a new family of genes responsible for autoinducer production. *Proceedings of the National Academy of Sciences USA* **96**, 1639–1644.
- Thampuran, N. and Surendran, P.K. (1996) Effect of chemical agents on swarming of *Bacillus* species. *Journal of Applied Bacteriology* **80**, 296–302.