

Inhibition of human LDL lipid peroxidation by phenol-rich beverages and their impact on plasma total antioxidant capacity in humans

Mauro Serafini,* João A.N. Laranjinha,[†] Leonor M. Almeida,[†] and G. Maiani*

*Antioxidant Research Laboratory, National Institute for Food and Nutrition Research, Rome, Italy and [†]Laboratório de Bioquímica, Faculdade de Farmácia, Universidade de Coimbra, Coimbra, Portugal

Mounting evidence shows that phenol-rich beverages exert strong antioxidant activity. However, in vivo evidence has produced conflicting results. In the present study, we studied the impact of the ingestion of 300 mL of black and green tea, alcohol-free red wine, alcohol-free white wine, or water on plasma total antioxidant capacity in five healthy volunteers. Red wine has the highest content of phenolics (3.63 ± 0.48 g QE/L), followed by green tea (2.82 ± 0.07 g QE/L), black tea (1.37 ± 0.15 g QE/L), and white wine (0.31 ± 0.01 g QE/L). Plasma total antioxidant capacity values of subjects who drank green tea rose at 30 min (P < 0.05). After black tea and red wine ingestion, the peaks were at 50 min (P < 0.05 and P < 0.01, respectively). No changes were observed in the control and white wine groups. Red wine and green tea were the most efficient in protecting low density lipoprotein from oxidation driven by peroxy and ferril radicals, respectively. Phenol-rich beverages are a natural source of antioxidants; however, the phenolic content alone cannot be considered an index of their in vivo antioxidant activity. (J. Nutr. Biochem. 11:585–590, 2000) © Elsevier Science Inc. 2000. All rights reserved.

Keywords: antioxidant capacity; LDL; wine; tea; phenolics; plasma; humans

Introduction

Advances in understanding the role of free radicals in the pathogenesis of degenerative diseases and in the process of aging suggest a potential health-promoting role of dietary antioxidants. Recent evidence suggests that many antioxidant compounds, commonly called “phytochemicals” and not yet identified as nutrients in plant foods, are bioactive.^{1,2} Phytochemicals include numerous substances such as tannins, phytoestrogens, tocotrienols, and phenolic compounds. Phenolic compounds comprise a considerable number of compounds (more than 4,000) present in large amounts in plant foods,^{3,4} tea, and wine.⁵

Epidemiological studies have shown that consumption of phenol-rich beverages (PRB) such as tea and wine correlates with reduced coronary heart disease mortality.^{6,7}

Nevertheless, it has been claimed that wine might, indeed, be one of the most active ingredients involved in the protective effect of the Mediterranean Diet and one of the possible explanations of the “French Paradox.”^{8,9} The putative role that PRB are thought to play in the prevention of oxidative stress-linked diseases is mainly based on their antioxidant properties.¹⁰ In vitro studies have consistently showed that wine and tea phenolics have significant antioxidant activity as measured by their ability to inhibit low density lipoprotein (LDL) oxidation^{11–13} and in neutralizing free radicals.^{10,14} However, experimental ex vivo and in vivo studies have produced contrasting results,^{15–22} suggesting that the in vivo situation cannot be extrapolated on the basis of the in vitro evidence (i.e., the bioavailability of phenolics is limited and depends upon such different variables as nutrient interactions, gut microflora, etc.).

In the present study, the in vitro and in vivo antioxidant properties of black tea (BT), green tea (GT), alcohol-free red wine (RW), and alcohol-free white wine (WW) have been studied in two different models: (1) evaluation of the in vitro total antioxidant capacity and effect on LDL

Address correspondence to Dr. Mauro Serafini, Antioxidant Research Laboratory, National Institute for Food and Nutrition Research, Via Ardeatina 546, 00178 Rome, Italy.

Received April 5, 2000; accepted August 15, 2000.

oxidized by a physiologically relevant oxidant (ferrylmyoglobin) and by a controlled flux of peroxy radicals, and (2) the impact on plasma total radical-trapping antioxidant parameter (TRAP) in humans.

Methods and materials

All chemicals were purchased from Sigma Chemical Co. (St. Louis, MO USA) unless otherwise stated. 6-Hydroxy-2,5,7,8-tetramethyl-chroman-2-carboxylic acid (Trolox) was purchased from Aldrich Chemical Co. (Milwaukee, MI USA), 2,2'-diazobis(2-amidinopropane) dihydrochloride (ABAP) was purchased from Wako Pure Chemical Industries Ltd. (Osaka, Japan), hydrogen peroxide was purchased from Merck (Darmstadt, Germany) as a 30% solution, and cis-Parinaric acid (PnA) was purchased from Molecular Probes (Eugene, OR USA). Solutions and reagents for TRAP assays were made using Milli-Q (Millipore, Bedford, MA USA) double-distilled water (resistance > 18 mW/cm²), and filtered through Chelex 100 resin Na⁺ form. Tea infusions were prepared using two commercially available brands of tea (Twinings Earl Grey and Birko Chinese green tea, KI). One hundred milliliters of boiling tap water was poured over 2.0 g of tea leaves, the mixture allowed to steep for 1.5 min, after which it was filtered. RW ("Chianti Classico Rocca Castagnoli" Farm, Siena, Italy), and WW ("Valdadige" bottled by Terre Freddo, Trento, Italy) were used in this study. To separate the nonalcoholic fraction rich in phenolics from the ethanol contained in the wine (a source of possible experimental confounds), the two wines were dealcoholized in a rotary evaporator at 25°C for 4 hr.

Total phenolic content

Total phenolic content of the tea and wine was measured according to a modification of the Folin-Ciocalteu colorimetric method.^{1,23} Five hundred microliters of the sample were added to 1 mL of 1.0 N HCL, and vigorously vortexed for 60 sec. After incubation at 37°C for 60 min, 1.0 mL of 2.0 mol/L NaOH in 75% methanol was added, and the resulting mixture vortexed for 3 min. Then, 1.0 mL of 10% (v/v) phosphoric acid was added to this mixture, and the sample was centrifuged at 1,500 × g for 10 min. The supernatant was removed and kept on ice in the dark, whereas the pellet was extracted again by adding 1.0 mL of a solution 1:1 (v/v) acetone: water and centrifuged for 10 min at 2,700 × g. The two supernatants were combined and filtered through a HV 0.45 μm filter (Millipore SJHV004NS). Two hundred microliters of sample were assayed for total polyphenols with the Folin-Ciocalteu reagent and expressed as Quercetin Equivalents (g QE/L).

TRAP assay

The method employed to assess the *in vitro* and *in vivo* peroxy-radical scavenging activities of PRB was developed in our laboratory.²⁴ It is based on the protection afforded by plasma or other substrate against the decay of a fluorescent target, R-Phycoerythrin (R-PE) during a peroxidation reaction. In the present study, samples were added to the reaction mixture made up to a 2.0-mL final volume and preincubated at 37°C for 5 min in 10-mm quartz fluorometer cuvettes. The oxidation reaction was started by adding ABAP at a final concentration of 5.0 mmol/L. The decay of R-PE fluorescence was monitored every 5 min for 90 min on a Perkin-Elmer (Norwalk, CT USA) LS-5 Luminescence Spectrometer equipped with a thermostatically controlled cell-holder. The monochromators operated at an excitation wavelength of 495 nm/5 nm slit width and an emission wavelength of 575 nm/5 nm slit width. The results were standardized using Trolox, a water-soluble analogue of α-tocopherol.

LDL Isolation and oxidation

LDL particles were isolated from fresh human plasma by density gradient ultracentrifugation in a Beckman L80 ultracentrifuge (Beckman Instruments, Inc., Palo Alto, CA USA) equipped with a Beckman 70.1 Ti fixed-angle rotor at 65,000 rpm at 15°C, following the procedure of Vieira et al.²⁵ The LDL fraction was then concentrated and simultaneously dialyzed for 45 min by ultrafiltration under nitrogen to obtain a fraction concentrated in LDL and free of water-soluble antioxidants.

PnA incorporated into LDL particles has been previously used as a probe to study LDL oxidation and its inhibition by antioxidants because it sensitively detects oxidant reactions within the LDL particles and allows for a quantitative characterization of the activity of the antioxidant under study.²⁶ The experimental conditions for PnA oxidation were set up as previously described²⁷ to ensure that the fluorescence intensity of incorporated PnA was linear with its concentration, and that LDL concentration was high enough to incorporate most of the probe, thus preventing PnA from interfering in the water phase. Changes in fluorescence were monitored with a Perkin-Elmer LS 50B luminescence spectrometer. The standard reaction mixture consisted in 2 mL phosphate buffer (110 mM NaCl, 20 mM phosphate, pH 7.4) containing 3 nmol PnA, 50 μM DTPA, and 120 μg LDL protein at 37°C with gentle stirring. Metmyoglobin was dialyzed against phosphate buffer (110 mM NaCl, 20 mM phosphate, pH 7.4) containing 50 μM EDTA and previously passed through Chelex 100 resin Na⁺ form. The oxidation reaction was initiated by addition of metmyoglobin (1 μM), followed 2 min later by hydrogen peroxide (1.5 μM) or AAPH (10 mM). Beverages were added 1 min before the peroxide or AAPH. The fluorescence of the probe was recorded in time using excitation and emission wavelengths of 324 and 413 nm, respectively (slit widths 3.5 nm). To minimize PnA fluorescence quenching due to the presence of hemoprotein, excitation and emission wavelengths were 310 and 425 nm, respectively. The effect of myoglobin redox transitions on PnA signal was checked.

In vivo study

Five healthy nonsmoking volunteers, ages 25–50 years, were selected after ascertaining that they had not been taking any vitamin supplements. On the first experimental day after an overnight fast, each subject was asked to ingest a bolus quantity (300 mL) of water. The same experiment was then repeated on different days using BT, GT, RW, and WW instead of water. The venous blood samples were obtained by antecubital venipuncture, with NaEDTA vacutainers, at times 0, 30, 50, and 120 min after the ingestion of the different beverages. Blood samples were immediately centrifuged at 12,000 × g for 3 min, plasma was separated and placed at –80°C, and assayed for TRAP within 3 hr. The study protocol was approved by the Human Ethics Committee of the National Institute of Nutrition, and written informed consent was obtained from each volunteer.

Statistics

Statistical analyses were performed using Statview II (Abacus Concepts Inc., Berkeley, CA USA) software for Macintosh. A *P*-value lower than 0.05 was considered statistically significant. Data are expressed as mean ± SEM or SD.

Results

In vitro study

Phenolic content and TRAP values of the beverages are shown in *Table 1*. RW had the highest amount of phenolics

Table 1 Total phenolic concentration and total radical-trapping antioxidant parameter (TRAP) of black tea, green tea, alcohol-free red wine, and white wine*

	Phenolics (g QE/L)	TRAP [†] (mmol/L)	Regression [‡] (r)
Black tea	1.37 ± 0.15	3.5 ± 0.20	0.994
Green tea	2.82 ± 0.07	17.8 ± 0.08	1.000
Alcohol-free red wine	3.63 ± 0.48	40.0 ± 0.10	0.999
Alcohol-free white wine	0.31 ± 0.01	1.9 ± 0.10	0.998

*Values are mean ± SD of three separate experiments.

[†]TRAP value is referred to 8 μ L of beverage, and represents the number of moles (mmol/L) of peroxy radicals trapped per liter of beverage.

[‡]Regression coefficient (r) of linear regression of lag-phase vs. 2, 4, and 8 μ L of undiluted black tea, diluted 1:200 (v/v) alcohol-free red wine, and diluted 1:2 (v/v) green tea, and alcohol-free white wine.

QE—Quercetin Equivalents.

(3.63 ± 0.48 g QE/L), followed by GT (2.82 ± 0.07 g QE/L) and BT (1.37 ± 0.15 g QE/L), whereas WW presented the lowest phenolic content (0.31 ± 0.01 g QE/L). The procedure adopted for the dealcoholization did not change the phenol content of the wines (data not shown). RW displayed the highest antioxidant capacity: twice that of GT and at least 10 and 20 times higher than BT and WW, respectively. There were no changes in wine TRAP values after removal of ethanol (data not shown). The correlation coefficient of the regression between increasing amounts of beverages (2, 4, and 8 μ L) and the length of lag-phase was 0.999 for RW, 0.998 for WW, 1 for GT, and 0.994 for BT.

The effect of phenol-rich beverages on LDL oxidation was determined in our study by PnA assay using different catalysts. These catalysts included AAPH, which generates peroxy radical at a fixed constant rate in the aqueous phase inducing the chain oxidation of human LDL by a free radical-mediated chain mechanism,²⁸ and ferrylmyoglobin, the product of metmyoglobin reaction with H₂O₂.²⁹ Ferrylmyoglobin is a strong oxidant able to initiate lipid peroxidation in several lipid systems, but its harmful effect can be counteracted by antioxidants reducing ferrylmyoglobin to the ferric form (for review, see Giulivi and Cadenas)³⁰. *Figure 1* shows a typical time course of PnA assay. When LDL was challenged with AAPH-derived peroxy radicals in the absence of external added antioxidants, PnA quickly underwent oxidative modification, as indicated by its rapid fluorescence decay. The presence of increasing amounts of wine (0.3, 0.5, and 0.7 μ L) in the medium protected PnA and LDL from oxidation in a dose dependent manner. The effect of the different beverages on peroxy radical-LDL oxidation is displayed in *Figure 2*. The results are expressed as percent of PnA oxidized with respect to a control without antioxidant after 20 min. RW is the most efficient in protecting LDL from oxidation, followed by GT and BT. At doses similar to the other beverages, WW failed to show any protective effects. *Figure 3* shows the effect of the different beverages on the oxidation of PnA-loaded LDL oxidized with metmyoglobin/H₂O₂. GT displayed the stronger protective effect, whereas RW and BT exhibited an intermediate grade of protection and WW was ineffective. When results were standardized for phenolics concentration




Figure 1 Effect of red wine on AAPH-induced oxidation of parinaric acid (PnA) incorporated in LDL. PnA oxidation rates in LDL particles in the absence (Line 1) and presence of 0.3 (Line 2), 0.5 (Line 3), and 0.7 (Line 4) μ L of red wine under conditions described in the Methods and Materials section.

(*Table 2*), the inhibitory effect of PRB on AAPH-LDL oxidation was related to the phenolic content (RW > GT > BT > WW). However, when LDL were oxidized with ferril radicals, the association between phenolic content and antioxidant activity was not observed for GT (*Table 2*).




Figure 2 Inhibition of AAPH-induced oxidation of parinaric acid (PnA) incorporated in LDL by black tea (Δ), green tea (\blacklozenge), red wine (\bullet), and white wine (\square). The amount of oxidized PnA was taken 20 min after starting the reaction with AAPH and expressed as % of control in the absence of beverages. The experimental conditions are described in the Methods and Materials section.




Figure 3 Inhibition of metmyoglobin/H₂O₂-induced oxidation of parinaric acid (PnA) incorporated in LDL by black tea (Δ), green tea (◆), red wine (●), and white wine (□). The amount of oxidized PnA was taken 20 min after starting the reaction with AAPH and expressed as % of control in the absence of beverages. The experimental conditions are described in the Methods and Materials section.

In vivo study

The overall antioxidant effect of acute beverage drinking was assessed by measuring plasma TRAP of the individual response throughout the entire period of observation. Table 3 describes the time trend of plasma TRAP levels after the ingestion of the different beverages: TRAP levels started to increase after 30 min and reached the peak at 50 min (*P* < 0.05) for BT. Base-line TRAP values immediately rose for subjects who drank GT (*P* < 0.05 at 30 min) and decreased after 50 min. RW raised the antioxidant capacity after 30 min (*P* < 0.05), peaked at 50 min (*P* < 0.01), and fell back to baseline values after 2 hr, whereas in the WW and control

Table 2 Inhibition of AAPH and metmyoglobin/H₂O₂-induced oxidation of parinaric acid (PnA) incorporated in LDL by the same amounts of alcohol-free red wine (RW), green tea (GT), black tea (BT), and white wine (WW) standardized for phenolic concentration*

	AAPH		Metmyoglobin/H ₂ O ₂	
	QE (µM)	PnA oxidized (%)	QE (µM)	PnA oxidized (%)
RW	3.0 ± 0.24	39.5 ± 0.3	0.6 ± 0.02	56.1 ± 0.6
GT	2.5 ± 0.03	60.4 ± 0.3	0.4 ± 0.02	49.9 ± 0.7
BT	1.1 ± 0.07	71.3 ± 0.2	0.2 ± 0.01	65.0 ± 0.5
WW	0.2 ± 0.01	98.0 ± 0.2	0.05 ± 0.01	99.0 ± 0.5

*Values are mean ± SD, *n* = 3. The same amount of RW, GT, BT, and WW has been used for the experiment with AAPH (0.5 µL) and metmyoglobin/H₂O₂ (0.1 µL). The amount of oxidized PnA was taken 20 min after starting the reaction with H₂O₂ and expressed as % of control in the absence of beverages. QE—Quercetin Equivalents. AAPH—2,2'-azobis(2-amidinopropane) hydrochloride.

groups, there were no changes in plasma TRAP values over the entire period of observation.

Discussion

Tea and wine, widely consumed beverages throughout the world, are extremely rich in phenolic compounds. Tea leaves contain more than 35% of their dry weight in phenolics, the nature of which depending upon the manufacturing procedure.³¹ GT differs from BT in phenolic composition: The former is rich in low molecular weight flavanols (a class of phenolics), the latter is rich in phenolic-condensation products such as theaflavins and thearubigens. Wine also contains flavanols but their chemical profile is different from that of teas.³² RW is rich in anthocyanins and in soluble and hydrolyzable tannins that represent the condensed form of phenolics, whereas WW does not contain anthocyanins. Ample evidence^{33,34} has shown that in vitro antioxidant activity of wine and tea is directly correlated with their phenolic content; however, data from in vivo and ex vivo studies have produced contrasting results.¹⁵⁻²²

In this study, we compared the antioxidant activity of different PRB in vitro with those in vivo through the measure of the total antioxidant capacity of human plasma after acute ingestion of PRB. The in vitro experiments showed that all of the PRB studied, except for WW, have a remarkably high antioxidant capacity and can inhibit LDL oxidation challenged with ferrylmyoglobin or AAPH with different efficiency. The ranking of the in vitro antioxidant efficiency mirrors the phenolic concentration of the beverages: RW with the higher concentration in phenolics displays the strongest antioxidant capacity, WW with the lower phenolic content has also the lower TRAP value. Phenolic concentration also provides a justification for RW higher efficiency in preventing peroxy-radical-induced LDL oxidation when compared to tea. However, RW is not so effective when LDL oxidation is driven by ferrylmyoglobin. Under these conditions, GT showed the best protection. These apparent contradictory results may be explained on the basis of differences in scavenging mechanisms necessary to neutralize AAPH-derived peroxy radicals, or to reduce the strong oxidant ferrylmyoglobin to the harmless metmyoglobin. Both mechanisms avoid LDL oxidation, and phenols are implicated in both. Polymerized phenols in BT and RW may exhibit free hydroxyl groups for peroxy radical scavenging, but more simple flavanols exhibiting a catechol structure and present in higher concentrations in GT may be better reductants of ferrylmyoglobin. The lack of any effect on LDL oxidation for the WW with both catalysts at the amount where the other beverages are effective may also be explained by its low phenolic content. WW begins to have a protective effect at doses 10 times higher (data not shown), suggesting that the phenolic content of the beverages play a crucial role in preserving LDL from oxidation.

The second part of our study was aimed at studying the effects of acute ingestion of BT, GT, RW, and WW by healthy volunteers on the total antioxidant capacity of plasma. Our results showed that both teas and RW contribute to raise plasma TRAP, but with different modalities. The different time of action for the three beverages mirrors their

Table 3 Plasma total antioxidant capacity* before and after drinking water, black tea, green tea, and alcohol-free red and white wine

Time (min)	TRAP				
	Water	Black tea	Green tea	Alcohol-free red wine	Alcohol-free white wine
T0	1318 ± 79	1158 ± 150	1303 ± 160	1151 ± 38	1386 ± 166
T30	1308 ± 68	1317 ± 129	1826 ± 185 ^a	1241 ± 48 ^a	1311 ± 94
T50	1343 ± 89	1761 ± 100 ^b	1570 ± 161 ^b	1395 ± 49 ^b	1267 ± 100
T120	1310 ± 93	1189 ± 131	1488 ± 125	1098 ± 60	1350 ± 68

*Total radical-trapping antioxidant parameter (TRAP) values are expressed as micromoles of peroxy radicals trapped per liter of plasma. Values are mean ± SEM, $n = 5$.

^{a,b}Values with different superscripts are significantly different from T0 value (Fisher PLSD's test): ^a $P < 0.05$, ^b $P < 0.01$.

phenolic composition: GT, richer in simple monomers units give the fastest response (30 min), whereas BT and RW, rich in polymeric units exerted their effect 20 min later. On the other hand, WW, with the lowest phenolics content, failed to show any protective effect in vivo. Again, the most likely explanation for the lack of effect of the WW is that phenolic concentrations in WW are too low to achieve an increase of phenolic plasma levels able to significantly raise antioxidant potential. These results are in agreement with previous evidence showing that WW drinking did not change phenolic levels¹ but increased peroxide plasma levels: when WW was added with 1 g/day of polyphenol powder, phenolic levels significantly increased with a parallel decrease in plasma peroxidation.¹⁶ However, on the basis of the phenolic content, RW would be expected to show the highest antioxidant in vivo activity, but the percent of increase with respect to base-line of TRAP levels was lower (21%) compared to the increase displayed by BT (52%) and GT (40%).

Alternatively, an important role might be played by the interaction between phenolics and other nutrients contained in the beverages; for instance, the interaction between protein and tannins (high molecular weight phenolics) can lead to the formation of insoluble complexes³⁵ that may affect phenolic bioavailability. Although the choice of using alcohol-free wine did not affect the phenolic content and the in vitro antioxidant capacity of the beverage, it may have had consequences on the order of magnitude of the in vivo results. Reduction of the alcohol content of wine has been shown to produce a linear increase in protein-tannin interaction and a parallel decrease in the antioxidant capacity of wine.³⁶ Moreover, Duthie et al.³⁷ have shown that more phenolics were absorbed from whisky than from wine, ascribing these results to the greater alcohol content of whisky. Along these lines, RW impact on plasma TRAP in our experimental conditions could have been higher if we had used non-dealcoholized wine.

On the other hand, BT displays (on the basis of its in vitro antioxidant activity) an unexpected high in vivo response, confirming previous evidence.¹⁹ BT contains phenolic condensation products such as theaflavins and thearubigins with molecular weights of about 500–3000,³¹ which display a lower in vitro antioxidant activity compared to the free phenolics of GT.³⁸ Moreover, Plumb et al.³⁹ have shown that polymerization decreases the ability of phenolics to prevent free radical damage in a lipid system. We

speculate that modifications in the molecular structure of BT polyphenols may have taken place after ingestion, restoring the antioxidant capacity lost as a consequence of the manufacturing process. We think that the condensed polyphenols might be broken down in the stomach by the acid gastric secretion. The monomer phenolics released by gastric hydrolysis would thus become available for absorption and for exerting their antioxidant activity.

The mixture of different phenolics with different red-ox potentials present in tea and wine represent a vast array of antioxidant compounds that might interact in a synergistic way, carrying out a transient protective in vivo effect through mechanisms still unexplored. In light of this experimental evidence, PRB consumption seems to represent a good natural source of compounds involved in the health-promoting effects of a diet rich in fruit and vegetables. However, our results also indicate that phenolic content itself cannot be considered a universal index of in vivo antioxidant activity. In fact, the different doses, combinations of different phenolics, interactions with nutrients, food matrix, and chemical nature of either the phenol or the oxidant must be taken into account when evaluating the in vivo antioxidant capacity of PRB.

In addition, our results give indirect evidence of the metabolic processes that may occur during phenolics absorption, the interactions with nutrients leading to different degrees of absorption, and of compounds with distinct physiological activity that need to be identified.

Acknowledgments

This research was partially supported by the Italian National Research Council short-term fellowship program.

References

- 1 Serafini, M., Maiani, G., and Ferro-Luzzi A. (1998). Alcohol-free red wine enhances plasma antioxidant capacity in humans. *J. Nutr.* **128**, 1003–1007
- 2 Lairon, M. and Amiot, M.J. (1999). Flavonoids in food and natural antioxidants in wine. *Curr. Opin. Lipidol.* **10**, 23–28
- 3 Velioglu, Y.S., Mazza, G., Gao, L., and Oomah, B.D. (1998). Antioxidant activity and total phenolics in selected fruits, vegetables, and grain products. *J. Agric. Food Chem.* **46**, 4113–4117
- 4 Vinson, J.A., Hao, Y., Su, X., and Zubik, L. (1998). Phenol antioxidant quantity and quality in foods: Vegetables. *J. Agric. Food Chem.* **46**, 3630–3634

- 5 Balentine, D.A., Wiseman, S.A., and Bouwens, L.C.M. (1997). The chemistry of tea flavonoids. *Crit. Rev. Food Sci. Nutr.* **37**, 693–704
- 6 Friedman, L.A. and Kimball, A.W. (1986). Coronary heart disease mortality and alcohol consumption in Framingham. *Am. J. Epidemiol.* **124**, 481–489
- 7 Knekt, P., Jarvinen, R., Reunanen, A., and Maatela, J. (1996). Flavonoid intake and coronary mortality in Finland: A cohort study. *Br. Med. J.* **312**, 478–481
- 8 Renaud, S. and De Lorgeril, M. (1992). Wine, alcohol, platelets and French paradox for coronary heart disease. *Lancet* **339**, 1523–1526
- 9 Frankel, E.N., Kanner, J., German, J.B., Parks, E., and Kinsella, J.E. (1993). Inhibition of oxidation of human low-density lipoprotein by phenolic substances in red wine. *Lancet* **341**, 454–457
- 10 Kanner, J., Frankel, E., Granit, R., German, B., and Kinsella, J.E. (1994). Natural antioxidants in grapes and wines. *J. Agric. Food Chem.* **42**, 64–69
- 11 Frankel, E.N., Waterhouse, A.L., and Teissedre, P.L. (1995). Principal phenolic phytochemicals in selected California wines and their antioxidant activity in inhibiting oxidation of human low-density lipoproteins. *J. Agric. Food Chem.* **43**, 1165–1169
- 12 Yen, G. and Chen, H. (1995). Antioxidant activity of various tea extracts in relation to their antimutagenicity. *J. Agric. Food Chem.* **43**, 27–32
- 13 Zhenhua, D., Yuan, C., Mei, Z., and Yunzhong, F. (1991). Inhibitory effect of China green tea polyphenol on the oxidative modification of low-density lipoprotein by macrophages. *Med. Sci. Res.* **19**, 767–768
- 14 Terao, J., Piskula, M., and Yao, Q. (1994). Protective effect of epicatechin, epicatechin gallate and quercetin on lipid peroxidation in phospholipid bilayers. *Arch. Biochem. Biophys.* **308**, 278–284
- 15 Cherubini, A., Beal, M.F., and Frei, B. (1999). Black tea increases the resistance of human plasma to lipid peroxidation in vitro, but not ex vivo. *Free Rad. Biol. Med.* **27**, 381–387
- 16 Nigdikar, S.V., Williams, N.R., Griffin, B.A., and Howard, A.N. (1998). Consumption of red wine polyphenols reduces the susceptibility of low-density lipoproteins to oxidation in vivo. *Am. J. Clin. Nutr.* **68**, 258–265
- 17 Whitehead, T.P., Robinson, D., Allaway, S., Syms, J., and Hale, A. (1995). Effect of red wine ingestion on the antioxidant capacity of serum. *Clin. Chem.* **41**, 32–35
- 18 Furrman, B., Lavy, A., and Aviram, M. (1995). Consumption of red wine with meals reduces the susceptibility of human plasma and low-density lipoprotein to lipid peroxidation. *Am. J. Clin. Nutr.* **61**, 549–554
- 19 Serafini, M., Ghiselli, A., and Ferro-Luzzi, A. (1996). In vivo antioxidant effect of green and black tea in man. *Eur. J. Clin. Nutr.* **50**, 28–32
- 20 McAnlis, G.T., McEneny, J., Pearce, J., and Young, I.S. (1998). Black tea consumption does not protect low density lipoprotein from oxidative modification. *Eur. J. Clin. Nutr.* **52**, 202–206
- 21 Maxwell, S. and Thorpe, G. (1996). Tea flavonoids have little short term impact on serum antioxidant activity. *Br. Med. J.* **313**, 229
- 22 de Rijke, Y.B., Demacker, N.M.P., Assen, N.A., Sloots, L.M., Katan, M.B., and Stalenhof, A.F.H. (1996). Red wine consumption does not affect oxidizability of low-density lipoproteins in volunteers. *Am. J. Clin. Nutr.* **63**, 329–334
- 23 Swain, T. and Hillis, E. (1959). The phenolic constituents of *Prunus Domestica*. The quantitative analysis of phenolic constituents. *J. Sci. Food Agric.* **10**, 63–68
- 24 Ghiselli, A., Serafini, M., Maiani, G., Azzini, E., and Ferro-Luzzi, A. (1995). A fluorescence based method for measuring total plasma antioxidant capability. *Free Rad. Biol. Med.* **18**, 29–36
- 25 Vieira, O., Laranjinha, J., Madeira, V., and Almeida, L. (1996). Rapid isolation of low density lipoproteins in a concentrated fraction free from water-soluble plasma antioxidants. *J. Lipid Res.* **37**, 2715–2721
- 26 Laranjinha, J., Almeida, V., and Madeira, V. (1994). Reactivity of dietary phenolic acids with peroxy radicals: Antioxidant activity upon low-density lipoprotein oxidation. *Biochem. Pharmacol.* **48**, 487–494
- 27 Laranjinha, J.A.N., Almeida, L.M., and Madeira, V.M.C. (1992). Lipid peroxidation and its inhibition in low-density lipoproteins: Quenching of cis-parinaric acid fluorescence. *Arch. Biochem. Biophys.* **297**, 147–155
- 28 Sato, K., Niki, E., and Shimasaki, H. (1990). Free radical-mediated chain oxidation of low density lipoprotein and its synergistic inhibition by vitamin E and vitamin C. *Arch. Biochem. Biophys.* **279**, 402–405
- 29 George, P. and Irvine, D.H. (1952). The reaction between metmyoglobin and hydrogen peroxide. *Biochem. J.* **52**, 511–517
- 30 Giulivi, C. and Cadenas, E. (1994). Ferrylmyoglobin: Formation and chemical reactivity toward electron-donating compounds. *Methods Enzymol.* **233**, 189–202
- 31 Balentine, A.D. (1992). Manufacturing and chemistry of tea. In *Phenolic Compounds in Food and Their Effect on Health I* (H. Chi-Tang, Y.L. Chang, and H. Mou-Tuan, eds.), pp. 103–117, ACS, Washington, DC, USA
- 32 Macheix, J.J., Fleurient, A., and Billot, J. (1990). *Fruit Phenolics*. CRC Press, Boca Raton, FL, USA
- 33 Vinson, J.A. and Hontz, B.A. (1995). Phenol antioxidant index: Comparative antioxidant effectiveness of red and white wines. *J. Agric. Food Chem.* **43**, 401–403
- 34 Yu-Li, L., I-Ming, J., Ying-Ling, C., Yu-Chih, L., and Jen-Kun, L. (1996). Composition of polyphenols in fresh tea leaves and associations of their oxygen-radical-absorbing capacity with antiproliferative actions in fibroblast cells. *J. Agric. Food Chem.* **44**, 1387–1394
- 35 Mehansho, H., Butler, L.G., and Carlson, D.M. (1987). Dietary tannins and salivary proline-rich proteins: Interactions, induction and defense mechanisms. *Annu. Rev. Nutr.* **7**, 423–440
- 36 Serafini, M., Maiani, G., and Ferro-Luzzi, A. (1997). Effect of ethanol on red wine tannin-protein (BSA) interactions. *J. Agric. Food Chem.* **45**, 3148–3151
- 37 Duthie, G.G., Pedersen, M.W., Gardner, P.T., Morrice, P.C., Jenkinson, A.M., McPhail, D.B., and Steele, G.M. (1998). The effect of whisky and wine consumption on total phenol content and antioxidant capacity of plasma from healthy volunteers. *Eur. J. Clin. Nutr.* **52**, 733–736
- 38 Gow-Chin, Y. and Hui-Yin, C. (1995). Antioxidant activity of various tea extracts in relation to their antimutagenicity. *J. Agric. Food Chem.* **43**, 27–32
- 39 Plumb, G.W., Pascual-Teresa, S., Santos-Buelga, C., Cheynier, V., and Williamson, G. (1998). Antioxidant properties of catechins and proanthocyanidins: Effect of polymerisation, galloylation and glycosylation. *Free Rad. Res.* **29**, 351–358