



# Inhibition of PI3K $\delta$ Enhances Poly I:C-Induced Antiviral Responses and Inhibits Replication of Human Metapneumovirus in Murine Lungs and Human Bronchial Epithelial Cells

Akitaka Fujita<sup>1†</sup>, Keiko Kan-o<sup>1,2\*†</sup>, Ken Tonai<sup>1,3</sup>, Norio Yamamoto<sup>1</sup>, Tomohiro Ogawa<sup>1</sup>, Satoru Fukuyama<sup>1</sup>, Yoichi Nakanishi<sup>1</sup> and Koichiro Matsumoto<sup>1</sup>

<sup>1</sup> Research Institute for Diseases of the Chest, Graduate School of Medical Sciences, Kyushu University, Fukuoka, Japan, <sup>2</sup> Department of Endoscopic Diagnostics and Therapeutics, Kyushu University Hospital, Fukuoka, Japan, <sup>3</sup> Division of Intensive Care, Department of Anesthesiology and Intensive Care Medicine, Jichi Medical University School of Medicine, Tochigi, Japan

## OPEN ACCESS

### Edited by:

Guochang Hu,  
University of Illinois at Chicago,  
United States

### Reviewed by:

Patrick Geraghty,  
SUNY Downstate Medical Center,  
United States  
Yang Jin,  
Boston University, United States

### \*Correspondence:

Keiko Kan-o  
hanamura@kokuu.med.kyushu-u.ac.jp

<sup>†</sup>These authors have contributed  
equally to this work

### Specialty section:

This article was submitted to  
Inflammation,  
a section of the journal  
Frontiers in Immunology

**Received:** 17 October 2019

**Accepted:** 25 February 2020

**Published:** 11 March 2020

### Citation:

Fujita A, Kan-o K, Tonai K,  
Yamamoto N, Ogawa T, Fukuyama S,  
Nakanishi Y and Matsumoto K (2020)  
Inhibition of PI3K $\delta$  Enhances Poly  
I:C-Induced Antiviral Responses and  
Inhibits Replication of Human  
Metapneumovirus in Murine Lungs  
and Human Bronchial Epithelial Cells.  
*Front. Immunol.* 11:432.  
doi: 10.3389/fimmu.2020.00432

Viral infections of the airway can exacerbate respiratory diseases, such as asthma or chronic obstructive pulmonary disease (COPD), and accelerate disease progression. Phosphoinositide 3-kinase (PI3K) $\delta$ , a class 1A PI3K, has been studied as a potential target for achieving anti-oncogenic and anti-inflammatory effects. However, the role of PI3K $\delta$  in antiviral responses is poorly understood. Using a synthetic double-stranded RNA poly I:C and a selective PI3K $\delta$  inhibitor IC87114, we investigated the role of PI3K $\delta$  signaling in poly I:C-induced expression of the T lymphocyte-inhibitory molecule programmed death 1 ligand 1 (PD-L1), inflammatory responses and antiviral interferon (IFN) responses. C57BL/6N mice were treated with IC87114 or vehicle by intratracheal (i.t.) instillation followed by i.t. administration of poly I:C. Poly I:C increased PD-L1 expression on epithelial cells, lymphocytes, macrophages, and neutrophils in the lungs and IC87114 suppressed poly I:C-induced PD-L1 expression on epithelial cells and neutrophils possibly via inhibition of the Akt/mTOR signaling pathway. IC87114 also attenuated poly I:C-induced increases in numbers of total cells, macrophages, neutrophils and lymphocytes, as well as levels of KC, IL-6 and MIP-1 $\beta$  in bronchoalveolar lavage fluid. Gene expression of IFN $\beta$ , IFN $\lambda_2$  and IFN-stimulated genes (ISGs) were upregulated in response to poly I:C and a further increase in gene expression was observed following IC87114 treatment. In addition, IC87114 enhanced poly I:C-induced phosphorylation of IRF3. We assessed the effects of IC87114 on human primary bronchial epithelial cells (PBECs). IC87114 decreased poly I:C-induced PD-L1 expression on PBECs and secretion of IL-6 and IL-8 into culture supernatants. IC87114 further enhanced poly I:C-induced increases in the concentrations of IFN $\beta$  and IFN $\lambda_{1/3}$  in culture supernatants as well as upregulated gene expression of ISGs in PBECs. Similar results were obtained in PBECs transfected with siRNA targeting the PI3KCD gene encoding PI3K p110 $\delta$ , and stimulated with poly I:C. In human metapneumovirus (hMPV) infection of PBECs, IC87114 suppressed hMPV-induced PD-L1 expression and

reduced viral replication without changing the production levels of IFN $\beta$  and IFN $\lambda_{1/3}$  in culture supernatants. These data suggest that IC87114 may promote virus elimination and clearance through PD-L1 downregulation and enhanced antiviral IFN responses, preventing prolonged lung inflammation, which exacerbates asthma and COPD.

**Keywords: poly I:C, phosphoinositide 3-kinase  $\delta$ , IC87114, programmed death 1 ligand 1, interferon, bronchial epithelial cells, human metapneumovirus**

## INTRODUCTION

Asthma and chronic obstructive pulmonary disease (COPD) are chronic respiratory diseases that are major causes of morbidity and mortality worldwide (1, 2). Airway infection by viruses including rhinovirus, respiratory syncytial virus, human metapneumovirus, and influenza virus is one of the leading causes of asthma and COPD exacerbation, which can lead to heightened inflammation and a general decline in health status (3, 4). These viruses possess single-stranded RNA genomes and generate double-stranded RNA (dsRNA) in host cells during replication (excluding influenza virus which has a unique 5'-triphosphate dsRNA panhandle structure). These dsRNAs, known as pathogen-associated molecular patterns, bind to Toll-like receptor (TLR) 3 in endosomes of antigen-presenting cells including bronchial epithelial cells, as well as to cytosolic RNA helicase retinoic acid-inducible gene-1 or melanoma differentiation-associated gene 5 (MDA5) to subsequently activate antiviral immune responses (5–7). A synthetic dsRNA analog, polyinosinic-polycytidylic acid (poly I:C), is recognized by TLR3 and MDA5 and subsequent signaling leads to activation of transcription factors including interferon regulatory factor-3 (IRF3) and nuclear factor kappa-light-chain-enhancer of activated B cells (NF- $\kappa$ B), resulting in secretion of type I and III interferons (IFNs) and proinflammatory cytokines and chemokines, respectively (8–11). Type I IFNs stimulate neighboring cells to express IFN-regulated genes (IRGs), inducing an antiviral state (12).

Adaptive antiviral immunity is initiated by interactions between antigen-presenting cells and T cells. These interactions include binding of an array of co-stimulatory or co-inhibitory molecules including B7 family members and their putative receptors (13). Programmed cell death 1 ligand 1 (PD-L1) is a B7-family co-inhibitory molecule, which is constitutively expressed on antigen presenting cells. Ligation of programmed cell death 1 (PD-1) by PD-L1 on activated T cells causes T cell impairment and exhaustion (14, 15). Some investigators reported that inhibition of the PD-1/PD-L1 pathway restored virus-specific CD8<sup>+</sup> T cell activity and reduced viral loads in murine models of acute or chronic viral infection, suggesting that PD-L1 upregulation on infected cells is responsible for CD8<sup>+</sup> T cell exhaustion and failure to achieve viral clearance (16–19). We have previously shown that poly I:C stimulation enhanced the expression of PD-L1 on human and murine bronchial epithelial cells; these cells are targeted by respiratory viruses for their replication (20–22). Given the high expression of PD-1 on virus-specific CD8<sup>+</sup> T cells, dsRNA-induced PD-L1 expression on

infected cells may prevent CD8<sup>+</sup> T cells from eliminating these cells and thereby promote the spread of virus to neighboring cells.

Phosphoinositide 3-kinases (PI3Ks) can be divided into three distinct classes (I, II, and III) according to their structures and substrate specificities. PI3Ks affect important signaling pathways via several downstream molecules such as Akt, 3'-phosphoinositide dependent kinase 1, the mechanistic target of rapamycin (mTOR), and glycogen synthase kinase 3 which control biological functions including cell proliferation, differentiation, migration, and metabolism (23, 24). There are three isoforms of class IA PI3Ks in mammalian cells:  $\alpha$ ,  $\beta$ , and  $\delta$ . Each class IA PI3K is composed of a catalytic subunit (p110 $\alpha$ , p110 $\beta$ , or p110 $\delta$ ) and a SH2 domain-containing regulatory subunit (p85 $\alpha$ , p55 $\alpha$ , p50 $\alpha$ , p85 $\beta$ , and p55 $\gamma$ ) (25). The catalytic subunits p110 $\alpha$  and p110 $\beta$  are broadly expressed, while p110 $\delta$  is predominantly expressed in cells of the myeloid and lymphoid lineages. Recently, a primary immunodeficiency caused by mutations in the *PIK3CD* or *PIK3R1* genes (encoding p110 $\delta$  and p85 $\alpha$ , respectively) was reported and called activated PI3K delta syndrome (APDS) (26–28). The *PIK3CD* or *PIK3R1* mutations increased PI3K $\delta$  activities and patients with APDS suffered from infectious complications such as recurrent bacterial respiratory infections and severe or persistent infections by herpesviruses, including Epstein-Barr virus, cytomegalovirus, and varicella-zoster virus (29). Although APDS phenotypes suggest that high PI3K $\delta$  activities aggravate viral infection, the roles of PI3K $\delta$  signaling in modulating co-inhibitory molecule (e.g., PD-L1) expression and antiviral IFN responses are poorly understood.

Human metapneumovirus (hMPV) discovered in 2001 has been identified as the secondary cause of acute lower respiratory infection in children, and moreover recently implicated in exacerbations of asthma and COPD (30, 31). There is currently no licensed vaccine or therapy available for hMPV. Previous several mouse studies have shown that hMPV is able to persist in respiratory infected cells by inhibiting innate immune responses and causing CD8<sup>+</sup> T cells impairment mediated by PD-1 (32, 33). While animal and human *in vivo* studies confirmed that airway epithelium can sustain hMPV infection and replication (34, 35), there have been limited studies describing hMPV infection of primary culture bronchial epithelial cells.

In this study, we evaluated the effects of the PI3K $\delta$  inhibitor IC87114 on PD-L1 expression, inflammatory responses, and antiviral IFN responses in lung cells following intratracheal (i.t.) administration of poly I:C in mice. We also assessed the effects of IC87114 on human primary bronchial epithelial cells (PBEs) stimulated by poly I:C or infected with hMPV.

## MATERIALS AND METHODS

### Animals

Male 8-week-old C57BL/6N mice were purchased from Japan SLC, Inc. (Shizuoka, Japan). Mice were housed under specific pathogen-free conditions. The study protocol was approved by the Kyushu University Animal Care and Use Committee (A19-021-1).

### Intratracheal Administration of Poly I:C

Mice were anesthetized intraperitoneally with 80 mg/kg ketamine and 16 mg/kg xylazine and their tracheas were cannulated using 22-gauge catheters. IC87114 (1 mg/kg body weight in 25  $\mu$ L total volume; BioVision, Milpitas, CA) or vehicle was administered by i.t. instillation followed by i.t. administration of poly I:C (3  $\mu$ g in 30  $\mu$ L total volume; Innaxon, Tewkesbury, United Kingdom) or vehicle. Mice were sacrificed 24 h following i.t. administration and lungs were removed following bronchoalveolar lavage. IC87114 was diluted to 1 mg/mL with 0.1% dimethyl sulfoxide. Poly I:C was diluted with 0.9% NaCl. Sham-treated mice were used as controls.

### Flow Cytometry

Lungs were minced and single-cell suspensions were prepared. Cells were suspended in 100  $\mu$ L of phosphate-buffered saline (PBS) containing 0.5% bovine serum albumin (BSA) and preincubated with anti-mouse CD16/CD32 antibody (BD Biosciences, San Jose, CA) for 15 min to prevent non-specific binding via the Fc $\gamma$  receptor. The cells were washed and suspended in 100  $\mu$ L of PBS containing 0.5% BSA and the following antibodies for 30 min: phycoerythrin (PE) anti-mouse CD326 (epithelial cell adhesion molecule; Ep-CAM) (BioLegend, San Diego, CA), peridinin-chlorophyll-protein (PerCP)/Cy5.5 anti-mouse CD45 (BioLegend), fluorescein isothiocyanate (FITC) anti-mouse CD11b (eBioscience, San Diego, CA), FITC anti-mouse CD11c (eBioscience), PE/Cy7 anti-mouse CD274 (PD-L1) (eBioscience), and allophycocyanin (APC)/Cy7 anti-mouse CD279 (PD-1) (BioLegend). Isotype controls were included as appropriate to facilitate gating of each population. Living cells were analyzed by addition of propidium iodide (PI) to samples in the first experiment and via the forward scatter (FSC) threshold in subsequent experiments. Cells were thoroughly washed and analyzed using a BD FACSVerser flow cytometer with FACSsuite software (Becton Dickinson, Franklin Lakes, NJ). One hundred thousand events were acquired in list mode. Epithelial cells, lymphocytes, macrophages, and neutrophils were identified as CD11b/CD11c<sup>low</sup> CD45<sup>low</sup> CD326 (Ep-CAM)<sup>high</sup>, CD45<sup>high</sup> CD11b/CD11c<sup>low</sup>, CD45<sup>high</sup> CD11b/CD11c<sup>high</sup> side scatter (SSC)-A<sup>low</sup>, and CD45<sup>high</sup> CD11b/CD11c<sup>high</sup> SSC-A<sup>high</sup>, respectively. In some experiments, cells were suspended in PBS containing 0.5% BSA with both APC anti-mouse CD3 (BioLegend) and PE anti-mouse CD19 (BioLegend) antibodies, or both APC anti-mouse Ly-6C (BD Biosciences) and PE anti-mouse Ly-6G (BioLegend) antibodies instead of PE anti-mouse CD326, PE/Cy7 anti-mouse CD274, and APC/Cy7 anti-mouse CD279 antibodies in order to validate gating strategy for lymphocytes, neutrophils and macrophages.

In some experiments, cells were sorted using a BD FACSAria SORP (Becton Dickinson) and centrifuged at 72 $\times$  g for 5 min. Pelleted cells were stained with Diff-Quick and observed using an optical microscope (Nikon Eclipse Ni-U, Nikon Corporation, Tokyo, Japan).

For *in vitro* experiments, PBECs or BEAS-2B were cultured to semi-confluence in 12-well plates. Cells were incubated in 100  $\mu$ L of PBS containing 0.5% BSA and PE anti-human PD-L1 (clone: MIH1; Invitrogen, Carlsbad, CA) at room temperature for 30 min. The cells were thoroughly washed and analyzed using a BD FACSVerser flow cytometer with FACSsuite software (Becton Dickinson). Ten thousand events were acquired in list mode with debris excluded by the FSC threshold.

For the analysis of cell viability, resuspended lung cells or PBECs were incubated with PI and non-viable cells stained with PI were counted in a flow cytometer with debris excluded by the FSC threshold.

### Culture and Treatment of Primary Bronchial Epithelial Cells (PBECs) and BEAS-2B

PBECs were obtained during routine fiberoptic bronchoscopy from never-smoker patients with normal lung function and pulmonary nodules. Bronchial brushings were performed from healthy lobes without a pulmonary nodule. All bronchial brushings were obtained from the same anatomical region (bronchial generations 4–7). The study protocol was approved by the Kyushu University Institutional Review Board for Clinical Research (29-170). All subjects provided written informed consent in accordance with the principles laid out in the Declaration of Helsinki.

Cells were cultured at 37°C/5% CO<sub>2</sub> under submerged conditions on flasks coated with collagen (Cell Applications, Inc. San Diego, CA) in supplemented bronchial epithelial growth medium (BEGM; Lonza, Basel, Swiss) and used within four passages. PBECs (~80% confluency) were pretreated with 10  $\mu$ M IC87114 or vehicle for 1 h, then stimulated by addition of 1  $\mu$ g/mL poly I:C to the culture medium. In virus infection experiments, 10  $\mu$ M IC87114 was added to cells for 1 h prior to hMPV infection, and then following infection. PBECs were cultured in BEGM without hydrocortisone for at least 24 h prior to stimulation or infection.

In some experiments, human bronchial epithelial cells, BEAS-2B, were cultured in DMEM/F12 (gibco, Thermo Fisher Scientific, San Diego, CA) with 10% FBS and 1% penicillin-streptomycin and incubated at 37°C/5% CO<sub>2</sub>. When cells reached semi-confluence, they were pretreated with 10 or 100 nM rapamycin (LC Laboratories, Woburn, MA) or vehicle for 1 h, then stimulated by addition of 3  $\mu$ g/mL poly I:C to the culture medium.

### Gene Expression in Mouse Lungs and Human Bronchial Epithelial Cells

Total RNA was isolated from mouse lungs, PBECs or BEAS-2B using TRIzol<sup>®</sup> Reagent (Thermo Fisher Scientific) or TRI Reagent<sup>®</sup> (Molecular Research Center, Inc, Cincinnati, OH). Reverse transcription was performed using Multiscribe Reverse

Transcriptase (Invitrogen). Real-time quantitative reverse-transcriptase PCR analyses were performed using SYBR Premix Ex Taq II (Takara, Shiga, Japan) and a Thermal Cycler Dice Real Time System II (Takara). Target gene expression levels were normalized to expression of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) or 18S rRNA. Primer sequences are provided in **Supplementary Table 1**.

## Western Blotting

Proteins were extracted from mouse lungs using TRIzol<sup>®</sup> Reagent (Thermo Fisher Scientific) according to the manufacturer's instructions. For *in vitro* experiments, PBECs cultured to semi-confluence in 6-well plates were lysed using Pierce<sup>®</sup> RIPA Buffer (Thermo Fisher Scientific) according to the manufacturer's instructions. Ten micrograms of the protein sample were denatured, separated by SDS-PAGE, and transferred to polyvinylidene difluoride membranes. Membranes were blocked with TBS-Tween (10 mM Tris, 150 mM NaCl, 0.05% Tween 20, pH 8.0) containing 5% skimmed milk for 1 h at room temperature. After blocking, blotting was performed with anti-I $\kappa$ B $\alpha$  (Cell signaling Technology, Beverly, MA), anti-phospho-I $\kappa$ B $\alpha$  (Ser32, Cell signaling Technology), anti-v-akt murine thymoma viral oncogene homolog (Akt) (Cell signaling Technology), anti-phospho-Akt (Ser473, Cell signaling Technology), anti-IRF3 (Cell signaling Technology), anti-phospho-IRF3 (Ser379, Cell signaling Technology), anti-PI3 Kinase p110 $\delta$  (Cell Signaling Technology), or anti- $\beta$ -actin (Santa Cruz Biotechnology, Dallas, TX) antibodies at 4°C overnight. Membranes were washed three times with TBS-Tween and then incubated with a horseradish peroxidase-conjugated secondary antibody for 60 min at room temperature. Membranes were washed three times with TBS-Tween, and specific bands were visualized using ImmunoStar<sup>®</sup> LD (Wako, Osaka, Japan) according to the manufacturer's instructions. All blots were imaged using the ChemiDoc<sup>™</sup> XRS+ system (Bio-Rad Laboratories, Inc. Hercules, CA). Densitometric analysis of band intensities was performed using Image J.

## Collection of Bronchoalveolar Lavage Fluid (BALF)

Mice were given lethal doses of pentobarbital and their lungs were gently lavaged once with 1 mL of 0.9% saline via the tracheal cannula. Total and differential cell counts in BALF were performed as described previously (36).

## Detection of Cytokines and Chemokines

BALF was centrifuged at 250  $\times$  g for 10 min, and the supernatants were stored at -80°C. Levels of mouse keratinocyte-derived chemokine (KC; detection limit 15.6–1,000 pg/mL), interleukin (IL)-6 (detection limit 7.8–500 pg/mL) and macrophage inflammatory protein-1 $\beta$  (MIP-1 $\beta$ ; detection limit 7.8–500 pg/mL) in BALF were measured by enzyme-linked immunosorbent assay (ELISA) (R&D systems, Inc. Minneapolis, MN). Levels of human IL-6 (detection limit 9.38–600 pg/mL), IL-8 (detection limit 31.2–2,000 pg/mL), IFN- $\beta$  (detection limit 7.81–500 pg/mL), and IFN- $\lambda_{1/3}$  (detection limit 62.5–4,000 pg/mL) in cell culture supernatants were determined using

ELISA (R&D systems, Inc.) Cell supernatants were diluted as appropriate for measurement.

## Preparation and Transfection of Small Interfering RNA (siRNA)

PBECs cultured in 6-well plates (60–80% confluence) were transiently transfected with 10 nM PIK3CD siRNA (Silencer<sup>®</sup> Select Validated siRNA, s10529; Ambion, Life Technologies, Carlsbad, CA) or 10 nM negative control (NC) siRNA (Silencer<sup>®</sup> Select Negative Control siRNA, 4390843; Ambion, Life Technologies) using Lipofectamine<sup>®</sup> RNAiMAX Reagent (Invitrogen) according to the manufacturer's instructions. Lipofectamine<sup>®</sup> RNAiMAX Reagent or siRNA were diluted in BEGM without hydrocortisone and antibiotics. Cells were used for experiments at 48 h after transfection.

## Virus Infection of Cultured Cells

The CAN97-83 strain of hMPV was used in most experiments, propagated in Vero E6 cells (CRL-1586; ATCC, Manassas, VA) and virus titer was determined by TCID<sub>50</sub> assay. In brief, virus was prepared in serum free MEM (gibco) containing 5  $\mu$ g/ml trypsin (gibco) by infecting cells in a 10-cm culture dish at a multiplicity of infection (MOI) of 0.05. After absorption for 2 h at 37°C/5%CO<sub>2</sub>, serum free MEM containing 5  $\mu$ g/ml trypsin was added. Serum free medium containing 5  $\mu$ g/ml trypsin was changed every other day during incubation and cells were incubated until 70–90% cytopathic effect was apparent, usually within 7 days. Cells and culture supernatant were collected and virus was obtained by freeze-thawing the cells. This strain of hMPV has been used in other studies (31) and we did not attempt further purification since this may change key properties of the virus (37). Virus was quantified by immunostaining assay. Briefly, semi-confluent Vero E6 monolayers in a 24 well plate were infected with 200  $\mu$ l of 10-fold serial dilutions. Five  $\mu$ g/ml trypsin was added in serum free MEM during infection. After absorption for 2 h at 37°C/5%CO<sub>2</sub>, 200  $\mu$ l of MEM containing 5% FBS was added and then cells were incubated. After 6 days, infected wells were identified by using mouse anti-hMPV monoclonal antibody (MAB8510; EMD Millipore, Temecula, CA) followed by a horseradish peroxidase-labeled goat anti-mouse secondary antibody (abcam, Cambridge, UK). Cells were visualized using DAB peroxidase substrate (Vector Laboratories, Inc. Burlingame, CA) and the TCID<sub>50</sub>/ml was determined using standard methods. The CAN97-83 strain of recombinant hMPV with green fluorescent protein (hMPV-GFP) was purchased from ViraTree (<http://www.viratree.com>). Control experiments using UV-inactivated hMPV or hMPV-GFP were included to confirm that responses were a result of hMPV infection and not from factors remaining in the culture media following hMPV propagation. To UV-irradiate virus, hMPV was exposed to a short-wavelength (254 nm) UV lamp at a distance of 5 cm for 15 min.

PBECs (~80% confluency) were washed with PBS and infected with hMPV or hMPV-GFP in BEGM without hydrocortisone for 1 h at a MOI of 0.1. Cell monolayers were washed, then incubated in BEGM without hydrocortisone for 24–72 h. Cells infected with hMPV-GFP were observed using fluorescence microscopy (BZ-X800, KEYENCE, Osaka, Japan).

The number of cells infected with hMPV-GFP were counted in four random microscopic fields (4 $\times$  objective lens) per well and the mean value was calculated.

Standard biosecurity and institutional safety procedures were adhered to during this study.

## Statistical Analyses

Unless otherwise stated, data were expressed as means  $\pm$  standard deviations (SDs). The Mann-Whitney U-test was used for comparisons between two groups. Comparisons of three or more groups were conducted using one-way analysis of variance (ANOVA) or two-way ANOVA followed by Tukey's multiple comparisons test. Correlations were examined using Spearman correlation. All statistical analyses were conducted using GraphPad Prism 8 software (GraphPad Software, San Francisco, CA). Differences were considered statistically significant at  $p < 0.05$ .

## RESULTS

### Poly I:C Upregulates PD-L1 Expression on Lung Cells in Mice

We have previously showed that PD-L1 on the epithelium in mouse lung was significantly upregulated 24 and 72 h following i.t. poly I:C administration, while numbers of neutrophils in the BALF increased 24 h and then diminished promptly 48 h following i.t. poly I:C administration (22). Therefore, we evaluated PD-L1 expression and enumerated cells in BALF 24 h following poly I:C administration.

We measured PD-L1 and PD-1 expression on epithelium, lymphocytes, macrophages, and neutrophils in the lungs of poly I:C-treated and untreated mice using the flow cytometry gating strategy outline in **Figure 1A**. To validate our method, lung cells were sorted and examined by microscopy, or further classified using specific antibodies (**Figure 1A**). Approximately 90% of identified lymphocytes were positive for only CD3 or CD19 and more than 90% of identified neutrophils were positive for Ly-6G. More than 90% of identified macrophages were negative for Ly-6G and the Ly-6G<sup>low</sup>/Ly-6C<sup>high</sup> population (inflammatory macrophages) was increased in poly I:C-treated group (**Figure 1A**). Under unstimulated conditions, PD-L1 was consistently expressed on epithelium, macrophages, and neutrophils, but not on lymphocytes (**Figure 1B**, histograms). Administration of poly I:C i.t. significantly increased the expression of PD-L1 on lung cells at 24 h post-stimulation (**Figure 1B**). By contrast, PD-1 was consistently expressed on lymphocytes and poly I:C stimulation had no effect on its expression (**Figure 1C**).

### A PI3K $\delta$ Inhibitor Attenuates Poly I:C-Induced PD-L1 Expression on Epithelium and Neutrophils in Mouse Lungs

We previously reported that poly I:C increased the expression of PD-L1 on BEAS-2B human bronchial epithelial cells


and that the selective PI3K $\delta$  inhibitor IC87114 partially suppressed poly I:C-induced PD-L1 expression (21, 38). Therefore, we investigated whether IC87114 attenuated poly I:C-induced PD-L1 expression *in vivo*. Treatment with IC87114 alone did not affect PD-L1 expression on mouse lung cells or PD-1 expression on lymphocytes (**Figures 2A,B**). Twenty-four hours following poly I:C administration, PD-L1 expression was significantly upregulated on epithelium, lymphocytes, macrophages, and neutrophils in mouse lungs (**Figure 2C**). Treatment with IC87114 significantly attenuated poly I:C-induced upregulation of PD-L1 on epithelium and neutrophils, but not its upregulation on lymphocytes and macrophages (**Figure 2C**). Cell viability was evaluated by PI exclusion assay 24 h following poly I:C administration. Treatment with IC87114 alone, poly I:C alone, or IC87114 plus poly I:C did not affect cell viability (**Figure 2D** and **Supplementary Figure 1A**).

### A PI3K $\delta$ Inhibitor Suppresses Poly I:C-Induced PD-L1 Expression via Inhibition of the Akt/mTOR Signaling Pathway


In our previous studies using BEAS-2B, the selective PI3K $\delta$  inhibitor IC87114 suppressed poly I:C-induced PD-L1 expression without affecting the phosphorylation of I $\kappa$ B $\alpha$  although the NF- $\kappa$ B pathway played an essential role in poly I:C-induced upregulation of PD-L1 (21, 38). Further experiments were conducted to explore the inhibitory mechanism of PD-L1 by IC87114 *in vitro* and *in vivo*. Poly I:C increased PD-L1 gene expression in human bronchial epithelial cells at 6 h following poly I:C stimulation and combined treatment with IC87114 and poly I:C did not alter the levels of PD-L1 gene expression, suggesting that PI3K $\delta$  provides translational control of PD-L1 (**Figure 3A**). Lastwika et al. reported that activation of the PI3K/Akt/mTOR pathway regulates translationally PD-L1 expression in bronchial epithelial cells (39). Similarly in our study, treatment with the mTOR inhibitor rapamycin suppressed poly I:C-induced PD-L1 on human bronchial epithelial cells 24 h following poly I:C stimulation (**Figure 3B**). *In vivo*, an hour following poly I:C administration, I $\kappa$ B $\alpha$  and Akt was phosphorylated in poly I:C-treated mouse lungs (**Figure 3C**). Treatment with IC87114 significantly suppressed Akt phosphorylation by poly I:C, but not I $\kappa$ B $\alpha$  phosphorylation (**Figure 3C**). These results indicate that PI3K $\delta$  may contribute translational induction of PD-L1 expression via the Akt/mTOR pathway in poly I:C-stimulated bronchial epithelial cells.

### A PI3K $\delta$ Inhibitor Suppresses Poly I:C-Induced Influx of Inflammatory Cells and Production of Pro-Inflammatory Chemokines and Cytokines in BALF


Next, we analyzed the effects of IC87114 on inflammatory cell infiltration and production of pro-inflammatory chemokines and cytokines in BALF. Total cell numbers in BALF were increased nearly 3-fold compared with controls



**FIGURE 1** | Poly I:C-induced upregulation of PD-L1 on mouse lung cells. **(A)** Flow cytometry gating strategy to distinguish epithelial cells, lymphocytes, macrophages, and neutrophils in mouse lungs. Sorted and pelleted cells were stained with Diff-Quick and observed using an optical microscope. Scale bar, 20  $\mu$ m. **(B,C)** Poly I:C or vehicle was administered intratracheally to mice and PD-L1 **(B)** or PD-1 **(C)** expression on lung cells was analyzed 24 h following administration using flow cytometry. Representative histograms are shown. SSC, side scatter; FSC, forward scatter; MFI, mean fluorescence intensity. All results are representative of at least three independent experiments. Data represent means  $\pm$  SDs ( $n = 6-14$  per group). \* $p < 0.01$ , \*\* $p < 0.001$  by the Mann-Whitney  $U$ -test.




**FIGURE 2** | A PI3K $\delta$  inhibitor attenuated poly I:C-induced upregulation of PD-L1 on mouse lung cells. **(A,B)** IC87114 was administered intratracheally (i.t.) to mice and PD-L1 **(A)** or PD-1 **(B)** expression on mouse lung cells was analyzed 24 h following treatment using flow cytometry. **(C,D)** IC87114 or vehicle was administered i.t. to mice followed by i.t. administration of poly I:C or vehicle. PD-L1 expression **(C)** on lung cells or cell viability **(D)** was analyzed 24 h following administration using flow cytometry. MFI, mean fluorescence intensity. All results are representative of at least three independent experiments. Data represent means  $\pm$  SDs ( $n = 8-16$  per group). \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  by the Mann-Whitney U-test or one-way ANOVA as appropriate.



24 h following poly I:C administration. Treatment with IC87114 suppressed poly I:C-induced influx of inflammatory cells to the same levels as controls (**Figure 4A**). Poly I:C

stimulation significantly increased the numbers of BALF macrophages, neutrophils and lymphocytes, but not the numbers of eosinophils (**Figure 4A** and data not shown).






Poly I:C-induced increases in numbers of macrophages, neutrophils and lymphocytes were attenuated by IC87114 and treatment with IC87114 alone did not affect cell numbers in BALF (Figure 4A).

Levels of KC, IL-6, and MIP-1 $\beta$  in BALF were analyzed as representative pro-inflammatory chemokines and cytokines induced by TLR3 and MDA5 signaling. Poly I:C significantly increased the concentration of KC, IL-6 and MIP-1 $\beta$  in BALF 24 h following administration, and treatment with

IC87114 attenuated poly I:C-induced increases in KC, IL-6 and MIP-1 $\beta$  (Figure 4B).

### A PI3K $\delta$ Inhibitor Enhances Poly I:C-Induced IFN Responses by Promoting IRF3 Phosphorylation in Mouse Lungs

Types I and III IFNs play important roles in immune responses to viral infection (12, 40). Thus, we examined the effects of




IC87114 on type I and type III IFNs responses in mouse lungs using real-time quantitative PCR. Six hours following poly I:C administration, expression of *IFN $\beta$*  and *IFN $\lambda_2$*  was significantly increased compared with controls and with mice treated with IC87114 alone (Figure 5A). Furthermore, combined treatment with IC87114 and poly I:C enhanced expression of *IFN $\beta$*  and *IFN $\lambda_2$*  to a greater degree than poly I:C treatment alone (Figure 5A). Expression of IRGs such as IFN-induced protein with tetratricopeptide repeats (*Ifit1*), *Ifit2*, and IFN-stimulated gene (*ISG15*) were analyzed and similar results were obtained. Significant upregulation of *Ifit1*, *Ifit2*, and *ISG15* expression was observed 24 h following poly I:C administration. Combined treatment with IC87114 and poly I:C greatly enhanced the expression of these IRGs compared with poly I:C-treated mice (Figure 5B).

Previous studies showed that the production of early IFNs results from the phosphorylation and translocation of available IRF3 into the nucleus and does not require production of new IRF proteins (41, 42). Therefore, we examined the effects of IC87114 on IRF3 phosphorylation in lungs by western blotting. Three hours following poly I:C administration,

IRF3 was phosphorylated in poly I:C-treated mice. Combined treatment with IC87114 and poly I:C significantly increased IRF3 phosphorylation levels compared with mice treated with poly I:C alone (Figure 6). Total IRF3 levels were not affected by poly I:C administration with or without IC87114 treatment (Figure 6).

### A PI3K $\delta$ Inhibitor Suppresses Poly I:C-Induced PD-L1 Expression and Enhances Antiviral IFN Responses in PBECS

Using bronchial epithelial cells collected from never-smoker patients with normal lung function, we investigated whether IC87114 had similar effects on poly I:C-induced PD-L1 expression, cytokine and chemokine production, and IFN responses as in mouse lung cells. PD-L1 expression on PBECS was significantly increased 24 h following poly I:C stimulation and treatment with IC87114 attenuated poly I:C-induced upregulation of PD-L1 (Figure 7A). At the same time point, no significant difference was observed in cell viability among four groups (Figure 7B and Supplementary Figure 1B). IC87114 also



decreased levels of IL-6 and IL-8 induced by poly I:C treatment in culture supernatants 24 h following stimulation (Figure 7C). By contrast, production of IFN $\beta$  and  $\lambda_{1/3}$  in culture supernatants was significantly enhanced by combined treatment with IC87114 and poly I:C compared with PBECs treated with poly I:C alone (Figure 7D). We assessed expression of ISG56, MxA and 2'5'OAS 6 h following poly I:C stimulation as representative IRGs in PBECs. Combined treatment with IC87114 and poly I:C significantly increased expression of ISG56, MxA and 2'5'OAS compared with treatment with poly I:C alone (Figure 7E).


### siRNA Knockdown of the PIK3CD Gene Suppresses Poly I:C-Induced Upregulation of PD-L1 and Enhances Antiviral IFN Responses in PBECs

The effect of knockdown of the *PIK3CD* gene encoding PI3K p110 $\delta$  was assessed in PBECs. PI3K p110 $\delta$  expression was significantly suppressed in PBECs treated with siPIK3CD 48 and 72 h following transfection compared with that in PBECs treated with siNC (Figure 8A). Next, PBECs transfected with siPIK3CD or siNC for 48 h were stimulated with poly I:C and PD-L1 expression and production of cytokine and IFNs in culture supernatants were analyzed 24 h following


stimulation. PD-L1 expression was significantly increased on PBECs treated with siNC and poly I:C compared with poly I:C-untreated controls although poly I:C-induced increases in PD-L1 expression were attenuated in PBECs treated with siPIK3CD and poly I:C (Figure 8B). IL-6 production levels in culture supernatants of PBECs treated with siPIK3CD and poly I:C were reduced compared with PBECs treated with siNC and poly I:C (Figure 8C). On the other hand, enhanced production of IFN $\beta$  and  $\lambda_{1/3}$  in culture supernatants were observed in PBECs treated with siPIK3CD and poly I:C (Figure 8D). At the same time point, cell viability was assessed and not affected by siRNA knockdown of PIK3CD and stimulation of poly I:C (Figure 8E and Supplementary Figure 1C).

### A PI3K $\delta$ Inhibitor Suppresses hMPV-Induced PD-L1 Expression and Inhibits Viral Replication in PBECs Without Changing the Production Levels of IFNs in Supernatants


Finally, we investigated the effects of IC87114 on hMPV-induced PD-L1 expression, IFN responses and viral replication in PBECs. At 48 and 72 hpi, PD-L1 expression was significantly upregulated on PBECs infected with hMPV at a MOI of 0.1



**FIGURE 7** | Effects of PI3K $\delta$  inhibitor on poly I:C-induced upregulation of PD-L1, production of cytokines and chemokines, and IFN responses in PBECs. PBECs were pretreated with IC87114 or vehicle for 1 h, then stimulated with poly I:C. Cell lysates for RNA extraction or cells and culture supernatants were collected 6 or 24 h following administration, respectively. **(A,B)** PD-L1 expression **(A)** on PBECs or cell viability **(B)** was analyzed using flow cytometry. Representative histograms are shown. MFI, mean fluorescence intensity. **(C)** Pro-inflammatory cytokine and chemokine levels in supernatants were measured by ELISA. **(D)** IFN levels in supernatants were measured by ELISA. The dotted line shows the lower limit of detection. # Levels in all samples were below the detection limit. **(E)** RNA was extracted and real-time quantitative reverse-transcriptase PCR was performed. Target gene expression levels were normalized to those of 18S rRNA. All results are representative of at least two independent experiments. Data represent means  $\pm$  SDs (n=6 per group) of three replicates from a minimum of two independent donors. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  by one-way ANOVA.



**FIGURE 8 |** Effects of siRNA targeting the PIK3CD gene on poly I:C-induced upregulation of PD-L1, production of cytokines and IFN responses in PBECs. **(A)** Representative western blots showing PI3K p110 $\delta$  in PBECs treated with PIK3CD siRNA or negative control (NC) siRNA for 48 or 72 h. Band intensity was quantitated using densitometry. **(B–D)** PBECs were transfected with PIK3CD siRNA or NC siRNA for 48 h, then stimulated with poly I:C or vehicle for 24 h. **(B)** PD-L1 expression on PBECs was analyzed using flow cytometry. MFI, mean fluorescence intensity. **(C)** IL-6 levels in supernatants were measured by ELISA. **(D)** IFN levels in supernatants were measured by ELISA. The dotted line shows the lower limit of detection. # Levels in all samples were below the detection limit. **(E)** Viable cells (PI negative) were identified using flow cytometry. All results are representative of at least two independent experiments. Data represent means  $\pm$  SDs ( $n = 6–9$  per group) of three replicates from a minimum of two independent donors. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  by one- or two-way ANOVA as appropriate.



**FIGURE 9** | A PI3K $\delta$  inhibitor attenuated hMPV-induced upregulation of PD-L1 on PBECs. **(A)** PBECs were infected with hMPV (MOI 0.1) and PD-L1 expression was analyzed at the indicated times using flow cytometry. **(B)** IC87114 or vehicle was added prior to and after hMPV infection (MOI 0.1). PD-L1 expression was analyzed at 48 hpi using flow cytometry. MFI, mean fluorescence intensity. All results are representative of at least two independent experiments. Data represent means  $\pm$  SDs ( $n = 6$  per group) of three replicates from a minimum of two independent donors. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  by one- or two-way ANOVA as appropriate.

compared with uninfected controls (Figure 9A). The increases in PD-L1 expression or hMPV nucleocapsid protein (*hMPV N*) and *IFN $\lambda_1$*  gene expression were absent if the virus was UV-irradiated thereby excluding non-hMPV induced responses (Supplementary Figures 2A,B). Treatment with IC87114 attenuated hMPV-induced upregulation of PD-L1 at 48 hpi (Figure 9B). Similarly to previous observations (31), hMPV did not induce IL-8 gene in PBECs and protein in culture supernatants (Supplementary Figure 3). IFN production following hMPV infection and viral replication were examined in PBECs treated with or without IC87114. Levels of IFN $\beta$  and IFN $\lambda_{1/3}$  in culture supernatants were significantly increased at 48 hpi and treatment with IC87114 did not affect production levels of IFN $\beta$  and IFN $\lambda_{1/3}$  (Figure 10A). However, *hMPV N* and *IFN $\lambda_1$*  gene expression induced by hMPV infection were significantly decreased in PBECs treated with IC87114 at 36 and 48 hpi, respectively (Figure 10B). There was a linear association between *hMPV N* and *IFN $\lambda_1$*  gene expression in cells treated with or without IC87114 (Figure 10B). IC87114 changed a slope of a regression line larger, indicating more efficient induction of IFN gene expression responsive to infections in IC87114-treated cells. Next, PBECs were infected with hMPV-GFP at a MOI of 0.1 and observed at 72 hpi by fluorescence microscopy. The number of hMPV-infected cells were significantly decreased and smaller syncytium formation was induced in IC87114-treated cells compared with untreated controls (Figure 10C and Supplementary Figure 4). After infection with UV-irradiated hMPV-GFP, GFP fluorescence and *hMPV N* gene expression was not induced in PBECs (Supplementary Figures 2C,D).


## DISCUSSION

PI3K $\delta$  has been extensively studied as a potential target for anti-oncogenic and anti-inflammatory therapies (27, 43). However, the role of PI3K $\delta$  signaling in antiviral responses is poorly understood. Using a murine model of airway viral infection, we investigated PD-L1 and PD-1 expression on lung cells following i.t. poly I:C administration and observed upregulation of PD-L1

but not PD-1 on epithelial cells, lymphocytes, macrophages, and neutrophils. A selective PI3K $\delta$  inhibitor, IC87114, suppressed poly I:C-induced PD-L1 expression on epithelial cells and neutrophils, but not on lymphocytes and macrophages. This suppressive effects on PD-L1 might be due to inhibition of the Akt/mTOR pathway by IC87114. IC87114 also attenuated poly I:C-induced increases in influx of inflammatory cells and secretion of pro-inflammatory chemokines and cytokines in BALF. Interestingly, IC87114 enhanced poly I:C-induced expression of IFNs and IRGs in the lung and increased the phosphorylation of IRF3. Finally, we assessed the effects of IC87114 in PBECs. Similarly to our *in vivo* results, IC87114 or siRNA knockdown of the *PIK3CD* gene decreased poly I:C-induced PD-L1 expression, production of pro-inflammatory chemokines and cytokines and enhanced antiviral IFN responses in PBECs. Furthermore, IC87114 treatment reduced replication of hMPV in PBECs.

Our *in vivo* experiments showed that the suppressive effects of IC87114 on poly I:C-induced PD-L1 expression depend on cell type. It has been well-established that the upregulation of PD-L1 on infected epithelium causes virus-specific CD8<sup>+</sup> T cells to fail to eliminate virus in the epithelium (19, 44). Upregulation of PD-L1 on neutrophils was reported to suppress lymphocyte proliferation, induce lymphocyte apoptosis and compromise anti-pathogen defense (45, 46). These previous reports suggested that the suppressive effects of IC87114 on PD-L1 expression by epithelial cells and neutrophils may restore T cell function and promote viral clearance. However, little is known about PD-L1 expression and regulation in macrophages. Loke et al. showed that PD-L1 expression can be induced on macrophages not only by classical activation stimuli such as poly I:C and lipopolysaccharide, but also through alternative activation via IL-4 (47). This alternative induction of PD-L1 may be one reason why IC87114 did not suppress poly I:C-induced PD-L1 expression on macrophages in our study. Additional studies are needed to investigate the functions of PD-L1 on macrophages during immune responses.

Respiratory virus infection precipitates the majority of exacerbations of asthma and COPD. Subsequent production of



**FIGURE 10** | A PI3K $\delta$  inhibitor reduced replication of hMPV in PBECs without changing the production level of IFNs in supernatants. IC87114 or vehicle was added prior to and after hMPV (MOI 0.1) (A,B) or hMPV-GFP (MOI 0.1) (C) infection. (A) Cell culture supernatants were collected at 48 hpi and IFN levels in supernatants were measured by ELISA. The dotted line shows the lower limit of detection. #Levels in all samples were below the detection limit. (B) Cell lysates for RNA extraction were collected at 24, 36, and 48 hpi and real-time quantitative reverse-transcriptase PCR was performed. Target gene expression levels were normalized to those of 18S rRNA. (C) Images of hMPV-GFP-infected cells at 72 hpi were obtained using fluorescence microscopy (10 $\times$  objective lens) and the number of cells infected with hMPV-GFP (4 $\times$  objective lens) was counted. Scale bar, 200  $\mu$ m. All results are representative of at least two independent experiments. Data represent means  $\pm$  SDs ( $n = 6$  per group) of three replicates from a minimum of two independent donors. \* $p < 0.05$ , \*\* $p < 0.01$ , \*\*\* $p < 0.001$  by one- or two-way ANOVA and spearman correlation as appropriate.

chemokines and cytokines in the bronchial epithelium induce inflammation in the lungs and excessive inflammatory responses may induce airway hyperresponsiveness and remodeling (48). Previous studies using pan-PI3K inhibitors have showed discrepant results regarding a role for PI3K in TLR-mediated inflammatory responses, suggesting that the effect of PI3K on NF- $\kappa$ B activation may depend on cell type (49, 50). In our study, the selective PI3K $\delta$  inhibitor IC87114 attenuated poly I:C-induced inflammatory responses *in vivo* and *in vitro*. Other investigators also have showed that IC87114 decreased neutrophilic inflammation in BALF to almost control level (51–53). These anti-inflammatory effects of IC87114 might contribute to prevention of asthma and COPD exacerbation. PI3K $\delta$  signaling is thought to play an important role in the pathogenesis of COPD and asthma, both respiratory diseases characterized by chronic lung inflammation (43, 54). In an ovalbumin-induced asthma model, IC87114 treatment or p110 $\delta$  inactivation in mice decreased type 2 cytokine responses, allergic airway inflammation and airway hyperresponsiveness (55, 56). To et al. reported that PI3K $\delta$  mRNA levels and activity were elevated in the peripheral lung tissues of COPD patients and that IC87114 restored corticosteroid resistance in cigarette smoke-exposed mice through inhibition of oxidative stress-dependent PI3K $\delta$  activation (57). Based on these studies, clinical trials with inhaled PI3K $\delta$  inhibitors in asthmatics and patients with COPD are currently being conducted to evaluate their efficacy in preventing exacerbation of these conditions and maintaining lung function (43, 58).

In this study, we showed that IC87114 significantly enhanced antiviral IFN responses and increased the phosphorylation of IRF3. Phosphorylation of IRF3 is critical for type I interferon production triggered by detection of dsRNA by pathogen recognition receptors (59). Subsequently, binding of secreted IFNs to ubiquitously-expressed type I IFN receptors triggers IFN signaling, followed by induction of IRG expression. A previous study using a pan-PI3K inhibitor showed that the PI3K pathway played an essential role in high concentration of poly I:C-induced phosphorylation of IRF3 (60). However, our results suggest that PI3K $\delta$  may negatively regulate antiviral IFN responses in the lungs. To the best of our knowledge, only two previous studies showed involvement of PI3K $\delta$  in antiviral IFN responses and suppressive effects of a selective PI3K $\delta$  inhibitor on antiviral IFN production. Guiducci et al. reported that IC87114 inhibited IRF7-mediated IFN $\alpha$  production in a dose-dependent manner in human plasmacytoid DCs stimulated by CpG-C, a synthetic TLR9 agonist (61). In another report from Aksoy, IC87114 decreased IFN $\beta$  production in LPS-stimulated BMDCs than in IC87114-untreated cells through inhibition of IRF3 phosphorylation (62). PI3K $\delta$  signaling may differentially regulate antiviral IFN responses in different cell types as shown in our previous study in which opposite effects of a pan-PI3K inhibitor on poly I:C-induced PD-L1 expression were observed among bronchial epithelial cells and BMDC (38). As mentioned above, it was reported that PI3K $\delta$  activity was elevated in the peripheral lung tissues of patients with COPD. In addition, a recent study showed that patients with COPD and a history of frequent exacerbations had reduced antiviral IFN responses

associated with increased secondary bacterial infections (63). Impaired antiviral IFN responses in asthmatic PBECs were also reported (64, 65). Synthesizing these previous reports and our results, susceptibility to infection may occur through exaggerated PI3K $\delta$  signaling, and PI3K $\delta$  may be targeted therapeutically in COPD and asthma to prevent development and aggravation of viral infection.

Lastly, we demonstrated the effects of IC87114 in PBECs. Bronchial epithelial cells act as the first line of defense against inhaled exogenous substances including viral pathogens recognized by pattern recognition receptors. The expression of PI3K $\delta$  is generally restricted to leukocytes although a few previous studies suggest PI3K $\delta$  expression and activity in non-leukocytes (66–70). Ge et al. showed that PI3K $\delta$  mRNA and protein levels were increased in the airway smooth muscle cells of asthmatics and patients with COPD, respectively (66). In another study, IC87114 decreased hypoxia-inducible factor-1 $\alpha$  protein levels in primary airway epithelial cells isolated from OVA-treated mice; this protein stimulates the expression of vascular endothelial growth factor that promotes vascular permeability in respiratory diseases (69). Tumor necrosis factor- $\alpha$  stimulation induced PI3K $\delta$  expression in human endothelial cells and synovial fibroblasts (70). Our results in PBECs are the first demonstration that PI3K $\delta$  signaling may play an important role in upregulation of PD-L1 and negative regulation of IFN responses, resulting in suppression of adaptive and innate responses to viral infections in PBECs. Most importantly, hMPV is able to replicate less efficiently in PBECs treated with IC87114. A limitation of our study was that poly I:C or single-cell experiments are unable to activate adaptive immune responses by presentation of viral antigen. Using live respiratory virus, further *in vivo* investigations are needed to evaluate whether PI3K $\delta$  inhibitors decrease viral loads in lungs and promote recovery from infection.

In summary, a PI3K $\delta$  inhibitor, IC87114, attenuated poly I:C-induced expression of the co-inhibitory molecule PD-L1 on epithelial cells and neutrophils in the lungs. IC87114 also suppressed increases in the influx of inflammatory cells into the lung and secretion of pro-inflammatory chemokines and cytokines. Furthermore, IC87114 enhanced poly I:C-induced antiviral IFN responses in the lungs. Similar results were obtained in *in vitro* experiments using PBECs treated with IC87114 or siRNA knockdown of the *PIK3CD* gene. Finally, experiments using hMPV confirmed that IC87114 inhibits viral replication in PBECs. Taken together, these results indicate that IC87114 may promote virus elimination and clearance, and prevent prolonged lung inflammation exacerbates asthma and COPD. Use of a selective PI3K $\delta$  inhibitor can potentially have significant benefits in virus-exacerbated respiratory diseases.

## DATA AVAILABILITY STATEMENT

The datasets generated for this study are available on request to the corresponding author.



## ETHICS STATEMENT

The studies involving human participants were reviewed and approved by the Kyushu University Institutional Review Board for Clinical Research. The patients/participants provided their written informed consent to participate in this study. The animal study was reviewed and approved by the Kyushu University Animal Care and Use Committee.

## AUTHOR CONTRIBUTIONS

KK, SF, YN, and KM conceived the study design and supervised the scientific work. AF, KK, KT, NY, and TO performed the experiments. AF and KK analyzed the data. All authors contributed to and approved the final manuscript.

## FUNDING

This work was supported by JSPS KAKENHI Grant Numbers JP24591132, JP15K09222, and JP18K15953. This work was supported (in part) by Center for Clinical and Translational Research of Kyushu University.

## ACKNOWLEDGMENTS

The authors would like to acknowledge Philip G. Bardin and Belinda J. Thomas (Centre for Innate Immunity and Infectious Diseases, Hudson Institute of Medical Research) for providing hMPV, and Yuta Shirogane (Department of Virology, Faculty of Medicine, Kyushu University) for giving technical advice on virus propagation. We also appreciate technical assistance from Mayu Matsuo, Mikiko Nakano, Kayoko Ono and The Research Support Center, Research Center for Human Disease Modeling, Kyushu University Graduate School of Medical Sciences. We thank Edanz Group ([www.edanzediting.com/ac](http://www.edanzediting.com/ac)) for editing a draft of this manuscript.

## REFERENCES

- D'amato G, Vitale C, Molino A, Stanzola A, Sanduzzi A, Vatrella A, et al. Asthma-related deaths. *Multidiscipl Respir Med.* (2016) 11:37. doi: 10.4081/mrm.2016.331
- Afonso ASM, Verhamme KMC, Sturkenboom MCJM, Brusselle GGO. COPD in the general population: prevalence, incidence and survival. *Respiratory Med.* (2011) 105:1872–84. doi: 10.1016/j.rmed.2011.06.012
- Jackson DJ, Sykes A, Mallia P, Johnston SL. Asthma exacerbations: origin, effect, and prevention. *J Allergy Clin Immunol.* (2011) 128:1165–74. doi: 10.1016/j.jaci.2011.10.024
- Linden D, Guo-Parke H, Coyle PV, Fairley D, Mcauley DF, Taggart CC, et al. Respiratory viral infection: a potential “missing link” in the pathogenesis of COPD. *Eur Respir Rev.* (2019) 28:180063. doi: 10.1183/16000617.0063-2018
- Alexopoulou L, Holt AC, Medzhitov R, Flavell RA. Recognition of double-stranded RNA and activation of NF-kappaB by Toll-like receptor 3. *Nature.* (2001) 413:732–8. doi: 10.1038/35099560
- Yoneyama M, Kikuchi M, Natsukawa T, Shinobu N, Imaizumi T, Miyagishi M, et al. The RNA helicase RIG-I has an essential function in double-stranded RNA-induced innate antiviral responses. *Nat Immunol.* (2004) 5:730–7. doi: 10.1038/nri1087
- Kato H, Takeuchi O, Sato S, Yoneyama M, Yamamoto M, Matsui K, et al. Differential roles of MDA5 and RIG-I helicases in the recognition of RNA viruses. *Nature.* (2006) 441:101–5. doi: 10.1038/nature04734
- Caskey M, Lefebvre F, Filali-Mouhim A, Cameron MJ, Goulet JP, Haddad EK, et al. Synthetic double-stranded RNA induces innate immune responses similar to a live viral vaccine in humans. *J Exp Med.* (2011) 208:2357–66. doi: 10.1084/jem.20111171
- Mccartney S, Vermi W, Gilfillan S, Cella M, Murphy TL, Schreiber RD, et al. Distinct and complementary functions of MDA5 and TLR3 in poly(I:C)-mediated activation of mouse NK cells. *J Exp Med.* (2009) 206:2967–76. doi: 10.1084/jem.20091181
- Kawai T, Akira S. Toll-like receptor and RIG-I-like receptor signaling. *Ann N Y Acad Sci.* (2008) 1143:1–20. doi: 10.1196/annals.1443.020
- Levy DE, Marie IJ, Durbin JE. Induction and function of type I and III interferon in response to viral infection. *Curr Opin Virol.* (2011) 1:476–86. doi: 10.1016/j.coviro.2011.11.001
- Sadler AJ, Williams BR. Interferon-inducible antiviral effectors. *Nat Rev Immunol.* (2008) 8:559–68. doi: 10.1038/nri2314

## SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fimmu.2020.00432/full#supplementary-material>

**Supplementary Figure 1** | Representative dot plots showing a PI profile in mouse lungs or human PBECs. **(A)** IC87114 or vehicle was administered i.t. to mice followed by i.t. administration of poly I:C or vehicle. Viable cells (PI negative) were identified 24 h following administration using flow cytometry. **(B)** PBECs were pretreated with IC87114 or vehicle for 1 h, then stimulated with poly I:C or vehicle. Viable cells (PI negative) were identified 24 h following stimulation using flow cytometry. **(C)** PIK3CD or negative control (NC) siRNA was transfected into PBECs for 48 h, then stimulated with poly I:C or vehicle. Viable cells (PI negative) were identified 24 h following stimulation using flow cytometry.

**Supplementary Figure 2** | UV-irradiated hMPV was unable to replicate in human bronchial epithelial cells and did not induce PD-L1 expression and IFN responses. **(A,B)** PBECs **(A)** or BEAS-2B **(B)** were infected with hMPV (MOI 0.1) or UV-irradiated hMPV (MOI 0.1). **(A)** PD-L1 expression was analyzed at 24 hpi using flow cytometry. **(B)** Cell lysates for RNA extraction were collected at 48 and 72 hpi and real-time quantitative reverse-transcriptase PCR was performed. **(C,D)** PBECs were infected with hMPV-GFP (MOI 0.1) or UV-irradiated hMPV-GFP (MOI 0.1). **(C)** Images of infected cells at 72 hpi obtained using fluorescence microscopy. Scale bar, 100  $\mu$ m. **(D)** Cell lysates for RNA extraction were collected at 120 hpi and real-time quantitative reverse-transcriptase PCR was performed. Target gene expression levels were normalized to those of 18S rRNA. Data represent means  $\pm$  SDs ( $n = 3$  per group). \* $p < 0.05$ , \*\* $p < 0.01$  by one- or two-way ANOVA as appropriate.

**Supplementary Figure 3** | hMPV did not induce IL-8 gene expression in PBECs and protein expression in supernatants. IC87114 or vehicle was added prior to and after hMPV (MOI 0.1) infection. **(A)** Cell lysates for RNA extraction were collected at 24, 36, and 48 hpi and real-time quantitative reverse-transcriptase PCR was performed. Target gene expression levels were normalized to those of 18S rRNA. **(B)** Cell culture supernatants were collected at 24 and 48 hpi and IL-8 levels in supernatants were measured by ELISA. All results are representative of at least two independent experiments. Data represent means  $\pm$  SDs ( $n = 6$  per group) of three replicates from a minimum of two independent donors. \* $p < 0.01$  by two-way ANOVA.

**Supplementary Figure 4** | Images of hMPV-GFP-infected cells. PBECs were cultured to semi-confluence in 12-well plates. IC87114 or vehicle was added prior to and after hMPV-GFP (MOI 0.1) infection, then cells were observed at 72 hpi using fluorescence microscopy.

**Supplementary Table 1** | Sequences of real-time PCR primers used in this study.

13. Ceeraz S, Nowak EC, Noelle RJ. B7 family checkpoint regulators in immune regulation and disease. *Trends Immunol.* (2013) 34:556–63. doi: 10.1016/j.it.2013.07.003
14. Pauken KE, Wherry EJ. Overcoming T cell exhaustion in infection and cancer. *Trends Immunol.* (2015) 36:265–76. doi: 10.1016/j.it.2015.02.008
15. Rogers MC, Williams JV. Reining in the CD8<sup>+</sup> T cell: respiratory virus infection and PD-1-mediated T-cell impairment. *PLoS Pathog.* (2019) 15:e1007387. doi: 10.1371/journal.ppat.1007387
16. Barber DL, Wherry EJ, Masopust D, Zhu B, Allison JP, Sharpe AH, et al. Restoring function in exhausted CD8 T cells during chronic viral infection. *Nature.* (2006) 439:682–7. doi: 10.1038/nature04444
17. Iwai Y, Terawaki S, Ikegawa M, Okazaki T, Honjo T. PD-1 inhibits antiviral immunity at the effector phase in the liver. *J Exp Med.* (2003) 198:39–50. doi: 10.1084/jem.20022235
18. Erickson JJ, Gilchuk P, Hastings AK, Tollefson SJ, Johnson M, Downing MB, et al. Viral acute lower respiratory infections impair CD8<sup>+</sup> T cells through PD-1. *J Clin Invest.* (2012) 122:2967–82. doi: 10.1172/JCI62860
19. McNally B, Ye F, Willette M, Flano E. Local blockade of epithelial PDL-1 in the airways enhances T cell function and viral clearance during influenza virus infection. *J Virol.* (2013) 87:12916–24. doi: 10.1128/JVI.02423-13
20. Tsuda M, Matsumoto K, Inoue H, Matsumura M, Nakano T, Mori A, et al. Expression of B7-H1 and B7-DC on the airway epithelium is enhanced by double-stranded RNA. *Biochem Biophys Res Commun.* (2005) 330:263–70. doi: 10.1016/j.bbrc.2005.02.161
21. Kan-O K, Matsumoto K, Inoue H, Fukuyama S, Asai Y, Watanabe W, et al. Corticosteroids plus long-acting beta2-agonists prevent double-stranded RNA-induced upregulation of B7-H1 on airway epithelium. *Int Arch Allergy Immunol.* (2013) 160:27–36. doi: 10.1159/000338430
22. Hamano S, Matsumoto K, Tonai K, Fukuyama S, Kan OK, Seki N, et al. Effects of corticosteroid plus long-acting beta2-agonist on the expression of PD-L1 in double-stranded RNA-induced lung inflammation in mice. *J Inflamm.* (2017) 14:2. doi: 10.1186/s12950-017-0149-4
23. Cantley LC. The phosphoinositide 3-kinase pathway. *Science.* (2002) 296:1655–7. doi: 10.1126/science.296.5573.1655
24. Vanhaesebroeck B, Leevers SJ, Ahmadi K, Timms J, Katso R, Driscoll PC, et al. Synthesis and function of 3-phosphorylated inositol lipids. *Annu Rev Biochem.* (2001) 70:535–602. doi: 10.1146/annurev.biochem.70.1.535
25. Okkenhaug K. Signaling by the phosphoinositide 3-kinase family in immune cells. *Annu Rev Immunol.* (2013) 31:675–704. doi: 10.1146/annurev-immunol-032712-095946
26. Angulo I, Vadas O, Garcon F, Banham-Hall E, Plagnol V, Leahy TR, et al. Phosphoinositide 3-kinase delta gene mutation predisposes to respiratory infection and airway damage. *Science.* (2013) 342:866–71. doi: 10.1126/science.1243292
27. Michalovich D, Nejentsev S. Activated PI3 kinase delta syndrome: from genetics to therapy. *Front Immunol.* (2018) 9:369. doi: 10.3389/fimmu.2018.00369
28. Lucas CL, Kuehn HS, Zhao F, Niemela JE, Deenick EK, Palendira U, et al. Dominant-activating germline mutations in the gene encoding the PI(3)K catalytic subunit p110delta result in T cell senescence and human immunodeficiency. *Nat Immunol.* (2014) 15:88–97. doi: 10.1038/n.i.2771
29. Coulter TI, Chandra A, Bacon CM, Babar J, Curtis J, Screaton N, et al. Clinical spectrum and features of activated phosphoinositide 3-kinase delta syndrome: a large patient cohort study. *J Allergy Clin Immunol.* (2017) 139:597–606 e594. doi: 10.1016/j.jaci.2016.06.021
30. Rudd PA, Thomas BJ, Zaid A, Macdonald M, Kan OK, Rolph MS, et al. Role of human metapneumovirus and respiratory syncytial virus in asthma exacerbations: where are we now? *Clin Sci.* (2017) 131:1713–21. doi: 10.1042/CS20160011
31. Kan-O, K, Ramirez R, Macdonald MI, Rolph M, Rudd PA, Spann KM, et al. Human metapneumovirus infection in chronic obstructive pulmonary disease: impact of glucocorticosteroids and interferon. *J Infect Dis.* (2017) 215:1536–45. doi: 10.1093/infdis/jix167
32. Wen SC, Schuster JE, Gilchuk P, Boyd KL, Joyce S, Williams JV. Lung CD8<sup>+</sup> T cell impairment occurs during human metapneumovirus infection despite virus-like particle induction of functional CD8<sup>+</sup> T cells. *J Virol.* (2015) 89:8713–26. doi: 10.1128/JVI.00670-15
33. Soto JA, Galvez NMS, Benavente FM, Pizarro-Ortega MS, Lay MK, Riedel C, et al. Human metapneumovirus: mechanisms and molecular targets used by the virus to avoid the immune system. *Front Immunol.* (2018) 9:2466. doi: 10.3389/fimmu.2018.02466
34. Kuiken T, Van Den Hoogen BG, Van Riel DA, Laman JD, Van Amerongen G, Sprong L, et al. Experimental human metapneumovirus infection of cynomolgus macaques (*Macaca fascicularis*) results in virus replication in ciliated epithelial cells and pneumocytes with associated lesions throughout the respiratory tract. *Am J Pathol.* (2004) 164:1893–900. doi: 10.1016/S0002-9440(10)63750-9
35. Boivin G, De Serres G, Hamelin ME, Cote S, Argouin M, Tremblay G, et al. An outbreak of severe respiratory tract infection due to human metapneumovirus in a long-term care facility. *Clin Infect Dis.* (2007) 44:1152–8. doi: 10.1086/513204
36. Fukuyama S, Matsunaga Y, Zhanghui W, Noda N, Asai Y, Moriwaki A, et al. A zinc chelator TPEN attenuates airway hyperresponsiveness and airway inflammation in mice *in vivo*. *Allergol Int.* (2011) 60:259–66. doi: 10.2332/allergolint.09-OA-0167
37. Foster MW, Gerhardt G, Robitaille L, Plante PL, Boivin G, Corbeil J, et al. Targeted proteomics of human metapneumovirus in clinical samples and viral cultures. *Anal Chem.* (2015) 87:10247–54. doi: 10.1021/acs.analchem.5b01544
38. Kan-O K, Matsumoto K, Asai-Tajiri Y, Fukuyama S, Hamano S, Seki N, et al. PI3K-delta mediates double-stranded RNA-induced upregulation of B7-H1 in BEAS-2B airway epithelial cells. *Biochem Biophys Res Commun.* (2013) 435:195–201. doi: 10.1016/j.bbrc.2013.04.082
39. Lastwika KJ, Wilson W 3rd, Li QK, Norris J, Xu H, Ghazarian SR, et al. Control of PD-L1 expression by oncogenic activation of the AKT-mTOR pathway in non-small cell lung cancer. *Cancer Res.* (2016) 76:227–38. doi: 10.1158/0008-5472.CAN-14-3362
40. Zhou JH, Wang YN, Chang QY, Ma P, Hu Y, Cao X. Type III interferons in viral infection and antiviral immunity. *Cell Physiol Biochem.* (2018) 51:173–85. doi: 10.1159/000495172
41. Levy DE, Marié I, Prakash A. Ringing the interferon alarm: differential regulation of gene expression at the interface between innate and adaptive immunity. *Curr Opin Immunol.* (2003) 15:52–58. doi: 10.1016/S0952-7915(02)00011-0
42. Demoullins T, Baron ML, Kettaf N, Abdallah A, Sharif-Askari E, Sekaly RP. Poly (I:C) induced immune response in lymphoid tissues involves three sequential waves of type I IFN expression. *Virology.* (2009) 386:225–36. doi: 10.1016/j.virol.2009.01.024
43. Stark AK, Srikantharajah S, Hessel EM, Okkenhaug K. PI3K inhibitors in inflammation, autoimmunity and cancer. *Curr Opin Pharmacol.* (2015) 23:82–91. doi: 10.1016/j.coph.2015.05.017
44. Telcian AG, Laza-Stanca V, Edwards MR, Harker JA, Wang H, Bartlett NW, et al. RSV-induced bronchial epithelial cell PD-L1 expression inhibits CD8<sup>+</sup> T cell nonspecific antiviral activity. *J Infect Dis.* (2011) 203:85–94. doi: 10.1093/infdis/jiq020
45. De Kleijn S, Langereis JD, Leentjens J, Kox M, Netea MG, Koenderman L, et al. IFN-gamma-stimulated neutrophils suppress lymphocyte proliferation through expression of PD-L1. *PLoS ONE.* (2013) 8:e72249. doi: 10.1371/journal.pone.0072249
46. Wang JF, Li JB, Zhao YJ, Yi WJ, Bian JJ, Wan XJ, et al. Up-regulation of programmed cell death 1 ligand 1 on neutrophils may be involved in sepsis-induced immunosuppression: an animal study and a prospective case-control study. *Anesthesiology.* (2015) 122:852–63. doi: 10.1097/ALN.0000000000000525
47. Loke P, Allison JP. PD-L1 and PD-L2 are differentially regulated by Th1 and Th2 cells. *Proc Natl Acad Sci USA.* (2003) 100:5336–41. doi: 10.1073/pnas.0931259100
48. Jackson DJ, Johnston SL. The role of viruses in acute exacerbations of asthma. *J Allergy Clin Immunol.* (2010) 125:1178–87; quiz 1188–79. doi: 10.1016/j.jaci.2010.04.021
49. Troutman TD, Bazan JF, Pasare C. Toll-like receptors, signaling adapters and regulation of the pro-inflammatory response by PI3K. *Cell Cycle.* (2012) 11:3559–67. doi: 10.4161/cc.21572
50. Hazeki K, Nigorikawa K, Hazeki O. Role of phosphoinositide 3-kinase in innate immunity. *Biol Pharm Bull.* (2007) 30:1617–23. doi: 10.1248/bpb.30.1617

51. Kang BN, Ha SG, Ge XN, Reza Hosseinkhani M, Bahaie NS, Greenberg Y, et al. The p110delta subunit of PI3K regulates bone marrow-derived eosinophil trafficking and airway eosinophilia in allergen-challenged mice. *Am J Physiol Lung Cell Mol Physiol*. (2012) 302:L1179–91. doi: 10.1152/ajplung.00005.2012
52. Kim HK, Lee GH, Bhattarai KR, Junjappa RP, Lee HY, Handigund M, et al. PI3Kdelta contributes to ER stress-associated asthma through ER-redox disturbances: the involvement of the RIDD-RIG-I-NF-kappaB axis. *Exp Mol Med*. (2018) 50:e444. doi: 10.1038/emm.2017.270
53. Lee KS, Park SJ, Kim SR, Min KH, Jin SM, Puri KD, et al. Phosphoinositide 3-kinase-delta inhibitor reduces vascular permeability in a murine model of asthma. *J Allergy Clin Immunol*. (2006) 118:403–9. doi: 10.1016/j.jaci.2006.04.041
54. Sriskantharajah S, Hamblin N, Worsley S, Calver AR, Hessel EM, Amour A. Targeting phosphoinositide 3-kinase delta for the treatment of respiratory diseases. *Ann N Y Acad Sci*. (2013) 1280:35–9. doi: 10.1111/nyas.12039
55. Park SJ, Lee KS, Kim SR, Min KH, Moon H, Lee MH, et al. Phosphoinositide 3-kinase delta inhibitor suppresses interleukin-17 expression in a murine asthma model. *Eur Respir J*. (2010) 36:1448–59. doi: 10.1183/09031936.00106609
56. Nashed BF, Zhang T, Al-Alwan M, Srinivasan G, Halayko AJ, Okkenhaug K, et al. Role of the phosphoinositide 3-kinase p110delta in generation of type 2 cytokine responses and allergic airway inflammation. *Eur J Immunol*. (2007) 37:416–24. doi: 10.1002/eji.200636401
57. To Y, Ito K, Kizawa Y, Failla M, Ito M, Kusama T, et al. Targeting phosphoinositide-3-kinase-delta with theophylline reverses corticosteroid insensitivity in chronic obstructive pulmonary disease. *Am J Respir Crit Care Med*. (2010) 182:897–904. doi: 10.1164/rccm.200906-0937OC
58. Yoo EJ, Ojiaku CA, Sunder K, Panettieri RA Jr. Phosphoinositide 3-kinase in asthma: novel roles and therapeutic approaches. *Am J Respir Cell Mol Biol*. (2017) 56:700–7. doi: 10.1165/rcmb.2016-0308TR
59. Jefferies CA. Regulating IRFs in IFN driven disease. *Front Immunol*. (2019) 10:325. doi: 10.3389/fimmu.2019.00325
60. Sarkar SN, Peters KL, Elco CP, Sakamoto S, Pal S, Sen GC. Novel roles of TLR3 tyrosine phosphorylation and PI3 kinase in double-stranded RNA signaling. *Nat Struct Mol Biol*. (2004) 11:1060–7. doi: 10.1038/nsmb847
61. Guiducci C, Ghirelli C, Marloie-Provost MA, Matray T, Coffman RL, Liu YJ, et al. PI3K is critical for the nuclear translocation of IRF-7 and type I IFN production by human plasmacytoid dendritic cells in response to TLR activation. *J Exp Med*. (2008) 205:315–22. doi: 10.1084/jem.20070763
62. Aksoy E, Taboubi S, Torres D, Delbauve S, Hachani A, Whitehead MA, et al. The p110delta isoform of the kinase PI(3)K controls the subcellular compartmentalization of TLR4 signaling and protects from endotoxic shock. *Nat Immunol*. (2012) 13:1045–54. doi: 10.1038/ni.2426
63. Singanayagam A, Loo SL, Calderazzo MA, Finney LJ, Trujillo Torralbo MB, Bakhsooliani E, et al. Anti-viral immunity is impaired in COPD patients with frequent exacerbations. *Am J Physiol Lung Cell Mol Physiol*. (2019) 317:L893–903. doi: 10.1152/ajplung.00253.2019
64. Wark PA, Johnston SL, Bucchieri F, Powell R, Puddicombe S, Laza-Stanca V, et al. Asthmatic bronchial epithelial cells have a deficient innate immune response to infection with rhinovirus. *J Exp Med*. (2005) 201:937–47. doi: 10.1084/jem.20041901
65. Edwards MR, Regamey N, Vareille M, Kieninger E, Gupta A, Shoemark A, et al. Impaired innate interferon induction in severe therapy resistant atopic asthmatic children. *Mucosal Immunol*. (2013) 6:797–806. doi: 10.1038/mi.2012.118
66. Ge Q, Moir LM, Trian T, Niimi K, Poniris M, Shepherd PR, et al. The phosphoinositide 3'-kinase p110delta modulates contractile protein production and IL-6 release in human airway smooth muscle. *J Cell Physiol*. (2012) 227:3044–52. doi: 10.1002/jcp.23046
67. Jude JA, Tirumurugan KG, Kang BN, Panettieri RA, Walseth TF, Kannan MS. Regulation of CD38 expression in human airway smooth muscle cells: role of class I phosphatidylinositol 3 kinases. *Am J Respir Cell Mol Biol*. (2012) 47:427–35. doi: 10.1165/rcmb.2012-0025OC
68. Farghaly HS, Blagbrough IS, Medina-Tato DA, Watson ML. Interleukin 13 increases contractility of murine tracheal smooth muscle by a phosphoinositide 3-kinase p110delta-dependent mechanism. *Mol Pharmacol*. (2008) 73:1530–7. doi: 10.1124/mol.108.045419
69. Kim SR, Lee KS, Park HS, Park SJ, Min KH, Moon H, et al. HIF-1alpha inhibition ameliorates an allergic airway disease via VEGF suppression in bronchial epithelium. *Eur J Immunol*. (2010) 40:2858–69. doi: 10.1002/eji.200939948
70. Whitehead MA, Bombardieri M, Pitzalis C, Vanhaesebroeck B. Isoform-selective induction of human p110delta PI3K expression by TNFalpha: identification of a new and inducible PIK3CD promoter. *Biochem J*. (2012) 443:857–67. doi: 10.1042/BJ20112214

**Conflict of Interest:** KK received research grants from MSD Life Science Foundation and Public Interest Incorporated Foundation that were unrelated to the submitted work.

The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2020 Fujita, Kan-o, Tonai, Yamamoto, Ogawa, Fukuyama, Nakanishi and Matsumoto. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.