

Protein family review

## Inhibitor of apoptosis proteins and their relatives: IAPs and other BIRPs

Anne M Verhagen, Elizabeth J Coulson and David L Vaux

Address: The Walter and Eliza Hall Institute of Medical Research, Post Office, Royal Melbourne Hospital, Victoria 3050, Australia.

Correspondence: Anne M Verhagen. E-mail: verhagen@wehi.edu.au

Published: 5 July 2001

*Genome Biology* 2001, **2**(7):reviews3009.1–3009.10

The electronic version of this article is the complete one and can be found online at <http://genomebiology.com/2001/2/7/reviews/3009>

© BioMed Central Ltd (Print ISSN 1465-6906; Online ISSN 1465-6914)

### Summary


Apoptosis is a physiological cell death process important for development, homeostasis and the immune defence of multicellular animals. The key effectors of apoptosis are caspases, cysteine proteases that cleave after aspartate residues. The inhibitor of apoptosis (IAP) family of proteins prevent cell death by binding to and inhibiting active caspases and are negatively regulated by IAP-binding proteins, such as the mammalian protein DIABLO/Smac. IAPs are characterized by the presence of one to three domains known as baculoviral IAP repeat (BIR) domains and many also have a RING-finger domain at their carboxyl terminus. More recently, a second group of BIR-domain-containing proteins (BIRPs) have been identified that includes the mammalian proteins Bruce and Survivin as well as BIR-containing proteins in yeasts and *Caenorhabditis elegans*. These Survivin-like BIRPs regulate cytokinesis and mitotic spindle formation. In this review, we describe the IAPs and other BIRPs, their evolutionary relationships and their subcellular and tissue localizations.

### Gene organization and evolutionary history

The inhibitor of apoptosis proteins (IAPs) were originally identified in baculoviruses, where they provide a mechanism for enhancing viral propagation through inhibition of defensive apoptosis by host insect cells [1,2]. Cellular IAPs were subsequently described in insects and vertebrates [3-9]. In recent times, it has become apparent that there is a second group of BIR-domain-containing proteins (BIRPs) carried by organisms such as *Caenorhabditis elegans* and yeasts as well as mammals and insects that can be distinguished from IAPs both by function and by structural features of their BIR domains (reviewed by Miller [10]). The domain structures of various BIRPs is shown in Figure 1a, and the relationship of different BIR domains to each other is illustrated by the phylogenetic tree in Figure 1b and an alignment of a selection of BIR domains is given in Figure 1c. Table 1 lists the chromosomal localizations of the genes encoding human BIRPs as well as their tissue expression patterns and the disease situations in which alterations of the genes have been observed.

The first group of BIRPs encompasses those that inhibit cell death; they are appropriately called IAPs. These proteins have between one and three BIR domains and also often have a RING-finger domain. IAPs have been identified in several multicellular organisms from *Drosophila* to mammals, but are not present in plants, yeasts, protozoans or *C. elegans*. The close relationship between baculoviral IAPs and insect IAPs suggests that baculoviral IAPs may have been acquired through gene transfer from host insect cells [11]. The BIR domains of IAPs can be grouped into several subtypes (Figure 1b). The three BIR domains - BIR1, BIR2 and BIR3 - of human XIAP, cIAP-1 and cIAP-2 fall into three different subgroups, suggesting gene duplication of an ancestral IAP gene encoding three BIR domains and a RING finger.

The gene encoding murine XIAP/MIHA/hILP/BIRC4 spans approximately 20 kilobases (kb) and the protein is encoded by six exons [12]. The initiation codon, the BIR1 and BIR2 domains and half of the BIR3 domain is encoded by exon 1.



**Figure 1**

The rest of the BIR3 domain is encoded by exons 2 and 3; exons 4 and 5 encode the following non-structural region and exon 6 encodes the carboxy-terminal RING-finger domain and stop codon. The structures of the genes for cIAP-1/MIHB/hiap2/BIRC2 and cIAP-2/MIHC/hiap1/BIRC3 are reportedly similar to that of XIAP (referred to as unpublished data by Farahani *et al.* [12]). Mammalian cIAP-1 and cIAP-2 are very similar to each other and their genes are tightly linked (about 12 kb apart), suggesting a relatively recent gene-duplication event [13]. Both proteins have three BIR domains, a caspase recruitment domain (CARD) and a RING finger. The recently identified IAP ML-IAP/LIVIN/KIAP/BIRC7 has only one BIR domain which is most highly related to the BIR3 domains of cIAP-1, cIAP-2 and XIAP, particularly in having an  $\alpha$ -helical extension carboxy-terminal to the BIR domain [14-16].

There are thought to be six tightly linked *NAIP* genes in mice and humans [17-19]. The BIR domains of NAIPs are more distantly related to the BIR domains of the other mammalian IAPs (see Figure 1b) and NAIPs do not have a RING-finger domain but do have a nucleotide-binding domain at their carboxyl terminus [5,20]. There are two *Drosophila* IAPs, DIAP1 and DIAP2, which have two or three BIR domains, respectively and which each have a carboxy-terminal RING-finger domain [4]. Another insect IAP, SfiAP from *Spodoptera frugiperda*, has also recently been described, with two BIR domains and a carboxy-terminal RING-finger domain [11]. There are several baculoviral IAPs; most have two BIR domains and a carboxy-terminal RING-finger domain [1,2].

The second group of BIRPs includes mammalian Survivin/BIRC5 and Bruce/BIRC6, *C. elegans* BIR-1 and BIR-2 (shown as CeBIR-1 and CeBIR-2 in Figure 1), yeast Spbir1P and ScBIR1P and *Drosophila* proteins d-Bruce and Deterin [21-25]. Apart from their BIR domains, these proteins are otherwise highly variable in size and structure. They have slightly larger BIR domains than those of the IAPs (see Figure 1c) and there is a conserved intron after the invariant

glycine-encoding codon in the BIR-domain-encoding region that is not present in IAP genes. The presence of the Survivin-like BIRPs in a wide range of organisms and their conserved function suggests that they represent the earliest BIRPs. It is possible that, following a gene-duplication event, the BIR domains in IAPs evolved to have a different function, namely to interact with and inhibit caspases. The genes for both murine and human Survivin have been described, and both comprise four exons, with exons 2 and 3 encoding the BIR domain [26-28].

### Characteristic structural features

BIR domains are characterized by a number of invariant amino acids, including three conserved cysteines and one conserved histidine residue within the sequence  $CX_2CX_{16}HX_{6-8}C$  (Figure 1c). Within IAPs, BIR domains are typically about 70 amino acids long, but they can be more than 100 amino acids long in other BIRPs. The structures of the cIAP-1 BIR3 domain and the XIAP BIR2 and BIR3 domains are very similar, indicating that BIR domains typically comprise a series of four or five  $\alpha$  helices and a three-stranded  $\beta$  sheet with a single zinc ion coordinated by the conserved cysteine and histidine residues [29-33].

RING fingers, a type of zinc finger, are present in diverse proteins. A carboxy-terminal RING finger domain is present in most of the IAPs and has, for XIAP and c-IAP-1, been shown to have ubiquitin protein ligase activity, directly regulating self-ubiquitination and degradation [34]. RING domains are characterized by the presence of a set of invariant metal-binding residues ( $C_3HC_4$ ) that coordinate two zinc ions [35]. The equine herpes virus protein RING has been shown to consist of an amphipathic  $\alpha$  helix next to a triple-stranded  $\beta$  sheet [36].

The c-IAP1 and c-IAP2 proteins have caspase recruitment domains (CARDs) between their three BIR domains and the RING-finger domain. The name relates to the ability of CARDs within adaptor proteins such as Apaf-1 to interact

### Figure 1

Structure and evolution of BIRPs. **(a)** Domain structures of BIRPs in mammals (humans), insects, nematodes and yeast. The length of each protein (in amino acids) is shown on the right. IAPs have type 1 BIR domains (dark blue), whereas type 2 BIR domains (light blue) are found in Survivin-like BIRPs. **(b)** The evolutionary relationship of BIR domains. A phylogenetic tree was generated using the parsimony option of the PHYLIP Phylogeny Inference Package (version 3.5c) [72], from BIR domains aligned by the Clustal W1.7 program. The accession numbers and regions used in the analysis are shown in Table 2. Default settings were used; the order of the sequences was jumbled 10 times and the degree of certainty for each node (the bootstrap value, shown on the Figure) was calculated from 100 random replicates. All bootstrap values greater than 50 and selected others are shown. The values for placement of ML-IAP with NAIP BIR3 and BIR2 domains are very low, and trees calculated using distance methods placed ML-IAP BIR with the BIR3 domains of XIAP, cIAP-1 and cIAP-2. Also, note that the alignment used to generate this tree does not include the carboxy-terminal  $\alpha$ -helical extension that is conserved between the BIR3 domains of XIAP, cIAP-1, cIAP-2 and ML-IAP but not NAIP. **(c)** Two types of BIR domain. An alignment of a selection of BIR domains illustrates the difference between BIR domains present within classical IAPs (top four sequences) and within other BIRPs (bottom four). Aligned regions are as given in Table 2. Amino acids that are conserved in all BIR domains are highlighted. Type 1 BIR domains are approximately 70 amino acids long, whereas type 2 BIR domains are longer.

**Table 1**

<b>Properties of human BIRPs</b>					
Human BIRPs and alternative names	Size (kDa)	Map position of gene	Expression of mRNA	Diseases associated with chromosomal rearrangements, gene amplification or deletion	References
XIAP/MIHA/hILP/BIRC4	57	Xq25	Detected in all adult tissues examined		[5-7,9,12]
cIAP-1/MIHB/HIAP2/BIRC2	70	11q22-q23	Detected in all adult tissues examined, with highest levels in thymus, testis and ovary		[3,5-7,13]
cIAP-2/MIHC/HIAP1/BIRC3	68	11q22-q23	Detected in all adult tissues examined, with highest levels in spleen and thymus	In marginal zone cell lymphomas of the mucosa-associated lymphoid tissue, <i>API2</i> , the gene encoding cIAP-2, is often rearranged together with the novel gene <i>MLT</i> at 18q21 resulting in an <i>API2-MLT</i> fusion	[3,5-7,13]
LIVIN/ML-IAP/KIAP/BIRC7	31	20q13.3	Detected in embryonic tissue and some adult tissues. Elevated levels in cancer cell lines, particularly melanoma cell lines	The region is frequently amplified in melanomas, colon, lung and colorectal malignancies	[14-16]
NAIP/BIRC1 (6 genes)	156	5q13.1	Detected only in liver and placenta by northern blot analysis but was detected in spinal cord mRNA by RT-PCR	Deletions of the <i>NAIP</i> genes often occur in individuals with spinal muscular atrophy but deletions in the nearby gene <i>SMN</i> are the most likely cause of disease	[5,20]
Survivin/BIRC5	17	17q25	Cell-cycle-dependent expression, restricted to tissues in which there are dividing cells; for example, embryonic tissue, solid tumors and transformed cell lines. Expression is absent from most normal adult tissue		[21,26-28]
BRUCE/BIRC6	528	2p21-p22	Expression of mRNA is detectable in most adult tissues, with high levels in brain and kidney		[22]

with CARDs within some initiator caspases (see later), such as caspase 9 [37]. The CARD fold is related to other protein-protein interaction domains found in proteins involved in cell death and elsewhere, such as the death domain, the death effector domain and the pyrin domain. The only IAPs described to date that possess a CARD are cIAP-1 and cIAP-2, and the function of this domain in these proteins is not known. Protein-protein interactions of cIAP-1 or cIAP-2 CARDs with CARDs of other proteins have not been described, nor have the CARDs been shown to self-associate. The structure of cIAP-1 and cIAP-2 CARDs have not yet been described, although CARDs from other proteins have been shown to comprise six or seven tightly packed anti-parallel  $\alpha$  helices, with both charged regions and hydrophobic regions mediating CARD-CARD interactions [38-42].

### Localization and function

#### BIRPs involved in inhibiting cell death: IAPs

The tissue and subcellular localization of mammalian IAPs varies (see Table 1) and may be important in determining the

relative contribution of different IAPs to cell-death regulation in different cell types and in response to different apoptotic stimuli.

XIAP appears to be widely expressed [5,6]. It has a cytoplasmic location and has been reported to inhibit cell death in response to a variety of apoptotic stimuli including ultraviolet irradiation, tumor necrosis factor (TNF), Fas ligand, and a number of toxic drugs (reviewed by LaCasse *et al.* [43]). XIAP very efficiently interacts with and inhibits active caspases 3, 7 and 9 [44-46]. Interaction of XIAP via its RING-finger domain with bone morphogenetic protein (BMP) type I receptors has also been reported, which would presumably enable some XIAP to localize to the plasma membrane [47]. A role for XIAP in regulating the BMP receptor signaling pathway by linking the receptor to downstream signaling molecule TAB1 has been proposed.

The cIAP-1 and cIAP-2 proteins were initially identified in a complex with TNF receptor 2, an indirect association resulting from direct interaction with TNF-receptor-associated

Table 2

## Accession numbers of BIRPs shown in Figure 1b and regions used in the phylogenetic analysis

Protein name	Accession number (GenBank)	Amino acid residues used for each BIR domain	Species
XIAP	P98170	BIR1 21-93, BIR2 159-230, BIR3 258-330	Human
cIAP-1	XP_006266	BIR1 41-113, BIR2 179-250, BIR3 264-336	Human
cIAP-2	XP_006267	BIR1 24-96, BIR2 164-235, BIR3 250-322	Human
ML-IAP	NP_071444	BIR 82-154	Human
NAIP	NP_004527	BIR1 55-127, BIR2 154-227, BIR3 273-345	Human
Survivin	XP_008623	BIR 10-87	Human
Bruce	NP_031592	BIR1 256-333	Human
DIAP1	AAF49548	BIR1 39-110, BIR2 221-293	<i>Drosophila</i>
DIAP2	Q24307	BIR1 4-76, BIR2 108-179, BIR3 207-279	<i>Drosophila</i>
dBRUCE	AAF54520	BIR1 243-320	<i>Drosophila</i>
Deterin	AAF55399	BIR1 22-100	<i>Drosophila</i>
SfiAP	AAF35285	BIR1 95-166, BIR2 205-277	<i>Spodoptera frugiperda</i>
OpIAP	NP_046191	BIR1 13-84, BIR2 106-178	<i>Orgyia pseudotsugata</i> nuclear polyhedrosis virus
CplIAP	P41436	BIR1 2-73, BIR2 103-175	<i>Cydia pomonella</i> granulosis virus
CiIAP	P47732	BIR1 32-108	Chilo iridescent virus
AcIAP	D36828	BIR1 24-96, BIR2 126-199	<i>Autographa californica</i> nuclear polyhedrosis virus
OpBIRP2	NP_046197	BIR1 19-91, BIR2 121-193	<i>Orgyia pseudotsugata</i> nuclear polyhedrosis virus
BmBIRP	NP_047432	BIR1 24-96, BIR2 125-199	<i>Bombyx mori</i> nuclear polyhedrosis virus
CeBIRP2/ CeBIR-2	U72208	BIR1 18-97, BIR2 162-240	<i>C. elegans</i>
CeBIRP1/ CeBIR-1	T37471	BIR 12-86	<i>C. elegans</i>
Spbir1P	T41649	BIR1 17-98, BIR2 122-193	<i>Schizosaccharomyces pombe</i>
ScBIR1P	NP_012622	BIR1 12-116, BIR2 145-240	<i>Saccharomyces cerevisiae</i>

factors (TRAFs) 1 and 2 involving the BIR and TRAF domains of the respective proteins [3]. The expression of cIAP-1 and cIAP-2 is increased following activation of the NF- $\kappa$ B transcription factor by the TNF receptor, and these IAPs may have a role in protecting cells from TNF-induced apoptosis by reducing the amount of caspase 8 activation [48]. Exactly how they do this is unclear, because caspase 8 can not be directly inhibited by cIAP-1 or -2 or by any other known IAPs [49]. Perhaps they act by inhibiting downstream caspases such as caspase 3, preventing the feedback loop involved in the further activation of upstream caspase 8. The cIAP-1 and cIAP-2 proteins have been shown directly to inhibit the activity of caspases 3 and 7 [49]. Although cIAP-1 and cIAP-2 can be found within the TNF-R complex, and they are presumably localized in part to the cell membrane, it is not clear what proportion of total cellular cIAP-1 or cIAP-2 this represents, and when transiently transfected into cells, cIAP-1 has a perinuclear localization [50].

The recently described mammalian IAP ML-IAP is detectable in embryonic tissue, selected adult tissues and several cancer cell lines, particularly in melanoma cell lines (see Table 1) [14-16]. One study reported expression

predominantly in the nucleus but also in filamentous structures in the cytoplasm, whereas only cytoplasmic expression was reported in another study. Although ML-IAP has only one BIR domain, it is reported to interact with and inhibit both the initiator caspase 9 and effector caspases 3 and 7; it inhibits cell death induced through death receptors, by over-expression of the cell-death pathway proteins FADD, Bax, RIP, RIP3 and DR3 or in response to various toxic drugs.

The first *NAIP* gene was identified as a candidate gene defective in spinal muscular atrophy [5,20]. It is now clear that deletion of a neighboring gene encoding the protein survival motor neuron (SMN) is the cause of the disease [51], but it is possible that loss of functional NAIP may contribute to the severity of the disease. NAIP is reported to inhibit cell death in response to serum withdrawal, menadione and TNF, but because the *NAIP* cDNA used in these experiments does not correspond to any of the *NAIP* genes and may represent sequences derived from a number of different *NAIP* genes, the exact activities of NAIP proteins are unclear.

The strongest evidence for regulation of developmental cell death by IAPs comes from studies in *Drosophila*, where loss


of DIAP1 results in extensive early embryonic cell death and a corresponding increase in caspase activity [52]. An equivalent study in mammals is still required to establish the role of these proteins in mammalian developmental cell death (see Frontiers section). A punctate perinuclear pattern of expression has been described for DIAP2 when overexpressed in insect cells [53].

**BIRPs involved in the regulation of cell division**


Although originally described as a protein that could inhibit cell death [21], Survivin functions primarily in the regulation of cell division, a role conserved in the yeast and *C. elegans* BIRPs, whose BIR domains closely resemble that of Survivin [23,24,54-59]. Survivin is expressed only during mitosis (see Table 1). Subcellular localization has shown that Survivin is a chromosome passenger protein: that is, it

is initially associated with the centromeres, but at the metaphase-anaphase transition it leaves the centromeres and remains in the spindle midzone [56-58,60]. It can be found in the mid-body at telophase, after which it is ubiquitinated and degraded. Consistent with its subcellular localization, elimination of Survivin by homologous recombination results in mouse embryos that are unable to survive beyond day 5 because of failure of cytokinesis [56].

The *C. elegans* proteins BIR-1 and BIR-2 are also involved in the regulation of cytokinesis [23,57]. BIR-1, like Survivin, is expressed only during cell division and is a chromosome passenger protein. The phenotype of BIR-1-deficient embryos is identical to that of embryos deficient for the Aurora-like kinase AIR-2, and BIR-1 is required for localization of AIR-2. Like Survivin and the *C. elegans* BIRPs, yeast



**Figure 2**  
 The role of IAPs in the regulation of cell death. Apoptosis induced by TNF or Fas ligand via death receptors (DR) involves adaptor-protein-mediated recruitment and activation of initiator caspases, such as caspase 8 (C8), and the subsequent activation of downstream caspases such as caspase 3 (C3) and 7 (not shown). In cell death induced via stress pathways such as irradiation or drugs, cytochrome c (Cyt c) and DIABLO are released from the mitochondria. Cytochrome c binds to the carboxyl terminus of the adaptor protein Apaf-1, allowing it to unfold, interact with inactive caspase 9 (Pro-C9) and promote its oligomerization and autoprocessing to give active 3 (C3); active caspase 9 then can activate downstream caspases. Pro-survival members of the Bcl-2 family inhibit cell death via stress pathways and can prevent release of cytochrome c and DIABLO from mitochondria. The downstream caspases cleave cellular substrates (such as poly (ADP-ribose) polymerase (PARP) and inhibitor of caspase-activated DNase (ICAD)), resulting in many morphological changes and culminating in cell death. IAPs prevent cell death by interacting with and inhibiting active caspases, whereas the IAP antagonist DIABLO can prevent these interactions.



**Figure 3**

Mechanism of caspase inhibition by IAPs and its prevention by the IAP antagonist DIABLO/Smac. **(a)** The amino terminus of the p10 subunit of caspase 9, revealed following autoprocessing, interacts tightly within a groove of the BIR3 domain of XIAP. **(b)** The linker region upstream of the BIR2 domain of XIAP interacts tightly with the catalytic site of caspase 3 (and caspase 7; C indicates the catalytic cysteine residue), resulting in effective caspase inhibition. A less significant interaction of the amino terminus of the small subunit of caspase 3 within a groove of the BIR2 domain has been proposed. **(c)** The amino terminus of DIABLO (red) is similar to that of caspase 9 and competes for the exact same interaction site within the BIR3 domain. **(d)** A similar groove within the BIR2 domain of XIAP is believed to mediate interaction with DIABLO and provides a mechanism for DIABLO to remove caspase 3 from XIAP.

proteins ScBIR1P and Spbir1P have a role in regulating cell division [24], and as yeasts do not have caspases, a role in cell-death regulation can be excluded.

Mammalian Bruce/Apollon has a punctate pattern of cellular expression and colocalizes with the Golgi marker protein TGN38. Expression in dendrites and axons of primary neurons has also been detected [22]. While Bruce has a BIR domain at its amino terminus, it also has ubiquitin-conjugating activity as a result of a ubiquitin-conjugating enzyme (UBC) domain at the carboxyl terminus of the protein. The BIR domain of Bruce is most similar to the BIR domains of Survivin and other BIRPs that are involved in regulating cell division, although the function of Bruce is unknown.

#### Mechanism of caspase inhibition

Caspases are produced as inactive zymogens that are processed into an active form following cell-death stimuli.

The active caspase is believed to be a heterotetrameric complex generated from two zymogen monomers. Cell-death pathways involve the sequential activation of initiator and effector caspases (see Figure 2). Activation of initiator caspases such as caspases 9 and 8 is facilitated by adaptor proteins such as Apaf-1 or FADD, and when activated, caspases 8 and 9 can process and activate downstream effector caspases such as caspases 3 and 7. IAPs can inhibit some active caspases, whereas the mammalian protein DIABLO antagonizes IAP function [61-63] (see Figures 2,3).

The BIR3 domain of XIAP interacts with and inhibits active processed caspase 9 [46,63,64]. The amino terminus of DIABLO and that of the p10 subunit of active caspase 9 - which is revealed only after autoprocessing - are similar and compete for exactly the same site within a groove of the BIR3 domain of XIAP [64-69] (Figure 3). The interactions of DIABLO and caspase 9 with XIAP are thus mutually exclusive.

The amino termini of the *Drosophila* proteins Grim, Reaper and Hid are similar to the amino terminus of DIABLO within the first four amino acids, and this region is responsible for interaction with IAPs [67-69].

The main point of contact between caspases 3 and 7 and XIAP involves the linker region immediately upstream of the BIR2 domain of XIAP [31-33,70] (Figure 3). The linker forms a direct contact with the catalytic site of the caspases, thereby blocking their activity. An additional interaction of the amino terminus of the processed p10 fragment of caspase 3 within the groove of the BIR2 domain has been proposed [33] and, although dispensable for caspase interaction and inhibition by XIAP, the IAP antagonist DIABLO/Smac is thought to compete for the same groove, presumably leveraging out caspase 3 and dislodging it from the IAP.

DIABLO is a dimeric protein with two amino termini that can potentially interact with BIR domains [65]. Although DIABLO is able to interact with individual BIR2 and BIR3 domains, albeit with higher affinity for the BIR3, the type of interaction favored with full-length XIAP is not known. For example, it is possible that a DIABLO dimer interacts simultaneously via its two amino termini with both the BIR2 and BIR3 domains on a single XIAP molecule, or that the two amino termini interact with BIR domains on different XIAP molecules with three different possible BIR domain combinations (two BIR2 domains, a BIR2 plus BIR3 domain or two BIR3 domains). Establishing this structurally may be difficult given that the generation of well-folded full-length XIAP has eluded crystallographers so far.

## Frontiers

Our understanding of IAPs and how they inhibit cell death has taken enormous strides in the last few months. At a structural level, we now know how IAPs interact with and inhibit caspases and how this interaction can be regulated by IAP antagonists such as DIABLO. Much research is still required, however, to define the different roles for different mammalian IAP proteins in the face of different cell-death stimuli. It is likely that there is some redundancy between family members. Indeed, MIHA-deficient mice are without phenotype, with the exception of apparently increased expression levels of cIAP-1 and cIAP-2, suggesting some compensation [71]. It may be necessary to generate mice deficient for more than one IAP in order to establish the role of IAPs in mammals. So far, only one mammalian IAP antagonist, DIABLO, has been described, but given that there are three such proteins in *Drosophila*, other mammalian IAP antagonists are almost certain to exist. The second group of BIRPs discussed in this review are clearly involved in regulating cytokinesis, although the exact mechanism is not known. Direct interactions of BIR domains from these proteins with other cellular proteins have not yet been reported but are likely to provide some clues to how these proteins function.

## Acknowledgements

We would like to acknowledge John Silke for helpful discussions.

## References

- Birnbaum MJ, Clem RJ, Miller LK: **An apoptosis inhibiting gene from a nuclear polyhedrosis virus encoding a polypeptide with cys/his sequence motif.** *J Virol* 1994, **68**:2521-2528.  
The first description of OplAP.
- Crook NE, Clem RJ, Miller LK: **An apoptosis inhibiting baculovirus gene with a zinc finger like motif.** *J Virol* 1993, **67**:2168-2174.  
The first description of an IAP; baculoviral CplAP was identified by its ability to prevent cell death of insect cells transfected with *Autographa californica* nuclear polyhedrosis virus DNA from which the anti-apoptotic gene p35 had been deleted.
- Rothe M, Pan MG, Henzel WJ, Ayres TM, Goeddel DV: **The TNFR2-TRAF signaling complex contains two novel proteins related to baculoviral-inhibitor of apoptosis proteins.** *Cell* 1995, **83**:1243-1252.  
The first description of mammalian cIAP-1 and cIAP-2, identifying them as proteins that interact with the TNF receptor via direct interaction with TRAF1 and TRAF2.
- Hay BA, Wassarman DA, Rubin GM: ***Drosophila* homologs of baculovirus inhibitor of apoptosis proteins function to block cell death.** *Cell* 1995, **83**:1253-1262.  
The first description of *Drosophila* IAPs and demonstration that overexpression of DIAP1 in the fly eye can prevent both normal cell death and that induced by overexpression of the pro-death proteins Reaper and Hid.
- Liston P, Roy N, Tamai K, Lefebvre C, Baird S, Chertonhorvat G, Farahani R, Mclean M, Ikeda JE, Mackenzie A, Korneluk RG: **Suppression of apoptosis in mammalian cells by NAIP and a related family of IAP genes.** *Nature* 1996, **379**:349-353.  
One of the first descriptions of mammalian IAPs cIAP-1, cIAP-2, XIAP and NAIP and of *Drosophila* DIAP1.
- Uren AG, Pakusch M, Hawkins CJ, Puls KL, Vaux DL: **Cloning and expression of apoptosis inhibitory protein homologs that function to inhibit apoptosis and/or bind tumor necrosis factor receptor-associated factors.** *Proc Natl Acad Sci USA* 1996, **93**:4974-4978.  
One of the first descriptions of mammalian cIAP-1, cIAP-2 and XIAP.
- Duckett CS, Nava VE, Gedrich RW, Clem RJ, Vandongen JL, Gilfillan MC, Shiels H, Hardwick JM, Thompson CB: **A conserved family of cellular genes related to the baculovirus iap gene and encoding apoptosis inhibitors.** *EMBO J* 1996, **15**:2685-2694.  
One of the first descriptions of *Drosophila* and mammalian IAPs (DIAP1 and XIAP).
- You MJ, Ku PT, Hrdlickova R, Bose HR: **Ch-iap1, a member of the inhibitor-of-apoptosis protein family, is a mediator of the antiapoptotic activity of the v-rel oncoprotein.** *Mol Cell Biol* 1997, **17**:7328-7341.  
A report of an IAP in chickens.
- Human Gene Nomenclature Database** [<http://www.gene.ucl.ac.uk/cgi-bin/nomenclature/searchgenes.pl>]  
Contains information on the new nomenclature for IAPs/BIRPs (BIRC1,2, and so on).
- Miller LK: **An exegesis of IAPs: salvation and surprises from BIR motifs.** *Trends Cell Biol* 1999, **9**:323-328.  
A review on the different types of BIR domains and the two emerging functions of BIRPs.
- Huang QH, Deveraux QL, Maeda S, Salvesen GS, Stennicke HR, Hammock BD, Reed JC: **Evolutionary conservation of apoptosis mechanisms: lepidopteran and baculoviral inhibitor of apoptosis proteins are inhibitors of mammalian caspase-9.** *Proc Natl Acad Sci USA* 2000, **97**:1427-1432.  
Describes SfIAP, a baculoviral IAP that is highly related to insect IAPs.
- Farahani R, Fong WG, Korneluk RG, Mackenzie AE: **Genomic organization and primary characterization of miap-3 - the murine homologue of human X-linked IAP.** *Genomics* 1997, **42**:514-518.  
Description of the murine XIAP gene.
- Rajcanseparovic E, Liston P, Lefebvre C, Korneluk RG: **Assignment of human inhibitor of apoptosis protein (iap) genes xiap, hiap-1, and hiap-2 to chromosomes Xq25 and 11q22-q23 by fluorescence in situ hybridization.** *Genomics* 1996, **37**:404-406.  
A chromosomal localization analysis of the human IAP genes.



14. Vucic D, Stennicke HR, Pisabarro MT, Salvesen GS, Dixit VM: **ML-IAP, a novel inhibitor of apoptosis that is preferentially expressed in human melanomas.** *Curr Biol* 2000, **10**:1359-1366.  
This paper and [15,16] describe ML-IAP/LIVIN/BIRC7, a mammalian IAP with a single BIR domain and a RING-finger motif.
15. Lin JH, Deng G, Huang QH, Morser J: **KIAP, a novel member of the inhibitor of apoptosis protein family.** *Biochem Biophys Res Comm* 2000, **279**:820-831.  
See [14].
16. Kasof GM, Gomes BC: **Livin, a novel inhibitor of apoptosis protein family member.** *J Biol Chem* 2001, **276**:3238-3246.  
See [14].
17. Yaraghi Z, Korneluk RG, Mckenzie A: **Cloning and characterization of the multiple murine homologues of NAIP (neuronal apoptosis inhibitory protein).** *Genomics* 1998, **51**:107-113.  
This paper and [18,19] report the six different loci of murine NAIP genes and their various transcripts.
18. Huang SD, Scharf JM, Growney JD, Endrizzi MG, Dietrich WF: **The mouse Naip gene cluster on Chromosome 13 encodes several distinct functional transcripts.** *Mamm Genome* 1999, **10**:1032-1035.  
See [17].
19. Yamamoto K, Sakai H, Hadano S, Gondo Y, Ikeda JE: **Identification of two distinct transcripts for the neuronal apoptosis inhibitory protein gene.** *Biochem Biophys Res Comm* 1999, **264**:998-1006.  
See [17].
20. Roy N, Mahadevan MS, Mclean M, Shutler G, Yaraghi Z, Farahani R, Baird S, Besnerjohnston A, Lefebvre C, Kang XL, et al.: **The gene for neuronal apoptosis inhibitory protein is partially deleted in individuals with spinal muscular atrophy.** *Cell* 1995, **80**:167-178.  
The first description of a mammalian NAIP.
21. Ambrosini G, Adida C, Altieri DC: **A novel anti-apoptosis gene, survivin, expressed in cancer and lymphoma.** *Nat Med* 1997, **3**:917-921.  
The first description of Survivin, a BIRP with a single BIR domain.
22. Hauser HP, Bardroff M, Pyrowolakis G, Jentsch S: **A giant ubiquitin-conjugating enzyme related to iap apoptosis inhibitors.** *J Cell Biol* 1998, **141**:1415-1422.  
The first description of Bruce, a large protein with an amino-terminal BIR domain and a UBC motif at the carboxyl terminus.
23. Fraser AG, James C, Evan GI, Hengartner MO: **Caenorhabditis elegans inhibitor of apoptosis protein (IAP) homologue BIR-1 plays a conserved role in cytokinesis.** *Curr Biol* 1999, **9**:292-301.  
The authors observed that worm embryos lacking BIR-1 are unable to complete cytokinesis.
24. Uren AG, Beilharz T, O'Connell MJ, Bugg SJ, van Driel R, Vaux DL, Lithgow T: **Role for yeast inhibitor of apoptosis (IAP)-like proteins in cell division.** *Proc Natl Acad Sci USA* 1999, **96**:10170-10175.  
Yeast deficient in BIRPs have defects in spore formation and/or division after germination.
25. Jones G, Jones D, Zhou L, Steller H, Chu YX: **Deterin, a new inhibitor of apoptosis from Drosophila melanogaster.** *J Biol Chem* 2000, **275**:22157-22165.  
First description of *Drosophila* Deterin, a single BIR-domain-containing protein.
26. Li FZ, Altieri DC: **The cancer antiapoptosis mouse survivin gene: characterization of locus and transcriptional requirements of basal and cell cycle-dependent expression.** *Cancer Res* 1999, **59**:3143-3151.  
Description of the exon structure and promoter of the mouse *survivin* gene.
27. Mahotka C, Wenzel M, Springer E, Gabbert HE, Gerharz CD: **Survivin-Delta Ex3 and survivin-2B: two novel splice variants of the apoptosis inhibitor survivin with different antiapoptotic properties.** *Cancer Res* 1999, **59**:6097-6102.  
Two different isoforms of human Survivin differ in the BIR-domain-containing region; Survivin-2B is less anti-apoptotic.
28. Conway EM, Pollefeyt S, Cornelissen J, DeBaere I, Steiner-Mosonyi M, Ong K, Baens M, Collen D, Schuh AC: **Three differentially expressed survivin cDNA variants encode proteins with distinct antiapoptotic functions.** *Blood* 2000, **95**:1435-1442.  
Description of mouse *survivin* gene and evidence for alternative splicing; only two splice forms inhibit caspase 3 activity.
29. Hinds MG, Norton RS, Vaux DL, Day CL: **Solution structure of a baculoviral inhibitor of apoptosis (IAP) repeat.** *Nat Struct Biol* 1999, **6**:648-651.  
The solution structure of cIAP-1 BIR3 domain.
30. Sun CH, Cai ML, Meadows RP, Xu N, Gunasekera AH, Herrmann J, Wu JC, Fesik SW: **NMR structure and mutagenesis of the third Bir domain of the inhibitor of apoptosis protein XIAP.** *J Biol Chem* 2000, **275**:33777-33781.  
The structure of the BIR3 domain of XIAP reveals that this domain is the minimal region required to inhibit caspase 9.
31. Chai JJ, Shiozaki E, Srinivasula SM, Wu Q, Dataa P, Alnemri ES, Shi YG: **Structural basis of caspase-7 inhibition by XIAP.** *Cell* 2001, **104**:769-780.  
Describes interaction of caspase 7 with an XIAP peptide comprising 18 amino acids from the linker region upstream of the BIR2 domain.
32. Huang YH, Park YC, Rich RL, Segal D, Myszkowski DG, Wu H: **Structural basis of caspase inhibition by XIAP: Differential roles of the linker versus the BIR domain.** *Cell* 2001, **104**:781-790.  
Demonstrates that caspase 7 interaction with XIAP involves the linker region upstream of the BIR2 domain rather than the BIR2 domain itself.
33. Riedl SJ, Renatus M, Schwarzenbacher R, Zhou Q, Sun CH, Fesik SW, Liddington RC, Salvesen GS: **Structural basis for the inhibition of caspase-3 by XIAP.** *Cell* 2001, **104**:791-800.  
This paper demonstrates that the main site of caspase 3 interaction with XIAP involves the linker region upstream of BIR2 domain, although an additional interaction of the amino terminus of the small caspase subunit within the DIABLO/Smac-binding pocket of the BIR2 domain is proposed.
34. Yang Y, Fang SY, Jensen JP, Weissman AM, Ashwell JD: **Ubiquitin protein ligase activity of IAPs and their degradation in proteasomes in response to apoptotic stimuli.** *Science* 2000, **288**:874-877.  
Demonstrating that the RING-finger domain of cIAP-1 and XIAP can catalyze self-ubiquitination of IAPs.
35. Freemont PS: **The RING finger. A novel protein sequence motif related to the zinc finger.** *Ann NY Acad Sci* 1993, **684**:174-192.  
A review of the properties of RING-finger domains.
36. Barlow PN, Luisi B, Milner A, Elliott M, Everett R: **Structure of the C3HC4 domain by 1H-nuclear magnetic resonance spectroscopy. A new structural class of zinc-finger.** *J Mol Biol* 1994, **237**:201-211.  
Describing the structure of a RING-finger domain.
37. Hofmann K, Bucher P, Tschoopp J: **The card domain - a new apoptotic signalling motif.** *Trends Biochem Sci* 1997, **22**:155-156.  
First observation of the CARD domain within adaptor proteins, caspases and IAPs.
38. Chou JJ, Matsuo H, Duan H, Wagner G: **Solution structure of the RAIDD CARD and model for CARD/CARD interaction in caspase-2 and caspase-9 recruitment.** *Cell* 1998, **94**:171-180.  
The structure of RAIDD CARD, the first structure of a CARD, is solved and shown to be similar to the Fas death domain.
39. Qin HX, Srinivasula SM, Wu G, Fernandes-Alnemri T, Alnemri ES, Shi YG: **Structural basis of procaspase-9 recruitment by the apoptotic protease-activating factor 1.** *Nature* 1999, **399**:549-557.  
This paper and [40-42] describe the structure of the Apaf-1 CARD, in the case of [39,40] interacting with the CARD domain from caspase 9.
40. Zhou P, Chou J, Olea RS, Yuan JY, Wagner G: **Solution structure of Apaf-1 CARD and its interaction with caspase-9 CARD: A structural basis for specific adaptor/caspase interaction.** *Proc Natl Acad Sci USA* 1999, **96**:11265-11270.  
See [39].
41. Vaughn DE, Rodriguez J, Lazebnik Y, Joshua-Tor L: **Crystal structure of Apaf-1 caspase recruitment domain: An alpha-helical Greek key fold for apoptotic signaling.** *J Mol Biol* 1999, **293**:439-447.  
See [39].
42. Day CL, Dupont C, Lackmann M, Vaux DL, Hinds MG: **Solution structure and mutagenesis of the caspase recruitment domain (CARD) from Apaf-1.** *Cell Death Diff* 1999, **6**:1125-1132.  
See [39].
43. LaCasse EC, Baird S, Korneluk RG, MacKenzie AE: **The inhibitors of apoptosis (IAPs) and their emerging role in cancer.** *Oncogene* 1998, **17**:3247-3259.  
A review of IAPs and their anti-apoptotic activities.

44. Deveraux QL, Takahashi R, Salvesen GS, Reed JC: **X-linked IAP is a direct inhibitor of cell-death proteases.** *Nature* 1997, **388**:300-304.  
The first description of an interaction between an IAP and a caspase.
45. Takahashi R, Deveraux Q, Tamm I, Welsh K, Assamunt N, Salvesen GS, Reed JC: **A single BIR domain of XIAP sufficient for inhibiting caspases.** *J Biol Chem* 1998, **273**:7787-7790.  
Demonstrates that the region of XIAP encompassing the BIR2 domain is able to inhibit caspases 3 and 7.
46. Deveraux QL, Leo E, Stennicke HR, Welsh K, Salvesen GS, Reed JC: **Cleavage of human inhibitor of apoptosis protein XIAP results in fragments with distinct specificities for caspases.** *EMBO J* 1999, **18**:5242-5251.  
Demonstrates that the BIR1-2 fragment is a specific inhibitor of caspase 3, whereas caspase 9 is inhibited by the BIR3-RING fragment.
47. Yamaguchi K, Nagai S, Ninomiya-Tsuji J, Nishita M, Tamai K, Irie K, Ueno N, Nishida E, Shibuya H, Matsumoto K: **XIAP, a cellular member of the inhibitor of apoptosis protein family, links the receptors to TAB1-TAK1 in the BMP signaling pathway.** *EMBO J* 1999, **18**:179-187.  
Describing interactions of XIAP with both TAB1, which lies upstream of a MAP kinase pathway, and the BMP receptor.
48. Wang CY, Mayo MW, Korneluk RG, Goeddel DV, Baldwin AS: **NF-kappa-B antiapoptosis - induction of TRAF1 and TRAF2 and c-IAP1 and c-IAP2 to suppress caspase-8 activation.** *Science* 1998, **281**:1680-1683.  
Levels of cIAP-1 and cIAP-2 increase following NF- $\kappa$ B activation.
49. Roy N, Deveraux QL, Takahashi R, Salvesen GS, Reed JC: **The cIAP-1 and cIAP-2 proteins are direct inhibitors of specific caspases.** *EMBO J* 1997, **16**:6914-6925.  
Demonstrating that cIAP-1 and cIAP-2 interact with and inhibit caspases 3 and 7.
50. McCarthy JV, Dixit VM: **Apoptosis induced by *Drosophila* reaper and grim in a human system - attenuation by inhibitor of apoptosis proteins (cIAPs).** *J Biol Chem* 1998, **273**:24009-24015.  
Demonstrating the pro-apoptotic effect of *Drosophila* proteins Reaper and Grim in human cells and inhibition by human IAPs.
51. Lefebvre S, Burglen L, Reboullet S, Clermont O, Burlet P, Viollet L, Benichou B, Cruaud C, Millaesseau P, Zeviani M, *et al.*: **Identification and characterization of a spinal muscular atrophy-determining gene.** *Cell* 1995, **80**:155-165.  
Identifying deletions or mutations of the SMN gene as the primary cause of spinal muscular atrophy.
52. Wang SL, Hawkins CJ, Yoo SJ, Muller HAJ, Hay BA: **The *Drosophila* caspase inhibitor DIAP1 is essential for cell survival and is negatively regulated by HID.** *Cell* 1999, **98**:453-463.  
Demonstrates that Hid promotes cell death by preventing inhibition of caspases by DIAP1.
53. Vucic D, Kaiser WJ, Harvey AJ, Miller LK: **Inhibition of reaper-induced apoptosis by interaction with inhibitor of apoptosis proteins (IAPs).** *Proc Natl Acad Sci USA* 1997, **94**:10183-10188.  
*Drosophila* and baculoviral IAPs can prevent Reaper-induced cell death. Their expression results in a shift from cytoplasmic to a punctate perinuclear pattern of Reaper expression, coinciding with the expression pattern of the IAPs.
54. Li FZ, Ambrosini G, Chu EY, Plescia J, Tognin S, Marchisio PC, Altieri DC: **Control of apoptosis and mitotic spindle checkpoint by survivin.** *Nature* 1998, **396**:580-584.  
First evidence for regulation of cell division by BIRPs.
55. Otaki M, Hatano M, Kobayashi K, Ogasawara T, Kuriyama T, Tokuhisa T: **Cell cycle-dependent regulation of TIAP/m-survivin expression.** *Biochim Biophys Acta* 2000, **7**:188-194.  
Description of murine Survivin and its promoter and regulation.
56. Uren AG, Wong L, Pakusch M, Fowler KJ, Burrows FJ, Vaux DL, Choo KHA: **Survivin and the inner centromere protein INCENP show similar cell-cycle localization and gene knockout phenotype.** *Curr Biol* 2000, **10**:1319-1328.  
Demonstrating the role of Survivin in the regulation of cytokinesis.
57. Speliotes EK, Uren A, Vaux D, Horvitz HR: **The survivin-like *C. elegans* BIR-1 protein acts with the aurora-like kinase AIR-2 to affect chromosomes and the spindle midzone.** *Mol Cell* 2000, **6**:211-223.  
Demonstrating the role of *C. elegans* BIRPs in regulating cytokinesis.
58. Skoufias DA, Mollinari C, Lacroix FB, Margolis RL: **Human survivin is a kinetochore-associated passenger protein.** *J Cell Biol* 2000, **151**:1575-1581.  
The authors overexpressed and showed that Survivin is a passenger protein; two mutants failed to localize to the kinetochores.
59. Li FZ, Flanary PL, Altieri DC, Dohleman HG: **Cell division regulation by BIR1, a member of the inhibitor of apoptosis family in yeast.** *J Biol Chem* 2000, **275**:6707-6711.  
Disruption of Spbir1p shows that BIRPs have a role in regulating cell division in *Schizosaccharomyces pombe*.
60. O'Connor DS, Grossman D, Plescia J, Li FZ, Zhang H, Villa A, Tognin S, Marchisio PC, Altieri DC: **Regulation of apoptosis at cell division by p34(cdc2) phosphorylation of survivin.** *Proc Natl Acad Sci USA* 2000, **97**:13103-13107.  
Survivin interacts with and is phosphorylated by the p34(cdc2)-cyclin B1 complex.
61. Verhagen AM, Ekert PG, Pakusch M, Silke J, Connolly LM, Reid GE, Moritz RL, Simpson RJ, Vaux DL: **Identification of DIABLO, a mammalian protein that promotes apoptosis by binding to and antagonizing IAP proteins.** *Cell* 2000, **102**:43-53.  
The first description (with [62]) of a mammalian IAP antagonist also known as DIABLO or Smac.
62. Du CY, Fang M, Li YC, Li L, Wang XD: **Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition.** *Cell* 2000, **102**:33-42.  
See [61].
63. Ekert PG, Silke J, Hawkins CJ, Verhagen AM, Vaux DL: **DIABLO promotes apoptosis by removing MIHA/XIAP from processed caspase 9.** *J Cell Biol* 2001, **152**:483-490.  
Demonstrating direct competition between DIABLO and caspase 9 for interaction with XIAP.
64. Srinivasula SM, Hegde R, Saleh A, Datta P, Shiozaki E, Chai J, Lee R, Robbins PD, Fernandes-Alnemri T, Shi Y, Alnemri ES: **A conserved XIAP-interaction motif in caspase-9 and Smac/DIABLO regulates caspase activity and apoptosis.** *Nature* 2001, **410**:112-116.  
The amino terminus of the small subunit of processed caspase 9 and the amino terminus of mature DIABLO are similar and compete for the same interaction site within the BIR3 domain of XIAP.
65. Chai JJ, Du CY, Wu JW, Kyin S, Wang XD, Shi YG: **Structural and biochemical basis of apoptotic activation by Smac/DIABLO.** *Nature* 2000, **406**:855-862.  
Describes the structure of DIABLO and demonstrates that DIABLO is a dimeric protein that interacts with IAPs through its amino terminus.
66. Srinivasula SM, Datta P, Fan XJ, Fernandes-Alnemri T, Huang ZW, Alnemri ES: **Molecular determinants of the caspase-promoting activity of Smac/DIABLO and its role in the death receptor pathway.** *J Biol Chem* 2000, **275**:36152-36157.  
Determines that the amino terminus of DIABLO is responsible for interaction with IAPs.
67. Liu ZH, Sun CH, Olejniczak ET, Meadows RP, Betz SF, Oost T, Herrmann J, Wu JC, Fesik SW: **Structural basis for binding of Smac/DIABLO to the XIAP BIR3 domain.** *Nature* 2000, **408**:1004-1008.  
Describing the solution structure of the XIAP BIR3 domain interacting with a nine-amino-acid peptide corresponding to the amino terminus of mature DIABLO.
68. Wu G, Chai JJ, Suber TL, Wu JW, Du CY, Wang XD, Shi YG: **Structural basis of IAP recognition by Smac/DIABLO.** *Nature* 2000, **408**:1008-1012.  
The crystal structure of DIABLO complexed with the BIR3 domain of XIAP.
69. Silke J, Verhagen AM, Ekert PG, Vaux DL: **Sequence as well as functional similarity for DIABLO/Smac and Grim, Reaper and Hid?** *Cell Death Differ* 2000, **7**:1275.  
Observation that the amino terminus of mature DIABLO is similar to the amino terminus of the *Drosophila* IAP antagonists Grim, Reaper and Hid.
70. Silke J, Ekert PG, Day CL, Hawkins CJ, Baca M, Chew J, Pakusch M, Verhagen AM, Vaux DL: **Direct inhibition of caspase 3 is dispensable for the anti-apoptotic activity of XIAP.** *EMBO J* 2001, **20**:3114-3123.  
Demonstrates the importance of residues within the linker region upstream of the BIR2 domain in XIAP inhibition of caspase 3.
71. Harlin H, Reffey SB, Duckett CS, Lindsten T, Thompson CB: **Characterization of XIAP-deficient mice.** *Mol Cell Biol* 2001, **21**:3604-3608.  
XIAP-deficient mice develop normally, with the only observed change being increased levels of cIAP-1 and -2 expression.
72. **PHYLIP: PHYLIP Phylogeny Inference Package** [<http://evolution.genetics.washington.edu/phylip.html>]  
Program used to generate the phylogenetic tree of IAPs (Figure 2).