

INJECTIVE REPRESENTATIONS OF QUIVERS

SANGWON PARK AND DERA SHIN

ABSTRACT. We prove that $M_1 \xrightarrow{f} M_2$ is an injective representation of a quiver $Q = \bullet \rightarrow \bullet$ if and only if M_1 and M_2 are injective left R -modules, $M_1 \xrightarrow{f} M_2$ is isomorphic to a direct sum of representation of the types $E_1 \rightarrow 0$ and $E_2 \xrightarrow{id} E_2$ where E_1 and E_2 are injective left R -modules. Then, we generalize the result so that a representation $M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} \dots \xrightarrow{f_{n-1}} M_n$ of a quiver $Q = \bullet \rightarrow \bullet \rightarrow \dots \rightarrow \bullet$ is an injective representation if and only if each M_i is an injective left R -module and the representation is a direct sum of injective representations.

1. Introduction

A quiver is just a directed graph. We allow multiple edges and edges going from a vertex back to the same vertex. Originally a representation of quiver assigned a vector space to each vertex - and a linear map to each edge (or arrow) - with the linear map going from the vector space assigned to the initial vertex to the one assigned to the terminal vertex. For example, a representation of the quiver $Q = \bullet \rightarrow \bullet$ is $M_1 \xrightarrow{f} M_2$. Then we can define a morphism of two representations of the same quiver. Now, instead of vector spaces we can use left R -modules and also instead of linear maps we can use R -linear maps. Representations of quivers were studied in ([1], [2]) and recently in [3] noetherian quivers were studied and in [4] projective representations of quivers were studied.

A left R -module E is injective if, for every left R -module B and every submodule A of B , every R -linear map $f : A \rightarrow E$ can be extended to an R -linear map $g : B \rightarrow E$. The diagram is

Received May 25, 2005. Revised November 16, 2005.

2000 Mathematics Subject Classification: 16E30, 13C11, 16D80.

Key words and phrases: module, quiver, representation of quiver, injective representation of quiver.

This paper was supported by Dong-A University Research Fund in 2005.

$$\begin{array}{ccccc}
 0 & \longrightarrow & A & \longrightarrow & B \\
 & & \downarrow f & \searrow \dot{g} & \\
 & & E & &
 \end{array}$$

DEFINITION 1.1. A representation $M_1 \xrightarrow{f} M_2$ of a quiver $Q = \bullet \rightarrow \bullet$ is called an injective representation if for any representation $N_1 \xrightarrow{g} N_2$ with a subrepresentation $S_1 \xrightarrow{s_2|g|s_1} S_2$ and morphisms

$$\begin{array}{ccc}
 S_1 & \xrightarrow{s_2|g|s_1} & S_2 \\
 h \downarrow & & \downarrow k \\
 M_1 & \xrightarrow{f} & M_2
 \end{array}$$

there exist $H \in \text{Hom}_R(N_1, M_1)$ and $K \in \text{Hom}_R(N_2, M_2)$ such that the following diagram

$$\begin{array}{ccc}
 N_1 & \xrightarrow{g} & N_2 \\
 H \downarrow & & \downarrow K \\
 M_1 & \xrightarrow{f} & M_2
 \end{array}$$

commutes and $H|_{S_1} = h$ and $K|_{S_2} = k$.

In other words, every diagram of representations

$$\begin{array}{ccccc}
 (0 \longrightarrow 0) & \longrightarrow & (S_1 \xrightarrow{s_2|g|s_1} S_2) & \longrightarrow & (N_1 \xrightarrow{g} N_2) \\
 & & \downarrow h & & \downarrow k \\
 & & (M_1 \xrightarrow{f} M_2) & &
 \end{array}$$

can be completed to a commutative diagram as follows :

$$\begin{array}{ccccc}
 (0 \longrightarrow 0) & \longrightarrow & (S_1 \xrightarrow{s_2|g|s_1} S_2) & \longrightarrow & (N_1 \xrightarrow{g} N_2) \\
 & & \downarrow & & \swarrow H \\
 & & & & (M_1 \xrightarrow{f} M_2) \\
 & & & & \nwarrow K
 \end{array}$$

LEMMA 1.2. If $M_1 \xrightarrow{f} M_2$ is an injective representation of a quiver $Q = \bullet \rightarrow \bullet$, then M_1 and M_2 are injective left R -modules.

PROOF. Let N be a left R -module, S be a submodule of N and $\alpha : S \rightarrow M_1$ be an R -linear map. Then, since $M_1 \xrightarrow{f} M_2$ is an injective representation, we can complete the diagram

$$\begin{array}{ccccc}
 (0 \longrightarrow 0) & \longrightarrow & (S \xrightarrow{id} S) & \longrightarrow & (N \xrightarrow{id} N) \\
 & & \downarrow \alpha & & \swarrow f \circ \alpha \\
 & & & & (M_1 \xrightarrow{f} M_2) \\
 & & & & \nwarrow f
 \end{array}$$

as a commutative diagram. Thus, M_1 is an injective left R -module.

Let $g : S \rightarrow M_2$ be an R -linear map. Then, since $M_1 \xrightarrow{f} M_2$ is an injective representation, we can complete the diagram

$$\begin{array}{ccccc}
 (0 \longrightarrow 0) & \longrightarrow & (0 \longrightarrow S) & \longrightarrow & (0 \longrightarrow N) \\
 & & \downarrow & & \swarrow h \\
 & & & & (M_1 \xrightarrow{f} M_2) \\
 & & & & \nwarrow g
 \end{array}$$

as a commutative diagram. Thus, M_2 is an injective left R -module. \square

LEMMA 1.3. If E is an injective left R -module, then a representation $E \rightarrow 0$ of a quiver $Q = \bullet \rightarrow \bullet$ is an injective representation.

PROOF. The lemma follows by completing the diagram

$$\begin{array}{ccccc}
(0 \longrightarrow 0) & \longrightarrow & (S_1 \xrightarrow{s_2|f|s_1} S_2) & \longrightarrow & (M_1 \xrightarrow{f} M_2) \\
& & \downarrow g & & \downarrow h \\
& & (E \longrightarrow 0) & &
\end{array}$$

as a commutative diagram. \square

REMARK 1.4. A representation $0 \rightarrow E$ of a quiver $Q = \bullet \rightarrow \bullet$ is not an injective representation if $E \neq 0$, because we cannot complete the diagram

$$\begin{array}{ccccc}
(0 \longrightarrow 0) & \longrightarrow & (0 \longrightarrow E) & \longrightarrow & (E \xrightarrow{id} E) \\
& & \downarrow & & \downarrow id \\
& & (0 \longrightarrow E) & &
\end{array}$$

as a commutative diagram.

LEMMA 1.5. *If E is an injective left R -module, then a representation $E \xrightarrow{id} E$ of a quiver $Q = \bullet \rightarrow \bullet$ is an injective representation.*

PROOF. Let $M_1, M_2, E, S_1(\subset M_1)$ and $S_2(\subset M_2)$ be an left R -modules and let $f : M_1 \rightarrow M_2$ be R -linear map. Let $g : S_2 \rightarrow E$ be an R -linear map and choose $g \circ f : S_1 \rightarrow E$ as an R -linear map. And consider the following diagram :

$$\begin{array}{ccccc}
(0 \longrightarrow 0) & \longrightarrow & (S_1 \xrightarrow{s_2|f|s_1} S_2) & \longrightarrow & (M_1 \xrightarrow{f} M_2) \\
& & \downarrow g \circ f & & \downarrow g \\
& & (E \xrightarrow{id} E) & &
\end{array}$$

Then, since E is an injective left R -module, there exists a map $h : M_2 \rightarrow E$. Now choose $h \circ f : M_1 \rightarrow E$ as an R -linear map. Then h and $h \circ f$ complete the above diagram as a commutative diagram. Therefore, $E \xrightarrow{id} E$ is an injective representation. \square

2. Direct sum of injective representations

THEOREM 2.1. *A representation $M_1 \xrightarrow{f} M_2$ is an injective representation of a quiver $Q = \bullet \rightarrow \bullet$ if and only if M_1, M_2 are injective left R -modules and $M_1 \xrightarrow{f} M_2$ is isomorphic to a direct sum of representations of the types $E_1 \rightarrow 0$ and $E_2 \xrightarrow{id} E_2$ where E_1 and E_2 are injective left R -modules.*

PROOF. Consider the following diagram :

$$\begin{array}{ccccc}
 (0 \longrightarrow 0) & \longrightarrow & (0 \longrightarrow M_2) & \longrightarrow & (M_2 \xrightarrow{id} M_2) \\
 & & \downarrow & & \downarrow id \\
 & & (M_1 \xrightarrow{f} M_2) & &
 \end{array}$$

Since $M_1 \xrightarrow{f} M_2$ is an injective representation, we can complete the above diagram as a commutative diagram as follows:

$$\begin{array}{ccccc}
 (0 \longrightarrow 0) & \longrightarrow & (0 \longrightarrow M_2) & \longrightarrow & (M_2 \xrightarrow{id} M_2) \\
 & & \downarrow & \nearrow id & \downarrow id \\
 & & (M_1 \xrightarrow{f} M_2) & \xleftarrow{id \circ g} &
 \end{array}$$

Thus, $f \circ g = id_{M_2}$. Therefore, $M_1 \cong M_2 \oplus \ker(f)$ and

$$(M_1 \xrightarrow{f} M_2) \cong (M_2 \xrightarrow{id} M_2) \oplus (\ker(f) \longrightarrow 0).$$

□

Now let $Q = \bullet \rightarrow \bullet \rightarrow \bullet \rightarrow \dots \rightarrow \bullet \rightarrow \bullet$ be a quiver with n vertices and $n - 1$ arrows. Then, we can easily generalize the results of Lemmas 1.3 and 1.5 as follows : the representations

$$\begin{array}{ccccccc}
 E & \rightarrow & 0 & \rightarrow & 0 & \rightarrow & \dots & \rightarrow & 0 & \rightarrow & 0 \\
 E & \xrightarrow{id} & E & \rightarrow & 0 & \rightarrow & \dots & \rightarrow & 0 & \rightarrow & 0 \\
 & & & & & & \vdots & & & & \\
 E & \xrightarrow{id} & E & \xrightarrow{id} & E & \xrightarrow{id} & \dots & \rightarrow & E & \xrightarrow{id} & E & \rightarrow & 0 \\
 E & \xrightarrow{id} & E & \xrightarrow{id} & E & \xrightarrow{id} & \dots & \rightarrow & E & \xrightarrow{id} & E & \xrightarrow{id} & E
 \end{array}$$

are all injective representations of a quiver $Q = \bullet \rightarrow \bullet \rightarrow \bullet \rightarrow \cdots \rightarrow \bullet \rightarrow \bullet$, if each E_i is an injective left R -module. We can also generalize Lemma 1.2 so that if $M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} M_n$ of a quiver $Q = \bullet \rightarrow \bullet \rightarrow \bullet \rightarrow \cdots \rightarrow \bullet \rightarrow \bullet$ is an injective representation, then each M_i is an injective left R -module.

THEOREM 2.2. *A representation $M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} M_3$ is an injective representation of a quiver $Q = \bullet \rightarrow \bullet \rightarrow \bullet$ if and only if M_1 , M_2 and M_3 are injective left R -modules,*

$$(M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} M_3) \\ \cong (E_1 \xrightarrow{id} E_1 \xrightarrow{id} E_1) \oplus (E_2 \xrightarrow{id} E_2 \rightarrow 0) \oplus (E_3 \rightarrow 0 \rightarrow 0).$$

PROOF. The diagram

$$(0 \longrightarrow 0 \longrightarrow 0) \longrightarrow (0 \longrightarrow M_2 \xrightarrow{id} M_2) \longrightarrow (M_2 \xrightarrow{id} M_2 \xrightarrow{id} M_2) \\ \downarrow \quad \quad \downarrow \quad \quad \downarrow \quad \quad \downarrow \\ (M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} M_3)$$

can be completed to a commutative diagram by $g_{21} : M_2 \rightarrow M_1$, $id : M_2 \rightarrow M_2$ and $g_{23} : M_2 \rightarrow M_3$. Then we can get $f_1 \circ g_{21} = id_{M_2}$ so that $M_1 \cong M_2 \oplus \ker(f_1)$. Now the diagram

$$(0 \longrightarrow 0 \longrightarrow 0) \longrightarrow (0 \longrightarrow 0 \longrightarrow M_3) \longrightarrow (0 \longrightarrow M_3 \xrightarrow{id} M_3) \\ \downarrow \quad \quad \downarrow \quad \quad \downarrow \quad \quad \downarrow \\ (M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} M_3)$$

can be completed to a commutative diagram by $i : 0 \rightarrow M_1$, $g_{32} : M_3 \rightarrow M_2$, $id : M_3 \rightarrow M_3$. Then, we can get $f_2 \circ g_{32} = id_{M_3}$ so that $M_2 \cong M_3 \oplus \ker(f_2)$. Therefore, $M_1 \cong M_3 \oplus \ker(f_2) \oplus \ker(f_1)$. Hence,

$$(M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} M_3) \\ \cong (E_1 \xrightarrow{id} E_1 \xrightarrow{id} E_1) \oplus (E_2 \xrightarrow{id} E_2 \rightarrow 0) \oplus (E_3 \rightarrow 0 \rightarrow 0).$$

This completes the proof. \square

Now, we can easily generalize Theorem 2.2 so that a representation $M_1 \xrightarrow{f_1} M_2 \xrightarrow{f_2} \cdots \xrightarrow{f_{n-1}} M_n$ of a quiver $Q = \bullet \rightarrow \bullet \rightarrow \bullet \rightarrow \cdots \rightarrow \bullet \rightarrow \bullet$

is an injective representation if and only if each M_i is an injective left R -module and the representation is the direct sum of the following injective representations:

$$\begin{array}{c} E \rightarrow 0 \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow 0 \\ E \xrightarrow{id} E \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow 0 \\ \vdots \\ E \xrightarrow{id} E \xrightarrow{id} E \xrightarrow{id} \cdots \rightarrow E \xrightarrow{id} E \rightarrow 0 \\ E \xrightarrow{id} E \xrightarrow{id} E \xrightarrow{id} \cdots \rightarrow E \xrightarrow{id} E \xrightarrow{id} E \end{array}$$

REMARK 2.3. The representations of a quiver $Q = \bullet \rightarrow \bullet \rightarrow \bullet \rightarrow \cdots \rightarrow \bullet \rightarrow \bullet$:

$$\begin{array}{c} 0 \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow E \\ 0 \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow E \xrightarrow{id} E \\ \vdots \\ 0 \rightarrow 0 \rightarrow \cdots \rightarrow 0 \rightarrow E \xrightarrow{id} E \xrightarrow{id} E \\ 0 \rightarrow E \xrightarrow{id} E \xrightarrow{id} \cdots \rightarrow E \xrightarrow{id} E \xrightarrow{id} E \end{array}$$

are not injective representations if $E \neq 0$.

References

- [1] E. Enochs, I. Herzog, and S. Park, *Cyclic quiver ring and polycyclic-by-finite group ring*, Houston J. Math. **25** (1999), no. 1, 1–13.
- [2] E. Enochs and I. Herzog, *A homotopy of quiver morphism with applications to representations*, Canad. J. Math. **51** (1999), no. 2, 294–308.
- [3] E. Enochs, J. R. Rozas, L. Oyonarte, and S. Park, *Noetherian quivers*, Quaestiones Mathematicae **25** (2002), no. 4, 531–538.
- [4] S. Park, *Projective representations of quivers*, Internart. J. Math. Math. Sci. **31** (2002), no. 2, 97–101.

Department of Mathematics
Dong-A University
Pusan 604-714, Korea
E-mail: swpark@donga.ac.kr
drshin@donga.ac.kr