

Innate and adaptive immunity in teleost fish: a review

C. URIBE¹, H. FOLCH², R. ENRIQUEZ¹, G. MORAN¹

¹Faculty of Veterinary Science, Universidad Austral de Chile, Valdivia, Chile

²Faculty of Medicine, Universidad Austral de Chile, Valdivia, Chile

ABSTRACT: The immune system of fish is very similar to vertebrates, although there are some important differences. Fish are free-living organisms from the embryonic stage of life in their aquatic environment. They have mechanisms to protect themselves from a wide variety of microorganisms. Consequently, fish rely on their innate immune system for an extended period of time, beginning at the early stages of embryogenesis. The components of the innate immune response are divided into physical, cellular and humoral factors and include humoral and cellular receptor molecules that are soluble in plasma and other body fluids. The lymphoid organs found in fish include the thymus, spleen and kidney. Immunoglobulins are the principal components of the immune response against pathogenic organisms. Immunomodulatory products, including nucleotides, glucans and probiotics, are increasingly used in aquaculture production. The use of these products reduces the need for therapeutic treatments, enhances the effects of vaccines and, in turn, improves the indicators of production. The aim of this review is to provide a review of the immune system in fish, including the ontogeny, mechanisms of unspecific and acquired immunity and the action of some immunomodulators.

Keywords: fish; immune response; immunomodulators

CRP = C-reactive protein; **ICM** = intermediate cell mass; **IHNV** = infectious hematopoietic necrosis virus; **IL** = interleukin; **INF** = interferon; **IPNV** = infectious pancreatic necrosis virus; **ISAV** = infectious salmon anaemia virus; **LPS** = lipopolysaccharide; **MHC** = Major histocompatibility complex; **MMCs** = melanomacrophage centers; **SAA** = serum amyloid protein; **TCR** = T cell receptor; **TLR** = toll-like receptor; **TNF** = tumour necrosis factor; **VHSV** = viral haemorrhage septicaemia virus

Contents

- | | |
|--|--|
| 1. Introduction | 5.10. Protease inhibitors |
| 2. Ontogeny of the immune system | 5.11. Lysozyme |
| 3. Lymphoid organs | 5.12. Natural antibodies |
| 3.1. Thymus | 5.13. Pentraxins |
| 3.2. Kidney | 5.14. Transferrin |
| 3.3. Spleen | 6. Specific immunity |
| 4. Cells involved in immune response | 6.1. Antibodies |
| 5. Nonspecific immunity | 6.2. Immunological memory |
| 5.1. Physical barriers | 6.3. Cellular cytotoxicity |
| 5.2. Nonspecific cellular cytotoxicity | 6.4. Cytokines involved in adaptive immunity |
| 5.3. Antimicrobial peptides | 7. Environmental effect on immune response |
| 5.4. Phagocytosis | 8. Compounds that modulate the immune system in fish |
| 5.5. Complement | 8.1. Nucleotides |
| 5.6. Tumor necrosis factor (TNF) | 8.2. Glucans |
| 5.7. Interferon (INF) | 8.3. Probiotics |
| 5.8. Interleukins (IL) | 9. Concluding remarks |
| 5.9. Others cytokines and chemokines | 10. References |

1. Introduction

The immune system of fish is physiologically similar to that of higher vertebrates, despite certain differences. In contrast to higher vertebrates, fish are free-living organisms from early embryonic stages of life and depend on their innate immune system for survival (Rombout et al., 2005). Nonspecific immunity is a fundamental defence mechanism in fish. In addition, it plays a key role in the acquired immune response and homeostasis through a system of receptor proteins. These receptor proteins identify molecular patterns that are typical of pathogenic microorganisms, including polysaccharides, lipopolysaccharide (LPS), peptidoglycan bacterial DNA, viral RNA and other molecules that are not normally on the surface of multicellular organisms. This response is divided into physical barriers and cellular and humoral immune response. These immunological parameters include growth inhibitors, lytic enzymes, the classic complement pathways, the alternative and lectin pathway, agglutinins and precipitins (opsonins and primary lectins), antibodies, cytokines, chemokines and antibacterial peptides. Various internal and external factors can influence innate immune response parameters. Temperature changes, stress management and density may have suppressive effects on this type of response, while several food additives and immunostimulants can enhance their efficiency (Magnadottir, 2006, 2010).

The aim of this review is to provide some background on the immune system of fish, describing the ontogeny and core components, the nonspecific and adaptive immune mechanisms and elements that promote their actions as immunomodulators.

2. Ontogeny of the immune system

The ontogeny of the immune system of teleost fish has been studied in various species, including rainbow trout (*Oncorhynchus mykiss*), catfish (*Ictalurus punctatus*), zebrafish (*Danio rerio*) and grouper (*Acanthistius brasilianus*). Teleosts have been described as primitive models of hematopoiesis. The first hematopoietic organ is called the intermediate cell mass (ICM). In kili (*Pseudepiplatys annulatus*) and rainbow trout, hematopoiesis exists for a short period in the yolk sac and then shifts to the ICM (Zapata et al., 2006). With respect to the development of the organs involved in the immune

response, the anterior kidney and thymus are considered to be complete before hatching occurs in rainbow trout and Atlantic salmon (*Salmo salar*) (Razquin et al., 1990); this has not been clarified at the end of incubation in many marine species, including bream (*Sparus aurata*), grouper and turbot (*Scophthalmus maximus*). Despite these differences, the sequence of lymphomyeloid organ development in marine teleosts is as follows: kidney, spleen and thymus (Mulero et al., 2007), but larval spleen has a greater erythropoietic function than lymphopoietic function (Nakanishi, 1986; Schroder et al., 1998). Cells of myeloid lineage can be found around the yolk sac from 24 hours post fertilization (hpf) but are not seen in the ICM. Myeloblasts and neutrophilic myelocytes are observed between 34 and 48 hpf. On the other hand, numerous erythrocytes circulate in the blood vessels, and mature granulocytes are found in the blood system in the connective tissue that surrounds the yolk sac at 34 hpf and adjacent to the first renal tubules at 48 hpf. In addition, young myelocytes and neutrophils migrate through blood vessel walls during the same period. Mature neutrophils are described in various tissues at 72 hpf (Willett et al., 1999).

On the other hand, the first appearance of IgM in lymphocytes varies considerably among fish species (Magnadottir et al., 2005). However, the first appearance of B-lymphocytes and immunoglobulins is late in marine species compared to fresh water species, and larvae have reached about 20–30 mm in length when IgM is first expressed (Chantanachookhin et al., 1991; Magnadottir et al., 2005). In addition, Dalmo (2005) mentioned that the transfer of maternal antibody to eggs and embryos has been demonstrated in several species. Further, Magnadottir et al. (2005) suggest that the primary role of maternal antibodies is to protect the eggs against vertical transfer of certain pathogens or that maternal IgM may aid phagocytosis or the activation of complement pathways in early developmental stages; IgM may even function simply as a nutritional yolk protein.

Finally, complement component C3 has been found in unfertilised eggs in the spotted wolfish (*Anarhichas minor* Olafsen) indicating maternal transfer (Ellingsen et al., 2005). Using immunohistochemistry techniques, C3 has been found in several different organs and tissues of developing cod and halibut (Lange et al., 2004a,b). These studies suggest that complement may play a role in the generation of different organs and not only in the defence against invading pathogens (Dalmo, 2005).

3. Lymphoid organs

The thymus, kidney (anterior and middle) and spleen are the largest lymphoid organs in teleosts (Zapata et al., 2006). In freshwater teleosts, the thymus is the first organ to become lymphoid, but before this event, the anterior kidney may contain hematopoietic progenitors, but not lymphocytes (Lam et al., 2004).

3.1. Thymus

This organ has two lobes, is homogeneous, and is represented by a thin sheet of oval lymphoid tissue that is arranged subcutaneously in the dorsal commissure of the operculum and is lined by mucous tissue of the pharyngeal epithelium (Ellis, 2001). The structure that characterizes the thymus of fish is a capsule that surrounds the lymphoid bark tissue. Basically, the thymus can be considered as an aggregation of macrophages that promote the encapsulated proliferation of T cells (Davis et al., 2002). The differentiation of the thymic structure is highly variable in teleosts, and in many species, it is not possible to observe a clear differentiation between the cortex and medulla that is found in higher vertebrates (Bowden et al., 2005). Furthermore, myeloid cells and eosinophilic granular cells can be found in this organ. Other studies described the appearance of focal epithelial nests, known as Hassal's corpuscles (Zapata et al., 2006). The thymus is responsible for the production of T cells. In zebrafish, thymus development involves neural crest cells derived from the neuroectoderm, which migrate to the third and fourth pharyngeal pouches and interact with the endoderm (Trede et al., 2001). The early development of the thymus has been studied in several teleost fish species, and development time differs between species according to the effects of temperature on growth. The relationship between growth and development is dynamic, with physiological age expressed as degree days, and is not a factor for differences in historical temperature (Bowden et al., 2005).

3.2. Kidney

The kidney in teleost fish is the equivalent of the bone marrow in vertebrates and is the largest site of haematopoiesis until adulthood (Zapata et al.,

2006). In rainbow trout, the kidney is well-developed after hatching, when it mainly produces red blood cells and granulocytes. Studies suggest the presence of lymphoid cells that release IgM between 12 and 14 days post fertilization (dpf) (Castillo et al., 1993) and have also shown two variants of IgM by ELISA in embryos eight days before the end of the incubation period (Sanchez et al., 1995). This finding suggests that a source of B cells exists before the end of the incubation period in the kidney or in other hematopoietic sites. Structurally, the anterior kidney is composed of a network of reticular fibres that provide support for lymph tissue and are found scattered among hematopoietic system cells that line the sinusoid reticuloepithelium (Press et al., 1994). The main cells found in the anterior kidney are macrophages, which aggregate into structures called melanomacrophage centers (MMCs), and lymphoid cells, which are found at all developmental stages and exist mostly as Ig⁺ cells (B cells) (Press et al., 1994). Reticular cells play an important role in supplying the interactions necessary for the function of lymphoid cells (Press et al., 1994) and endothelial cells of sinusoids. The latter system is the main component for blood filtering due to its ability to perform endocytosis (Danneving et al., 1994).

3.3. Spleen

The spleen is composed of a system of splenic ellipsoids, MMCs and lymphoid tissue. In most species, ellipsoids are clustered together and are organized around the other two components (Ferguson, 1989). The ellipsoids are thick-walled capillaries that open in the pulp and result from the division of the splenic arterioles. The cells along the walls are actively involved in the macrophage phagocytosis of antigens. Usually in the form of antibodies or metabolic products, antigens may be detained for long periods of time, which has an important role in immunological memory. The spleen in zebrafish remains a small organ that contains large amounts of eritroblast at 30 days post fertilization. At three months, when lymphoblasts are evident in the spleen, emerging ellipsoids are involved in the capture of antigen. A similar developmental pattern has been described for other teleost spleen, such as that of the Atlantic salmon, grouper and catfish (Dos Santos et al., 2000; Petrie-Hanson and Ainsworth, 2000).

4. Cells involved in immune response

Fish possess lymphocyte populations that are analogous to T cells, B cells, cytotoxic cells (similar to natural killer cells), macrophages and polymorphonuclear leukocytes. The immune system of teleosts has sub-populations of T lymphocytes that exhibit differential responses to mitogens, B cell acute allograft reactions, mixed leukocyte reactions and cooperative interactions between T cells, B cells and macrophages that are essential for the production of antibodies. Moreover, elasmobranch and teleost fish are the most primitive groups that possess the Major Histocompatibility Complex (MHC) and T cell receptor (TCR) (Manning and Nakanishi, 1996).

5. Nonspecific immunity

In fish, the innate response has been considered an essential component in combating pathogens due to limitations of the adaptive immune system, their poikilothermic nature, their limited repertoire of antibodies and the slow proliferation, maturation and memory of their lymphocytes (Whyte, 2007). It is commonly divided into three compartments: the epithelial/mucosal barrier, the humoral parameters and the cellular components. The epithelial and mucosal barrier of the skin, gills and alimentary tract is an extremely important disease barrier in fish, being constantly immersed in media containing potentially harmful agents (Magnadottir, 2010). This type of response requires a series of mechanisms that involve humoral factors, cell and tissue, antimicrobial peptides and complement factors. Humoral factors may be cellular receptors or molecules that are soluble in plasma and other body fluids (Magnadottir, 2006; Subramanian et al., 2007; Subramanian et al., 2008)

5.1. Physical barriers

Flakes, skin mucus and gills act as the first barrier to infection (Ingram, 1980; Shepard, 1994; Ellis, 2001). The mucus of fish contains lectins, pentraxins, lysozymes, complement proteins, antibacterial peptides and immunoglobulin M (IgM), which have an important role in inhibiting the entry of pathogens (Alexander and Ingram, 1992; Rombout et al., 1993; Aranishi and Nakane, 1997; Boshra et al., 2006; Saurabh and Sahoo, 2008). In addition, the epidermis

is able to react to different attacks (thickening and cellular hyperplasia), and its integrity is essential for osmotic balance and to prevent the entry of foreign agents (Hibiya, 1994). On the other hand, defending cells are present, such as lymphocytes, macrophages and eosinophilic granular cells (Sveinbjornsson et al., 1996; Ellis, 2001; Fischer et al., 2006).

5.2. Nonspecific cellular cytotoxicity

In mammals, nonspecific response situations are mainly executed by cytotoxic cells, known as natural killer cells. Although the nonspecific cytotoxic cells of catfish are morphologically distinct from the large granular lymphocytes of mammals, they are suggested to be functionally similar (Evans and Jaso-Friedmann, 1992). These cells are able to eliminate a range of spontaneously xenogeneic targets, including parasites in fish and traditional targets of natural killer cells in mammals (Hasegawa et al., 1998). Unlike the natural killer cells of mammals, the nonspecific cytotoxic cells of catfish are agranular, small lymphocytes that are commonly found in lymphoid tissues, such as the anterior kidney and spleen, but are rarely found in the blood (Shen et al., 2002). In addition to the case in catfish, nonspecific cytotoxic cells have shown activity in other fish species, including rainbow trout (Greenle et al., 1991), common carp (*Cyprinus carpio*) (Hinuma et al., 1980; Suzumura et al., 1994), damselfish (*Dascyllus albisella*) (McKinney and Achmale, 1994) and tilapia (*Oreochromis* spp.) (Faisal et al., 1989).

5.3. Antimicrobial peptides

Studies of the integument and integument secretions of fish (Hellio et al., 2002) have demonstrated an important role for this system in host defence against viruses and bacteria (Ellis, 2001; Chinchar et al., 2004; Maier et al., 2008). These peptides have been found in the mucus, liver and gill tissue of teleost fish (Birkemo et al., 2003). These low molecular weight polypeptides have the ability to break down bacterial walls (Ellis, 2001).

5.4. Phagocytosis

Phagocytosis is one of the most important processes in poikilothermic animals because it is

the process that is least influenced by temperature (Blazer, 1991; Lange and Magnadottir, 2003; Magnadottir et al., 2005). The main cells involved in phagocytosis in fish are neutrophils and macrophages (Secombes and Fletcher, 1992). These cells remove bacteria mainly by the production of reactive oxygen species during a respiratory burst. In addition, neutrophils possess myeloperoxidase in their cytoplasmic granules, which in the presence of halide and hydrogen peroxide kills bacteria by halogenation of the bacterial cell wall. Moreover, these cells have lysozymes and other hydrolytic enzymes in their lysosomes (Fischer et al., 2006). Similarly, macrophages can produce nitric oxide in mammals and can be as potent as antibacterial agents, peroxy-nitrites and hydroxyl groups (Secombes and Fletcher, 1992).

5.5. Complement

The complement system in teleosts, as well as that in higher vertebrates, can be activated in three ways: the classical pathway, which is triggered by antibody binding to the cell surface (Holland and Lambris, 2002), the alternative pathway, which is independent of antibodies and is activated directly by foreign microorganisms, and the lectin pathway, which is activated by the binding of a protein complex consisting of mannose/mannan-binding lectin in bacterial cells (Sakai, 1992). However, the mechanisms and molecules involved in this system in teleosts are not well understood, with the exception of the genetic sequence of the mannose-binding lectin protease that is associated with serum (Matsushita et al., 1998; Nikoskelainen et al., 2002). Studies suggest that the alternative complement pathway is of great importance in the innate immune response in teleost fish (Yano, 1996). The activity of complement-opsonized teleosts has been documented in a variety of species, such as common carp (Matsuyama et al., 1992), catfish (Jenkins and Ourth, 1993) and salmonids (Lammens et al., 2000). Salmonid antibodies, in the presence of complement proteins, are able to neutralize enveloped viruses, including the infectious hematopoietic necrosis virus (IHNV) and the viral haemorrhagic septicaemia virus (VHSV) (Lorenzen and La Patra, 1999). In bacterial infections, complement activation by lipopolysaccharide found in the cell wall of gram-negative pathogenic bacteria stimulates C5a factor production, a pow-

erful chemokine to macrophages and neutrophils that have receptors for C3, the central complement molecule which is a part of all three pathways, and which remains attached to the bacteria, favouring phagocytosis (Jenkins and Ourth, 1993). Teleost C3 is composed of a disulfide-linked two-chain (a and b) glycoprotein containing a thioester bond comparable to C3 from several vertebrate species (Magdanottir et al., 2005).

5.6. Tumor necrosis factor (TNF)

Several studies in fish have provided direct evidence suggesting that TNF- α and - β are important activators of macrophages. Studies in rainbow trout, turbot, sea bream (*Sparus aurata*), goldfish (*Carassius auratus*) and catfish have shown that TNF causes the activation of macrophages, leading to increased respiratory activity, phagocytosis and nitric oxide production (Yin et al., 1997; Mulero and Meseguer, 1998; Tafalla et al., 2001).

5.7. Interferon (INF)

INF α and β are cytokines with a nonspecific antiviral function that is based on the inhibition of nucleic acid replication within infected cells. INF plays an important role in the defence against viral infection in vertebrate host cells, which secrete INF α / β upon recognition of viral nucleic acid (Robertsen, 2006). These INFs protect other cells from viral infection by binding to different receptors, which results in the induction of several hundred genes that are stimulated by INF (ISGs). Some of these genes encode antiviral proteins, such as protein MX (MX), protein kinase dsRNA activated (PKR) and 2,5-oligoadenylate synthetase (OAS) (De Veer et al., 2001; Samuel, 2001). Two interferons (INF α -1 and INF α -2) have been cloned from the Atlantic salmon and were characterized with respect to their sequence, gene structure, promoters and induction of antiviral activity of interferon-stimulated genes (ISG) (Killeng et al., 2007; Rokenes et al., 2007). The INF α -1 Atlantic salmon induces the expression of MX and ISGs, and both have similar properties to INF α / β and INF γ in mammals (Rokenes et al., 2007). Furthermore, INF α -1 in Atlantic salmon induces a potent antiviral activity against infectious pancreatic necrosis virus (IPNV) in cells. However, this cytokine does not protect Atlantic salmon cells

against infectious salmon anaemia virus (ISAV), suggesting that this virus must have developed mechanisms to counteract the antiviral activity of INF (Killeng et al., 2007). Moreover, INF α -2 expressed recombinantly in Atlantic salmon was also confirmed to confer antiviral activity against IPNV (Ooi et al., 2008). Some authors suggest that rainbow trout have at least three INFs (Zou et al., 2007). Rainbow trout (INF1 (rtl INF1) and INF2 (rtl INF2) show sequences similar to INF α -1 and INF-2 in Atlantic salmon. Rainbow trout INF 3 (rtl INF 3) is related to INF α in mammals and fish, as shown by the conservation of four cysteine residues. The rtl INF1 and 2 proteins have been shown to regulate the production of MX, inhibiting the proliferation of VHSV in cell lines from the gonad of rainbow trout (GTR 2). Moreover, rtl INF 3 has been proven to be a poor inducer of antiviral activity. It is worth noting that these three INFs in rainbow trout show differential expression across cells and tissues, suggesting that they have different functions in the fish immune system (Zou et al., 2007).

5.8. Interleukins (IL)

IL-1 in mammals is comprised of 10 ligands and 10 protein receptor molecules and plays an important role in inflammation and host defence (Dinarello, 1997). IL-1 β has been detected in 13 teleost fish species and is involved in the regulation of immunity through the stimulation of T cells. The function of IL-1 β in these fish species is analogous to mammalian IL-1 β (Mathew et al., 2002; Magnadottir, 2010). In teleost fish, IL-1 receptors have been cloned and sequenced from the rainbow trout and Atlantic salmon. The expression of the IL-1 receptor in salmon appears to be constitutive in all tissues tested and is regulated in the anterior kidney, spleen, liver and gills after stimulation with LPS and TNF- α , suggesting a role for the IL-1 receptor in regulating IL-1 β during the inflammatory response (Sangrador-Vegas et al., 2000; Subramaniam et al., 2002).

5.9. Others cytokines and chemokines

As described above, TNF- α and IL-1 β are cytokines involved in the induction of inflammatory responses to Gram negative bacteria in fish. In addition, it has been demonstrated that IL-6 is

also involved in the cascade leading to an inflammatory response for this type of bacteria (Savan and Sakai, 2006). Further, cytokines involved in leukocyte differentiation, including granulocyte colony stimulating factor (CSF) (Arma et al., 2004), macrophage-CSF (Hanington et al., 2007) and IL-7 (Kono et al., 2008), are all identified in fish. On the other hand, chemokines are also present and in some fish species considerable numbers of genes have now been identified (DeVries et al., 2006); however, many have no clear homologues in other vertebrate groups and their function remains to be determined (Secombes, 2008).

5.10. Protease inhibitors

Several protease inhibitors are present in the serum and other body fluids of fish (Bowden et al. 1997). The main function of protease inhibitors is to maintain body fluid homeostasis. These molecules are involved in acute phase reactions and defence against pathogens that secrete proteolytic enzymes (Magdanottir, 2010). The most widely studied of the protease inhibitors is the α -2 macroglobulin, which has a high specificity for inhibiting the physical encapsulation of protease (Armstrong and Quigley, 1999).

5.11. Lysozyme

Lysozyme is a bacteriolytic enzyme that is widely distributed throughout the body and is part of the nonspecific defence mechanisms in most animals. In salmonids, lysozyme has been detected in serum, secretions, mucous membranes and tissues rich in leucocytes, mainly the kidney and intestine (Grinde et al., 1988; Lie et al., 1989). Apparently, the main sources of lysozyme are monocytes/macrophages and neutrophils. However, recent studies have detected this enzyme in the granules of the eosinophilic granular cells of the intestine (Sveinbjornsson et al., 1996). The bactericidal action of this enzyme involves the hydrolyzation of the peptidoglycan of bacterial cell walls resulting in cell lysis. Lysozyme was initially associated with the defence against Gram-positive bacteria, but has been found to lyse Gram-negative bacteria as well. Furthermore, this enzyme is known to trigger an opsonin of the complement system and phagocytic cells (Magnadottir, 2006).

5.12. Natural antibodies

Natural antibodies are produced in fish at a level that is regulated in the absence of antigenic stimulation of cells that are equivalent to B1 cells (Boes, 2000). These natural antibodies are found in high levels in the serum of fish, where they provide immediate and broad protection against bacterial and viral pathogens, making these factors key components of nonspecific immunity. Natural antibodies are also linked to adaptive immunity. Teleost fish are capable of generating specific IgM-type natural antibodies against various antigens. The intensity of this response, however, has been shown to vary between different species and environmental conditions (Whyte, 2007).

5.13. Pentraxins

C-reactive protein (CRP) and serum amyloid protein (SAA) are present in the body fluids of vertebrates and invertebrates and are commonly associated with the acute phase response of inflammation (Bayne and Gerwick, 2001; Wu et al., 2003). The expression of CRP has been reported in several teleost fish species, including rainbow trout (Hoover et al., 1998), catfish (Szalai et al., 1992), Atlantic salmon, common cod (*Gadus morhua*), halibut (*Hippoglossus hippoglossus*) and dog fish (*Hoplias malabaricus*) (Lund and Olafsen, 1998). The levels of these proteins are increased upon tissue injury, trauma or infection. These proteins play an active role in the immune system (Cook et al., 2003), activation of the classical complement pathway (De Haas et al., 2000) and the removal of apoptotic cells (Nauta et al., 2003). Different stimuli, such as tissue damage, trauma or infection, have been shown to generate various patterns of CRP production in teleost fish, in which either the level of CRP is decreased in serum (negative acute phase protein) (Szalai et al., 1994; Liu et al., 2004) or the level of CRP is increased in serum (positive acute phase protein) (Kodama et al., 2004). Although the pentraxins of teleosts have a recognized role in defence mechanisms, studies performed by Cook et al. (2003, 2005) have demonstrated their role in immune function through the activation of complement-mediated lysis in sheep erythrocytes. These authors note that the phagocytosis of beads soaked with pentraxins possibly occurs through the action of a surface receptor expressed in phagocytes.

5.14. Transferrin

Iron is an essential element in the establishment of infection by many pathogens, but the availability of iron in the tissue fluids of vertebrates is extremely low due to its high affinity for the blood protein transferrin. Only bacteria with high affinity systems for iron absorption are able to maintain sufficient iron levels to grow *in vivo* (Ellis, 2001; Stafford and Belosevic, 2003). Transferrin is a globular glycoprotein with a high iron chelator activity. This protein is the major iron ion transport protein in animals and plants. Transferrin has a high degree of genetic polymorphism in all species and is found in the serum and secretions of all vertebrates. Pathogenic bacteria have developed a set of mechanisms to obtain iron from the host, even from transferrin. However, because of the high degree of genetic polymorphism and the ability of certain bacterial pathogens to obtain iron from transferrin, this protein may be restricted to certain genotypes. In coho salmon (*Oncorhynchus kisutch*), three genotypes have been identified with respect to transferrin: AA, AC and CC. In some strains of coho salmon that originate in Oregon, increasing resistance against bacterial kidney disease has been associated with the C allele of transferrin (Suzumoto et al., 1997). However, the latter has not been confirmed for two strains of coho salmon that are native to British Columbia (Withler and Evelyn, 1990).

6. Specific immunity

The specific immune response occurs through mechanisms that involve a complex network of specialized cells, proteins, genes and biochemical messages that provide the means necessary for the body to respond specifically to antigens, antibodies and effector cells with high specificity and affinity.

6.1. Antibodies

The predominant immunoglobulin in teleosts is a tetramer of the IgM class and contains eight antigenic combining sites (Acton et al., 1971). Some teleosts have a monomer of IgM in their serum, although the factors leading to its expression are still unknown (Wilson and Warr, 1992). The binding affinities of monomeric and tetrameric IgM in rainbow trout are similar, but tetrameric IgM activates

complement more effectively than the monomeric form due to a structural difference in the Fc portion of the molecule (Elcombe et al., 1985). IgD was the second immunoglobulin isotype identified in fish, specifically catfish, due to sequence similarity with IgD in mammals, its location immediately under the IgM gene and its expression in B cells (Wilson et al., 1997). Moreover, the concentration of IgM in the serum of salmonids is extremely low compared to that of other teleosts such as Japanese eel (*Anguilla japonica*) (Uchida et al., 2000), cyprinids (Vilain et al., 1984) and some Perciformes (Scapigliati et al., 1997). However, the amounts of IgM in the serum of brown trout (*Salmo trutta*) and rainbow trout that are infected or acclimated to high temperature (19 °C) reach values similar to those of the common cod and haddock (*Melanogrammus aeglefinus*). In addition, IgM levels in salmon and atlantic cod (*Gadus morhua*) vary with size (Sanchez et al., 1993; Magnadottir et al., 1999), temperature (Sanchez et al., 1993) and water quality season (Olesen and Jorgensen, 1986; Magnadottir et al., 2001). Teleost antibodies are found in the skin (Hatten et al., 2001), intestine (Rombout et al., 1986), gill mucus (Lumsden et al., 1993), bile (Jenkins et al., 1994) and systemically in the plasma. The immune response of the skin and gills is important because these organs are in direct contact with the environment. Specific antibodies can be generated in the skin (Cain et al., 2000), intestine, (Jones et al., 1999) and gills (Lumsden et al., 1993) without necessarily generating a systemic response.

6.2. Immunological memory

Fish develop a memory response before a second exposure to an antigen (Arkoosh and Kaattari, 1991; Whittington et al., 1994). Rainbow trout respond to suboptimal doses of both T lymphocytes in an antigen-dependent and independent manner after an initial exposure to the same antigen (Arkoosh and Kaattari, 1991). It is remarkable that it takes two exposures before the fish responds to the second administration of T-dependent antigens, whereas T-independent antigen requires only one exposure. Additionally, while response is faster and of larger magnitude than the primary response, the number of antigen-specific B cells in the spleen is directly proportional to the frequency of B-cell specific antigen precursors. This finding suggests that the secondary response is caused by the expansion of the pool of memory B cells (Kattari, 1992) and not a specific

difference in the antibodies. However, the response to suboptimal levels of antigen suggests that perhaps B cells with high affinity receptors were selected as memory B cells (Morrison and Nowak, 2002).

6.3. Cellular Cytotoxicity

The leukocytes of fish are capable of generating cellular cytotoxicity reactions, which has been demonstrated in several fish species. However, the cells that are responsible for cell-mediated cytotoxicity in fish are difficult to characterize due to the lack of appropriate tools for cellular and molecular recognition (Fisher et al., 2006). In mammals, the adaptive immune response implemented by CD8+ cytotoxic T lymphocytes has been shown to be critical in combating various viral infections. These lymphocytes recognize and kill cells with virus-derived peptides using class I MHC molecules. Several studies indicate that this mechanism of cell death also exists in fish. The sequencing of fish homologues of MHC class I and CD3+ T cells suggests that CD8+ MHC class I presentation is similar to that which occurs in higher vertebrates (Fisher et al., 2006). Furthermore, we found a direct monoclonal antibody against the recombinant protein Onmy-UBA*501, the MHC class I molecule of rainbow trout, which are expressed in the same cell types as classical mammalian MHC mammalian class I molecules (Dijkstra et al., 2003). Like other vertebrates, fish acquire immunocompetency during ontogeny. In sea bass (*Dicentrarchus labrax* L.), T cells appear early in larval development, at least five days after the start of incubation (Dos Santos et al., 2000). Fisher et al. (2006) have demonstrated in rainbow trout that transcripts of CD8+ T cells and MHC class I are detectable one week after insemination, while TCR mRNA in T cells appears one week after insemination. MHC class I molecules and CD8 are expressed from the larval stage, suggesting that very young fish can be vaccinated to develop cellular immunity. In fact, up to 14 days after the end of incubation young trout can destroy skin allografts, a process accompanied by infiltration of cytotoxic T lymphocytes (Tatner and Manning, 1983).

6.4. Cytokines involved in adaptive immunity

Cytokines have been described in adaptive immunity in fish and with the recent discovery of CD4

in teleosts in it seems likely these cytokines will drive the activation and differentiation of T helper cell subsets to release different cytokine repertoires (Secombes, 2008). As stated above, type I and type II IFN are present that could potentially drive Th1 cell differentiation, with IFN- γ a potential effector of Th1 responses altogether with IL-12 (Yoshiura et al., 2003; Nascimento et al., 2007), IL-15 (Wang et al., 2007) and IL-18 (Huising et al., 2004; Zou et al., 2004). In relation to Th2-type cytokines, a molecule which is claimed to be a homologue of IL-4 has been found recently; however, it has no obvious similarity to known vertebrate IL-4 genes (Li et al., 2007). However, the presence of the transcription factor that drives Th2 cell differentiation, GATA3, suggests that Th2 responses may exist in fish, although this remains to be proven (Secombes, 2008). Finally, both IL-10 (Zou et al., 2003; Pinto et al., 2006) and TGF- β (Hardie et al., 1998) are present in fish, and thus may form the effectors of potential T regulatory cells.

7. Environmental effect on immune response

The immune system and response of fish can be greatly influenced by various external factors like temperature, light, water quality, salinity and different stress inducers (Magnadottir, 2010).

Decreases in temperature, which are important due to the poikilothermic nature of fish, affect the rate of their physiological functions (Hayward et al., 2009; e.g., enzymatic activity). In the rainbow trout, photoperiod has also been shown to impact the immune response through a reduction in the number of circulating leukocytes that is caused by increasing daylight hours; in turn, leukocytes are increased to reduce the hours of exposure to this factor. An increase in the photoperiod can also generate increases in the activity of lysozyme and the circulating levels of IgM. Furthermore, oxygen levels in the environment may modulate the immune response; hypoxia depresses the respiratory burst activity of macrophages and lowers the levels of circulating antibodies, which in turn, are elevated by hyperoxia (Watts et al., 2001; Bowden, 2008). On the other hand, increased levels of suspended solids in the fish environment help to raise hematocrit levels to compensate for the diminished ability of the gills to capture oxygen. This factor also increases the activity of lysozyme and the levels

of circulating IgM as a response to elevated levels of pathogens that are expected in an environment rich in particles. Increased salinity also affects immune parameters by increasing the activity of lytic enzymes, the respiratory burst of macrophages and the circulating levels of IgM. However, the reason why these changes are associated with salinity is currently unknown, but some authors suggest that increases in salinity levels cause an increased burden of pathogens from the environment to generate a more favourable osmotic environment (Bowden, 2008). Changes in environmental pH levels show conflicting results for immune system parameters, such as levels of lysozyme and IgM in the circulation. Finally, stress in fish as a result of population density associated with cultivation and production management can increase circulating cortisol levels, generating a decrease in specific and non-specific immunity and, therefore, making the fish prone to opportunistic pathogens (Brydges et al., 2009; Ramsay et al., 2009).

8. Compounds that modulate the immune system in fish

Another approach to immunoprophylactic control is the use of probiotics and immunostimulants including specific dietary manipulation (Magnottir, 2010). These compounds are defined as chemicals, drugs, stress generators or other actions that raise nonspecific defence mechanisms or the specific immune response (Anderson, 1992) and include nucleotides, glucans and probiotics. These treatments are primarily aimed at enhancing the innate system and therefore are of value as general preventive measures in aquaculture (Magnadottir, 2010).

8.1. Nucleotides

Nucleotides are composed of a purine or pyrimidine base, a ribose or deoxyribose sugar and one or more phosphate groups. The term nucleotide in this context refers not only to a specific form of the compound, but also to all forms that contain purine or pyrimidine bases (Rudolph, 1994). These compounds are essential in many physiological and biochemical functions, among which are included the encoding and decoding of genetic information, the mediating of energy metabolism and cell signals, and also forming components of coenzymes, al-

losteric effectors and cellular agonists (Carver et al., 1995; Cosgrove, 1998). Nucleotide supplementation of fish was introduced by the studies of Burrells et al. (2001a,b), which indicated that dietary supplementation with nucleotides confers increased resistance on salmonids to viral, bacterial and parasitic diseases and improves the effectiveness of vaccination and the ability to osmoregulate. Further, the binding of these compounds to Toll-like receptor 9 (TLR-9) is thought to stimulate the activity of intracellular bacterial DNA and trigger innate and adaptive responses (Dalmo, 2005). With respect to the innate immune response, it is known that dietary nucleotides may influence the activity of macrophages (Gil, 2002) and natural killer cell (Carver et al., 1990). Sakai et al. (2001) demonstrated in common carp that exogenous nucleotides may influence the humoral and cellular components of the innate immune system by increasing the complement lytic activity of enzymes and the production of the superoxide anion by anterior kidney phagocytes. Li et al. (2004) reported that hybrid striped bass (*Morone saxatilis*) that were fed diets supplemented with nucleotides had a higher production of oxidative radicals from the neutrophils in their blood compared to fish that were fed a normal diet. In addition, nucleotides also influence the activity of lymphocytes and immunoglobulin production (Jyonouchi et al., 1993; Navarro et al., 1996). Ramadan et al. (1994), were the first to observe the effect of dietary supplementation with nucleotides, and described a stimulatory effect on the humoral and cellular immune response in tilapia after intramuscular injection or bath with formalin-killed *Aeromonas hydrophila*. Antibody titres and lymphocyte mitogenic responses of fish supplemented with nucleotides were significantly higher in fish that were fed the basal diet. Similar phenomena have been observed in rainbow trout (Burrells et al., 2001b; Leonardi et al., 2003), hybrid striped bass (Li et al., 2004) and Atlantic salmon (Burrells et al., 2001b). However, the mechanisms of action of these compounds are still not well understood.

8.2. Glucans

One of the most used substances in immunostimulation experiments in fish are various forms of β -glucans from different sources, normally introduced in the feed but also by intraperitoneal injection or as a vaccine adjuvant (Dalmo and Bogwald,

2008). Glucan-linked glucans β 1-3 and β 1-6 are the major components of the cell walls of yeasts and mycelial fungi. These polysaccharides are not antigenic to animals (Ballou, 1982), but have been proven to be potent activators of nonspecific mechanisms of antibacterial defences in fish (Di Luzio, 1985; Sherwood et al., 1987). The injection of glucans generates improvements in the fish innate immune response by increasing the activity of macrophages, complement activation and levels of lytic enzymes (Jeney and Anderson, 1993; Jorgensen et al., 1993; Ai et al., 2007). Furthermore, several studies have shown that injection of glucans improves antibacterial defences and efficiency of vaccination (Yano et al., 1991; Rorstad et al., 1993). Similarly, when administered orally, these compounds are capable of stimulating the immune response and disease resistance (Siwicki et al., 1994). Engstad and Robertsen (1995) inoculated Atlantic salmon with an *Aeromonas salmonicida* vaccine that contained glucans in its formulation; six weeks after inoculation, there was a greater antibody response in the vaccine-inoculated fish compared to the fish that were vaccinated with a preparation that contained only the bacteria.

8.3. Probiotics

Probiotics are organisms or substances that contribute to the intestinal microbial balance. Fuller (1989) defined probiotics as live microbial feed supplements which exert beneficial effects on the host animal by improving its intestinal microbial balance. Research on the use of probiotics in aquatic animals has increased the demand for sustainable aquaculture (Gatesoupe, 1999). Many of the probiotics used in aquaculture belong to the Lactobacillus, the genus *Bacillus*, photosynthetic bacteria or yeasts, although other genera or species have also been employed, including nitrifying bacteria, streptococci and Roseobacter (Wang et al., 2008). These have been tested in aquaculture with generally good results (Panigrahi et al., 2005; Aly et al., 2008; Ma et al., 2009). The benefits of supplements include improvements in feed values, contribution to enzymatic digestion, inhibition of pathogenic microorganisms, anti-mutagenic and anti-carcinogenic activity, growth-promoting factors and an augmentation of the immune response (Wang and Xu, 2006; Wang, 2007). Studies have shown that the use of *Bacillus* sp. in cropping

systems has improved the water quality, survival, growth rate and health status of juvenile giant prawn (*Penaeus monodon*) and also reduced the amount of pathogenic vibrios (Dalmine et al., 2001). Moreover, probiotics may also stimulate the innate defence through TLR, other cellular receptors or humoral factors (Magnadottir, 2010).

9. Concluding remarks

In the last years, great progress has been made in understanding the immune response of fish which has benefitted the growing aquaculture industry worldwide. One critical element of this area of research has been established: the immunomodulation of larval fish. The production of fish larvae is often hampered by high mortality rates and economic loss due to infectious diseases. As is well known, in fish the innate response has been considered an essential component in combating pathogens due to limitations of the adaptive immune system. Therefore, the immunomodulation of larval fish has been proposed as a potential method for improving larval survival by increasing the innate responses of developing animals until their adaptive immune response is sufficiently developed to mount an effective response to the pathogen (Dalmo, 2005). This is a crucial point in further research that is currently focused on increasing the survival of larval fish. Finally, Dalmo (2005) suggest that the development of strategies to control pathogen load and immuno-prophylactic measures must be implemented further to realise the full potential of the production of marine fish larvae and thus improve the overall production of adult fish

REFERENCES

- Acton RT, Weinheimer PF, Hall SJ, Niedermeier W, Shelton E, Bennett JC (1971): Tetrameric immune macroglobulins in three orders of bony fishes. *Proceedings of the National Academy of Sciences of the United States of America* 68, 107–111.
- Ai Q, Mai K, Zhang L, Tan B, Zhang W, Xu W, Li H (2007): Effects of dietary [β]-1, 3 glucan on innate immune response of large yellow croaker, *Pseudosciaena crocea*. *Fish and Shellfish Immunology* 22, 394–402.
- Alexander JB, Ingram GA (1992): Noncellular nonspecific defense mechanisms of fish. *Annual Review of Fish Disease* 2, 249–279.
- Aly SM, Abdel-Galil Ahmed Y, Abdel-Aziz Ghareeb A, Mohamed MF (2008): Studies on *Bacillus subtilis* and *Lactobacillus acidophilus*, as potential probiotics, on the immune response and resistance of tilapia nilotica (*Oreochromis niloticus*) to challenge infections. *Fish and Shellfish Immunology* 25, 128–136.
- Anderson DP (1992): Immunostimulants, adjuvants and vaccine carriers in fish: applications to aquaculture. *Annual Review of Fish Disease* 2, 281–307.
- Aranishi F, Nakane M (1997): Epidermal proteases of the Japanese eel. *Fish Physiology and Biochemistry* 16, 471–478.
- Arkoosh MR, Kaattari SL (1991): Development of immunological memory in rainbow trout (*Oncorhynchus mykiss*). I. An immunochemical and cellular analysis of the B cell response. *Developmental and Comparative Immunology* 15, 279–293.
- Arma NR, Hirono I, Aoki T (2004): Characterization of expressed genes in kidney cells of Japanese flounder *Paralichthys olivaceus* treatment with ConA/PMA and LPS. *Fish Pathology* 39, 189–196.
- Armstrong PB, Quigley JP (1999): α -2-Macroglobulin: an evolutionarily conserved arm of the innate immune system. *Developmental and Comparative Immunology* 23, 375–390.
- Ballou CE (1982): Yest cell wall and cell surface. In: Strathern JN, Jones EW, Broach JR (eds.): *The Molecular Biology of the Yeast Saccharomyces: Metabolism and Gene Expression*. Cold Spring Harbor Laboratories, New York, USA. 335–360.
- Bayne CJ, Gerwick L (2001): The acute phase response and innate immunity of fish. *Developmental and Comparative Immunology* 25, 725–743.
- Birkemo GA, Luders T, Andersen O, Nes IF, Nissen-Meyer J (2003): Hipposin, a histone-derived antimicrobial peptide in Atlantic halibut (*Hippoglossus hippoglossus* L.). *Journal of Biochemistry, Molecular Biology and Biophysics* 1646, 207–215.
- Blazer VS (1991): Piscine macrophage function and nutritional influences: A review. *Journal of Aquatic Animal Health* 3, 77–86.
- Boes M (2000): Role of natural and immune IgM antibodies in immune responses. *Molecular Immunology* 37, 1141–1149.
- Boshra H, Li J, Sunyer JO (2006): Recent advances on the complement system of teleost fish. *Fish and Shellfish Immunology* 20, 239–262.
- Brydges NM, Boulcott P, Ellis T, Braithwaite VA (2009): Quantifying stress responses induced by different handling methods in three species of fish. *Applied Animal Behaviour Science* 116, 295–301.

- Bowden TJ (2008): Modulation of the immune system of fish by their environment. *Fish and Shellfish Immunology* 25, 373–383.
- Bowden TJ, Butler R, Bricknell IR, Ellis AE (1997): Serum trypsininhibitory activity in five species of farmed fish. *Fish and Shellfish Immunology* 7, 377–385.
- Bowden TJ, Cook P, Rombout JHWM (2005): Development and function of the thymus in teleosts. *Fish and Shellfish Immunology* 19, 413–427.
- Burrells C, William PD, Forno PF (2001a): Dietary nucleotides: a novel supplement in fish feeds. 1. Effects on resistance to diseases in salmonids. *Aquaculture* 199, 159–169.
- Burrells C, William PD, Southage PJ, Wadsworth SL (2001b): Dietary nucleotides: a novel supplement in fish feeds. 2. Effects on vaccination, salt water transfer, growth rate and physiology of Atlantic salmon. *Aquaculture* 199, 171–184.
- Cain KD, Jones DR, Raison RL (2000): Characterisation of mucosal and systemic immune responses in rainbow trout (*Oncorhynchus mykiss*) using surface plasmon resonance. *Fish and Shellfish Immunology* 10, 651–666.
- Carver JD, Cox WI, Barness LA (1990): Dietary nucleotide effects upon murine natural killer cell activity and macrophage activation. *Journal of Parenteral and Enteral Nutrition* 14, 18–22.
- Castillo A, Sanchez C, Dominguez J, Kaattari SL, Villena AJ (1993): Ontogeny of IgM and IgM-bearing cells in rainbow trout. *Developmental and Comparative Immunology* 17, 419–424.
- Chantanachookhin C, Seikai T, Tanaka M (1991): Comparative study of the ontogeny of the lymphoid organs in three species of marine fish. *Aquaculture* 99, 143–155.
- Chinchar VG, Bryan L, Silphadaung U, Noga E, Wade D, Rollins-Smith L (2004): Inactivation of viruses infecting ectothermic animals by amphibian and piscine antimicrobial peptides. *Virology* 323, 268–275.
- Cook MT, Hayball PJ, Birdseye L, Bagley C, Nowak BF, Hayball JD (2003): Isolation and partial characterization of a pentraxin-like protein with complement-fixing activity from snapper (*Pagrus auratus*, Sparidae) serum. *Developmental and Comparative Immunology* 27, 579–588.
- Cook MT, Hayball PJ, Nowak BF, Hayball JD (2005): The opsonising activity of a pentraxin-like protein isolated from snapper (*Pagrus auratus*, Sparidae) serum. *Developmental and Comparative Immunology* 29, 703–712.
- Cosgrove M (1998): Nucleotides. *Nutrition* 14, 748–751.
- Dalmo RA (2005): Ontogeny of the fish immune system. *Fish and Shellfish Immunology* 19, 395–396.
- Dalmin G, Kathiresan K, Purushothaman A (2001): Effect of probiotics on bacterial population and health status of shrimp in culture pond ecosystem. *Indian Journal of Experimental Biology* 39, 939–942.
- Dalmo RA, Bogwald J (2008): β -glucans as conductors of immune symphonies. *Fish and Shellfish Immunology* 25, 384–396.
- Danneving BH, Lauve A, McPress LC, Landsverk T (1994): Receptor-mediated endocytosis and phagocytosis by rainbow trout head kidney sinusoidal cells. *Fish and Shellfish Immunology* 4, 3–18.
- Davis JM, Clay H, Lewis JL, Ghori N, Herbomel P, Ramakrishnan L (2002): Real-time visualization of mycobacterium-macrophage interactions leading initiation of granuloma formation in zebrafish embryos. *Immunity* 17, 693–702.
- De Haas CJ, van Leeuwen EMM, van Bommel T, Verhoef J, van Kessel KPM, van Strijp JAG (2000): Serum amyloid P component bound to Gram-negative bacteria prevents lipopolysaccharide-mediated classical pathway complement activation. *Infection and Immunity* 68, 1753–1759.
- De Veer MJ, Holko M, Frevel M, Walker E, Der S, Paranjape JM, Silverman RH, Williams BRG (2001): Functional classification of interferon-stimulated genes identified using microarrays. *Journal of Leukocyte Biology* 69, 912–920.
- DeVries ME, Kelvin AA, Xu L, Ran L, Robinson J, Kelvin DJ (2006): Defining the origins and evolution of the chemokine/chemokine receptor system. *Journal of Immunology* 176, 401–415.
- Dijkstra JM, Kollner B, Aoyagi K, Sawamoto Y, Kuroda A, Ototake M (2003): The rainbow trout classical MHC class I molecule Onmy-UBA 501 is expressed in similar cell types as mammalian classical MHC class I molecules. *Fish and Shellfish Immunology* 14, 1–23.
- Di Luzio NR (1985): Update on the immunomodulating activities of glucans. *Seminars in Immunopathology* 8, 387–400.
- Dinareello CA (1997): Interleukin-1. *Cytokine Growth Factor Review* 8, 253–265.
- Dos Santos NM, Romano N, de Sousa M, Ellis AE, Rombout JH (2000): Ontogeny of B and T cells in sea bass (*Dicentrarchus labrax*, L.). *Fish and Shellfish Immunology* 10, 583–596.
- Elcombe BM, Chang RJ, Taves CJ, Winkelhake JL (1985): Evolution of antibody structure and effector functions: comparative hemolytic activities of monomeric and tetrameric IgM from rainbow trout, *Salmo gairdnerii*. *Comparative Biochemistry and Physiology* 80, 697–706.
- Ellingsen T, Strand C, Monsen E, Bogwald J, Dalmo RA (2005): The ontogeny of complement component C3

- in the spotted wolfish (*Anarhichas minor* Olafsen). *Fish and Shellfish Immunology* 18, 351–358.
- Ellis AE (2001): Innate host defence mechanism of fish against viruses and bacteria. *Developmental and Comparative Immunology* 25, 827–839.
- Engstad RE, Robertsen B (1995): Effects of structurally different yeast B glucans on immune responses in Atlantic salmon (*Salmo salar*). *Journal of Marine Biotechnology* 3, 203–207.
- Evans D, Jaso-Friedmann L (1992): Nonspecific cytotoxic cells as effectors of immunity in fish. *Annual Review of Fish Disease* 2, 109.
- Faisal M, Ahmed G, Peters II, Cooper EL (1989): Natural cytotoxicity of tilapia leukocytes. *Disease of Aquatic Organisms* 7, 17–22.
- Ferguson HW (1989): *Systemic Pathology of Fish. A text and atlas comparative tissue response in diseases of teleost.* Iowa state University Press. Ames. Iowa, USA. 5–103.
- Fischer U, Utke K, Somamoto T, Kollner B, Ototake M, Nakanishi T (2006): Cytotoxic activities of fish leukocytes. *Fish and Shellfish Immunology* 20, 209–226.
- Fuller R (1989): Probiotic in man and animals. *Journal of Applied Bacteriology* 66, 365–378.
- Gatesoupe FJ (1999): The use of probiotics in aquaculture. *Aquaculture* 180, 147–165.
- Gil A (2002): Modulation of the immune response mediated by dietary nucleotides. *European Journal of Clinical Nutrition* 56, S1–S4.
- Greenlee AD, Brown RA, Ristlow SS (1991): Nonspecific cytotoxic cells of Rainbow trout (*Oncorhynchus mykiss*) kill Yac-1 targets by both necrotic and apoptotic mechanisms. *Developmental and Comparative Immunology* 15, 153–164.
- Grinde B, Lie O, Poppe T, Salte R (1988): Species and individual variation in lysozyme activity in fish of interest in aquaculture. *Aquaculture* 68, 299–304.
- Hanington PC, Wang T, Secombes CJ, Belosevic M (2007): Growth factors of lower vertebrates: characterization of goldfish (*Carassius auratus* L.) macrophage colony-stimulating factor-1. *Journal of Biological Chemistry* 282, 31865–31872.
- Hardie LJ, Laing KJ, Daniels GD, Grabowski PS, Cunningham C, Secombes CJ (1998): Isolation of the first piscine transforming growth factor β gene: analysis reveals tissue specific expression and a potential regulatory sequence in rainbow trout (*Oncorhynchus mykiss*). *Cytokine* 10, 555–563.
- Hasegawa S, Nakayasu C, Yoshitomi T, Nakanishi T, Okamoto N (1998): Specific cell-mediated cytotoxicity against an allogeneic target cell line in isogenic ginbuna crucian carp. *Fish and Shellfish Immunology* 8, 303–313.
- Hatten F, Fredriksen A, Hordvik I, Endresen C (2001): Presence of IgM in cutaneous mucus, but not in gut mucus of Atlantic salmon, *Salmo salar*. Serum IgM is rapidly degraded when added to gut mucus. *Fish and Shellfish Immunology* 11, 257–268.
- Hayward CJ, Bott NJ, Nowak BF (2009): Seasonal epizootics of sea lice, *Caligus* spp., on southern bluefin tuna, *Thunnus maccoyii* (Castelnau), in a long-term farming trial. *Journal of Fish Disease* 32, 101–106.
- Hellio C, Pons AM, Beaupoil C, Bourgougnon N, Gal YL (2002): Antibacterial, antifungal and cytotoxic activities of extracts from fish epidermis and epidermal mucus. *International Journal of Antimicrobial Agents* 20, 214–2199.
- Hibiya T (ed.) (1994): *An Atlas of Fish Histology. Normal and Pathological Features.* Gustav Fischer Verlag, Stuttgart, Germany. 5–125.
- Hinuma S, Abo T, Kumagai K, Hata M (1980): The potent activity of fresh water fish kidney cells in cell-killing. I. Characterization and species distribution of cytotoxicity. *Developmental and Comparative Immunology* 4, 653–666.
- Holland MC, Lambris JD (2002): The complement system of teleosts. *Fish and Shellfish Immunology* 12, 399–420.
- Hoover GJ, El-Mowafi A, Simko E, Kocal TE, Ferguson HW, Hayes MA (1998): Plasma proteins of rainbow trout (*Oncorhynchus mykiss*) isolated by binding to lipopolysaccharide from *Aeromonas salmonicida*. *Comparative Biochemistry and Physiology* 120, 559–569.
- Huising MO, Stet RJM, Savelkoul HFJ, Verburg-van Kemenade BML (2004): The molecular evolution of the interleukin-1 family of cytokines; IL-18 in teleost fish. *Developmental and Comparative Immunology* 28, 395–413.
- Ingram GA (1980): Substances involved in the natural resistance of fish to infection a review. *Journal of Fish Biology* 16, 23–60.
- Jeney G, Anderson DP (1993): Glucan injection or bath exposure given alone or in combination with a bacteria enhance the nonspecific defense mechanisms in rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 116, 315–329.
- Jenkins JA, Ourth DD (1993): Opsonic effect of the alternative complement pathway on channel catfish peripheral blood phagocytes. *Veterinary Immunology Immunopathology* 39, 447–459.
- Jenkins PG, Wrathmell AB, Harris JE, Pulsford AL (1994): Systemic and mucosal immune response to enterically delivered antigen in *Oreochromis mossambicus*. *Fish and Shellfish Immunology* 4, 255–271.

- Jones DR, Hannan CM, Russel-Jones GJ, Raison RL (1999): Selective B cell non-responsiveness in the gut of the rainbow trout (*Oncorhynchus mykiss*). *Aquaculture* 172, 29–39.
- Jorgensen JB, Sharp GJ, Secombes CJ, Robertsen B (1993): Effect of yeast cell wall glucan on the bacterial activity of rainbow trout macrophages. *Fish and Shellfish Immunology* 3, 267–277.
- Jyonouchi, H, Zhang L, Tomita Y (1993): Studies of immunomodulating actions of RNA/nucleotides: RNA/Nucleotides enhance in vitro immunoglobulin production by human peripheral blood mononuclear cells in response to T-dependent stimuli. *Pediatric Research* 33, 458–465.
- Kaattari SL (1992): Fish B lymphocytes: Defining their form and function. In: Faisal M, Hetrick FM (eds.): *Annual Review of Fish Diseases*. Vol. 2. Pergamon, Tarrytown, NY, USA. 161–180.
- Kileng O, Brundtland MI, Robertsen B (2007): Infectious salmon anemia virus is a powerful inducer of key genes of the type I interferon system of Atlantic salmon, but is not inhibited by interferon. *Fish and Shellfish Immunology* 23, 378–389.
- Kodama H, Matsuoka Y, Tanaka Y, Liu YC, Iwasaki T, Watarai S (2004): Changes of C-reactive protein levels in rainbow trout (*Oncorhynchus mykiss*) sera after exposure to anti-ectoparasitic chemicals used in aquaculture. *Fish and Shellfish Immunology* 16, 589–597.
- Kono T, Bird S, Sonoda K, Savan R, Secombes CJ, Sakai M (2008): Characterization and expression analysis of an interleukin-7 homologue in the Japanese pufferfish, *Fugu rubripes*. *FEBS Journal* 275, 1213–1226.
- Lam SH, Chua HL, Gong Z, Lam TJ, Sin Y (2004): Development and maturation of the immune system in zebrafish, *Danio rerio*: a gene expression profiling, in situ hybridization and immunological study. *Developmental and Comparative Immunology* 28, 9–28.
- Lammens M, Decostere A, Haaesebrouck F (2000): Effects of *Flavobacterium psychrophilum* strains and their metabolites on the oxidative activity of rainbow trout *Oncorhynchus mykiss* phagocytes. *Disease of Aquatic Organisms* 41, 173–179.
- Lange S, Magnadottir B (2003): Spontaneous haemolytic activity of Atlantic halibut (*Hippoglossus hippoglossus* L.) and sea bass (*Dicentrarchus labrax*) serum. *Comparative Biochemistry and Physiology, Molecular Biology* 136, 99–106.
- Lange SR, Bambir S, Dodds AW, Magnadottir B (2004a): The ontogeny of complement component C3 in Atlantic cod (*Gadus morhua* L.) an immunohistochemical study. *Fish and Shellfish Immunology* 16, 359–367.
- Lange SR, Bambir S, Dodds AW, Magnadottir B (2004b): An immunohistochemical study on complement component C3 in juvenile Atlantic halibut (*Hippoglossus hippoglossus* L.). *Developmental and Comparative Immunology* 28, 593–601.
- Leonardi M, Sandino AM, Klempau A (2003): Effect of a nucleotide-enriched diet on the immune system, plasma cortisol levels and resistance to infectious pancreatic necrosis (IPN) in juvenile rainbow trout (*Oncorhynchus mykiss*). *Bulletin of the European Association of Fish Pathologists* 23, 52–59.
- Li P, Lewis DH, Gatlin DM (2004): Dietary oligonucleotide from yeast RNA influences immune responses and resistance of hybrid striped bass (*Morone chrysops* × *M. saxatilis*) to *Streptococcus iniae* infection. *Fish and Shellfish Immunology* 16, 561–569.
- Li JH, Shao JZ, Xiang LX, Wen Y (2007): Cloning, characterization and expression analysis of pufferfish interleukin-4 cDNA: the first evidence of Th2-type cytokine in fish. *Molecular Immunology* 44, 2078–2086.
- Lie O, Evensen O, Sorensen A, Froysadal E (1989): Study on lysozyme activity in some fish species. *Disease of Aquatic Organism* 6, 1–5.
- Liu Y, Iwasaki T, Watarai S, Kodama H (2004): Effect of turpentine oil on C-reactive protein (CRP) production in rainbow trout (*Oncorhynchus mykiss*). *Fish and Shellfish Immunology* 17, 203–210.
- Lorenzen N, LaPatra SE (1999): Immunity to rhabdovirus in rainbow trout: the antibody response. *Fish and Shellfish Immunology* 9, 345–360.
- Lumsden JS, Ostland VE, Byrne PJ, Ferguson HW (1993): Detection of a distinct gill-surface antibody response following horizontal infection and bath challenge of brook trout *Salvelinus fontinalis* with *Flavobacterium branchiophilum*, the causative agent of bacterial gill disease. *Disease of Aquatic Organism* 16, 21–27.
- Lund V, Olafsen JA (1998): A comparative study of pentraxin-like proteins in different fish species. *Developmental and Comparative Immunology* 22, 185–194.
- Ma CW, Cho YS, Oh KH (2009): Removal of pathogenic bacteria and nitrogens by *Lactobacillus* spp. JK-8 and JK-11. *Aquaculture* 287, 266–270.
- Maier VH, Dorn KV, Gudmundsdottir BK, Gudmundsson GH (2008): Characterisation of cathelicidin gene family members in divergent fish species. *Molecular Immunology* 45, 3723–3730.
- Magnadottir B (2006): Innate immunity of fish (overview). *Fish and Shellfish Immunology* 20, 137–151.
- Magnadottir B (2010): Immunological control of fish diseases. *Journal of Marine Biotechnology* 12, 361–379.
- Magnadottir B, Jonsdottir H, Helgason S, Bjornsson B, Jorgensen TO, Pilstrom L (1999): Humoral immune

- parameters in Atlantic cod (*Gadus morhua* L.). II. The effects of size and gender under different environmental conditions. *Comparative Biochemistry and Physiology* 122B, 181–188.
- Magnadottir B, Jonsdottir H, Helgason S, Bjornsson B, Jorgensen TO, Pilstrom L (2001): Immuneparameters of immunised cod (*Gadus morhua* L.). *Fish and Shellfish Immunology* 10, 75–89.
- Magnadottir B, Lange S, Gudmundsdottir S, Bogwald J, Dalmo RA (2005): Ontogeny of humoral immune parameters in fish. *Fish and Shellfish Immunology* 19, 429–439.
- Manning MJ, Nakanishi T (1996): Cellular defenses. In: Iwama GK, Nakanishi T (eds.): *Fish Physiology*. XV. The fish immune system. Academic press. London, England. 159–205.
- Mathew JA, Guo YX, Goh KP, Chan J, Verburg van Kemnade BM, Kwang J (2002): Characterisation of a monoclonal antibody to carp IL-1beta and the development of a sensitive capture ELISA. *Fish and Shellfish Immunology* 13, 85–95.
- Matsushita M, Endo Y, Nonaka M, Fujita T (1998): Complement-related serine proteases in tunicates and vertebrates. *Current Opinion in Immunology* 10, 29–35.
- Matsuyama H, Yano T, Yamakawa T, Nakao M (1992): Opsonic effect of the third complement component (C3) of carp (*Cyprinus carpio*) on phagocytosis by neutrophils. *Fish and Shellfish Immunology* 2, 69–78.
- McKinney EC, Achmale MC (1994): Damsel fish with neurofibromatosis exhibit cytotoxicity in toward tumor targets. *Developmental and Comparative Immunology* 18, 305–313.
- Morrison RN, Nowak B (2002): The antibody response of teleost fish. *Seminars in Avian and Exotic Pet Medicine* 11, 46–54.
- Mulero V, Meseguer J (1998): Functional characterisation of a macrophage-activating factor produced by leucocytes of gilthead seabream (*Sparus aurata* L.). *Fish and Shellfish Immunology* 8, 143–156.
- Mulero I, Garcia Ayala A, Meseguer J, Mulero V (2007): Maternal transfer of immunity and ontogeny of autologous immunocompetence of fish: A mini review. *Aquaculture* 268, 244–250.
- Nakanishi T. (1986): Seasonal changes in the humoral immune response and the lymphoid tissues of the marine teleost, *Sebastes marmoratus*. *Veterinary Immunology and Immunopathology* 12, 213–221.
- Nascimento DS, do Vale A, Toma's AM, Zou J, Secombes CJ, dos Santos NMS (2007): Cloning, promoter analysis and expression in response to bacterial exposure of sea bass (*Dicentrarchus labrax* L.) interleukin-12 p40 and p35 subunits. *Molecular Immunology* 44, 2277–2791.
- Nauta AJ, Daha M, van Kooten C, Roos A (2003): Recognition and clearance of apoptotic cells: a role for complement and pentraxins. *Trends in Immunology* 24, 148–154.
- Navarro J, Ruiz-Bravo A, Jimenez-Varela M, Gil A (1996): Modulation of antibody-forming cell and mitogen-driven lymphoproliferative responses by dietary nucleotides in mice. *Immunology Letters* 53, 141–145.
- Nikoskelainen S, Lehtinen J, Lilius EM (2002): Bacteriolytic activity of rainbow trout (*Oncorhynchus mykiss*) complement. *Developmental and Comparative Immunology* 26, 797–804.
- Olesen NJ, Jorgensen PEV (1986): Quantification of serum immunoglobulin in rainbow trout *Salmo gairdneri* under various environmental conditions. *Disease of Aquatic Organism* 1, 183–189.
- Ooi EL, Verjan N, Hirono I, Nochi T, Kondo H, Aoki T, Kiyono H, Yuki Y. (2008): Biological characterisation of a recombinant Atlantic salmon type I interferon synthesized in *Escherichia coli*. *Fish and Shellfish Immunology* 24, 506–513.
- Panigrahi A, Kiron V, Puangkaew J, Kobayashi T, Satoh S, Sugita H (2005): The viability of probiotic bacteria as a factor influencing the immune response in rainbow trout *Oncorhynchus mykiss*. *Aquaculture* 243, 241–254.
- Petrie-Hanson L, Ainsworth AJ (2000): Differential cytochemical staining characteristics of channel catfish leukocytes identify cell populations in lymphoid organs. *Veterinary Immunology and Immunopathology* 73, 129–144.
- Pinto RD, Nascimento DS, Reis MIR, do Vale A, dos Santos NMS (2006): Molecular characterization, 3D modelling and expression analysis of sea bass (*Dicentrarchus labrax*) interleukin-10. *Molecular Immunology* 44, 2066–2075.
- Press CM, Dannevig BH, Landsverk T (1994): Immune and enzyme histochemical phenotypes of lymphoid and nonlymphoid cells within the spleen and head kidney of Atlantic salmon (*Salmo salar* L.). *Fish and Shellfish Immunology* 4, 79–93.
- Ramadan A, Afifi NA, Moustafa M, Samy AM (1994): The effect of ascogen on the immune response of tilapia fish to *Aeromonas hydrophila* vaccine. *Fish and Shellfish Immunology* 5, 159–165.
- Ramsay JM, Feist GW, Varga ZM, Westerfield M, Kent ML, Schreck CB (2009): Whole-body cortisol response of zebrafish to acute net handling stress. *Aquaculture* 297, 157–162.
- Razquin BE, Castillo A, Lopez-Fierro P, Alvarez F, Zapata A, Villena A (1990): Ontogeny of IgM-producing cells in the lymphoid organs of rainbow trout, *Salmo gairdneri*

- Richardson: an immunological and enzyme-histochemical study. *Journal of Fish Biology* 36, 159–173.
- Robertsen B (2006): The interferon system of teleost fish. *Fish and Shellfish Immunology* 20, 172–191.
- Rokenes TP, Larsen R, Robertsen B (2007): Atlantic salmon ISG15: expression and conjugation to cellular proteins in response to interferon, double-stranded RNA and virus infections. *Molecular Immunology* 44, 950–959.
- Rombout JH, Blok LJ, Lamers CHJ (1986): Immunization of carp (*Cyprinus carpio*) with a *Vibrio anguillarum* bacterin: Indications for a common mucosal immune system. *Developmental and Comparative Immunology* 10, 341–351.
- Rombout JH, Taverne N, van de Kamp M, Taverne-Thiele AJ (1993): Differences in mucus and serum immunoglobulin of carp (*Cyprinus carpio* L.). *Developmental and Comparative Immunology* 17, 309–317.
- Rombout JH, Huttenhuis HBT, Picchietti S, Scapigliati S (2005): Phylogeny and ontogeny of fish leucocytes. *Fish and Shellfish Immunology* 19, 441–455.
- Rorstad G, Aasjord PM, Robertsen B (1993): Adjuvant effect of a yeast glucan in vaccines against furunculosis in Atlantic salmon (*Salmo salar*). *Fish and Shellfish Immunology* 3, 179–190.
- Rudolph FB (1994): The biochemistry and physiology of nucleotides. *Journal of Nutrition* 124, 124S–127S.
- Sakai DK (1992): Repertoire of complement in immunological defense mechanisms of fish. *Annual Review of Fish Disease* 2, 223–247.
- Sakai M, Taniguchi K, Mamoto K, Ogawa H, Tabata M (2001): Immunostimulant effects of nucleotide isolated from yeast RNA on carp, *Cyprinus carpio* L. *Journal of Fish Disease* 24, 433–438.
- Samuel CE (2001): Antiviral actions of interferons. *Clinical Microbiology Review* 14, 778–809.
- Sanchez C, Babin M, Tomillo J, Ubeira FM, Dominguez J (1993): Quantification of low levels of rainbow trout immunoglobulin by enzyme immunoassay using two monoclonal antibodies. *Veterinary Immunology and Immunopathology* 36, 65–74.
- Sanchez C, Alvarez A, Castillo A, Zapata A, Villena A, Dominguez J (1995): Two different subpopulations of Ig-bearing cells in lymphoid organs of rainbow trout. *Developmental and Comparative Immunology* 19, 79–86.
- Sangrador-Vegas A, Martin SA, O’Dea PG, Smith TJ (2000): Cloning and characterization of the rainbow trout (*Oncorhynchus mykiss*) type II interleukin-1 receptor cDNA. *European Journal of Biochemistry* 267, 7031–703.
- Saurabh S, Sahoo PK (2008): Lysozyme: an important defence molecule of fish innate immune system. *Aquaculture Research* 39, 223–239.
- Savan R, Sakai M (2006): Genomics of fish cytokines. *Comparative Biochemistry and Physiology D* 1, 9–101.
- Scapigliati G, Chausson F, Cooper EL, Scalia D, Mazzini (1997): Qualitative and quantitative analysis of serum immunoglobulins of four Antarctic fish species. *Polar Biology* 18, 209–213.
- Schroder MB, Villena AJ, Jorgensen TO (1998): Ontogeny of lymphoid organs and immunoglobulin producing cells in Atlantic cod (*Gadus morhua* L.). *Developmental and Comparative Immunology* 22, 507–517.
- Secombes CJ (2008): Will advances in fish immunology change vaccination strategies? *Fish and Shellfish Immunology* 25, 409–416.
- Secombes CJ, Fletcher TC (1992): The role of phagocytes in the protective mechanisms of fish. *Annual Review of Fish Disease* 2, 53–71.
- Shen L, Stuge TB, Zhou H, Khayat M, Barker KS, Quinloun SM, Wilson M, Bengten E, Chinchar VG, Clem LW, Miller NW (2002): Channel catfish cytotoxic cells: a mini-review. *Developmental and Comparative Immunology* 26, 141–149.
- Shephard KL (1994): Functions for fish mucus. *Reviews in Fish Biology Fisheries* 4, 401–429.
- Sherwood ER, Williams DL, McNamee RB, Jones EL, Browder IW, Di Luzio NR (1987): Enhancement of interleukin 2 production by soluble glucan. *Journal of Immunopharmacology* 9, 261–267.
- Siwicki AK, Anderson DP, Rumsey GL (1994): Dietary intake of immunostimulants by rainbow trout affects non specific immunity and protects against furunculosis. *Veterinary Immunology and Immunopathology* 41, 125–139.
- Stafford JL, Belosevic M (2003): Transferrin and the innate immune response of fish: identification of a novel mechanism of macrophage activation. *Developmental and Comparative Immunology* 27, 539–554.
- Subramaniam S, Stansberg C, Olsen L, Zou J, Secombes CJ, Cunningham C (2002): Cloning of a *Salmo salar* interleukin-1 receptor-like cDNA. *Developmental and Comparative Immunology* 26, 415–431.
- Subramaniam S, Mackinnon SL, Ross NW (2007): A comparative study on innate immune parameters in the epidermal mucus of various fish species. *Comparative Biochemistry and Physiology B* 148, 256–263.
- Subramaniam S, Ross NW, Mackinnon SL (2008): Comparison of antimicrobial activity in the epidermal mucus extracts of fish. *Comparative Biochemistry and Physiology B* 150, 85–92.
- Suzumura E, Kurata O, Okamoto N, Ykeda Y (1994): Characteristics of natural killer cells in carp. *Fish Pathology* 29, 199–203.
- Suzumoto BK, Schreck CB, McIntyre JD (1997): Relative resistances of three transferrin genotypes of coho

- salmon (*Oncorhynchus kisutch*) and their hematological responses to bacterial kidney disease. Journal of the Fisheries Research Board of Canada 34, 1–8.
- Sveinbjornsson B, Olsen R, Paulsen S (1996): Immunocytochemical localization of lysozyme in intestinal eosinophilic granule cells (EGCs) of Atlantic salmon, *Salmo salar* L. Journal of Fish Disease 19, 349–355.
- Szalai AJ, Norcum MT, Bly JE, Clem LW (1992): Isolation of an acute-phase and phosphorylcholine-reactive pentraxin from channel catfish (*Ictalurus punctatus*). Comparative Biochemistry and Physiology B 102, 535–543.
- Szalai AJ, Bly JE, Clem LW (1994): Changes in serum concentrations of channel catfish (*Ictalurus punctatus*) phosphorylcholine-reactive protein (PRP) in response to inflammatory agents, low temperature-shock and infection by the fungus *Saprolegnia* sp. Fish and Shellfish Immunology 4, 323–336.
- Tafalla C, Figueras A, Novoa B (2001): Viral hemorrhagic septicemia virus alters turbot *Scophthalmus maximus* macrophage nitric oxide production. Disease of Aquatic Organism 47, 101–107.
- Tatner MF, Manning MJ (1983): The ontogeny of cellular immunity in the rainbow trout, *Salmo gairdneri* Richardson, in relation to the stage of development of the lymphoid organs. Developmental and Comparative Immunology 7, 69–75.
- Trede NS, Zapata A, Zon LI (2001): Fishing for lymphoid genes. Trends in Immunology 22, 302–307.
- Uchida D, Hirose H, Chang PK, Aranishi F, Hirayabu E, Mano N, Mitsuya T, Prayitno SB, Natori M (2000): Characterization of Japanese eel immunoglobulin M and its level in serum. Comparative Biochemistry and Physiology B 127, 525–532.
- Vilain C, Wetzel MC, Du Pasquier L, Charlemagne J (1984): Structural and functional analysis of spontaneous antinitrophenyl antibodies in three cyprinid fish species: carp (*Cyprinus carpio*), goldfish (*Carassius auratus*) and tench (*Tinca tinca*). Developmental and Comparative Immunology 8, 611–622.
- Wang YB (2007): Effect of probiotics on growth performance and digestive enzyme activity of the shrimp *Penaeus vannamei*. Aquaculture 269, 259–264.
- Wang YB, Xu ZR (2006): Effect of probiotics for common carp (*Cyprinus carpio*) based on growth performance and digestive enzyme activities. Animal Feed Science and Technology 127, 283–292.
- Wang T, Holland J, Bols N, Secombes CJ (2007): Molecular and functional characterisation of interleukin-15 in rainbow trout *Oncorhynchus mykiss*: a potent inducer of interferon-gamma expression in spleen leucocytes. Journal of Immunology 179, 1475–1488.
- Wang YB, Li JR, Lin J (2008): Probiotics in aquaculture: challenges and outlook. Aquaculture 281, 1–4.
- Watts M, Munday BL, Burke CM (2001): Immune responses of teleost fish. Australian Veterinary Journal 79, 570–574.
- Whittington RJ, Munday BL, Akhlaghi M, Reddacliff GL, Carson J (1994): Humoral and peritoneal cell responses of rainbow trout (*Oncorhynchus mykiss*) to ovalbumin, *Vibrio anguillarum* and Freund's complete adjuvant following intraperitoneal and bath immunisation. Fish and Shellfish Immunology 4, 475–488.
- Whyte SK (2007): The innate immune response of finfish e A review of current knowledge. Fish and Shellfish Immunology 23, 1127–1151.
- Wilson MR, Warr GW (1992): Fish immunoglobulins and the genes that encode them. Annual Review of Fish Disease 2, 201–221.
- Wilson MR, Bengten E, Miller N, Clem LW, Du Pasquier L, Warr GW (1997): A novel chimeric Ig heavy chain from a teleost fish shares similarities to IgD. Proceedings of the National Academy of Sciences of the United States of America 94, 4593–4597.
- Willett C, Cortes A, Zuasti A, Zapata A. (1999): Early hematopoiesis and developing lymphoid organs in the zebrafish. Developmental Dynamics 214, 323–336.
- Withler RE, Evelyn TP (1990): Genetic variation in resistance to bacterial kidney disease within and between two strains of coho salmon from British Columbia. Transactions of the American Fisheries Society 119, 1003–1009.
- Wu Y, Wang S, Peng X (2003): Serum acute phase response (APR)-related proteome of loach to trauma. Fish and Shellfish Immunology 16, 381–389.
- Yano T (1996): The nonspecific immune system: Humoral defence. In: Iwama G, Nakanishi T (eds.): The Fish Immune System: Organism, Pathogen and Environment. Academic Press, San Diego. 105–157.
- Yano T, Matsuyama H, Mangindaan REP (1991): Polysaccharide induced protection of carp *Cyprinus carpio* against bacterial infection. Journal of Fish Disease 14, 577–582.
- Yin Z, Lam TL, Sin YM (1997): Cytokine-mediated antimicrobial immune response of catfish, *Clarias gariepinus*, as a defence against *Aeromonas hydrophila*. Fish and Shellfish Immunology 7, 93–104.
- Yoshiura Y, Kiryu I, Fujiwara A, Suetake H, Suzuki Y, Nakanishi T (2003): Identification and characterization of *Fugu orthologues* of mammalian interleukin-12 subunits. Immunogenetics 55, 296–306.
- Zapata A, Diez B, Cejalvo T, Gutierrez-De Frias C, Cortes A (2006): Ontogeny of the immune system of fish. Fish and Shellfish Immunology 20, 126–136.

Zou J, Clarke M, Secombes CJ (2003): Characterisation, expression and promoter analysis of an interleukin 10 homologue in the puffer fish, *Fugu rubripes*. Immunogenetics 55, 325–335.

Zou J, Bird S, Truckle J, Bols N, Horne M, Secombes CJ (2004): Identification and expression analysis of an IL-18 homologue and its alternatively spliced form in rainbow trout *Oncorhynchus mykiss*. European Journal of Biochemistry 271, 1913–1923.

Zou J, Tafalla C, Truckle J, Secombes CJ (2007): Identification of a second group of type I IFNs in fish sheds light on IFN evolution in vertebrates. Journal of Immunology 179, 3859–3871.

Received: 2010–10–27

Accepted after corrections: 2011–11–02

Corresponding Author:

Gabriel Moran, Universidad Austral de Chile, Faculty of Veterinary Science, Department of Pharmacology, Valdivia, Chile

Tel. +56 63 221936, E-mail: gmoran@uach.cl
