

Insights into the Evolution of Vitamin B₁₂ Auxotrophy from Sequenced Algal Genomes

Katherine E. Helliwell,¹ Glen L. Wheeler,^{2,3} Kyriacos C. Leptos,⁴ Raymond E. Goldstein,⁴ and Alison G. Smith^{*,1}

¹Department of Plant Sciences, University of Cambridge, Cambridge, United Kingdom

²Plymouth Marine Laboratory, Plymouth, United Kingdom

³Marine Biological Association of United Kingdom, The Laboratory, Plymouth, United Kingdom

⁴Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Cambridge, United Kingdom

*Corresponding author: E-mail: as25@cam.ac.uk.

Associate editor: Charles Delwiche

Abstract

Vitamin B₁₂ (cobalamin) is a dietary requirement for humans because it is an essential cofactor for two enzymes, methylmalonyl-CoA mutase and methionine synthase (METH). Land plants and fungi neither synthesize or require cobalamin because they do not contain methylmalonyl-CoA mutase, and have an alternative B₁₂-independent methionine synthase (METE). Within the algal kingdom, approximately half of all microalgal species need the vitamin as a growth supplement, but there is no phylogenetic relationship between these species, suggesting that the auxotrophy arose multiple times through evolution. We set out to determine the underlying cellular mechanisms for this observation by investigating elements of B₁₂ metabolism in the sequenced genomes of 15 different algal species, with representatives of the red, green, and brown algae, diatoms, and coccolithophores, including both macro- and microalgae, and from marine and freshwater environments. From this analysis, together with growth assays, we found a strong correlation between the absence of a functional METE gene and B₁₂ auxotrophy. The presence of a METE unitary pseudogene in the B₁₂-dependent green algae *Volvox carteri* and *Gonium pectorale*, relatives of the B₁₂-independent *Chlamydomonas reinhardtii*, suggest that B₁₂ dependence evolved recently in these lineages. In both *C. reinhardtii* and the diatom *Phaeodactylum tricornutum*, growth in the presence of cobalamin leads to repression of METE transcription, providing a mechanism for gene loss. Thus varying environmental conditions are likely to have been the reason for the multiple independent origins of B₁₂ auxotrophy in these organisms. Because the ultimate source of cobalamin is from prokaryotes, the selective loss of METE in different algal lineages will have had important physiological and ecological consequences for these organisms in terms of their dependence on bacteria.

Key words: algae, cobalamin, pseudogene, methionine synthase, vitamin B₁₂, vitamin auxotrophy, RT-PCR.

Introduction

Vitamins are essential for all organisms because they provide the precursors to enzyme cofactors important for metabolism. Animals must obtain these organic micronutrients in their diet, but plants and microorganisms generally synthesize de novo the cofactors they need (Smith et al. 2007). It is thus perhaps surprising that, despite their photosynthetic lifestyle, many algae exhibit vitamin auxotrophy, which is the inability to synthesize an organic nutrient essential for growth. Over half of all microalgal species require an exogenous supply of vitamin B₁₂ (cobalamin), whilst just over 20% require vitamin B₁ (thiamine) and a smaller proportion (5%) require biotin (vitamin B₇) (Croft et al. 2006). However, vitamin auxotrophy is not phylogenetically related but instead distributed throughout the algal clades, implying that it has arisen multiple times throughout evolution (Croft et al. 2005). Biotin and thiamine auxotrophy are the result of the loss of one or more of the biosynthetic enzymes to make the vitamin (Croft et al. 2006), but for cobalamin, this does not appear to be the case. Those algae that are not dependent on B₁₂

do not contain it unless it is supplied in the medium (Croft et al. 2005). In fact, B₁₂ biosynthesis appears to be confined to prokaryotes, and there is no evidence that any eukaryote is capable of producing the vitamin (Warren et al. 2002). Interestingly, not all bacteria make B₁₂—some, such as *Escherichia coli*, will use it if it is available, whereas other species have no enzymes that use B₁₂. A notable example is *Pelagibacter rubique*, a member of the highly abundant SAR11 clade of marine bacteria (Giovannoni et al. 2005).

Cobalamin is a complex Co²⁺-containing modified tetrapyrrole that acts as a cofactor for enzymes involved in C1 metabolism and certain radical reactions. In prokaryotes, there are over 20 enzymes that have a cobalamin cofactor (Marsh 1999), three of which have been found in eukaryotes. In humans, there are just two B₁₂-dependent enzymes, methylmalonyl-CoA mutase (MCM), involved in odd-chain fatty acid metabolism in the mitochondria, and methionine synthase (METH), which catalyzes the C1 transfer from methyltetrahydrofolate to homocysteine to make methionine. An alternative B₁₂-independent form of methionine synthase (METE) is found in land plants and fungi, so they

have no requirement for the cofactor and do not synthesize it. *E. coli* possesses both forms of methionine synthase and can utilize METH if B₁₂ is available in the environment, otherwise it uses METE. Similarly, there are both B₁₂-independent and B₁₂-dependent forms of ribonucleotide reductase (RNR) (type I & III RNR are B₁₂ independent, type II is B₁₂ dependent), involved in deoxyribose biosynthesis (Hamilton 1974; Carell and Seeger 1980). Many prokaryotes encode both types of enzyme and again appear to switch between them depending on the availability of cobalamin in the environment. All eukaryotes have the type I isoform, whilst *Euglena gracilis*, an excavate protist, is reported to have B₁₂-dependent RNR II as well (Hamilton 1974).

An initial search for the presence of genes encoding B₁₂-dependent enzymes in four sequenced algal genomes suggested that B₁₂ auxotrophy may be related to the form of methionine synthase present (Croft et al. 2005). The green alga *Chlamydomonas reinhardtii* and the diatom *Phaeodactylum tricorutum* have both METE and METH and are B₁₂ independent, whereas another diatom *Thalassiosira pseudonana*, which contains METH only, is B₁₂ dependent. The B₁₂-independent red alga *Cyanidioschyzon merolae* has METE only. However, algae are an extremely diverse group of organisms, so a much broader study is necessary to allow definitive conclusions to be drawn.

The dominant algal lineages have arisen from a complex succession of endosymbiotic events. Over a billion years ago, engulfment of a photosynthetic cyanobacterium by a heterotrophic protist gave rise to the basal algal groups: green and red algae and the glaucocystophytes (Delwiche 1999; Reyes-Prieto et al. 2007; Gould et al. 2008). Secondary endosymbiotic events then followed, in which one or more heterotrophic organisms engulfed either a red or a green alga, giving rise to algae with complex plastids (Moreira and Philippe 2001; Gould et al. 2008). These include the chromalveolate supergroup, which encompasses important marine phytoplankton such as diatoms, coccolithophores, and dinoflagellates, as well as brown algae (as defined in Hackett et al. 2007).

Here, we perform a comprehensive survey of B₁₂-dependent enzymes and associated proteins in 15 algal species with completed genome sequences that span the algal kingdom. Our results provide strong evidence that the major determinant for the B₁₂ requirements of algae relates to the isoform(s) of methionine synthase that they possess and that multiple independent losses of a functional METE is the principal factor underlying the evolution of B₁₂ auxotrophy.

Materials and Methods

Strains and Growth Conditions

The algal strains used were either from culture collections or gifts from colleagues (see [supplementary table 1, Supplementary Material](#) online, for more information and details of growth conditions). *Salmonella enterica* (AR3621) was a gift from Professor Martin Warren (University of Kent, UK) and was grown as described by Raux et al. (1996).

Sequence Similarity Searches and Putative Sequence Analysis and Verification

BlastP and TblastN (Altschul et al. 1990) sequence similarity searches were performed to assess the presence of each protein in the described algal genomes. The organisms and sequence IDs of the proteins that were used to perform these searches are as follows: *C. reinhardtii* METE (XP_001702934) and METH (76715), *Homo sapiens* MTRR (AAF16876.1), MCM (AAA59569.1), and CBLB (AAH11831.1), *E. coli* CBLA (P27254.2), *Eu. gracilis* RNR II (Q2PDF6.1) (protein IDs of all putative hits can be found in [supplementary table 2, Supplementary Material](#) online). The genome version of each species in which the searches were carried out are as follows: *C. reinhardtii* (v4) (Merchant et al. 2007), *Volvox carteri* f. *nagariensis* (v1) (Prochnik et al. 2010), *Coccomyxa* sp. C-169 (v1) (http://genome.jgi-psf.org/Coc_C169_1/Coc_C169_1.home.html), *Chlorella variabilis* NC64A (v1) (Blanc et al. 2010), *Micromonas pusilla* (CCMP 1545) (v2) (Worden et al. 2009), *Ostreococcus tauri* (OTH 95) (v2), *O. lucimarinus* (v2) (Palenik et al. 2007), and *Ostreococcus* sp. RCC809 (v2) (http://genome.jgi-psf.org/OstRCC809_2/OstRCC809_2.home.html), *P. tricorutum* (CCAP 1055/1) (v2) (Armbrust et al. 2004; Bowler et al. 2008), *Fragilariopsis cylindrus* (CCMP 1102) (v1) (<http://genome.jgi-psf.org/Fracy1/Fracy1.home.html>), *T. pseudonana* (CCMP 1335) (v3) (Armbrust et al. 2004), *Aureococcus anophagefferens* (CCMP 1984) (v1) (<http://genome.jgi-psf.org/Auran1/Auran1.home.html>), *Emiliania huxleyi* (CCMP 1516) (v1) (<http://genome.jgi-psf.org/Emihu1/Emihu1.home.html>). All of the above can be found at <http://genome.jgi-psf.org/>. The *Ectocarpus siliculosus* (CCAP 1310) (v1) sequence (Cock et al. 2010) can be found at <http://bioinformatics.psb.ugent.be/genomes/view/Ectocarpus-siliculosus> and that for *Cy. merolae* (Matsuzaki et al. 2004) at <http://merolae.biol.s.u-tokyo.ac.jp/>.

To verify the identity of all putative orthologous proteins, we employed the following techniques: 1) multiple sequence alignment using Jalview (Waterhouse et al. 2009) to check visually each gene model, 2) Genewise 2.0 analysis (Birney et al. 2004) to identify missing introns in poor gene models, and 3) Pfam (24.0) analysis (Finn et al. 2008) to identify conserved functional domains in each protein (METE: PF01717, PF08267; METH: PF02965, PF02965, PF02607, PF00809, and PF02574; RNR II: PF09747, PF02867; MCM: PF01642, PF02310; MTRR: PF00258, PF0667, and PF00175; CBLA: PF3308, PF00875; and CBLB: PF01923).

Treatment of Algae Prior to B₁₂ Growth Assays

Bacteria are a potential source of B₁₂, so prior to the growth assays, cultures were either obtained axenic (*P. tricorutum* and *F. cylindrus*) or treated to remove contaminating bacteria. *V. carteri* EVE was made axenic by several serial washings of spheroids in six-well microplates using a sterile pipette. The final washed spheroids were tested for the presence of bacterial contamination on lysogeny broth (LB) plates. *Gonium pectorale* was made axenic by serial streak

outs on standard Volvox medium agar plates followed by incubation at 28 °C. This was done multiple times until bacterial colonies were growing distinctly from algal colonies. Finally, an isolated algal colony was used to generate an axenic stock. *Ostreococcus* sp. RCC809, *O. tauri* (OTH 95), *Em. huxleyi*, and *M. pusilla* were treated with antibiotics. Cultures were grown for 1 week with penicillin (1 mg/ml), kanamycin (25 mg/ml), and neomycin (20 mg/ml). An additional subculture without antibiotics for approximately 6 days was then grown, before inoculating the experimental vitamin B₁₂ growth assays. For *Em. huxleyi*, before the antibiotic treatment, cells were filtered on a 1 micron filter (to attempt to remove external bacteria) and then decalcified to reduce the possibility that bacteria may have been present underneath the coccoliths (calcifying form only). This was achieved by addition of sterile hydrochloric acid to pH 4, followed by restoration to pH 8.5 by the addition of sterile sodium hydroxide. Cultures were stained with the nucleic acid-specific stain 4',6-diamidino-2-phenylindole (DAPI) (1 ng/ml, 5 min at 20 °C), placed in a microscopy dish and viewed under epifluorescent illumination (excitation 330–380 nm, emission above 420 nm) using an Eclipse E1000 microscope (Nikon, Tokyo, Japan). Bacteria were clearly visible in nontreated control cultures but were not seen in the antibiotic treated cultures used in this study.

Assessment of Vitamin B₁₂ Requirements of Algal Species

The vitamin B₁₂ requirements of the algal species (treated as above) were assessed by growing each alga in the appropriate medium, plus or minus B₁₂. The concentration of B₁₂ used in the plus condition was as described in the media appropriate to each alga (see [supplementary table 1, Supplementary Material](#) online). Aliquots (1 ml) of culture were then transferred to fresh media within 5–21 days depending on the speed of growth, which varies between species. Up to 5 subcultures were carried out, or until the algae had died in the experimental condition (minus B₁₂) and not in the control (plus B₁₂) as described in [Croft et al. 2005](#). Three biological replicates were carried out for each condition. If the cultures were still viable after B₁₂ starvation, they were visually checked with DAPI and/or plated onto LB plates (for freshwater species) or high salt LB (for marine species).

B₁₂ Bioassay

B₁₂ bioassays on cell lysates of *Em. huxleyi* were carried out as described by [Raux et al. \(1996\)](#), using *S. enterica* (AR3621). For preparation of the algal cell lysates, cells were washed 3 times and resuspended in 0.9% NaCl. The cells were then subjected to three freeze–thaw cycles and placed at 100 °C for 15 min before cooling to room temperature.

Molecular Methods

DNA was extracted from algal cells grown in liquid culture using the phenol–chloroform method described in [Newman et al. 1990](#). Total RNA was extracted as described by [Witman et al. \(1972\)](#). RNA quality was assessed using a Nanodrop

Spectrophotometer (ND-1000). A 5 µg sample of RNA was treated with RNase-free DNase (Promega) for 1 h at 37 °C to remove contaminating DNA. Reverse transcription (RT) was performed using Superscript II transcriptase (Invitrogen) with oligo dT primers according to the manufacturer's instructions. Polymerase chain reaction (PCR) was used to amplify sequences from genomic DNA and cDNA using Taq DNA polymerase (Bioline) (see [supplementary table 3, Supplementary Material](#) online, for details of primers). For the semiquantitative RT-PCR analysis in *P. tricornutum*, PCR amplification was carried out for 26–38 cycles, to determine the optimal cycle number for each primer pair and allow comparisons between samples ([supplementary fig. 1, Supplementary Material](#) online). The following cycle numbers for each primer set was used throughout: *METE* (35 cycles), *METH* (33 cycles), and *HISTONE H4* (33 cycles).

Phylogenetic Analysis of Methionine Synthases

Methionine synthase sequences were selected from both prokaryotes and other nonalgal eukaryotes. In order to ensure that the data set included appropriate prokaryote sequences, BlastP searches of the GenBank nonredundant database using methionine synthase genes identified in algal genomes were used to select the most similar sequences in prokaryotes. An alignment of *METE* or *METH* was constructed using MUSCLE ([Edgar 2004](#)) and manually corrected using BioEdit ([Hall 1999](#)) where appropriate, to ensure only unambiguous residues were compared. Maximum likelihood phylogenetic analysis was performed using PhyML within the Bosque software package ([Ramirez-Flandes and Ulloa 2008](#)), based on the Whelan and Goldman substitution matrix ([Whelan and Goldman 2001](#)). One hundred bootstrap replicates were run for each analysis.

Results

Survey of B₁₂-Dependent Enzymes in Sequenced Algal Genomes

We examined 15 algal genomes, including one or more representatives from the Rhodophyta (red algae), Chlorophyta (green algae), diatoms, Haptophyta, and brown macroalgae, for the presence of enzymes that require vitamin B₁₂. We also searched for accessory proteins involved in supplying the cofactor to these B₁₂-dependent enzymes. MCM utilizes the adenosylated form of cobalamin (AdoCbl) and requires two accessory proteins, CBLA for vitamin B₁₂ transport into the mitochondria and CBLB for AdoCbl synthesis ([Dobson, Wai, Leclerc, Kadir, et al. 2002](#); [Dobson, Wai, Leclerc, Wilson, et al. 2002](#)) (see [fig. 1](#)). Conversely, *METH* uses methylcobalamin (MeCbl) and requires the molecular chaperone methionine synthase reductase (MTRR) to regenerate the methylated cofactor ([Dobson, Wai, Leclerc, Kadir, et al. 2002](#); [Yamada et al. 2006](#)). Blast searches were performed with the protein sequences as queries (as described in Materials and Methods) using default parameters. All hits were verified by multiple sequence alignment and Pfam analyses, (see [supplementary table 2, Supplementary Material](#) online, for accession number details).




Fig. 1. Schematic diagram illustrating the intracellular metabolism of B_{12} in eukaryotes. The figure shows the B_{12} -dependent enzymes methionine synthase (*METH*) and methylmalonyl-CoA mutase (*MCM*), and their accessory proteins (in dashed boxes). *METE* catalyzes the same reaction as *METH* but does not require B_{12} as a cofactor. Both use methyltetrahydrofolate as the methyl donor, but for *METH*, the methyl group is transferred to the substrate via cobalamin. AdoCbl, adenosylcobalamin; MeCbl, methylcobalamin; Cbl I, Cbl II refers the oxidation state of the cobalt ion contained within vitamin B_{12} .

We found that all species have the B_{12} -dependent methionine synthase, *METH*, except *Coccomyxa* sp. C-169 and *Cy. merolae*, (table 1). Although partial sequences were found for *A. anophagefferens* and *F. cylindrus*, these appear to be the result of gaps in the genome assembly. All genomes contained the gene encoding *MTRR*, the accessory protein for *METH*, even those that did not encode this B_{12} -dependent isoform of methionine synthase. In contrast, the gene encoding the vitamin B_{12} -independent isoform, *METE*, was present in only seven of the genomes and absent altogether from another seven. For *V. carteri*, a gene was identified that had sequence similarity to *METE* of *C. reinhardtii*. However, although the genome assembly was complete in this region, when the predicted amino acid sequence was obtained using GeneWise 2.0, there were numerous frameshifts and regions of missing sequence, implying that this was a pseudogene. This is considered in more detail below.

MCM was found in the genomes of all species representing the Chromalveolata, but only in one of the species with simple plastids, the green alga *Ch. variabilis* (NC64A). However, the distribution of the accessory proteins was

not identical. *A. anophagefferens* does not appear to encode *CBLA*, indicating that *MCM* would not function in this species, and *CBLB* was found in several of the green algae that do not encode *MCM*, suggesting that this gene was retained after the loss of the associated B_{12} -dependent enzyme. The third B_{12} -dependent enzyme RNR II, previously reported in *Eu. gracilis* (Hamilton 1974), was present only in *Ec. siliculosus*. This alga also encodes both subunits of B_{12} -independent RNR I (Cock et al. 2010).

Assessment of Algal Requirements for Vitamin B_{12}

To interpret this genomic data set, the B_{12} requirement of each alga must be established. For eight species, this information was already available in existing literature. For the other algal species, we performed standard vitamin B_{12} growth assays to examine growth with or without this cofactor. In order to avoid false positive results, the assays were designed to limit all potential sources of B_{12} . Artificial seawater was used where possible (see supplementary table 1, Supplementary Material online) to exclude a potential source of B_{12} from natural seawater, and all cultures used were either obtained axenic or treated with an antibiotic cocktail prior to the growth assay. Cultures were visually inspected using DAPI-staining and epifluorescence microscopy to look for the presence of bacteria.

After 2–4 subcultures, it became clear that *V. carteri*, *Ostreococcus* sp. RCC809, *O. tauri*, and *M. pusilla* were B_{12} dependent (fig. 2A–D). The dramatic decrease in growth within two subcultures suggests our precautions to avoid contaminating sources of B_{12} were sufficient and that false positives were unlikely. In contrast, for the Arctic diatom *F. cylindrus* and the haptophyte *Em. huxleyi*, no difference was seen in growth with or without B_{12} .

Methionine Synthase Isoform Determines B_{12} Dependence

An analysis of the B_{12} -dependent enzymes in the six algal species that require B_{12} for growth (*V. carteri*, *Ostreococcus* sp. RCC809, *O. tauri*, *M. pusilla*, *T. pseudonana*, and *A. anophagefferens*) indicates that *METH* is present in all species, whereas only two species possess *MCM*. Moreover, *METE* is absent in each of these B_{12} -dependent algal species, indicating that *METH* would be the sole source of methionine synthase activity. Conversely, with the exception of *Em. huxleyi*, *METE* is universally present in the B_{12} -independent algae, indicating that *METE* enables the organism to live in the absence of the cofactor. Although *Ec. siliculosus* contains type II RNR, it also encodes B_{12} -independent RNR I, explaining the observation that the alga itself does not require B_{12} for growth (Boalch 1961). From these observations, we conclude that the isoform of methionine synthase is the key factor determining the nutritional requirement for B_{12} in algae. However, two of our surveyed species warrant closer scrutiny. Firstly, it must be determined whether the partial *METE* sequence in *V. carteri* is likely to encode an active gene product, and secondly we need to understand how *Em. huxleyi* is able to grow in the absence of both *METE* and exogenous B_{12} .

Table 1. Analysis of 15 Sequenced Algal Genomes for Genes Encoding the Three B₁₂-Dependent Proteins (and their accessory proteins) Known to Occur in Eukaryotes.

Group	Species	Genes Encoding B ₁₂ -Dependent Enzymes				Genes Encoding Accessory Proteins			B ₁₂ Dependent? ⁴	
		<i>METE</i>	<i>METH</i>	<i>RNR II</i>	<i>MCM</i>	<i>MTRR</i>	<i>CBLA</i>	<i>CBLB</i>		
Chlorophyta	<i>Chlamydomonas reinhardtii</i> [2]	✓ ¹	✓	×	×	✓	×	✓	×	[3,4]
	<i>Volvox carteri f. nagariensis</i> EVE [5]	Pseudo ²	✓	×	×	✓	×	✓	✓	[1]
	<i>Micromonas pusilla</i> (CCMP 1545) [6]	×	✓	×	×	✓	×	✓	✓	[1]
	<i>Ostreococcustauri</i> (OTH95)	×	✓	×	×	✓	×	×	✓	[1]
	<i>Ostreococcus lucimarinus</i> [7]	×	✓	×	×	✓	×	✓	?	
	<i>Ostreococcus</i> sp. RCC809 [7]	×	✓	×	×	✓	×	✓	✓	[1]
	<i>Coccomyxa</i> sp. C-169	✓	×	×	×	✓	×	×	×	[8]
	<i>Chlorella</i> NC64A	✓	✓	×	✓	✓	✓	✓	?	
	Chromalveolata	<i>Emiliania huxleyi</i> (CCMP1516)	×	✓	×	✓	✓	✓	✓	×
<i>Phaeodactylum tricornutum</i> (CCAP1055/1) [11]		✓	✓	×	✓	✓	✓	✓	×	[12]
<i>Fragilariopsis cylindrus</i> (CCMP 1102) [13]		✓	Partial ³	×	✓	✓	✓	✓	×	[1]
<i>Thalassiosira pseudonana</i> [14]		×	✓	×	✓	✓	✓	✓	✓	[15]
<i>Ectocarpus siliculosus</i> [16]		✓	✓	✓	✓	✓	✓	✓	×	[17]
<i>Aureococcus anophagefferens</i>		×	Partial	×	✓	✓	×	✓	✓	[18]
<i>Cyanidioschyzon merolae</i> [19]		✓	×	×	×	✓	×	×	×	[15]

^a A tick indicates that a complete gene was present; an × indicates no hit obtained.

^b An incomplete sequence for *METE* in *V. carteri* was verified by resequencing and was subsequently shown to be a pseudogene (see text for details).

^c The *METH* results for *F. cylindrus* and *A. anophagefferens* were not complete but were in regions with gaps in the genome assembly and so are indicated as “partial.”

^d The B₁₂ requirements for each of the algal species were either established in this study [1] or taken from the literature: [2] Merchant et al. (2007); [3] Levin (1958); [4] Provasoli (1958); [5] Prochnik et al. (2010); [6] Worden et al. (2009); [7] Palenik et al. (2007); [8] Shihira (1965); [9] Guillard (1963); [10] Pintner and Provasoli (1968); [11] Bowler et al. (2008); [12] Hutner (1948); [13] Mock T, Parker M, Armbrust V, Valentin K, Bowler C, unpublished data; [14] Armbrust et al. (2004); [15] Croft et al. (2005); [16] Cock et al. (2010); [17] Boalch (1961); [18] Mahoney (2005); and [19] Matsuzaki et al. (2004).

Characterization of *METE* Genes in *V. carteri* and Related Algae

As explained above, sequence similarity searches identified a putative *METE* sequence in *V. carteri*, but it appeared to be a pseudogene. Closer inspection showed that elements of the N-terminus are missing, there is an in-frame stop codon (white asterisk, [fig. 3A](#)) and multiple deletions leading to seven frameshifts (dashed white lines, [fig. 3A](#)). The C-terminus of *METE* contains conserved Zn²⁺-binding residues required for catalytic activity (Gonzalez et al. 1992; Pejchal and Ludwig 2005). These residues, equivalent to His660, Cys662, Glu684, and Cys749 in *C. reinhardtii*, are conserved in all of the algal *METE* sequences we have obtained (black dots, [fig. 3B](#)), with the exception of that from *V. carteri*, in which the two cysteines are replaced by arginine and proline, respectively. We resequenced this region of the *V. carteri* genome and verified that the assembled genomic sequence information was correct. In combination, these features indicate that *METE* in *V. carteri* is no longer functional.

In order to determine whether the *V. carteri* *METE* gene is expressed, we performed an RT-PCR analysis (See [supplementary table 3, Supplementary Material](#) online, for primer sequences). As *C. reinhardtii* *METE* is repressed in the presence of vitamin B₁₂, we depleted *V. carteri* cells of B₁₂, taking samples from each subsequent subculture, up to the third subculture in which the cells failed to grow as a result of B₁₂ depletion. Two sets of primers were designed, targeted to predicted exons within conserved regions of the *METE* gene. *METH* was found to be expressed in both B₁₂-replete and B₁₂-depleted treatments ([fig. 3C](#)), as is the case in *C. reinhardtii* (Croft et al. 2005). However, despite being able to amplify a *METE* product from genomic DNA, we could not do the same with either primer set using

cDNA as the template, indicating that the gene is not expressed.

V. carteri and *C. reinhardtii* are members of the chlorophycean order of Volvocales (Herron et al. 2009). Examination of the B₁₂ requirements within this order ([fig. 4A](#)) indicates that B₁₂ dependence may have arisen on multiple occasions even within this lineage. We therefore investigated another B₁₂-dependent member of the Volvocales, *G. pectorale*, for further evidence of recent *METE* gene loss. We confirmed *G. pectorale* is B₁₂ dependent using the B₁₂ growth assay ([fig. 4B](#)). Using degenerate primers targeted against a conserved region of the *C. reinhardtii* and *V. carteri* *METE* sequences, we were able to amplify a *METE* transcript from *G. pectorale* cDNA of the correct size (237 bp) ([fig. 4C](#)), although only in cells grown in the absence of vitamin B₁₂; like *C. reinhardtii*, the gene appears to be repressed by cobalamin. The larger band is likely to be genomic DNA contamination. The sequence of the 237 bp band exhibited high similarity to that of *C. reinhardtii* ([supplementary fig. 2, Supplementary Material](#) online). However, closer examination of these sequences revealed a significant deletion in *G. pectorale* *METE* that results in a frameshift and a premature stop codon, so that, like *V. carteri*, *G. pectorale* *METE* would not be able to produce a functional enzyme.

Taken together, these results indicate that in both *V. carteri* and *G. pectorale* *METE* has become a pseudogene, resulting in B₁₂ dependence in these algae. Given that *C. reinhardtii* diverged from *V. carteri* and *G. pectorale* approximately 250 Ma (Herron et al. 2009), this suggests a recent gene loss and uniquely captures the evolution of B₁₂ auxotrophy in action. As several other members of the Volvocales are B₁₂ independent, we suggest that the two pseudogenes in *V. carteri* and *G. pectorale* represent independent gene loss events.




Fig. 2. Assessment of B₁₂ requirements of selected algae. Species were grown in the appropriate liquid medium (supplementary table 1, Supplementary Material online) with and without B₁₂ in batch culture over several subcultures, or until the cells had died. (A) *Volvox carteri*, (B) *Ostreococcus* sp. RCC809, (C) *O. tauri*, (D) *Micromonas pusilla*, (E) *Fragilariopsis cylindrus*, and (F) *Emiliana huxleyi*. filled bars: +B₁₂, White: no B₁₂. The following measurements were used to quantify growth: spheroid number (A), optical density (B, C, and D), and cell count (E and F). Three replicate cultures were used for each treatment. Error bars denote standard errors.

The *V. carteri* *METE* gene represents a unitary pseudogene as there is no functional duplicate gene, and thus, there is an accompanying phenotype. This is presumably also the case in *G. pectorale* because it too is B₁₂ dependent. Another notable example of a unitary pseudogene resulting in vitamin auxotrophy is that of L-gulonolactone oxidase involved in the biosynthesis of vitamin C in animals (Nishikimi et al. 1992, 1994). As a consequence of losing this gene, guinea pigs and primates are no longer able to synthesize the vitamin themselves and instead must acquire it in their diet.

Reassessing the B₁₂ Requirements of *Em. huxleyi*

The haptophyte *Em. huxleyi* does not contain *METE* within its genome, but it can survive in the absence of exogenous B₁₂ (fig. 2F), confirming earlier reports (Guillard 1963; Carlucci and Bowes 1970; Haines and Guillard 1974). Assuming that methionine synthase activity is essential for growth in *Em.*

huxleyi, either *METE* is present in the *Em. huxleyi* genome but absent from the current genome assembly or the growth assays did not reveal the true B₁₂ requirements of *Em. huxleyi*.

The first suggestion is unlikely. *METE* is not present in either the *Em. huxleyi* (CCMP 1516) genome assembly, the unassembled genomic reads or in the *Em. huxleyi* expressed sequence tag collections (Von Dassow et al. 2009). Additionally, using broad specificity degenerate primers (Huang et al. 2005), we were unable to amplify *METE* from genomic DNA of *Em. huxleyi* (supplementary fig. 3, Supplementary Material online). Although this is not conclusive evidence for the absence of the gene, our combined data do not support the presence of *METE* in the *Em. huxleyi* genome.

We therefore reassessed the B₁₂ requirements of *Em. huxleyi*. We treated cultures of both calcifying and noncalcifying *Em. huxleyi* (CCMP 1516) using extended methods to minimize bacterial contamination (see Materials and Methods).




FIG. 3. *Volvox carteri* contains a unitary *METE* pseudogene. (A) Schematic diagram of the gene model for *Chlamydomonas reinhardtii* (*C. r*) *METE* (upper) compared with that of *V. carteri* (*V. c*; lower) indicating that *V. carteri* gene is truncated with multiple frameshifts and an in-frame stop codon (boxes: exons, arrows + dashed white line: frameshift, *in-frame stop codon). (B) Multiple sequence alignment (ClustalW) of *METE* protein sequences from eight algae for the region spanning the functionally important conserved residues required for binding Zn²⁺ (indicated by a black dot). The two cysteines are absolutely conserved except in *V. carteri*. (C) PCR amplification of *METE*, *METH*, and actin (*ACTA*) in *V. carteri*, using two different subcultures (Sub 1 & 2) grown with and without B₁₂. RNA was reversed transcribed and then used with two sets of primers (F1/R1 and F2/R2) targeted to different conserved regions of *METE* and avoiding intron/exon boundaries in *C. reinhardtii*. However, no *METE* transcripts were detected, although a product of the expected size was amplified from genomic DNA (right-hand boxes), and the *METH* and *ACTA* transcripts were readily detected.

Both strains were able to grow in the absence of B₁₂ for up to five subcultures (supplementary fig. 4, Supplementary Material online). However, using a bioassay that measures B₁₂, we were able to detect up to 11.7 ng/ml of B₁₂ in lysed cell extracts of the *Em. huxleyi* cells grown for four subcultures in medium without B₁₂. In contrast, Croft et al. (2005) showed that cultures of several other B₁₂-independent algae: *Haematococcus pluvialis*, *Cyanidium caldarium*, *Tetraselmis verrucosa*, and *C. reinhardtii* have no detectable B₁₂ after just two subcultures. The source of B₁₂ in *Em. huxleyi* is unlikely to be de novo synthesis because only two of the total of 20 enzymes required for aerobic B₁₂ biosynthesis (Warren et al. 2002) were identified in the *Em. huxleyi* genome. These genes exhibited weak similarity to *cobB* and *cobN*.

DAPI-staining of the *Em. huxleyi* cultures did not reveal any associated bacteria. However, when the cultures were plated on high salt LB media, growth of bacteria was observed. Analysis of the 16S ribosomal RNA gene established that the contaminant for the noncalcifying strain was a member of the *Oceanicaulis* genus, whereas for the calcifying strain, it was of the *Aurantimonas* genus, within the order Rhizobiales, whose members are known to synthesize cobalamin. As the vitamin was not depleted in *Em. huxleyi*,

this result supports the hypothesis that algal species lacking *METE* require B₁₂ for growth. The apparent very close association of these bacteria with *Em. huxleyi*, which could not be disrupted by antibiotic treatment, suggests an intimate relationship that warrants further attention

Phylogenetic Analysis of *METE* and *METH*

To examine the evolutionary relationships between the requirement for B₁₂ and the presence of methionine synthase isoforms, we generated a schematic phylogenetic tree (fig. 5) (Keeling and Palmer 2008), with representatives of the major eukaryote lineages, including land plants, mammals, and fungi. It is clear that the B₁₂ requirement correlates with the absence of a functional *METE*.

A more detailed phylogenetic analysis of *METE* and *METH* sequences from a broad sample of organisms was then performed to enable a distinction to be made between whether multiple independent losses of *METE* resulted in B₁₂ auxotrophy, or conversely that multiple independent gains of *METE* may have led to the evolution of B₁₂ independence. Aside from the major endosymbiotic gene transfer events, abundant horizontal gene transfers (HGTs) have been proposed to encode novel metabolic capacities in some algae, for example in the




Fig. 4. B₁₂ requirements in the Volvocales (A) Schematic tree displaying phylogenetic relationships between Volvocalean species (based on Herron et al. 2009) for which B₁₂ requirements are already known: *Chlamydomonas reinhardtii* (Lewin 1958), *Gonium pectorale* (Stein 1966), *G. multicoccum* (Saito 1972), *Volvox globata*, *Eudorina elegans* (Dusi 1940), *V. tertius* (Pintner and Provasoli 1959), *Pandorina morum* (Palmer and Starr 1971), and *V. aureus* (Pringsheim 1970), or for *V. carteri* determined in this study. (B) Confirmation of B₁₂ dependence of *G. pectorale* with a B₁₂ growth assay. (C) RT-PCR analysis of *METE* transcripts in *G. pectorale* grown with and without B₁₂. A band of the correct size (237 bp) is obtained in the latter case and sequencing confirmed it to be from *METE*.

diatom *P. tricornutum* (Bowler et al. 2008). Our results indicate that the phylogeny of eukaryote *METH* is well resolved, with strong support for the monophyly of *METH* in the chlorophyte, chromalveolate, and opisthokont (animal and fungal) lineages (fig. 6A). This argues against multiple HGT events in these lineages and supports the view that the absence of *METH* in the *Cy. merolae* and *Coccomyxa* sp. C-169 (table 1) and in land plants reflects independent gene loss.

In contrast, phylogenetic analysis of *METE* sequences indicates a more complex evolutionary history. The majority of algal *METE* sequences forms a well-supported monophyletic clade with cyanobacterial homologues (fig. 6B), suggesting that it was probably acquired during the primary cyanobacterial endosymbiosis, as proposed before for *C. reinhardtii*




Fig. 5. Schematic tree showing phylogenetic relationships between organisms and their B₁₂ requirements. The phylogeny is based on Keeling and Palmer (2008). In addition to the algae species used in the current study for which complete genome sequences are known, two prokaryotes are shown (the cyanobacterium *Synechocystis* sp. strain PCC6803 and *E. coli*), as well as members of the land plants, animals, and fungi. The organisms fall into the following distinct phylogenetic groups indicated by shading from top to bottom: Prokaryota, Chromalveolata, Rhodophyta, Chlorophyta, Streptophyta, and the Opisthokonta (in dashed box). Ticks indicate that a species can grow independently without an exogenous source of B₁₂, whereas an asterisk (*) is shown for *Synechocystis* sp. strain PCC6803. This can synthesize cobalamin (Raux et al. 1998) and therefore does not require an external source of the vitamin, even though it lacks *METE*. Black squares indicate the presence of a gene, and partial black boxes indicate a partial hit. In eukaryotes, a strong correlation between the requirement for B₁₂ and the absence of a complete *METE* can be observed.

METE (Moustafa and Bhattacharya 2008; Maruyama et al. 2009). This observation argues against multiple HGT events in these lineages, and thus, the most likely explanation for the absence of *METE* in representatives of the Chlorophyta and Chromalveolata is as a result of multiple independent losses. This is in accordance with our discovery of *METE* pseudogenes in *V. carteri* and *G. pectorale*.

Surprisingly, *METE* sequences from land plants and the red alga *Cy. merolae* do not fall into this algal clade, although *METE* from another rhodophyte, *Galdieria sulphuraria*, clusters with the green algal sequences. A more recent HGT event must therefore be considered as a possible origin for *METE* in land plants and *Cy. merolae*. An alternative hypothesis is that *METE* may have originated from the eukaryote ancestor prior to the cyanobacterial primary endosymbiosis. Therefore, unlike the majority of algae that have retained the cyanobacterial copy of *METE*, the land plants may have retained the ancestral eukaryote gene. This implies that the ancestral members of the Archaeplastida (the eukaryotic supergroup including the Rhodophyta, Chlorophyta, and Glaucocystophyta, as well as the land plants, see Parfrey et al. (2006), for review) retained both forms of *METE* (ancestral eukaryotic and cyanobacterial), at least until the streptophytes diverged from the




Fig. 6. Phylogenetic analysis of methionine synthase isoforms. (A) Maximum likelihood consensus tree constructed for METH using PhyML. 100 bootstraps were run, values above 50% are shown. (B) Maximum likelihood consensus tree for METE, constructed as described above.

chlorophytes. Although we have identified no extant Archaeplastida species with both forms of the *METE* gene to support this hypothesis, coexistence followed by multiple gene loss has been proposed to explain the complex distribution of other genes in the green algal lineage, such as *EFL* and *EF-1 α* (Noble et al. 2007). It is worth noting that *Arabidopsis* has three different copies of *METE* (one plastidic and two cytosolic) (Ravanel et al. 2004), all of which belong to the land plant clade. Gene duplication and divergence may therefore have also played a role in the replacement of *METH* or the cyanobacterial *METE* in higher plants.

Effect of B₁₂ on Expression of *METE* in *P. tricornutum*

A comparison of *E. coli* methionine synthase isoforms found that METH has a much greater catalytic activity (ca. 100-fold) than METE (Gonzalez et al. 1992). This suggests that METH would be used preferentially in the presence of B₁₂, and indeed in *E. coli*, the *metE* gene is repressed by cobalamin in the medium. In agreement with this, *METE* expression is also strongly repressed by B₁₂ in *C. reinhardtii* (Croft et al. 2005), and we made a similar observation for *G. pectorale* (fig. 4C). Thus, a prolonged continuous supply of B₁₂ could

provide an opportunity for the accumulation of mutations in *METE*, which in time could lead to the loss of gene function. However, if this is to explain *METE* loss more broadly, it is important to determine whether B₁₂ represses *METE* expression in other algal species. Therefore, we assessed *METE* regulation in the chromalveolate diatom *P. tricornutum*, which like *C. reinhardtii* has both methionine synthase isoforms.

We used semiquantitative RT-PCR to analyze the expression of *METE* and *METH* from *P. tricornutum* cells grown in the presence or absence of B₁₂. *METH* transcript is detectable in both treatments, whereas *METE* is not expressed in the presence of vitamin B₁₂ (fig. 7A and supplementary fig. 1, Supplementary Material online), and the repression occurs within 14 h after the addition of the vitamin (fig. 7B). Experiments presented in this paper, alongside those conducted on *C. reinhardtii* (Croft et al. 2005), thus indicate that *METH* is used preferentially in organisms with both methionine synthase isoforms, providing a potential mechanism for *METE* gene loss.

Discussion

The advent of whole genome sequencing has provided an unparalleled opportunity for comparisons between different organisms, in particular to test evolutionary questions. However, it is essential that sequence analysis is coupled with experimental approaches to provide independent verification of proposed events. We have combined growth assays and expression data with bioinformatics searches to investigate the reason for B₁₂ auxotrophy within the algal kingdom and the possible mechanism by which it arose. Our results indicate strongly that the major determinant for the B₁₂ requirement of algae relates to the isoform(s) of methionine synthase they possess, and the available evidence supports multiple evolutionary losses of *METE* as a key factor in the distribution of B₁₂ auxotrophy amongst photosynthetic eukaryotes. In contrast, it is unlikely that B₁₂-dependent MCM causes the auxotrophy because it is found in B₁₂-independent algal species such as *P. tricornutum*, indicating that the cellular functions of MCM activity may not be vital and thus could be dispensed with in the absence of B₁₂. Similarly, the absence of MCM in species that have an absolute requirement for B₁₂ (e.g., *V. carteri*, *M. pusilla*) supports the inference that this enzyme is not a determining factor.

Nonetheless, if we are to gain insight into the factors leading to the loss of *METE*, a fuller understanding of the functional roles of methionine synthase isoforms in different lineages is required. Methionine synthase is responsible both for de novo synthesis of methionine, and the regeneration of methionine following transmethylation reactions. In land plants, at least, it is proposed that these functions occur in different subcellular compartments (the chloroplast and cytosol, respectively) (Ravanel et al. 2004). In certain organisms with both isoforms, *METE* and *METH* could conceivably perform different functions and/or exist in different subcellular compartments. For example, *METE* is also an abundant flagella protein in *C. reinhardtii* and has been implicated in protein methylation




FIG. 7. *METE* expression is repressed by B₁₂ in the diatom *Phaeodactylum tricornutum*. (A) RT-PCR analysis of transcripts for *METE* and *METH* following growth +B₁₂ or -B₁₂. (B) Twenty-four hour time-course of *METE* and *METH* expression following the addition of vitamin B₁₂ at time 0 h. H4, constitutively expressed histone H4 gene. The following cycle number for each primer set was used throughout: *METE* (35 cycles), *METH* (33 cycles), and *HISTONE H4* (33 cycles), see supplementary figure 4, Supplementary Material online.

during flagella resorption (Schneider et al. 2008; Sloboda and Howard 2009). Thus, in *C. reinhardtii*, there may be a selective pressure to maintain *METE* due to its specialized role in flagella function, although repression of *METE* by B₁₂ would suggest this is not an essential role or that *METH* can function in its place. This explanation presumably also applies to flagellated algae such as *M. pusilla* and *V. carteri*, which lack *METE*. Differences in substrate availability may further influence the selective pressure to retain methionine synthase isoforms because evidence from bacteria, plants, and yeast indicate that *METE* is unable to utilize the monoglutamate form of methyltetrahydrofolate as a methyl donor (Burton and Sakami 1969; Whitfield and Weissbach 1970; Ravanel et al. 2004).

Exogenous sources of methionine may be available in the ocean, as one of the most abundant marine prokaryotes, SAR11 clade *Pelagibacter* lacks methionine synthase (Tripp et al. 2008). However, in eukaryotic algae, an exogenous source of methionine would be insufficient to compensate for a lack of methionine synthase activity. In all cells, the most important metabolic function for this enzyme is its role as a component of the methylation cycle, and when methionine synthase activity is limiting a phenomenon known as folate trapping is observed (Scott 1999). This is clearly demonstrated by the fact that methionine is insufficient to support the growth of *Lobomonas rostrata*, a B₁₂-dependent green alga, in the absence of B₁₂. It is necessary to supplement with folate in addition (Croft et al. 2005). Given these cellular considerations, we consider that the most

important environmental factor promoting the selective loss of *METE* in certain algal lineages is a reliable and readily available source of B₁₂ in the environment, which is produced exclusively by prokaryotes (Warren et al. 2002). Croft et al. (2005) proposed that algae may obtain B₁₂ via a symbiotic relationship with bacteria, although a direct symbiosis involving B₁₂ has not yet been demonstrated in natural populations. However, there are many documented examples of close associations between algae and bacteria (Liu et al. 2008; Amin et al. 2009) and, regardless of the initial reason for their formation, these interactions may result in an abundant supply of B₁₂ for the algal partners. Of particular interest in this regard are the reports of bacterial symbionts associated with *Volvox*, including possible endosymbionts (Hamburger 1958) as well as the tightly associated bacteria we found associated with *Em huxleyi*. Moreover, a recent study showed that vitamin auxotrophy for B₁₂ (and indeed thiamine and biotin) in harmful algal bloom species (HABs) such as dinoflagellates was much more prevalent than that for non-HAB species, (95% for B₁₂, 74% for thiamine, and 37% for biotin) (Zhong Tang et al. 2010). Many HABs are phagotrophic (e.g., Jeong et al. 2005) and therefore probably obtain these micronutrients heterotrophically from their prey.

It is also worth considering the consequences of standard algal stock maintenance practices in the laboratory in which cultures are grown with a recommended 1–10 nM exogenous vitamin B₁₂ (amongst other vitamins) regardless of whether they require it or not. Recent measurements indicate the lower range of vitamin B₁₂ concentrations in the Southern Ocean and North Atlantic Ocean approach potentially limiting values, with the lowest values being 0.4 and 0.2 pM, respectively (Panzeca et al. 2009), whereas in our hands, a minimum of 10 pM is required to support algal growth in batch culture. However, the requirement for a direct symbiosis between bacteria and algae for the delivery of B₁₂ has been disputed. Chemostat experiments indicate that the growth saturation constants for B₁₂ in phytoplankton reside in the subpicomolar range, suggesting that the concentration of B₁₂ released through bacterial cell lysis is sufficient to support algal growth, even in open ocean environments (Droop 2007). Nevertheless, there are several reports of B₁₂ stimulation of natural phytoplankton assemblages suggesting this cofactor may be limiting in certain areas (Sanudo-Wilhelmy et al. 2006; Bertrand et al. 2007). Although it remains to be determined whether B₁₂ limits phytoplankton productivity in a wider context, one would certainly expect selective pressure to retain *METE* to be higher in environments such as the open ocean where B₁₂ supply may be low or unreliable. In this regard, a metagenomic study of marine prokaryote communities indicated that methionine and cobalamin-dependent pathways varied significantly with environmental features (Gianoulis et al. 2009). An important factor limiting the ability of bacteria to synthesize cobalamin is the concentration of cobalt (Panzeca et al. 2008), so in turn this may have a significant influence on the distribution of B₁₂-auxotrophy in oligotrophic environments. Given that algae alone are responsible for the fixation of 50% of the world's carbon dioxide (Field

et al. 1998) and have a vital role in biogeochemical cycling in both marine and freshwater systems, it will be important to understand the extent to which B₁₂ availability influences both the physiology and the ecology of these organisms.

Supplementary Material

Supplementary tables 1–3 and figures 1–4 are available at *Molecular Biology and Evolution* online (<http://www.mbe.oxfordjournals.org/>).

Acknowledgments

We would like to thank Herve Moreau (Oceanological Observatory of Banyuls-ser-Mer, France) for donation of the strains *O. tauri* (OTH95), *O. lucimarinus*, and *M. pusilla*. We are grateful to Thomas Mock and Jan Strauss (University of East Anglia, UK) for providing us with the culture of *F. cylindrus* and for their helpful advice. Thanks also go to the Joint Genome Institute, and all those contributing to each genome project referred to in this paper. Specifically to the coordinators for specific projects who made data available before publication: Thomas Mock (*F. cylindrus*), Betsy Read (California State University, USA) (*Em. huxleyi*), James L. Van Etten (Nebraska Center for Virology, University of Nebraska-Lincoln, USA) (*Coccomyxa* sp. C-169 and *Ch. variabilis* NC64A), and Christopher Gobler (School of Marine and Atmospheric Sciences, Stony Brook University, NY, USA) (*A. anophagefferens*). We would also like to thank Karen Weynberg (Plymouth Marine Laboratory, UK) for helping us to grow the *Ostreococcus* strains. Finally, our appreciation goes to Roger Sloboda (Dartmouth College, Hanover, USA) for his helpful insight and discussions. We acknowledge funding from the Biotechnology and Biological Sciences Research Council (BBSRC) of the United Kingdom with grant reference BB/F021844/1, and a studentship for KEH.

References

- Altschul S, Gish W, Miller W, Myers E, Lipman D. 1990. Basic local alignment search tool. *J Mol Biol.* 215:403–410.
- Amin SA, Green DH, Hart MC, Kupper FC, Sunda WG, Carrano CJ. 2009. Photolysis of iron-siderophore chelates promotes bacterial-algal mutualism. *Proc Natl Acad Sci U S A.* 106:17071–17076.
- Armbrust EV, Berges JA, Bowler C, et al. (46 co-authors). 2004. The genome of the diatom *Thalassiosira pseudonana*: ecology, evolution, and metabolism. *Science.* 306:79–86.
- Bertrand EM, Saito MA, Rose JM, Riesselman CR, Lohan MC, Noble AE, Lee PA, DiTullio GR. 2007. Vitamin B₁₂ and iron co-limitation of phytoplankton growth in the Ross Sea. *Limnol Oceanogr.* 52:1079–1093.
- Birney E, Clamp M, Durbin R. 2004. Genewise and genomewise. *Genome Res.* 14:988–995.
- Blanc G, Duncan G, Agarkova I, et al. (15 co-authors). 2010. The *Chlorella variabilis* NC64A genome reveals adaptation to photosymbiosis, coevolution with viruses, and cryptic sex. *Plant Cell.* 22:2943–2955.
- Boalch GT. 1961. Studies on *Ectocarpus* in culture. 2. Growth and nutrition of a bacteria-free culture. *J Mar Biol Assoc U K.* 41:287.
- Bowler C, Allen AE, Badger JH, et al. (77 co-authors). 2008. The *Phaeodactylum* genome reveals the evolutionary history of diatom genomes. *Nature* 456:239–244.

- Burton EG, Sakami W. 1969. The formation of methionine from the mono-glutamate form of methyltetrahydrofolate by higher plants. *Biochem Biophys Res Commun.* 36:228–234.
- Carrell EF, Seeger JW Jr. 1980. Ribonucleotide reductase activity in vitamin B₁₂ deficient *Euglena gracilis*. *Biochem J.* 188:573–576.
- Carlucci AF, Bowes PM. 1970. Production of vitamin B₁₂ thiamine and biotin by phytoplankton. *J Phycol.* 6:351–357.
- Cock JM, Sterck L, Rouze P, et al. (77 co-authors). 2010. The *Ectocarpus* genome and the independent evolution of multicellularity in brown algae. *Nature* 465:617–621.
- Croft MT, Lawrence AD, Raux-Deery E, Warren MJ, Smith AG. 2005. Algae acquire vitamin B₁₂ through a symbiotic relationship with bacteria. *Nature* 438:90–93.
- Croft MT, Warren MJ, Smith AG. 2006. Algae need their vitamins. *Eukaryot Cell.* 5:1175–1183.
- Delwiche CF. 1999. Tracing the thread of plastid diversity through the tapestry of life. *Am Nat.* 154:S164–S177.
- Dobson CM, Wai T, Leclerc D, Kadir H, Narang M, Lerner-Ellis JP, Hudson TJ, Rosenblatt DS, Gravel RA. 2002. Identification of the gene responsible for the cblB complementation group of vitamin B₁₂-dependent methylmalonic aciduria. *Hum Mol Genet.* 11:3361–3369.
- Dobson CM, Wai T, Leclerc D, Wilson A, Wu XC, Dore C, Hudson T, Rosenblatt DS, Gravel RA. 2002. Identification of the gene responsible for the cblA complementation group of vitamin B₁₂-responsive methylmalonic acidemia based on analysis of prokaryotic gene arrangements. *Proc Natl Acad Sci U S A.* 99:15554–15559.
- Droop MR. 2007. Vitamins, phytoplankton and bacteria: symbiosis or scavenging? *J Plankton Res.* 29:107–113.
- Dusi H. 1940. Culture of bacteriologiquement pure et nutrition autotrophe d'*Eudorina elegans*. Role du fer pour la formation des colonies. *Ann Inst Pasteur.* 64:340–343.
- Edgar RC. 2004. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics.* 5:1–19.
- Field CB, Behrenfeld MJ, Randerson JT, Falkowski P. 1998. Primary production of the biosphere: integrating terrestrial and oceanic components. *Science* 281:237–240.
- Finn RD, Tate J, Mistry J, et al. (11 co-authors). 2008. The pfam protein families database. *Nucleic Acids Res.* 36:D281–D288.
- Gianoulis TA, Raes J, Patel PV, et al. (12 co-authors). 2009. Quantifying environmental adaptation of metabolic pathways in metagenomics. *Proc Natl Acad Sci U S A.* 106:1374–1379.
- Giovannoni SJ, Tripp HJ, Givan S, et al. (11 co-authors). 2005. Genome streamlining in a cosmopolitan oceanic bacterium. *Science* 309:1242.
- Gonzalez JC, Banerjee RV, Huang S, Sumner JS, Matthews RG. 1992. Comparison of cobalamin-independent and cobalamin-dependent methionine synthases from *Escherichia coli*—2 solutions to the same chemical problem. *Biochemistry* 31:6045–6056.
- Gould SB, Waller RR, McFadden GI. 2008. Plastid evolution. *Annu Rev Plant Biol.* 59:491–517.
- Guillard RRL. 1963. Organic sources of nitrogen for marine centric diatoms. *Symp Mar Microbiol.* 93–104.
- Hackett JD, Su Yoon H, Li S, Reyes-Prieto A, Rummele E, Bhattacharya D. 2007. Phylogenomic analysis supports the monophyly of cryptophytes and haptophytes and the association of rhizaria with chromalveolates. *Mol Biol Evol.* 24(8):1702–1713.
- Haines KC, Guillard RRL. 1974. Growth of vitamin B₁₂-requiring marine diatoms in mixed laboratory cultures with vitamin B₁₂ producing marine bacteria. *J Phycol.* 10:245–252.
- Hall TA. 1999. Bioedit: a user-friendly biological sequence alignment editor and analysis program for windows. *Nucleic Acids Symp Ser.* 41:95–98.
- Hamburger B. 1958. Bakteriensymbiose bei *Volvox aureus* ehrenberg. *Arch Mikrobiol.* 29:291–310.
- Hamilton FD. 1974. Ribonucleotide reductase from *Euglena gracilis*. a 5 deoxyadenosylcobalamin-dependent enzyme. *J Biol Chem.* 249:4428–4434.
- Herron MD, Hackett JD, Aylward FO, Michod RE. 2009. Triassic origin and early radiation of multicellular volvocine algae. *Proc Natl Acad Sci U S A.* 106:3254–3258.
- Huang L, Li DY, Wang SX, Zhang SM, Chen JH, Wu XF. 2005. Cloning and identification of methionine synthase gene from *Pichia pastoris*. *Acta Biochim Biophys Sin.* 37:371–378.
- Hutner SH. 1948. Essentiality of constituents of sea water for growth of a marine diatom. *Trans N Y Acad Sci.* 10:136–141.
- Jeong HJ, Yoo YD, Park JY, Song JY, Kim ST, Lee SH, Kim KY, Yih WH. 2005. Feeding by phototrophic red-tide dinoflagellates: five species newly revealed and six species previously known to be mixotrophic. *Aquat Microb Ecol.* 40:133–150.
- Keeling PJ, Palmer DP. 2008. Horizontal gene transfer in eukaryotic evolution. *Nat Rev Genet.* 9:605–618.
- Lewin JC. 1958. Vitamin-bezonog de algoj. *Sciencaj studoj. Copenhagen (Denmark): P. Neergaard.* p. 187–192.
- Liu JQ, Lewitus AJ, Brown P, Wilde SB. 2008. Growth promoting effects of a bacterium on raphidophytes and other phytoplankton. *Harmful Algae.* 7:1–10.
- Mahoney J. 2005. Modification of chemically-defined medium asp12 for picoplankter *Aureococcus anophagefferens*, with limited comparison of physiological requirements of New York and New Jersey isolates. Northeast Fisheries Science Centre Reference Document 05-17, Woods Hole, Massachusetts.
- Marsh EN. 1999. Coenzyme B12 (cobalamin)-dependent enzymes. *Essays Biochem.* 34:139154.
- Maruyama S, Matsuzaki M, Misawa K, Nozaki H. 2009. Cyanobacterial contribution to the genomes of the plastid-lacking protists. *BMC Evol Biol.* 9:197.
- Merchant SS, Prochnik SE, Vallon O, et al. (101 coauthors). 2007. The *Chlamydomonas* genome reveals the evolution of key animal and plant functions. *Science.* 318:245–251.
- Moreira D, Philippe H. 2001. Sure facts and open questions about the origin and evolution of photosynthetic plastids. *Res Microbiol.* 152:771–780.
- Moustafa A, Bhattacharya D. 2008. Phylosort: a user-friendly phylogenetic sorting tool and its application to estimating the cyanobacterial contribution to the nuclear genome of *Chlamydomonas*. *BMC Evol Biol.* 8:3.
- Newman SM, Boynton JE, Gillham NW, Randolph-Anderson BL, Johnson AM, Harris EH. 1990. Transformation of chloroplast ribosomal-RNA genes in *Chlamydomonas*—molecular and genetic-characterization of integration events. *Genetics* 126:875–888.
- Nishikimi M, Fukuyama R, Minoshima S, Shimizu N, Yagi K. 1994. Cloning and chromosomal mapping of the human nonfunctional gene for L-gulonono-gamma-lactone oxidase, the enzyme for L-ascorbic-acid biosynthesis missing in man. *J Biol Chem.* 269:13685–13688.
- Nishikimi M, Kawai T, Yagi K. 1992. Guinea pigs possess a highly mutated gene for L-gulonono-gamma-lactone oxidase, the key enzyme for L-ascorbic acid biosynthesis missing in this species. *J Biol Chem.* 267:21967–21972.
- Noble GP, Rogers MB, Keeling PJ. 2007. Complex distribution of EFL and EF-1 alpha proteins in the green algal lineage. *BMC Evol Biol.* 7:82.
- Palenik B, Grimwood J, Aerts A, et al. (38 co-authors). 2007. The tiny eukaryote *Ostreococcus* provides genomic insights into the paradox of plankton speciation. *Proc Natl Acad Sci U S A.* 104:7705–7710.

- Palmer EG, Starr RC. 1971. Nutrition of *Pandorina morum*. *J Phycol.* 7:85–89.
- Panzeca C, Beck AJ, Tovar-Sanchez A, Segovia-Zavala J, Taylor GT, Gobler CJ, Sanudo-Wilhelmy SA. 2009. Distributions of dissolved vitamin B₁₂ and Co in coastal and open-ocean environments. *Estuar Coast Shelf Sci.* 85:223–230.
- Panzeca C, Taylor GT, Sanudo-Wilhelmy SA, Leblanc K, Taylor GT, Hutchins DA, Sanudo-Wilhelmy SA. 2008. Potential cobalt limitation of vitamin B₁₂ synthesis in the North Atlantic Ocean. *Global Biogeochem Cycles.* 22:GB2029.
- Parfrey LW, Barbero E, Lasser E, Dunthorn M, Bhattacharya D, Patterson DJ, Katz LA. 2006. Evaluating support for the current classification of eukaryotic diversity. *PLoS Genet.* 2(12):e220.
- Pejchal R, Ludwig ML. 2005. Cobalamin-independent methionine synthase (metE): a face-to-face double barrel that evolved by gene duplication. *PLoS Biol.* 3:254–265.
- Pintner IJ, Provasoli L. 1959. The nutrition of *Volvox globator* and *V. tertius*. *Proc IX Int Bot Congr Montreal Abstr.* 9e:300–301.
- Pintner IJ, Provasoli L. 1968. Heterotrophy in sub-dyed light of 3 *Chrysochromulina* species. *Bull Misaki Mar Biol Inst Kyoto Univ.* 12:25–31.
- Pringsheim EG. 1970. Identification and cultivation of European *Volvox* sp. *Antonie Leeuwenhoek.* 36:33–43.
- Prochnik S, Umen J, Nedelcu A, et al. (28 co-authors). 2010. Genomic analysis of organismal complexity in the multicellular green alga *Volvox carteri*. *Science* 329:223–226.
- Provasoli L. 1958. Nutrition and ecology of protozoa and algae. *Annu Rev Microbiol.* 12:279–308.
- Ramirez-Flandes S, Ulloa O. 2008. Bosque: integrated phylogenetic analysis software. *Bioinformatics* 24:2539–2541.
- Raux E, Lanois A, Levillaye F, Warren M, Brody E, Rambach A, Thermes C. 1996. *Salmonella typhimurium* cobalamin (vitamin B₁₂) biosynthetic genes: functional studies in *S. typhimurium* and *Escherichia coli*. *J Bacteriol.* 178:753–767.
- Raux E, Lanois A, Warren M, Rambach A, Thermes C. 1998. Cobalamin (vitamin B₁₂) biosynthesis: identification and characterization of a *Bacillus megaterium* *cobI* operon. *Biochem J.* 335:159–166.
- Ravanel S, Block MA, Rippert P, Jabrin S, Curien G, Rebeille F, Douce R. 2004. Methionine metabolism in plants—chloroplasts are autonomous for de novo methionine synthesis and can import S-adenosylmethionine from the cytosol. *J Biol Chem.* 279:22548–22557.
- Reyes-Prieto A, Weber APM, Bhattacharya D. 2007. The origin and establishment of the plastid in algae and plants. *Annu Rev Genet.* 41:147–168.
- Saito S. 1972. Growth of *Gonium multicocum* in synthetic media. *J Phycol.* 8:169–175.
- Sanudo-Wilhelmy SA, Gobler CJ, Okbamichael M, Taylor GT. 2006. Regulation of phytoplankton dynamics by vitamin B₁₂. *Geophys Res Lett.* 33:L04604.
- Schneider MJ, Ulland M, Sloboda RD. 2008. A protein methylation pathway in *Chlamydomonas* flagella is active during flagellar resorption. *Mol Biol Cell.* 19:4319–4327.
- Scott J. 1999. Folate and vitamin B₁₂. *Proc Nutr Soc.* 58:441–448.
- Shihira I, Krauss RW. 1965. *Chlorella*: physiology and taxonomy of 41 isolates. Vol. 97. College Park, MD: University of Maryland.
- Sloboda RD, Howard L. 2009. Protein methylation in full length *Chlamydomonas* flagella. *Cell Motil Cytoskeleton.* 66:650–660.
- Smith AG, Croft MT, Moulin M, Webb ME. 2007. Plants need their vitamins too. *Curr Opin Plant Biol.* 10:266–275.
- Stein JR. 1966. Growth and mating of *Gonium pectorale* (Volvocales) in defined media. *J Phycol.* 2:27–28.
- Tripp HJ, Kitner JB, Schwalbach MS, Dacey JW, Wilhelm LJ, Giovannoni SJ. 2008. SAR11 marine bacteria require exogenous reduced sulphur for growth. *Nature* 452:741–744.
- Von Dassow P, Ogata H, Probert I, Wincker P, Da Silva C, Audic S, Claverie JM, de Vargas C. 2009. Transcriptome analysis of functional differentiation between haploid and diploid cells of *Emiliania huxleyi*, a globally significant photosynthetic calcifying cell. *Genome Biol.* 10(10):R114.
- Warren MJ, Raux E, Schubert HL, Escalante-Semerena JC. 2002. The biosynthesis of adenosylcobalamin (vitamin B₁₂). *Nat Prod Rep.* 19:390–412.
- Waterhouse A, Procter J, Martin D, Clamp M, Barton GJ. 2009. Jalview version 2—a multiple sequence alignment editor and analysis workbench. *Bioinformatics* 25(9):1189–1191.
- Whelan S, Goldman N. 2001. A general empirical model of protein evolution derived from multiple protein families using a maximum-likelihood approach. *Mol Biol Evol.* 18:691–699.
- Whitfield C, Weissbach H. 1970. Binding of folate substrate to 5-methyltetrahydropteroyltrimethylglutamate-homocysteine transmethylase. *J Biol Chem.* 245:402.
- Witman GB, Rosenbaum J, Berliner J, Carlson K. 1972. *Chlamydomonas* flagella. I. Isolation and electrophoretic analysis of microtubules, matrix, membranes, and mastigonemes. *J Cell Biol.* 54:507.
- Worden AZ, Lee JH, Mock T, et al. (48 co-authors). 2009. Green evolution and dynamic adaptations revealed by genomes of the marine picoeukaryotes *Micromonas*. *Science.* 325:147–147.
- Yamada K, Gravel RA, Toraya T, Matthews RG. 2006. Human methionine synthase reductase is a molecular chaperone for human methionine synthase. *Proc Natl Acad Sci U S A.* 103:9476–9481.
- Zhong Tang Y, Koch F, Gobler C. 2010. Most harmful algal bloom species are vitamin B₁ and B₁₂ auxotrophs. *Proc Natl Acad Sci U S A.* 107:20756–20761.