

Instanton Induced Neutrino Majorana Masses in CFT Orientifolds with MSSM-like spectra

L.E. Ibáñez¹ A.N. Schellekens² and A. M. Uranga³

¹ Departamento de Física Teórica C-XI and Instituto de Física Teórica C-XVI,
Universidad Autónoma de Madrid, Cantoblanco, 28049 Madrid, Spain

² NIKHEF, Kruislaan 409, 1009DB Amsterdam, The Netherlands;
IMAPP, Radboud Universiteit, Nijmegen, The Netherlands;
Instituto de Matemáticas y Física Fundamental, CSIC,
Serrano 123, Madrid 28006, Spain.

³ PH-TH Division, CERN CH-1211 Geneva 23, Switzerland
(On leave from IFT, Madrid, Spain)

Abstract

Recently it has been shown that string instanton effects may give rise to neutrino Majorana masses in certain classes of semi-realistic string compactifications. In this paper we make a systematic search for supersymmetric MSSM-like Type II Gepner orientifold constructions admitting boundary states associated with instantons giving rise to neutrino Majorana masses and other L- and/or B-violating operators. We analyze the zero mode structure of D-brane instantons on general type II orientifold compactifications, and show that only instantons with $O(1)$ symmetry can have just the two zero modes required to contribute to the 4d superpotential. We however discuss how the addition of fluxes and/or possible non-perturbative extensions of the orientifold compactifications would allow also instantons with $Sp(2)$ and $U(1)$ symmetries to generate such superpotentials. In the context of Gepner orientifolds with MSSM-like spectra, we find no models with $O(1)$ instantons with just the required zero modes to generate a neutrino mass superpotential. On the other hand we find a number of models in one particular orientifold of the Gepner model $(2, 4, 22, 22)$ with $Sp(2)$ instantons with a few extra uncharged non-chiral zero modes which could be easily lifted by the mentioned effects. A few more orientifold examples are also found under less stringent constraints on the zero modes. This class of $Sp(2)$ instantons have the interesting property that R-parity conservation is automatic and the flavour structure of the neutrino Majorana mass matrices has a simple factorized form.

1 Introduction

String Theory, as the leading candidate for a unified theory of Particle Physics and Gravity, should be able to describe all observed particle phenomena. One the most valuable experimental pieces of information obtained in the last decade concerns neutrino masses. Indeed the evidence from solar, atmospheric, reactor and accelerator experiments indicates that neutrinos are massive. The simplest explanation of the smallness of neutrino masses is the see-saw mechanism [1]. The SM gauge symmetry allows for two types of operators bilinear on the neutrinos (with dimension ≤ 4) :

$$\mathcal{L}_\nu = M_{ab}\nu_R^a\nu_R^b + h_{ab}\nu_R^a\bar{H}L^b \quad (1.1)$$

where ν_R is the right-handed neutrino, L is the left-handed lepton doublet and \bar{H} is the Higgs field. In supersymmetric theories, this term arise from a superpotential with the above structure, upon replacing fields by chiral superfields. If M_{ab} is large, the lightest neutrino eigenvalues have masses

$$M_\nu = \langle \bar{H} \rangle^2 h^T M^{-1} h \quad (1.2)$$

For $M \sim 10^{10} - 10^{13}$ GeV and Dirac neutrino masses of order charged lepton masses, the eigenvalues are consistent with experimental results.

What is the structure of neutrinos and their masses in string theory? In specific compactifications giving rise to the MSSM spectra singlet fields corresponding to right-handed neutrinos ν_R generically appear. Dirac neutrino masses are also generically present but the required Majorana ν_R masses are absent. This is because most MSSM-like models constructed to date have extra $U(1)$ symmetries, under which the right-handed neutrinos are charged, which hence forbid such masses. In many models, such symmetries are associated to a $U(1)_{B-L}$ gauge boson beyond the SM. In order to argue for the existence of ν_R masses, string model builders have searched for non-renormalizable couplings of the type $(\nu_R\nu_R\bar{N}_R\bar{N}_R)$ with extra singlets N_R . Once the latter fields get a vev, $U(1)_{B-L}$ is broken and a Majorana mass appears for the ν_R . Although indeed such couplings (or similar ones with higher dimensions) exist in some semi-realistic compactifications, such a solution to the neutrino mass problem in string theory has two problems: 1) The typical ν_R masses so generated tend to be too small due to the higher dimension of the involved operators and 2) The vevs for the N_R fields breaks spontaneously R-parity so that dimension 4 operators potentially giving rise to fast proton decay are generated. This is in a nutshell the neutrino problem in string compactifications (see [2] for a recent discussion in heterotic setups).

In [3] (see also [4]) two of the present authors pointed out that there is a built-in mechanism in string theory which may naturally give rise to Majorana masses for right-handed neutrinos. It was pointed out that string theory instantons may generate such masses through operators of the general form

$$M_{string} e^{-U} \nu_R \nu_R \quad . \quad (1.3)$$

Here U is a linear combination of closed string moduli whose imaginary part gets shifted under a $U(1)_{B-L}$ gauge transformation in such a way that the operator is fully gauge invariant. The exponential factor comes from the semi-classical contribution of a certain class of string instantons. This is a pure stringy effect distinct from the familiar gauge instanton effects which give rise to couplings violating anomalous global symmetries like $(B+L)$ in the SM. Here also $(B-L)$ (which is anomaly-free) is violated.

This operator is generated due to existence of instanton fermionic zero modes which are charged under $(B-L)$ and couple to the ν_R chiral superfield. Although the effect can take place in different constructions, the most intuitive description may be obtained for the case of Type IIA CY orientifold compactifications with background D6-branes wrapping 3-cycles in the CY. In the simplest configurations one has four SM stacks of D6-branes labeled **a**, **b**, **c**, **d** which correspond to $U(3)$, $SU(2)$ (or $U(2)$), $U(1)_R$ and $U(1)_L$ gauge interactions respectively, which contain the SM group. One can construct compactifications with the MSSM particle spectrum in which quarks and leptons lie at the intersections of those SM D6-branes. Then the relevant instantons correspond to euclidean D2-branes wrapping 3-cycles in the CY (satisfying specific properties so as to lead to the appropriate superspace interaction). The D2-D6 intersections lie the additional fermionic zero modes which are charged under $(B-L)$. For instantons with the appropriate number of intersections with the appropriate D6-branes, and with open string disk couplings among the zero modes and the ν_R chiral multiplet (see fig.(3.1)), the operator in (1.3) is generated.

The fact that the complex modulus U transforms under $U(1)_{B-L}$ gauge transformations indicates that the $U(1)_{B-L}$ gauge boson gets a mass from a Stückelberg term. So a crucial ingredient in the mechanism to generate non-perturbative masses for the ν_R 's is that there should be massless $U(1)_{B-L}$ gauge boson which become massive by a Stückelberg term. It turns out that not many semi-realistic models with $U(1)_{B-L}$ mass from Stückelberg couplings have been constructed up to date. In the literature there are two main classes of RR tadpole free models with massive B-L. The first class are non-susy type IIA toroidal orientifold models first constructed in [5]. The second class are the type II Gepner orientifold models constructed by one of the present authors and

collaborators [6, 7]. The former were already considered in [3]. In the present paper we will concentrate on the RCFT Gepner model constructions, which lead to a large class of MSSM like models, more representative of the general Calabi-Yau compactifications (for a recent discussion of instanton-induced neutrino masses in a model with no RR tadpole cancellation, see [8]).

The class of constructions in [6, 7] start with any of the 168 Type II compactifications obtained by tensoring $N = 2$ SCFT minimal models. In addition one can choose a number of modular invariant partition functions (MIPF), leading to a total of 5403. Then different consistent orientifold projections are performed on the different models. This yields a total of 49304 Type II orientifolds. The open string sector of the theory is defined in terms of the boundary states of the theory. Intuitively, they play the same role as D-branes wrapping cycles in the geometrical settings. Thus one associates boundary states **a, b, c, d** to the gauge groups giving rise to the SM. Different choices for the SM boundary states lead to different spectra. In the present paper we will make use of the data in [6] which contains 211634 different MSSM-like spectra (including also different hidden sectors). Although this number is huge, most of these models are really extensions of the MSSM, since they have either an extra $U(1)_{B-L}$ or $SU(2)_R \times U(1)_{B-L}$ group factor beyond the SM group. As we said, we are actually only interested in models in which the $U(1)_{B-L}$ gets a Stückelberg mass. Then we find that the number of MSSM-like models with these characteristics is dramatically reduced: only 0.18 percent of the models (391) have a massive $U(1)_{B-L}$.

As we said, in the geometrical setting of IIA orientifolds with intersecting D6-branes [9, 10] (see [11] for reviews and [12, 13] for the IIB counterparts), instantons are associated to D2-branes wrapping 3-cycles, like the background D6-branes do. Analogously, in the RCFT setting the same class of boundary states appearing in the SM constructions are the ones corresponding to instantons. The zero modes on the instanton is computable from the overlaps of instanton brane boundary states (zero modes uncharged under the 4d gauge group) or of instanton and 4d spacefilling brane boundary states (zero modes charged under the corresponding gauge factor). We find that the criteria for a non-perturbative superpotential to be generated [14] are only fulfilled if the Chan-Paton (CP) symmetry of the instantons is $O(1)$. For instantons with CP symmetry ¹ $Sp(2)$ or $U(1)$ we find that there are a few extra uncharged fermionic zero modes which would preclude the formation of the searched superpotentials. On the other hand we argue that the addition of fluxes and/or possible non-perturbative

¹We adopt the convention that the fundamental representation of $Sp(m)$ is m -dimensional.

extensions of the orientifold compactifications would allow also instantons with $Sp(2)$ and $U(1)$ symmetries to generate such superpotentials. We thus include all $O(1)$, $Sp(2)$ and $U(1)$ instantons ² in our systematic search. The computation of charged and uncharged fermion zero modes may be easily implemented as a routine in a systematic computer search for instanton zero modes in Gepner MSSM-like orientifolds. Results of such a systematic computer search are presented in this article.

We find that out of the 391 models with massive $U(1)_{B-L}$, there are very few admitting instantons with the required minimal $O(1)$ CP symmetry, and in fact none of them without additional vector-like zero modes. On the other hand we do find 32 models admitting $Sp(2)$ symmetric instantons with just the required charged zero mode content (and the minimal set of non-chiral fermion zero modes). In fact they are all variations of the same orientifold Gepner model based on the tensor product $(k_1, k_2, k_3, k_4) = (2, 4, 22, 22)$. These models all in fact correspond to the same MIPF and orientifold projection, they only differ on which particular boundary states corresponding to the four **a, b, c, d** SM ‘stacks’. All models have the same chiral content, exactly that of the MSSM, with extra vectorlike chiral fields which may in principle become massive in different points of the CY moduli space. They have no hidden sector, i.e., the gauge group is just that of the SM. For each of the models there are 8 instantons with $Sp(2)$ CP symmetry with just the correct charged zero mode structure allowing for the superpotential coupling (1.3) giving rise to ν_R Majorana masses. As we said, they have extra uncharged fermion zero modes beyond the two required to generate a superpotentials. However one would expect that these unwanted zero modes might be lifted in more generic situations in which e.g. NS/RR fluxes are added.

We thus see that, starting with a ‘large’ landscape of 211634 MSSM-like models, and searching for instantons inducing neutrino masses, we find there are none admitting the $O(1)$ instantons with exactly the required zero mode structure, and only few (32) examples with $Sp(2)$ instantons with next-to-minimal uncharged zero mode structure (and exactly the correct charged zero modes). Let us emphasize though that it is the existence of massive $U(1)_{B-L}$ models which is rare. Starting with the subset of models with a massive $U(1)_{B-L}$, finding models with correct instanton charged zero modes within that class is relatively frequent, 10 percent of the cases. Furthermore, we will see that there are further models beyond those 32 which contain extra non-chiral instanton zero modes and which may also be viable if these modes get massive by some

²We refer to the different kinds of instanton by their Chan-Paton symmetry on their volume. Since we are not interested in gauge theory instantons, this notation should not be confusing.

effect (like e.g. the presence of RR/NS fluxes).

Instantons may generate some other interesting superpotential couplings in addition to ν_R masses, some possibly beneficial and others potentially dangerous. In particular we find that in the models which contain $Sp(2)$ instantons which might induce ν_R masses, there are also other instantons which would give rise directly to the Weinberg operator [15]

$$\mathcal{L}_W = \frac{\lambda}{M}(L\overline{H}L\overline{H}) \quad (1.4)$$

Once the Higgs field gets a vev, this gives rise directly to left-handed neutrino masses. Thus we find that in that class of models both the see-saw mechanism (which also gives rise to a contribution to the Weinberg operator) and an explicit Weinberg operator might contribute to the physical masses of neutrinos. Which effect dominates will depend on the relative size of the corresponding instanton actions as well as on the size of the string scale. Among potentially dangerous operators which might be generated stand the R-parity violating operators of dimension < 5 , which might give rise e.g. to fast proton decay. We make a study of the possible generation of those, and find that in all models in which ν_R masses might be generated R-parity is exactly conserved. This is a very encouraging result.

A natural question to ask is whether one can say something about the structure of masses and mixings for neutrinos. As argued in [3] generically large mixing angles are expected, however to be more quantitative we also need to know the structure of Yukawa couplings for leptons. In principle those may be computed in CFT but in practice this type of computation has not yet been developed for CFT orientifolds. Nevertheless we show that, in the case of instantons with $Sp(2)$ CP symmetry, a certain factorization of the flavor structure takes place, which could naturally give rise to a hierarchical structure of eigenvalues for neutrino masses.

The structure of this article is as follows. In the next section we present a discussion of instanton induced superpotentials in Type II orientifolds. This discussion will apply both to Type IIA and Type IIB CY orientifolds as well as to more abstract CFT orientifolds. We discuss the structure of both uncharged and charged instanton zero modes. In particular we show that only instantons with $O(1)$ CP symmetry have the appropriate uncharged zero mode content to induce a superpotential contribution. We also discuss how $Sp(2)$ and $U(1)$ might still generate superpotential contributions if extra ingredients are added to the general setting. In section 3 we apply that discussion to the specific case of the generation of ν_R Majorana masses, showing what is the required zero mode structure in this case. We show how the flavor structure of the

Majorana mass term factorizes in the case of instantons with $Sp(2)$ CP symmetry, leading potentially to a hierarchical structure of eigenvalues. We further discuss the generation of other B/L-violating operators including the generation of the Weinberg operator as well as R-parity violating couplings. In section 4 we review the RCFT Type II orientifold constructions in [6, 7]. A general discussion of zero fermion modes for instantons in RCFT orientifolds is presented in section 5.

The results of our general search for instantons generating ν_R masses are presented in section 6. We provide a list of all Gepner orientifolds which admit instanton configurations potentially giving rise to ν_R Majorana masses. We describe the structure of the models with $Sp(2)$ instantons having the required charged zero modes for that superpotential to be generated. We also describe the boundary states of the corresponding instantons and provide the massless spectrum of the relevant MSSM-like models. Other orientifolds with zero mode structure close to the minimal one are also briefly discussed. A brief discussion about the possible generation of R-parity violating superpotentials is included. We leave section 7 for some final comments. Some notation on the CFT orientifold constructions, and a discussion of the CFT symmetries in the $Sp(2)$ examples are provided in two appendices.

As this paper was ready for submission, we noticed [16, 17], whose discussion of instanton zero modes partially overlaps with our analysis in Section 2.2.

2 Instanton induced superpotentials in Type II orientifolds

In this Section we review the generation of superpotentials involving 4d charged fields via D-brane instantons in type II compactifications. The discussion applies both to type IIA and IIB models, and to geometrical compactification as well as to more abstract internal CFT's. For recent discussions on D-brane instantons, see [4, 3, 25, 8] .

Before starting, a notational remark is in order. Our notation is adapted to working in the covering theory, namely the type II compactification, and orientifolding in a further step. Thus we describe the brane configurations as a system of branes (described by boundary states for abstract CFT's), labeled k , and their orientifold images labeled k' . Similarly, we denote M the brane / boundary state corresponding to the instanton brane, and M' its orientifold image. If a brane is mapped to itself under the orientifold action, we call it a 'real' brane / boundary, and 'complex' otherwise.

2.1 D-brane instantons, gauge invariance and effective operators

A basic feature of type II orientifold compactifications with D-branes is the generic presence of Stückelberg couplings between the $U(1)$ gauge fields on the D-branes, and certain 4d RR closed string 2-forms. These couplings are required by the Green-Schwarz mechanism when the $U(1)$'s have non-zero triangle contributions to mixed anomalies [18, 19], but can also exist for certain non-anomalous $U(1)$'s [5, 20]. These couplings make the $U(1)$ gauge bosons massive, but the symmetry remains as a global symmetry exact in perturbation theory. Since the closed string moduli involved are scalars (0-forms) in the RR sector, the natural candidate non-perturbative effects to violate these $U(1)$ symmetries are instantons arising from euclidean D-branes coupling to these fields.

In computing the spacetime effective interaction mediated by the instanton, one needs to integrate over the instanton zero modes. In the generic case (and in particular for the case of our interest) there are no bosonic zero modes beyond the universal ones (namely, the four translational bosonic zero modes associated to the position of the instanton). On the other hand, the structure of fermion zero modes will be crucial. Since we are interested in models with non-trivial 4d gauge group, arising from a set of background 4d spacetime filling branes, we consider separately fermion zero modes which are uncharged under the 4d gauge group and those which are charged. In this paper we restrict our discussion to 4d $N = 1$ supersymmetric models, and this will simplify the analysis of zero modes.

Fermion zero modes which are uncharged under the 4d gauge group determine the kind of 4d superspace interaction which is generated by the instanton. We are interested in generating superpotential interactions, which, as is familiar, requires the instanton to have two fermion zero modes to saturate the $d^2\theta$ superspace integration. For this, a necessary (but not sufficient!) condition is that the D-branes are half-BPS, so these fermion zero modes are the Goldstinos of the two broken supersymmetries. In the string description, uncharged zero modes arise from open strings in the MM sector (in our notation, the one leading to adjoint representations), which in particular contain these Goldstinos, and the MM' sector (in our notation, the one leading to two-index symmetric or antisymmetric tensors). Note that both are the same for 'real' branes. Hence we are primarily interested in D-branes whose MM sector contains just two fermion zero modes, and whose MM' sector (for 'complex' branes) does not contain additional fermion zero modes.

In analogy with the familiar case of gauge theory instantons [21], charged fermion zero modes determine the violation of perturbative global symmetries by the instanton-induced interaction. Namely, in order to saturate the integration over the charged fermions zero modes, the spacetime interaction must contain insertions of fields charged under the 4d gauge symmetry, and in particular under the global $U(1)$ factors, which are thus violated by the D-brane instanton. Notice that this holds irrespectively of the number of uncharged fermion zero modes, namely of the kind of superspace interaction induced by the instanton. Restricting to superpotential interactions, the resulting operator in the 4d effective action has roughly the form

$$W_{inst} = e^{-U} \Phi_1 \dots \Phi_n \quad (2.1)$$

Here the fields Φ_1, \dots, Φ_n are 4d $N = 1$ chiral multiplets charged under the 4d gauge group, and in particular also under the $U(1)$ symmetries. Note also that the instanton amplitude contains a prefactor (which in general depends on closed and open string moduli) arising from the Gaussian path integral over (massive) fluctuations of the instanton (hence described by an open string annulus partition function, see [22, 23] for related work), which we can ignore for our purposes in this paper.

For D-brane instantons, U is the closed string modulus to which the euclidean D-brane couples. In the D-brane picture, instanton fermion zero modes charged under the gauge factor carried by the k^{th} stack of 4d space-filling branes (and its image k') arise from open strings in the Mk and Mk' sectors, transforming as usual in the $(\square_M, \bar{\square}_k)$ and (\square_M, \square_k) representations, respectively (with both related in the case of ‘real’ branes). The (net) number of instanton fermion zero modes with such charges is given by certain multiplicities ³ $I_{Mk}, I_{Mk'}$.

A D-brane instanton, irrespectively of the superspace structure of the 4d interactions it may generate, violates $U(1)_k$ charge conservation by an amount $I_{Mk} - I_{Mk'}$ for ‘complex’ branes and I_{Mk} for ‘real’ branes. In particular, this is the total charge of the field theory operator $\Phi_1 \dots \Phi_n$ in (2.1). From the Stückelberg couplings, it is possible to derive [3] (see [23, 24, 25] for related work, also [26])

³In geometric type IIA compactifications with 4d spacetime-filling branes and instanton branes given by D6- and D2-branes wrapped on Special Lagrangian 3-cycles, I_{Mk} corresponds to the intersection number between the 3-cycles corresponding to the k^{th} D6- and the D2-brane M (and similarly for $I_{Mk'}$). In geometric type IIB compactifications, it corresponds to the index of a suitable Dirac operator. In general (even for abstract CFT’s) it can be defined as the Witten index for the 2d theory on the open string with the boundary conditions corresponding to the two relevant branes. We will often abuse language and refer to this quantity as intersection number, even in Section 6 where we work in the non-geometric regime of type IIB compactifications.

that for ‘complex’ instantons, gauge transformations of the $U(1)_k$ vector multiplets $V_k \rightarrow V_k + \Lambda_k$, transform U as

$$U \rightarrow U + \sum_k N_k (I_{Mk} - I_{Mk'}) \Lambda_k \quad (2.2)$$

For ‘real’ brane instantons, which were not considered in [3], the shift is given by ⁴

$$U \rightarrow U + \sum_k N_k I_{Mk} \Lambda_k \quad (2.3)$$

(this new possibility will be an important point in our instanton scan in Section 6).

The complete interaction (2.1) is invariant under the $U(1)$ gauge symmetries. However, from the viewpoint of the 4d low-energy effective field theory viewpoint, it leads to non-perturbative violations of the perturbative $U(1)$ global symmetries, by the amount mentioned above.

In the string theory construction there is a simple microscopic explanation for the appearance of the insertions of the 4d charged fields (related to the mechanism in [27]). The instanton brane action in general contains cubic terms $\alpha \Phi \gamma$, involving two instanton fermions zero modes α in the $(\square_M, \bar{\square}_k)$ and γ in the $(\square_p, \bar{\square}_M)$ coupling to the 4d spacetime field Φ in the $(\square_k, \bar{\square}_p)$ of the 4d gauge group ⁵. Performing the Gaussian path integral over the instanton fermion zero modes leads to an insertion of Φ in the effective spacetime interaction. In general, and for a ‘complex’ instanton, there are several fermion zero modes α_i, γ_i in the fundamental (resp. antifundamental) of the instanton gauge group, coupling to a 4d spacetime chiral operators \mathcal{O}_{ij} (which could possibly be elementary charged fields, or composite chiral operators). Gaussian integration over the fermion zero modes leads to an insertion of the form $\det \mathcal{O}$ (for ‘real’ brane instantons, $\det \mathcal{O}$ should be interpreted as a Pfaffian). It is straightforward to derive our above statement on the net charge violation from this microscopic mechanism.

Note that the above discussions show that instantons in different topological sectors (namely with different RR charges, and different intersection numbers with the 4d spacefilling branes) contribute to different 4d spacetime operators. In particular, multiwrapped instantons, if they exist as BPS objects, contribute to operators different

⁴Equivalently, one may use (2.2), but must include an additional factor of 1/2 from the reduction of the volume for a real brane (which is invariant under the orientifold action).

⁵ Although there is no chirality in 0+0 dimension, the fermion zero modes α and γ are distinguished by their chirality with respect to the $SO(4)$ global symmetry of the system (which corresponds to rotations in the 4d spacetime dimensions, longitudinal to the space-filling branes, and transverse to the instanton brane). Supersymmetry of the instantons constrains the couplings on the instanton action (such as the cubic couplings above) to have a holomorphic structure.

from the singly wrapped instanton. This implies that the instanton expansion for a fixed operator is very convergent, and could even be finite.

Another important implication of the above discussion is that, in order to generate a specific operator via an instanton process, a necessary condition is that the instanton has an appropriate number and structure of charged zero modes. However, this is not sufficient. Insertions of 4d fields appear only if the fields couple to the instanton fermion zero modes via terms at most quadratic in the zero modes. In equivalent terms, only zero modes appearing in the Gaussian part of the instanton action can be saturated by insertions of 4d fields (those to which they couple). The requirement that the zero modes have appropriate couplings to the 4d fields may be non-trivial to verify in certain constructions. This is the case for the Gepner model orientifolds in coming sections, whose couplings are computable in principle, but unknown in practice. In such cases we will assume that any coupling which is not obviously forbidden by symmetries will be non-vanishing. Unfortunately there are no arguments to estimate the actual values of such non-vanishing couplings, hence we can argue about the existence of certain instanton induced operators, but not about the coefficients of such terms.

2.2 Zero mode structure for D-brane instantons

In this section we describe more concretely different kinds of instantons and the structure of interesting and unwanted zero modes. Our discussion will be valid for general D-brane models in perturbative type II orientifolds without closed string fluxes, although we also make some comments on more general F-theory vacua and the effects of fluxes. A more specific discussion is presented in Section 5.

2.2.1 Uncharged zero modes

We start discussing zero modes uncharged under the 4d gauge group. These are crucial in determining the kind of superspace interaction induced by the instanton on the 4d theory. In particular, we are interested in instantons contributing to the 4d superpotential, namely those which contain just two fermion zero modes in this sector. We are also interested in instantons which may contain additional fermion zero modes, and the possible mechanisms that can be used to lift them. Let us discuss ‘real’ and ‘complex’ brane instantons in turn.

- **Real brane instantons**

Real brane instantons correspond to branes which are mapped to themselves by the

orientifold action, hence $M = M'$. Uncharged zero modes arise from the MM open string sector. As discussed in Section 5, there is a universal sector of zero modes, in the sense that it is present in any BPS D-brane instanton, which we now describe. Before the orientifold projection, we have a gauge group $U(n)$ on the volume of n coincident instantons. Notice that, although there are no gauge bosons in $0 + 0$ dimensions, the gauge group is still well-defined, since it acts on charged states (open string ending on the instanton brane). There are four real bosonic zero modes and four fermion zero modes in the adjoint representation. For the minimal $U(1)$ case, the four bosons are the translational Goldstones. The four fermions arise as follows. This sector is insensitive to the extra 4d spacefilling branes, and feels an accidental 4d $N = 2$ supersymmetry. The BPS D-brane instanton breaks half of this, and leads to four Goldstinos, which are the described fermions ⁶.

The orientifold projection acts on this universal sector as follows (see Section 5 for further discussion). The gauge group is projected down to orthogonal or symplectic. Hence instanton branes with symplectic gauge group must have even multiplicity (a related argument, in terms of the orientifold action on Chan-Paton indices, is given in Section 5). For instantons with $O(n)$ gauge symmetry, the orientifold projects the four bosonic zero modes and two fermion zero modes (related by the two supercharges of 4d $N = 1$ supersymmetry broken by the instanton) to the two-index symmetric representation, and the other two fermion zero modes (related by the other two supercharges of the accidental 4d $N = 2$ in this sector) to the antisymmetric representation. Hence for $O(1)$ instantons (namely instantons with $O(1)$ gauge group on their volume), we have just two fermion zero modes, which are the Goldstinos of 4d $N = 1$ supersymmetry, and the instanton can in principle contribute to the superpotential (if no additional zero modes arise from other non-universal sectors). For instantons with $Sp(n)$ gauge symmetry, the orientifold projects the four bosonic zero modes and two fermion zero modes to the two-index antisymmetric representation, and the other two fermion zero modes to the symmetric representation. Hence, even for the minimal case of $Sp(2)$ instantons, we have just two fermion zero modes in the triplet representation, in addition to the two 4d $N = 1$ Goldstinos. Hence $Sp(2)$ instantons cannot contribute to the superpotential in the absence of additional effects which lift these zero modes (see later) ⁷.

⁶We thank F. Marchesano for discussions on this point.

⁷ For D-brane instantons corresponding to gauge instantons, the additional fermion zero modes in the universal sector couple to the boson and fermion zero modes from open strings stretched between the instanton and the 4d spacefilling brane. They act as Lagrange multipliers which impose

In addition to this universal sector, there exist in general additional modes, whose presence and number depends on the detailed structure of the branes. Namely, on the geometry of the brane in the 6d compact space in geometric compactifications, or on the boundary state of the internal CFT in more abstract setups. They lead to a number of boson and fermion zero modes in the symmetric or antisymmetric representation. The computation of these multiplicities in terms of the precise properties of the instanton branes is postponed to Section 5. In order to generate a superpotential, one must require these modes to be absent, except for the case of antisymmetrics of $O(1)$ instantons, which are actually trivial.

An important point is that extra fermion zero modes (including the extra triplet fermion zero modes in the universal sector of $Sp(2)$ instantons, and any two-index tensor fermion zero mode in the non-universal sectors) are in principle not protected against acquiring non-zero masses once the model is slightly modified. In other words, such fermions are non-chiral, in terms of the $SO(4)$ chirality in footnote 5. One such modification is motion in the closed string moduli space, which can lift the non-universal modes if there are non-trivial couplings between them and closed string moduli (unfortunately, such correlators are difficult to compute, even in cases where the CFT is exactly solvable, like the Gepner models). Note that extra zero modes in the universal sector of $Sp(2)$ instantons cannot be lifted by this effect, since it does not break the accidental 4d $N = 2$ in this sector. A second possible modification which in general can lift extra zero modes is the addition of fluxes, generalizing the results for D3-brane instantons in geometric compactifications [29] (for non-geometric CFT compactifications, we also expect a similar effect, once fluxes are introduced following [30]). Note that fluxes can lift extra zero modes in the universal sector as well, since they can break the accidental 4d $N = 2$ susy in this sector. A last mechanism arising in more general F-theory compactifications and discussed below for complex instantons, is valid for real instanton branes as well.

The bottom line is that in the absence of such extra effects, only $O(1)$ instantons can contribute to superpotential terms. However, in modifications of the model such extra effects can easily lift the extra fermion zero modes. Hence, this kind of extra vector-like zero modes will not be considered catastrophic, and real instantons (including the $O(1)$ and $Sp(2)$ cases) with such zero modes are considered in our scan in Section 6.

the fermionic constraints in the ADHM construction [28], and may not spoil the generation of a superpotential.

- **Complex brane instantons**

Zero modes uncharged under the 4d gauge group can arise from the MM and MM' open string sectors. Notice that the orientifold action maps the MM sector to the $M'M'$, hence we simply discuss the former and impose no projection. The discussion of the MM sector is as for real brane instantons before the orientifold projection. The universal sector leads to a $U(n)$ gauge symmetry, and four bosonic and four fermionic zero modes in the adjoint representation. The bosons are translational Goldstones, while the fermions are Goldstinos of the accidental 4d $N = 2$ present in this sector. Hence, even in the minimal case of $U(1)$ brane instantons there are two extra fermion zero modes, beyond the two fermion zero modes corresponding to the 4d $N = 1$ Goldstinos. Hence $U(1)$ instanton (except for those corresponding to gauge instantons, see footnote 7) cannot contribute to superpotential terms in the absence of additional effects, like closed string fluxes. However, keeping in mind the possibility of additional effects lifting them in modifications of the model, we include them in the discussion. Also, in what follows we will use the $U(n)$ notation for the different fields to keep track of the Chan-Paton index structure.

The above statement would seem in contradiction⁸ with computations of non-perturbative superpotentials [14] induced by M5-branes instantons in M-theory compactifications on Calabi-Yau fourfolds, which are dual to D3-brane instantons (with world-volume $U(1)$ gauge group) on type IIB compactifications. The resolution is that the M5-branes that contribute are intersected by different (p, q) degenerations of the elliptic fiber. This implies that $U(1)$ D3-brane instanton only contribute if they are intersected by mutually non-local (p, q) 7-branes. The two extra fermion zero modes exist locally on the D3-brane volume, but cannot be defined globally due to the 7-brane monodromies. Hence such effect can take place only on non-perturbative type IIB compactifications including (p, q) 7-branes. Note that in perturbative compactifications, namely IIB orientifolds, the (p, q) 7-branes are hidden inside orientifold planes [31] with their monodromy encoding the orientifold projection; hence the only branes that can contribute to the superpotential are real branes, for which the projection/monodromy removes the extra fermion zero modes, as discussed above.

In addition to this universal sector, the MM sector may contain a non-universal set of fermions and bosons, in the adjoint representation (hence uncharged under $U(1)$). They depend on the specific properties of the brane instanton, and will be discussed in Section 5. These additional zero modes should be absent in order for the instanton

⁸We thank S. Kachru for discussions on the ideas in this paragraph.

to induce a non-trivial superpotential. Notice however that these zero modes are uncharged under any gauge symmetry, and hence vector-like. Thus, there could be lifted in modifications of the model, as discussed for real instantons.

The MM' sector is mapped to itself under the orientifold action. Hence it leads to a number of bosons and fermions in the two index symmetric or antisymmetric representations. Notice that the two-index antisymmetric representation is trivial for $U(1)$, so these modes are actually not present. On the other hand, fermion zero modes in the two-index symmetric representation are chiral and charged under the brane instanton gauge symmetry. Hence they cannot be lifted by any of the familiar mechanisms, and thus spoil the appearance of a non-perturbative superpotential, even if the model is modified. Such fermion zero modes are considered catastrophic and we will look for models avoiding them in our scan in Section 6.

2.2.2 Charged fermion zero modes

• Real brane instantons

Instanton zero modes charged under the 4d gauge group arise from Mk open string sectors (and their image Mk'). In the generic case, there are no scalar zero modes in these sectors. This is because in mixed Mk open string sectors the 4d spacetime part leads to DN boundary conditions, which already saturate the vacuum energy in the NS sector. Only in the special case where the internal structure of the spacetime filling brane k and the instanton brane are the same, there may be NS ground states of the internal CFT which do not contribute extra vacuum energy, hence leading to massless scalars. However, this corresponds to brane instantons which can be interpreted as instantons of the 4d gauge theory on the 4d space-filling branes. The instantons we are interested in for the generation of neutrino Majorana mass terms are not of this kind [3] (see e.g. [32, 33, 28, 25, 34] for discussions on gauge theory instantons from D-brane instantons).

Hence we focus on charged fermion zero modes, which are generically present in any mixed Mk sector. Let us define L_{Mk} , $L_{Mk'}$ the (positive by definition) number of left-handed chiral fermion zero modes in the representations (\square_M, \square_k) , $(\square_M, \bar{\square}_k)$, respectively. The net number of chiral fermion zero modes in the $(\square_M, \bar{\square}_k)$ is given by $I_{Mk} = L_{Mk'} - L_{Mk}$. This controls the violation of the $U(1)_a$ global charge by the instanton. Namely, such fermion zero modes in the Mk , Mp sectors lead (if suitable couplings are present) to the insertion of 4d charged fields Φ_{kp} and/or $\bar{\Phi}_{kp'}$.

In addition, there are $P_{Mk} = \min(L_{Mk'}, L_{Mk})$ vector-like pairs of fermion zero

modes. Since they are vector-like, they may be lifted by slight modifications of the model, like moving in the closed string moduli space, or by introducing additional ingredients, like fluxes. In addition, they may be lifted by moving in the open string moduli space of the 4d spacefilling branes, as follows. The zero modes may lead to insertions of 4d fields Φ_{kk} , if the kk sector contains such 4d chiral multiplets (or to insertions of composite 4d operators in the adjoint of the k^{th} 4d gauge factor), and if they couple to the zero modes. Although this may not be generically not the case, many of our models in coming section contain such adjoint fields. Hence, a non-trivial vev for the latter can lift these extra vector-like zero modes, hence leading to instanton generating the superpotential of interest. Given these diverse mechanisms to lift these zero modes, their presence of such zero modes is thus unwanted, but again not necessarily catastrophic.

One last comment, related to the concrete kind of instanton search we will be interested in. Namely, we will be searching for instantons leading to a specific operator, carrying non-trivial charges under a specific set of 4d gauge factors. Postponing the detailed discussion to Sections 3.1, 4.2, let us denote **a**, **b**, **c**, **d** the set of branes leading to a field theory sector, denoted ‘observable’ (and which reproduces the SM in our examples). We will require the instanton to have a prescribed number of chiral fermion zero modes charged under these branes, namely we require the intersection numbers of the instanton with these branes $I_{M\mathbf{a}}, \dots, I_{M\mathbf{d}}$ to have specific values (as mentioned above, in the most restrictive scan we forbid vector-like pairs of zero modes under these branes). In addition, the model in general contains an additional sector of branes, denoted ‘hidden’ (since there is zero net number of chiral multiplets charged under both sectors) and labeled h_i , required to fulfill the RR tadpole cancellation conditions. In general there may be instanton fermion zero modes from e.g. the Mh_1 , h_2M sectors, which would contribute to insertions of the 4d fields in the h_1h_2 sector if there are suitable cubic couplings. These extra insertions could be avoided if such 4d fields in the hidden sector acquire vevs (note that vevs for the (vector-like) fields charged under the visible and hidden sectors would typically break hypercharge, and should be avoided), and hence lift the zero modes. Equivalently, from the 4d perspective, the unwanted extra h_1h_2 field insertion is replaced by its vev. However, this renders the discussion very model dependent. Moreover, the possibility of hidden brane recombination was not included in the search for SM-like models in [6, 7] (namely, the possibility of allowing for chiral fields charged under the observable and hidden gauge groups, which may become non-chiral and possibly massive upon hidden brane

recombination). Hence we will consider these chiral fermion zero modes as unwanted (as usual, non-chiral modes are unwanted but not catastrophic, hence they are allowed for in a more relaxed search).

• **Complex brane instantons**

The discussion of ‘complex’ brane instantons is somewhat analogous to the previous one, with the only complication that the brane M and its image M' lead to independent modes, leading to a more involved pattern of fermion zero modes. Instanton zero modes charged under the 4d gauge group arise from the Mk, Mk' and related sectors. As for ‘real’ brane instantons, there are generically no scalars in these sectors (and certainly not in our case of interest). Hence we focus on charged fermion zero modes, which are generically present in any mixed sector.

In contrast with ‘real’ brane instantons, a net combination of fermion zero modes in the $(\square_M, \bar{\square}_k) + (\square_M, \square_k)$ is not vector-like, but chiral under the instanton gauge symmetry. Such a pair cannot therefore be lifted even by modifications of the theory. In general, if the instanton has a mismatch in the total numbers n_α, n_γ of fermion zero modes α_i in the \square_M and γ_j in the $\bar{\square}_M$, the instanton amplitude automatically vanishes. Namely, the matrix of operators \mathcal{O}_{ij} coupling to the zero modes necessarily has rank at most $\min(n_\alpha, n_\gamma)$. That is, if $n_\alpha > n_\gamma$ there are linear combinations of the α_i which do not couple, and cannot lead to insertions. Moreover, they are not liftable by the familiar mechanisms⁹, thus in our instanton search in Section 6 such excess zero modes are forbidden even in relaxed scans.

Let us thus discuss a sector of fermion zero modes with equal number $n_\alpha = n_\gamma$. Considering a given 4d space-filling brane k , let us denote $L_{Mk}, L_{M'k'}, L_{Mk'}, L_{M'k}$ the (positive by definition) number of left-handed chiral fermion zero modes in the representations $(\square_M, \square_k), (\bar{\square}_M, \bar{\square}_k), (\square_M, \bar{\square}_k)$, and $(\bar{\square}_M, \square_k)$ respectively. The net number of chiral fermion zero modes in the $(\square_M, \bar{\square}_k)$ and (\square_M, \square_k) is given by $I_{Mk} = L_{Mk'} - L_{M'k'}$ and $I_{Mk'} = L_{Mk} - L_{M'k}$, respectively. This net number of fermions zero modes controls the violation of the $U(1)_a$ global charge by the instanton. Namely, such fermion zero modes in the Mk, Mp, Mk', Mp' sectors lead (if suitable couplings are present) to the insertion of 4d charged fields Φ_{kp} and/or $\Phi_{kp'}$.

⁹Note that such a mismatch is always correlated with the existence of extra chiral zero modes in the MM' sectors, discussed above. Denoting $\vec{Q}_a, \vec{Q}_{\text{orient}}$ the vector of RR charges of the 4d space-filling branes and orientifold planes, they satisfy the RR tadpole conditions $\sum_a N_a \vec{Q}_a + \sum_{a'} N_{a'} \vec{Q}_{a'} + \vec{Q}_{\text{orient}} = 0$. By taking the ‘intersection’ bilinear with the RR charges \vec{Q}_M of the brane instanton, we have $I_{Ma} + I_{Ma'} + I_{M,\text{orient}} = 0$. This implies that the number of fundamentals minus anti-fundamentals of the instanton gauge group is related to the number of two-index tensors.

The remaining fields in this sector are vector-like pairs, in the $(\square_M, \bar{\square}_k) + (\bar{\square}_M, \square_k)$ or the $(\square_M, \square_k) + (\bar{\square}_M, \bar{\square}_k)$, which in principle lead to the vanishing of the instanton amplitude, but which can be lifted by additional effects (motion in closed or open string moduli space, or addition of fluxes), in a way consistent with the gauge symmetries in 4d spacetime and on the instanton.

Just like for ‘real’ brane instantons, we conclude by commenting on our concrete instanton search in models with a set of visible branes **a**, **b**, **c**, **d** and a set of hidden branes h_i . The requirement that the instanton leads to an operator with specific charges under the visible branes fixes the values of the quantities $I_{M\mathbf{a}} - I_{M\mathbf{a}'}$, etc. As we described for real branes, one may still have fermion zero modes charged under the hidden sector branes, but they lead to additional insertions, hence we rather focus on instantons with $I_{Mh_i} - I_{Mh'_j} = 0$. The two kinds of conditions, on intersection numbers with visible and hidden branes, still leave the possibility of combinations of fermion zero modes of the kind $(\square_M, \bar{\square}_k) + (\bar{\square}_M, \square_k)$, which do not contribute to I_{Mk} , or of the kind $(\square_M, \square_k) + (\bar{\square}_M, \bar{\square}_k)$, which does not contribute to $I_{Mk'}$. Such combinations are automatically vector-like, and thus may be lifted in modifications of the theory. But the condition also allow combinations like $(\square_M, \bar{\square}_k) + (\square_M, \square_k)$, which exploit a cancellation between I_{Mk} and $I_{Mk'}$ (as also does $(\bar{\square}_M, \square_k) + (\bar{\square}_M, \bar{\square}_k)$). Such combinations are chiral by themselves, and in general imply a mismatch of modes in the \square_M and the $\bar{\square}_M$. The total mismatch can be arranged to vanish using combinations of the kind $(\square_M, \bar{\square}_k) + (\square_M, \square_k)$ and $(\bar{\square}_M, \square_p) + (\bar{\square}_M, \bar{\square}_p)$ for different branes. However, the only way to lift these pairs is by breaking the gauge symmetry on the 4d space-filling branes k and p . This can be done without damage to the visible sector if these are hidden branes, but this corresponds to the recombination of hidden branes that, as mentioned already, we are not going to consider. Hence only vector-like pairs with respect to *each* brane are considered to be liftable in simple modifications of the theory. In our instanton search, these are the only additional fermion zero modes which we allow in relaxed scans (but they are clearly not allowed for in restricted scans)

3 Instanton induced Majorana neutrino masses

In this Section we discuss the possible physical effects of D-brane instantons in string models with SM-like spectrum. In particular we describe the conditions to generate right-handed neutrino Majorana masses. We also comment on other possible B and/or L violating operators that can be generated by instantons. In this section we will again




Figure 1: Quarks and leptons at intersecting branes

use the geometrical language of IIA intersecting D-branes but it should be clear that our discussion equally applies to general CFT orientifolds like the ones presented in the next section.

3.1 The MSSM on the branes

Let us now specify the discussion in the previous section to the case of the generation of a right-handed neutrino mass term. In order to do that we need some realistic orientifold construction with the gauge group and fermion spectrum of the Standard Model (SM). In the context of Type II orientifolds perhaps the most economical brane configuration leading to a SM spectrum is the one first considered in [5]. This consists of four stacks, labelled **a**, **b**, **c**, **d**. The gauge factor on branes **a** is $U(3)$, and contains QCD and baryon number. The **d** factor is $U(1)_d$, and corresponds to lepton number. Stack **b** contains $SU(2)_{\text{Weak}}$ either embedded in $U(2)$ or $Sp(2)$. Finally brane **c** can either provide a $U(1)$ or an $O(2)$ factor. In the brane intersection language, the chiral fermions of the SM live at the intersections of these branes, as depicted in Fig. 3.1.

The $U(1)_Y$ factor of the standard model is embedded in the Chan-Paton factors of branes **a**, **c** and **d** as

$$Y = \frac{1}{6}Q_a - \frac{1}{2}Q_c - \frac{1}{2}Q_d = \frac{1}{2}(Q_{B-L} - Q_R) \quad (3.1)$$

where Q_x denotes the generator of the $U(1)$ of brane stack **x** (in case the Chan-Paton factor of brane **c** is $O(2)$ one should use the properly normalized $O(2)$ generator). Note that in this convention the Q_d generator appears with sign opposite to other conventions in the literature, e.g. in [3]. In addition to Y these models have two

additional $U(1)$ gauge symmetries:

$$\begin{aligned} Q_{\text{anom}} &= 3Q_{\mathbf{a}} + Q_{\mathbf{d}} = 9Q_B + Q_L \\ Y' &= \frac{1}{3}Q_{\mathbf{a}} + Q_{\mathbf{c}} - Q_{\mathbf{d}} = Q_{B-L} + Q_R \end{aligned} \quad (3.2)$$

The first is anomalous whereas the second, which we will call $B - L$ (with a slight abuse of language, since it is in fact a linear combination of $B - L$ and hypercharge), is anomaly free. In models in which the electroweak gauge group is embedded in $U(2)$, rather than in $Sp(2)$, there is a second anomalous $U(1)_{\mathbf{b}}$. The charges of the SM particles under these $U(1)$ symmetries are given in table 1.

Intersection	$D = 4$ fields/ zero modes		$Q_{\mathbf{a}}$	$Q_{\mathbf{c}}$	$Q_{\mathbf{d}}$	Y	Q_M
$(\mathbf{ab}), (\mathbf{ab}')$	Q_L	$3(3, 2)$	1	0	0	1/6	0
(\mathbf{ca})	U_R	$3(3, 1)$	-1	1	0	-2/3	0
$(\mathbf{c}'\mathbf{a})$	D_R	$3(3, 1)$	-1	-1	0	1/3	0
$(\mathbf{db}), (\mathbf{db}')$	L	$3(1, 2)$	0	0	1	-1/2	0
$(\mathbf{c}'\mathbf{d})$	E_R	$3(1, 1)$	0	-1	-1	1	0
(\mathbf{cd})	ν_R	$3(1, 1)$	0	1	-1	0	0
(\mathbf{Mc})	α_i	$2(0, 0)$	0	-1	0	1/2	1
(\mathbf{dM})	γ_i	$2(0, 0)$	0	0	1	-1/2	-1

Table 1: Standard model spectrum and $U(1)$ charges of particles and zero modes. Q_M stands for the world-volume gauge symmetry in the case of $U(1)$ complex instantons.

The $U(1)_k$ gauge symmetries have couplings with the RR 2-forms B_r of the model, as follows

$$S_{BF} = \sum_{k,r} N_k (p_{kr} - p_{k'r}) \int_{Ad} B_r \wedge F_k \quad (3.3)$$

where $p_{kr}, p_{k',r}$ are given by the RR charges of the D-branes. These imply that under a $U(1)_k$ gauge transformation $A_k \rightarrow A_k + d\Lambda_k$ the scalar a_r dual to B_r transforms as

$$a_r \rightarrow a_r + \sum_k N_k (p_{kr} - p_{k',r}) \Lambda_k \quad (3.4)$$

This has two effects: 1) The linear combination of axion fields $\sum_r (p_{kr} - p_{k',r}) a_r$ is eaten up by the $U(1)_k$ massless gauge boson, making it massive. 2) For anomalous $U(1)_k$, the anomalies cancel through a 4d version of the Green-Schwarz mechanism. This works due to the existence of appropriate $a_r F \wedge F$ couplings, involving the gauge fields in the theory.

It is obvious that all anomalous $U(1)$'s become massive by this mechanism. However it is important to realize [5] that gauge bosons of anomaly-free symmetries like $U(1)_{B-L}$ may also become massive by combining with a linear combination of axions. This is

interesting since it provides a mechanism to reduce the gauge symmetry of the model without needing explicit extra Higgsing. In the models in which $U(1)_{B-L}$ becomes massive in this way, the gauge group left over is purely that of the SM. Moreover, we will see that having (B-L) massive by this Stückelberg mechanism is crucial to allow the generation of instanton-induced Majorana neutrino masses.

Note that the $B \wedge F$ couplings may also be potentially dangerous, since in principle they could also exist for hypercharge, removing $U(1)_Y$ from the low-energy spectrum. As we will see in our RCFT examples later on, having massless $U(1)_Y$ but massive $U(1)_{B-L}$ turns out to be a strong constraint in model building.

3.2 Majorana mass term generation

As discussed in the previous section, string instantons can give rise to non-perturbative superpotentials breaking explicitly the perturbative global $U(1)$ symmetries left-over from $U(1)$ gauge bosons made massive through the Stückelberg mechanism. The kind of operator we are interested in has the form

$$W \simeq e^{-S_{ins}} \nu_R \nu_R \quad (3.5)$$

where ν_R is the right-handed neutrino superfield¹⁰. Here S_{ins} transforms under both $U(1)_{B-L}$ and $U(1)_R$ in such a way that the overall operator is gauge invariant. This operator may be created if the mixed open string sectors lead to fermionic zero modes α_i, γ_i , $i = 1, 2$, appropriately charged under the 4d gauge factors. As we discussed in the previous section, to generate a superpotential one needs instanton with $O(1)$ Chan-Paton symmetry, in order to lead to two uncharged fermion zero modes to saturate the $d^2\theta$ 4d superspace integration. On the other hand, as we argued, instantons with $Sp(2)$ or $U(1)$ CP symmetries may also induce the required superpotentials if there is some additional dynamics getting rid of the extra uncharged zero modes which in principle appear in instantons with these symmetries. We thus consider all $O(1)$, $Sp(2)$ and $U(1)$ instantons in our discussion.

In order to get a ν_R bilinear, the intersection numbers of instanton M and d, c branes are as follows

$$Sp(2) \text{ case :} \quad I_{M\mathbf{c}} = 1 ; I_{M\mathbf{d}} = -1 \quad (3.6)$$

(since there is an extra multiplicity from the two branes required to produce $Sp(2)$)

$$O(1) \text{ case :} \quad I_{M\mathbf{c}} = 2 ; I_{M\mathbf{d}} = -2 \quad (3.7)$$

¹⁰ Actually we denote by ν_R the left-handed ν_L^c field following the usual (a bit confusing) convention.




Figure 2: Disk amplitude coupling two charged zero modes to ν_R in the geometrical Type IIA intersecting brane approach.

$$U(1) \text{ case : } \quad I_{M\mathbf{c}} = 2 ; I_{M\mathbf{d}} = -2 \text{ or } I_{M\mathbf{d}'} = 2 ; I_{M\mathbf{c}'} = -2 \quad (3.8)$$

Furthermore there must be cubic couplings involving the right-handed neutrino superfield ν^a in the a^{th} family and the fermionic zero modes α_i, γ_j

$$L_{cubic} \propto d_a^{ij} (\alpha_i \nu^a \gamma_j) , a = 1, 2, 3 \quad (3.9)$$

In type IIA geometric compactifications, this coupling arises from open string disk instantons, see Fig. 3.1. In general type IIA models (resp. IIB models), the coefficients d_a^{ij} depend on the Kähler (resp. complex structure) moduli, and possibly on open string moduli. In simple CFT models (like e.g. in toroidal cases) these quantities may be in principle explicitly computed.

These trilinear couplings appear in the instanton action and after integration of the fermionic zero modes α_i, γ_i one gets a superpotential coupling proportional to

$$\int d^2\theta \int d^2\alpha d^2\gamma e^{-d_a^{ij} (\alpha_i \nu^a \gamma_j)} = \int d^2\theta \nu_a \nu_b (\epsilon_{ij} \epsilon_{kl} d_a^{ik} d_b^{jl}) \quad (3.10)$$

yielding a right-handed neutrino mass term. This term is multiplied by the exponential of the instanton euclidean action so that the final result for the right-handed neutrino mass (up to a 1-loop prefactor) has the form

$$M_{ab}^R = M_s (\epsilon_{ij} \epsilon_{kl} d_a^{ik} d_b^{jl}) \exp\left(-\frac{V_{\Pi_M}}{g_s} + i \sum_r q_{M,r} a_r\right) \quad (3.11)$$

For geometric compactifications V_{Π_M} is roughly related to the wrapped volume. We keep the same notation to emphasize that the effect is non-perturbative in g_s . In supersymmetric models the term in the exponential is the linear combination U of complex structure moduli to which the instanton D-brane couples, as described in the previous section. As explained, the gauge $U(1)_{\mathbf{c}}, U(1)_{\mathbf{d}}$ transformation of the bilinear

piece and the $e^{-S_{D2}}$ factor nicely cancel. Note that from the viewpoint of the 4d SM effective field theory, the instanton has generated a Majorana neutrino mass violating $B - L$. Notice also that since this symmetry is non-anomalous, its violation cannot be associated to a gauge instanton, hence this is a pure string theory instanton effect.

3.3 Flavor and the special case of $Sp(2)$ instantons

In order to extract more specific results for the flavor structure of the obtained Majorana mass operator, one needs to know more details about the quantities d_a^{ij} coming from the disk correlators. However in the particular case of $Sp(2)$ instantons, the labels i, j are $Sp(2)$ doublet indices, and the symmetry requires $d_a^{ij} = d_a \epsilon^{ij}$. The mass matrix for the three neutrinos is given by $M_{ab}^R = 2M_s d_a d_b \exp(-U)$, so that the flavour dependence on $a, b = 1, 2, 3$ factorizes. More generally, as we will see in our RCFT search in Section 6, there are typically several different instantons contributing to the amplitude, so that we actually have a result for the mass

$$M_{ab}^R = 2M_s \sum_r d_a^{(r)} d_b^{(r)} e^{-U_r} \quad (3.12)$$

where the sum goes over the different contributing instantons. One thus has a structure of the form

$$M^R \sim \sum_r e^{-U_r} \text{diag}(d_1^{(r)}, d_2^{(r)}, d_3^{(r)}) \cdot \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix} \cdot \text{diag}(d_1^{(r)}, d_2^{(r)}, d_3^{(r)}) \quad (3.13)$$

This structure is very interesting. Indeed, each instanton makes one particular (instanton-dependent) linear combination of the neutrinos massive, leaving two linear combinations massless. Hence, for three or more instantons, one generically has a matrix with three non-zero eigenvalues. It is easy to imagine a hierarchical structure among the three eigenvalues if e.g. the exponential suppression factors $\exp(-\text{Re } U_r)$ are different for each instanton.

3.4 Other $B-$ and $L-$ violating operators

Our main focus in this paper is on the generation of right-handed neutrino Majorana masses. However instantons may induce other $L-$ and $B-$ violating operators which we briefly summarize in this subsection.

3.4.1 The Weinberg operator

A right-handed neutrino Majorana mass term is not the only possible operator violating lepton number. Instanton effects may also give rise to dimension 5 operators not involving ν_R . Specifically, the Weinberg operator

$$\mathcal{L}_W = \frac{\lambda}{M} (L\bar{H}L\bar{H}) . \quad (3.14)$$

might be generated. Once Higgs fields get a vev v this operator gives rise directly to Majorana masses for the left-handed neutrinos of order $\simeq v^2/M$. Indeed, it is easy to check that in this case the required instanton M must verify

$$Sp(2) \text{ case :} \quad I_{M\mathbf{c}} = -1 ; I_{M\mathbf{d}} = 1 \quad (3.15)$$

$$O(1) \text{ case :} \quad I_{M\mathbf{c}} = -2 ; I_{M\mathbf{d}} = 2 \quad (3.16)$$

$$U(1) \text{ case :} \quad I_{M\mathbf{c}} = -2 ; I_{M\mathbf{d}} = 2 \text{ or } I_{M\mathbf{c}'} = 2 ; I_{M\mathbf{d}'} = -2 \quad (3.17)$$

(here we are assuming $SU(2)_{weak}$ to be embedded in an $Sp(2)$). Note that these intersection numbers are different to those giving rise to ν_R mass terms. In particular they lead to a transformation under $B - L$ opposite to that of ν_R mass operators¹¹. In the present case there are altogether four fermionic zero modes α_i, γ_i corresponding to the intersections of the instanton M with the branes \mathbf{c}, \mathbf{d} . These zero modes can have couplings involving the left-handed leptons L and the u-type Higgs multiplet \bar{H}

$$L_{disk} \propto c_a^{ij} (\alpha_i (L^a \bar{H}) \gamma_j) . \quad (3.18)$$

Upon integration over the fermionic zero modes one recovers the Weinberg operator. In the present case the scale M of the Weinberg operator will be the string scale M_s and the coupling $\lambda \simeq \exp(-S_{ins})$. Again, in the particular case of $Sp(2)$ instantons the situation simplifies ($c_a^{ij} = c^a \epsilon^{ij}$) and one gets left-handed neutrino Majorana masses

$$M_{ab}^L = \frac{\langle \bar{H} \rangle^2}{M_s} \sum_r 2c_a^{(r)} c_b^{(r)} e^{-S_r} \quad (3.19)$$

where r runs over the different contributing instantons and S_r is their corresponding action. The flavour structure of this left-handed neutrino mass matrix is the same as in eq.(3.13) and again may potentially lead to a hierarchical structure of left-handed neutrino masses, as is experimentally observed.

¹¹Instantons with these intersection numbers will be denoted with a plus sign in the instanton search later on

In a given model both this kind of instanton and the one giving rise to right-handed neutrino masses (which is different) may be present. This contribution to the left-handed neutrino Majorana mass is in principle sub-leading compared to the see-saw contribution

$$M_{ab}^L(\text{see-saw}) = \frac{\langle \overline{H} \rangle^2}{2M_s} h_D^T \left(\sum_r d_a^{(r)} d_b^{(r)} e^{-S_r} \right)^{-1} h_D \quad (3.20)$$

where is the ordinary Yukawa coupling constant $h_D^{ab}(\nu_R^a \overline{H} L^b)$. In principle the former is doubly suppressed both by $1/M_s$ and the exponential factor. On the other hand if the exponential suppression is not too large this mechanism involving directly the Weinberg operator may be the most relevant source of neutrino masses. This is because the see-saw contribution coming from ν_R exchange is proportional to the square of the ordinary Yukawa couplings h_D^{ab} which could be small. One could even think of having just the Weinberg operator as the unique source of the observed left-handed neutrino masses. Note however that in string vacua like this, in which the ν_R 's are present and massless at the perturbative level, having just the Weinberg operator would not be phenomenologically correct, and instantons of the first class are still needed so that the ν_R 's get a sufficiently large mass.

3.4.2 R-parity violating operators

In the case of $N = 1$ SUSY models like the MSSM there might be operators of dimension 3 and 4 violating lepton and/or baryon number. These are the superpotential couplings

$$W_{R_p} = \mu_a^L L^a \overline{H} + \lambda_{abc} Q^a D^b L^c + \lambda'_{abc} U^a D^b D^c + \lambda''_{abc} L^a L^b E^c \quad (3.21)$$

in standard notation. Unlike the neutrino operators mentioned above, these operators violate $B - L$ in one unit (rather than 2). It is well known that the standard R-parity of the MSSM may be identified with a Z_2 subgroup of $U(1)_{B-L}$, so these terms are odd under R-parity. The simultaneous presence of all these couplings is phenomenologically unacceptable. Indeed, the third coupling violates baryon number, and the other three violate lepton number. Together they lead to proton decay at an unacceptably large rate. On the other hand couplings violating *either* B *or* L are phenomenologically allowed.

It is an interesting question whether any of these operators may be induced by string instanton effects. A first point to note is that instantons with $Sp(2)$ Chan-Paton symmetry can never generate operators of this type. The reason is that all charged zero modes will necessarily come in $Sp(2)$ doublets and hence the charged operators induced will always involve an even number of charged $D = 4$ fields and R-parity is

automatically preserved. On the other hand $O(1)$ and $U(1)$ instantons may generate R-parity violating operators. In particular, the LH bilinear is essentially the square root of the Weinberg operator, and may be induced if a $U(1)$ or $O(1)$ instanton M exists with

$$I_{M\mathbf{c}} = -1 ; I_{M\mathbf{d}} = 1 \quad \text{or} \quad I_{M\mathbf{c}'} = 1 ; I_{M\mathbf{d}'} = -1 . \quad (3.22)$$

(in the $O(1)$ case the second option is not independent from the first). Again, if the appropriate disk couplings are non-vanishing a term with $\mu_L^a \sim M_s \exp(-S_{ins})$ is generated. The rest of the operators in W_{R_p} may also be generated. Possible instanton zero modes which may induce them are shown in table 2. For example, the QDL operator may be induced if a $U(1)$ instanton M with intersection numbers

$$I_{M\mathbf{b}} = -1 ; I_{M\mathbf{c}'} = 1 ; I_{M\mathbf{d}} = 1 \quad (3.23)$$

is present and in addition couplings

$$L_{disk} \propto c_{ab} (\alpha(U^a Q_j^b) \gamma^j) + c'_a (\beta L_j^a \gamma_j) \quad (3.24)$$

exist. Here α, β, γ are zero modes corresponding to $(M\mathbf{c}')$, $(M\mathbf{d})$ and (bM) intersections and $a, b(j)$ are flavor($SU(2)_L$) indices. Analogous trilinear or quartic disk amplitudes involving two charged zero modes should exist to generate the rest of the R-parity violating amplitudes in table 2.

$D = 4$ Operator	$I_{M\mathbf{a}}$	$I_{M\mathbf{a}'}$	$I_{M\mathbf{b}}$	$I_{M\mathbf{c}}$	$I_{M\mathbf{c}'}$	$I_{M\mathbf{d}}$	$I_{M\mathbf{d}'}$
$\nu_R \nu_R$	0	0	0	2	0	-2	0
$LHLH$	0	0	0	-2	0	2	0
LH	0	0	0	-1	0	1	0
QDL	0	0	-1	0	1	1	0
UDD	-1	0	0	1	2	0	0
LLE	0	0	-1	0	1	1	0
$QQQL$	1	0	-2	0	0	1	0
$UUDE$	-1	0	0	2	2	-1	0

Table 2: Zero modes required to generate Lepton/Baryon-number violating superpotential operators. $Sp(2)$ instantons cannot give rise to R-parity violating operators whereas $O(1), U(1)$ instantons may in principle contribute to all of them. In the case of $U(1)$ instantons there are additional zero mode possibilities which are obtained by exchanging $I_{M\mathbf{x}} \leftrightarrow -I_{M\mathbf{x}'}$.

3.4.3 Dimension 5 proton decay operators

There are also superpotential dimension-5 operators violating B and L which may be constructed from the MSSM matter superfields. Indeed the dimension 5 operators

$$\left(\frac{1}{M}\right)QQQL \quad ; \quad \left(\frac{1}{M}\right)UUDE \quad (3.25)$$

are in fact the leading source of proton decay in SUSY GUT models with R-parity. Unlike the other operators considered here these ones preserve $B - L$ (hence R-parity) but not $B + L$. These operators do not contribute directly to a proton decay but need to be 'dressed' by a one loop exchange of some fermionic SUSY particle. This makes that, even although they are suppressed only by one power of the relevant fundamental scale, the loop factor and the corresponding couplings make the overall rate in SUSY-GUTS (barely) consistent with present experimental bounds for M of order the GUT scale or larger.

These dimension 5 operators may also be induced in D-brane models of the class here considered by the presence of instantons with appropriate intersection numbers. For instance, the first operator may be induced through $O(1)$ or $U(1)$ instantons M with

$$I_{M\mathbf{b}} = I_{M\mathbf{b}'} = -2 \quad ; \quad I_{M\mathbf{a}} = 1 \quad ; \quad I_{M\mathbf{d}} = 1 \quad (3.26)$$

Again $Sp(2)$ instantons cannot induced this operator, since the the Ma intersection would yield 6 (rather than 3) colored fermionic zero modes. The proton decay rate obtained from these operators depend on the ratio $\exp(-S_{\text{ins}}) \times 1/M_s$. For M_s of order 10^{16} GeV, the rate is consistent with present bounds if $\exp(-S_{\text{ins}})$ provides a suppression of a few orders of magnitude. On the other hand, models with a low string scale may be in danger unless the exponential suppression is sufficiently large (or such particular instantons are absent).

As a general conclusion, these phenomenological aspects of instanton induced operators very much depend on the action of the instanton, e.g. the volume of the wrapped D2-instanton in the intersecting D-brane constructions. In any event it is clear that the instantons here considered may indeed induce proton decay at a model-dependent rate. However in certain models R-parity will be preserved and prevent too rapid proton decay. Indeed, this is what we find in our instanton search in Gepner orientifolds. As we said $Sp(2)$ instantons automatically preserve R-parity. More generally, models that violate R-parity are rare, and the corresponding instantons actually generate very high dimensional operators, so R-parity breaking effects seems quite suppressed. In

fact in our search within MSSM-like models in Gepner model orientifolds we do not find instantons with just the correct charged zero modes to generate the low dimensional couplings discussed above. So, at least within our class of RCFT constructions, R-parity preservation is quite a common feature.

4 CFT orientifolds

In this section we describe the 4d string models we consider, namely orientifolds of type IIB Gepner model compactifications. This is a very large class, on which one can carry out large scans for certain desired properties. And moreover at present the only known class of (SUSY) models with massive $B - L$.

4.1 Construction of the models

In general, RCFT orientifolds are orientifold projections of closed string theories constructed using rational conformal field theory. Although this includes in principle rational tori and orbifolds, the real interest lies in cases where the two-dimensional CFT is interacting, because such theories are hard to access by other methods. A disadvantage of the use of RCFT is that this method is algebraic, and not geometric in nature, so that one cannot easily explore small deformations of a certain string theory. It is best thought of as a rational scan of moduli spaces.

The most easily accessible examples are the orientifolds of tensor products of minimal $N = 2$ conformal field theories (“Gepner models”) forming a type IIB closed string theory. During the last decade, examples in this class have been studied by many authors (see [35][36][37][38][39][40][41][42]), and searched systematically in [6] and [7]. Although the Gepner models form only a small subset of RCFT’s, they already offer a large number of possibilities. The total number of tensor products with the required central charge $c = 9$ is 168. On top of this, one can choose a large number of distinct modular invariant partition functions on the torus. The orientifold formalism is not available for all of them, but it has been completely worked out [43] for all simple current invariants (based on the charge conjugation invariant). This yields a total of 5403 distinct MIPFs. On top of this, we may choose various orientifold projections. Here the only known possibilities are a class of simple-current based choices [44][45][46][47]. This then yields a total of 49304 orientifolds.

For each orientifold choice, the full open string partition function is

$$\frac{1}{2} \left(\sum_{a,b,i} N_a N_b A_{ab}^i \chi_i \left(\frac{\tau}{2} \right) + \sum_{a,i} N_a M_a^i \hat{\chi}_i \left(\frac{\tau}{2} + \frac{1}{2} \right) \right) \quad (4.1)$$

Here A_{ab}^i are the annulus coefficients, M_a^i the Moebius coefficients, N_a the Chan-Paton multiplicities and $\chi(\tau)$ are the closed string characters, and $\hat{\chi}_i(\tau) = T^{-1/2} \chi_i(\tau)$. The set of integers i is simply the set of primary fields of the closed string CFT, and depends only on the tensor product. The integers a, b are the boundary labels; this set depends on the MIPF. Our notation and labelling conventions for these CFT quantities are explained in Appendix A. The integers A_{ab}^i and M_a^i depend in addition also on the orientifold choice; in the case of A_{ab}^i the latter dependence is very simple: all distinct annuli can be written as $A_{ab}^{\Omega,i} = \sum_c A_a^i {}^c C_{cb}^{\Omega}$, where Ω is the orientifold choice (which we usually do not specify explicitly) and C_{cb}^{Ω} is the boundary conjugation matrix, which acts as an involution on the set of boundaries.

Suppressing some details (which can be found in [43]) we may write these integers as

$$A_{ab}^{\Omega,i} = \sum_{m,J,K} \frac{S_m^i R_{a,(m,J)} g_{JK}^{\Omega,m} R_{b,(m,K)}}{S_{0m}} \quad (4.2)$$

$$M_a^{\Omega,i} = \sum_{m,J,K} \frac{P_m^i R_{a,(m,J)} g_{JK}^{\Omega,m} U_{(m,K)}^{\Omega}}{S_{0m}} \quad (4.3)$$

Here m is the label of an Ishibashi-state (the set of states that propagates in the transverse (or closed string) channel of the the annulus or Moebius diagrams). It is a subset of the set of closed string labels i , but in general there are degeneracies, so that more than one distinct Ishibashi state belongs to a given closed string label. These degeneracies are distinguished by the labels J, K (see Appendix A). The complex numbers R and U are respectively the boundary and crosscap coefficients. Note that the latter depend on the orientifold choice, but the former do not. The only dependence of the annulus coefficients on the orientifold choice is through the Ishibashi metric g_{JK}^{Ω} , which is a matrix on each Ishibashi degeneracy space, and which can be a sign if there are no degeneracies. Finally, the matrix P is given by $P = \sqrt{T} S T^2 S \sqrt{T}$, where S and T are the generators of the modular group of the torus. Similar expressions exist for the Klein bottle multiplicities defining the unoriented closed sector, but they will not be needed in this paper.

The boundary labels a, b, \dots refer to all boundaries that respect the bulk symmetries of the CFT. This includes the individual $N = 2$ chiral algebras of the factors in the tensor product, the *alignment currents*¹² that ensure the proper definition of world-sheet supersymmetry and the space-time supersymmetry generator that imposes a generalized GSO-projection on the spectrum. The latter implies that all characters χ_i respect (at least) $N = 1$ space-time supersymmetry. By construction, the boundary states are then supersymmetric as well. Both conditions (boundary and bulk space-time supersymmetry) can in principle be relaxed within the formalism, but this leads to a much larger set of bulk and boundary states. The precise labelling of the boundaries is explained in Appendix A and involves a subset of the closed string labels i and a degeneracy label, distinct from the one used for the Ishibashi states. The set of boundary labels is complete in the sense of [45]. This means that no additional boundary states exist that respect all the aforementioned symmetries. It also means that the matrices R are square matrices (although their rows and columns are defined in terms of different index sets). It is in principle possible to write down additional boundary states that break some of the world-sheet symmetries. This is an important possibility to keep in mind, but we will not consider it here.

The massless spectrum is obtained by restricting the characters χ_i to massless states. Since the characters are supersymmetric those massless states are either vector multiplets or chiral multiplets. The latter can be restricted to one chirality (*e.g.* left-handed); the other choice merely produces the CPT conjugates. Boundaries are called real if $a = a'$, where the conjugate boundary a' is defined by $C_{a,a'}^\Omega = 1$, and complex otherwise. The Chan-Paton multiplicities N_a give rise to gauge groups $U(N_a)$ for complex boundaries and $SO(N_a)$ or $Sp(N_a)$ for real ones. In the latter case N_a must be even. To count bi-fundamentals we define

$$L_{ab} \equiv \sum_i A_{ab}^i \chi_i\left(\frac{\tau}{2}\right)_{\text{massless}, L} . \quad (4.4)$$

Note that because of the factor $\frac{1}{2}$ in (4.1) and the fact that L_{ab} is symmetric, the value of L_{ab} is indeed precisely the number of bi-fundamentals in the representation (N_a, N_b) .

¹²These are spin-3 currents consisting of products of the world-sheet supercurrents of the factors in the tensor product, including the NSR space-time factor.

It is convenient to introduce the intersection matrix¹³

$$I_{ab} \equiv L_{ab'} - L_{a'b} , \quad (4.5)$$

which is manifestly antisymmetric in a and b . Note that for a pair of complex boundaries a, b with conjugates a', b' one can define four quantities that are relevant for the massless spectrum, two of which are chiral, namely I_{ab} and $I_{ab'}$.

It is often convenient to associate a geometric picture to these integers. Thus we will often refer to the boundary labels and their multiplicities as “stacks of branes”, and view the integers I_{ab} as brane intersection numbers. This is only done for convenience and does not imply a concrete brane realization; indeed, it does not make sense to say that a given boundary label corresponds to a Dp -brane for some give p . Such an interpretation might be valid in a large radius limit, assuming such a limit exists.

In general, for a choice of Chan-Paton multiplicities N_a there will be tadpoles in the one-point closed string amplitudes on the disk and the crosscap. These have to be cancelled in order to make the theory consistent (since we work with supersymmetric strings we do not have the option of cancelling RR and NS-NS tadpoles separately). This leads to a condition on the Chan-Paton multiplicities:

$$\sum_a N_a R_{a,(m,J)} = 4\eta_m U_{m,J} \quad (4.6)$$

where $\eta_0 = 1$ and all other η 's are -1 ; there is such a condition for any Ishibashi label (m, J) that leads to a massless scalar in the transverse channel. The one for $m = 0$ (which is non-degenerate) is the dilaton tadpole condition. It has the special feature that all coefficients R_{a0} are real and positive. The crosscap coefficient U_0 is also real and can be chosen positive (in the CFT both signs are acceptable). If $U_0 \neq 0$ (4.6) limits the Chan-Paton multiplicities; if $U_0 = 0$ the only solution is $N_a = 0$ for all a , which rules out any realization of the Standard Model. This reduces the number of usable orientifolds to 33012.

Tadpole cancellation condition implies cancellation of RR-charges coupling to long-range fields, and absence of local anomalies. There is a second condition that has

¹³Note that L_{ab} is a symmetric matrix giving the number of chiral multiplets in the (\square_a, \square_b) bi-fundamental. This is a natural quantity in unoriented CFT's, where a symmetric definition for the annulus amplitude exists. In oriented CFT the annulus is, in general, not symmetric, but on the other hand it is possible to choose the branes in such a way that only $(\square, \bar{\square})$ bi-fundamentals appear. This has become the customary way of counting states in the intersecting brane literature, even for orientifold models. The quantity I_{ab} is defined in such a way that it is anti-symmetric in a and b . This is why boundary conjugations appear in the right hand side. This has the additional advantage of making I a more familiar quantity for readers used to the standard intersection brane conventions.

to be taken into account, which has to do with \mathbf{Z}_2 charges that do not couple to long-range fields, usually referred to as “K-theory charges” in geometric constructions. Uncancelled K-theory charges may lead to global anomalies in symplectic factors of the gauge group. But even if this symptom is absent, the disease may still exist. A much more general way to probe for uncancelled K-theory charges is to require the absence of global anomalies not only in the Chan-Paton gauge group but also on all symplectic brane-anti-brane pairs that can be added to it as “probe-branes” [48]. Presently this is the most general constraint that be imposed in these models, but it is not known if additional ones are required. This probe brane constraint leads to a large number of mod-2 constraint and is potentially very restrictive, but almost harmless in practice [49]. It is satisfied by all models we consider in the present paper.

4.2 Search for SM-like models

The complete set of solutions to these conditions is finite but huge, but the vast majority is of no phenomenological interest. In the last few years systematic searches have been carried out for models that contain the Standard Model. The models that were considered have the property that the set of Chan-Paton labels can be split into two subsets, the observable and the hidden sector. The former has been limited, for practical reasons, to at most four complex brane stacks, required to contain the Standard Model gauge group and the right intersections to yield three families of quarks and leptons, plus (in general) some non-chiral (vector-like) additional matter. The hidden sector is only constrained by the requirement that there be no net number of chiral multiplets charged under both the observable and hidden sector, and by practical computational limitations. The main purpose of the hidden sector in these models is to provide variables that can be used to satisfy the tadpole and global anomaly conditions, since the multiplicities in the observable sector are already fixed. In some cases the observable sector already satisfies the constraints by itself, and there is no hidden sector.

The observable sector can be realized in many different ways if one only imposes the constraint that the standard model should be contained in it. These possibilities were recently explored in [7]. We will focus on the realization described in Section 3.1, first considered in [5]. There are four stacks, namely **a** (containing QCD and baryon number as $U(3)$), **b** (containing electroweak $SU(2)$ embedded as $U(2)$ or $Sp(2)$), **c** (providing a $U(1)$ or an $O(2)$ factor¹⁴, and **d** (providing another $U(1)$ factor).

¹⁴In [6] also $Sp(2)$ was considered, but this requires an additional Higgs mechanism.

The standard model hypercharge generator is , defined in (3.1):

$$Y = \frac{1}{6}Q_{\mathbf{a}} - \frac{1}{2}Q_{\mathbf{c}} - \frac{1}{2}Q_{\mathbf{d}} \quad (4.7)$$

where $Q_{\mathbf{x}}$ denotes the generator of the $U(1)$ of brane stack \mathbf{x} ; in case the Chan-Paton factor of brane \mathbf{c} is $O(2)$ one should use the properly normalized $O(2)$ generator. In addition to Y these models have two or three additional $U(1)$ gauge symmetries (the latter case if electroweak $SU(2)$ arises from $U(2)$). These (except the combination $B - L$) are anomalous, with anomaly cancelled by the Green-Schwarz mechanism, implying the existence of a $B \wedge F$ coupling making them massive. In fact, as already mentioned, such Stückelberg couplings may be present for non-anomalous $U(1)$'s as well. We are interested in models where the hypercharge gauge boson does not have such couplings (otherwise the model would be phenomenologically unacceptable), but where the $B - L$ gauge boson is massive by such couplings (both in order that the gauge group reduces to the SM one, and that neutrino Majorana masses may be induced by string instantons, as discussed in previous sections).

The combined requirements of having a massive $B - L$ and a massless Y turn out to be difficult to satisfy. In fact, if the group on brane \mathbf{c} is $O(2)$ they are impossible to satisfy simultaneously, because the $O(2)$ component of the vector boson does not couple to any axions, and hence the $B - L$ and Y bosons have the same mass. But even in models with a $U(1)$ group on brane \mathbf{c} it happens rather rarely that both constraints are satisfied simultaneously, at least in the searches that have been done so far.

We will make use here of the data presented in [6, 7], which are available in slightly improved form on the website www.nikhef.nl/~t58/filtersols.php. This database consist of 211634 distinct spectra. Here “distinct” means that they are physically different for a given MIPF¹⁵ if the hidden sector is ignored. Hence the differences can be the number of vector-like states of various kinds or the dilaton couplings of branes \mathbf{a} , \mathbf{b} , \mathbf{c} , \mathbf{d} . Geometrically, these spectra may originate from the same moduli space, but then in any case from different points on this moduli space. The improvements in comparison with the data presented in [6] consist of taking into account the full global anomaly conditions from probe branes. In some cases this required nothing more than checking these conditions for an existing solution of the tadpole conditions, but in other cases a new solution had to be found. As a result, a few models disappeared from the original database, but due to improved algorithms a few new ones could be added. The net re-

¹⁵Rare cases of identical spectra and couplings originating from different MIPFs are treated as distinct.

sult is some small but inconsequential changes in the total number of models of various kinds. The numbers we will mention below are based on the improved database.

The total number of models in that database with a Chan-Paton group $U(3) \times Sp(2) \times U(1) \times U(1)$ is 10587. Of these, 391 (about 4%) have a massive $B - L$ vector boson. For $U(3) \times U(2) \times U(1) \times U(1)$ these numbers are, respectively, 51 and 0. Hence no examples of the latter type were found, although they were found with 1,2 and 4 families (in a limited search), in a few percent of the total number of models. It seems therefore reasonable to expect that $U(3) \times U(2) \times U(1) \times U(1)$ with massive $B - L$ do exist, and that their absence is just a matter of statistics. Just for comparison, the total number of $U(3) \times Sp(2) \times O(2) \times U(1)$ models is 56627.

5 Fermion zero modes for instantons on RCFT's

In this section we discuss D-brane instantons for general compactifications, including abstract CFT ones. We also provide the spectrum of zero modes on an instanton brane, using the information about their internal structure i.e. in the compactified dimension in geometric models, or of the internal CFT in more abstract setups like in previous section. We will be interested in the latter case.

A first question that should be addressed is what this internal structure is. For instance, in type IIA geometric compactifications, it corresponds to a supersymmetric (i.e. special lagrangian) 3-cycle. Notice that these are the same kind of 3-cycles already used to wrap the D6-branes that give rise to the 4d gauge symmetry of such models. For general CFT's, D-branes are described as boundary states. To describe instantons, one can simply use the same boundary state of the internal CFT to describe the 4d space-filling branes present in the model and the instanton branes. The only difference is that boundaries satisfy Neumann conditions in the 4d space-filling case, and Dirichlet in the instanton case. This exploits the fact that whenever a boundary state of the internal CFT, and with Neumann boundary conditions in the 4d space is an acceptable state of the full CFT, the same boundary state of the internal CFT, combined with Dirichlet boundary conditions in the 4d space also gives an acceptable state of the full CFT. For geometric compactifications this is related to Bott periodicity of the K-theory classes associated to the D-brane charges, but it is possible to show it in general.

Since instanton D-branes can thus be naturally associated to the boundary states of 4d space-filling branes, it is convenient to express the spectrum of zero modes of the former in terms of the massless states of the latter. This is particularly useful, since

the computation of the spectra on 4d space-filling branes for Gepner model orientifolds has already been described (although the arguments below are valid also for geometric compactifications). Hence, let us denote by \mathcal{M} a 4d space-filling brane associated with the same boundary state of the internal CFT as the instanton brane M of interest. Note that the 4d space-filling brane \mathcal{M} is an auxiliary tool, and need not be (and, for our instantons of interest, will not be) one of the 4d space-filling branes present in the model.

‘Real’ brane instantons

Let us first consider the case of ‘real’ brane instantons. Consider a set of m 4d space-filling branes \mathcal{M} , and focus first on the massless spectrum in the $\mathcal{M}\mathcal{M}$ sector. Before the orientifold projection, it leads to a universal 4d $N = 1$ $U(m)$ vector multiplet, and a number $L_{\mathcal{M}\mathcal{M}}$ of adjoint chiral multiplets. The orientifold operation maps this sector to itself, acting on the Chan-Paton with a matrix $\gamma_{\Omega, \mathcal{M}}$. This matrix satisfies

$$\gamma_{\Omega, \mathcal{M}}^T \gamma_{\Omega, \mathcal{M}}^{-1} = \pm \mathbf{1}_m \quad (5.1)$$

The two possibilities can be chosen to correspond to $\gamma_{\Omega, \mathcal{M}} = \mathbf{1}_m$ or $\gamma_{\Omega, \mathcal{M}} = \epsilon_m$, with $\epsilon_m = \begin{pmatrix} 0 & \mathbf{1}_r \\ -\mathbf{1}_r & 0 \end{pmatrix}$, and $m = 2r$ hence necessarily even in the latter case. They correspond to the SO and Sp projections, respectively.

The orientifold projection on the $N = 1$ vector multiplet Chan-Paton matrices is given by

$$\lambda = -\gamma_{\Omega, \mathcal{M}} \lambda^T \gamma_{\Omega, \mathcal{M}}^{-1} \quad (5.2)$$

and leads to $SO(m)$ or $Sp(m)$ vector multiplets for the SO or Sp projection (hence the name). Concerning the $N = 1$ chiral multiplets, they fall in two classes of p_- , p_+ (with $p_- + p_+ = L_{\mathcal{M}\mathcal{M}}$) which suffer the projections

$$\lambda = \pm \gamma_{\Omega, \mathcal{M}} \lambda^T \gamma_{\Omega, \mathcal{M}}^{-1} \quad (5.3)$$

For the SO projection, this leads to p_+ , p_- chiral multiplets in the \square, \square representation. For the Sp projection, there are p_+ , p_- chiral multiplets in the \square, \square representation.

The sectors $\mathcal{M}a$ (where a is a 4d space-filling branes present in the model) are mapped to sectors $\mathcal{M}a'$, so it is enough to focus on the former. After the orientifold projection one gets $L_{\mathcal{M}a}$, $L_{\mathcal{M}a'}$ chiral multiplets in the $(\square_{\mathcal{M}}, \square_a)$, $(\square_{\mathcal{M}}, \square_a)$.

Let us now obtain the zero modes for a set of m instanton branes M in terms of the above spectrum. The MM sector is closely related to the $\mathcal{M}\mathcal{M}$ sector, by changing the NN boundary conditions in 4d spacetime to DD boundary conditions

(which can be done in a covariant formalism, but not in the light-cone gauge). Before the orientifold projection, one obtains the same set of states (since moddings for NN and DD boundary conditions are identical, both in the NS and R sector), but with different world-volume interpretation. Also, the change in boundary conditions implies that some polarization states which are unphysical for the 4d spacefilling brane are physical in the instanton brane. Hence, the $U(m)$ gauge bosons on the 4d space-filling brane \mathcal{M} correspond to four adjoint real scalars in the instanton brane M . Similarly, the 4d spinors in \mathcal{M} , correspond to four fermion zero modes on M , transforming as two spinors of opposite chiralities $\theta^\alpha, \tilde{\theta}_{\dot{\alpha}}$ of the $SO(4)$ rotation group in transverse space. The orientifold projection maps the MM sector to itself, acting on Chan-Paton indices with a matrix $\gamma_{\Omega,M}$. In close analogy with the argument in [50] for the familiar D5-D9-brane system in type I (see [51, 52] for related derivations), one can show that the condition (5.1) flips sign upon changing four NN boundary conditions to DD, hence

$$\gamma_{\Omega,M}^T \gamma_{\Omega,M}^{-1} = \mp \mathbf{1}_m \quad (5.4)$$

Namely, the instanton brane has $Sp(m)$ gauge group when the 4d space-filling brane (with same internal boundary state) has gauge group $O(m)$, and vice-versa. We still refer to these projections as SO and Sp , hoping no confusion arises. Note that, as mentioned in Section 2.2, although there are no gauge bosons in $0+0$ dimensions, the gauge group is present on the instantons in that it acts on open string endpoints.

Let us consider the effect of the orientifold projection on the MM states, as compared with the effect on $\mathcal{M}\mathcal{M}$ states. Again, following arguments familiar in the D5-D9 brane system in type I, one can show that the signs in conditions like (5.2), (5.3) remain unchanged upon changing four NN dimensions to DD, except for bosonic modes polarized along the directions longitudinal to these four dimensions (and for fermions related to them by the unbroken susy of the total system). To be concrete, considering the four MM adjoint bosons, and two MM adjoint fermions θ^α associated to the universal $\mathcal{M}\mathcal{M}$ vector multiplets, they suffer the projection

$$\lambda = +\gamma_{\Omega,M} \lambda^T \gamma_M^{-1} \quad (5.5)$$

Hence they transform in the \square of $Sp(m)$ for the SO projection, and in the \square of $SO(m)$ for the Sp projection. On the other hand, for the two fermion zero modes $\tilde{\theta}_{\dot{\alpha}}$, the projection is

$$\lambda = -\gamma_{\Omega,M} \lambda^T \gamma_M^{-1} \quad (5.6)$$

and leads to two fermion zero modes in the \square of $Sp(m)$ for the SO projection, and in the $\bar{\square}$ of $SO(m)$ for the Sp projection.

This implies that in order to obtain two fermion zero modes from this universal multiplet, in order to generate a superpotential, one should consider instantons with orthogonal gauge group and multiplicity one ($O(1)$ instantons). For instantons with symplectic gauge group and multiplicity two ($Sp(2)$ instantons), there are two additional fermion zero modes in the triplet representation. As mentioned, we will continue to consider such instantons in our relaxed scan. Multiple instantons, i.e. boundary states with higher multiplicity, lead to a larger amount of additional fermion zero modes (due to the larger gauge representations for the fermions), and do not contribute to superpotentials; we will not consider such cases even in relaxed scans, since they also very often lead to too many charged fermion zero modes and cannot contribute to the operators of interest (except possibly for $O(2)$ and $U(2)$ instantons with low intersections, which are kept in our scan as a curiosity).

Similarly, for the p_{\pm} sets of MM scalars and fermions associated to the MM 4d chiral multiplets, the projection is

$$\lambda = \pm \gamma_{\Omega, M} \lambda^T \gamma_{\Omega, M}^{-1} \quad (5.7)$$

with the same sign choice as in (5.3). The different structure of γ_{Ω} implies that, for the SO projection we get p_+ , p_- sets of scalars and fermions in the $\bar{\square}$, \square , while for the Sp projection there are p_+ , p_- sets of scalars and fermions in the \square , $\bar{\square}$.

This concludes the discussion of the MM sector. Let us not consider the Ma sectors, from the information from the $\mathcal{M}a$ sectors. Notice that this implies changing four NN boundary conditions to DN, which have different moddings. Hence the states are different in both situations, but the information on the multiplicities is preserved. Specifically, in the NS sector the DN boundary condition introduce an additional vacuum energy which generically makes all states massive. Hence there are no massless scalar zero modes in generic $\mathcal{M}a$ sectors. In the R sector, the change in the moddings reduces the dimension of the massless ground state, leading to a single (chiral) fermionic degree of freedom. Since the orientifold action maps the Ma sector to Ma' sectors, there are no subtleties in the orientifold projection. The end result is $L_{\mathcal{M}a}$, $L_{\mathcal{M}a'}$ fermion zero modes in the $(\square_{\mathcal{M}}, \square_a)$, $(\square_{\mathcal{M}}, \bar{\square}_a)$. The net number of chiral fermion zero modes in the $(\square_{\mathcal{M}}, \bar{\square}_a)$ is given by $I_{\mathcal{M}a} = L_{\mathcal{M}a'} - L_{\mathcal{M}a}$, i.e. the net number of chiral multiplets in the related $\mathcal{M}a$ sector.

The results for orientifold projections for real branes are shown in table 3.

Proj.	Multiplet in \mathcal{M}	\mathcal{M} (before orient.)	\mathcal{M} (after orient.)	M (after orient.)
SO	$N = 1$ vect. mult.	$U(m)$	$O(m)$	$Sp(m)$ $2\boxed{f} + 2\boxed{f} + 4\boxed{b}$
	$N = 1$ ch. mult.	$(p_+ + p_-)$ Ad	$p_+\boxed{f} + p_-\boxed{f}$	$2p_+(\boxed{f} + \boxed{b}) +$ $2p_-(\boxed{f} + \boxed{b})$
Sp	$N = 1$ vect. mult.	$U(m)$	$Sp(m)$	$O(m)$ $2\boxed{f} + 2\boxed{f} + 4\boxed{b}$
	$N = 1$ ch. mult.	$(p_+ + p_-)$ Ad	$p_+\boxed{f} + p_-\boxed{f}$	$2p_+(\boxed{f} + \boxed{b}) +$ $2p_-(\boxed{f} + \boxed{b})$
Any	$N = 1$ ch. mult.	$L_{\mathcal{M}a'}(\square_{\mathcal{M}}, \bar{\square}_a) +$ $L_{\mathcal{M}a}(\square_{\mathcal{M}}, \bar{\square}_a)$	$L_{\mathcal{M}a'}(\square_{\mathcal{M}}, \bar{\square}_a) +$	$L_{\mathcal{M}a'}(\square_{\mathcal{M}}, \bar{\square}_a)_f$
			$L_{\mathcal{M}a}(\square_{\mathcal{M}}, \square_a)$ net $I_{\mathcal{M}a}(\square_{\mathcal{M}}, \bar{\square}_a)$	$L_{\mathcal{M}a}(\square_{\mathcal{M}}, \square_a)_f$ net $I_{\mathcal{M}a}(\square_{\mathcal{M}}, \bar{\square}_a)_f$

Table 3: Orientifold projection for real branes: Massless modes of the 4d space-filling branes \mathcal{M} (before and after the orientifold projection) and zero modes on the instanton branes M (denoted with sub-indices b, f for bosonic and fermionic modes)

Complex brane instantons

We now consider the case of complex brane instantons. The arguments are very similar, hence the discussion is more sketchy. Consider m 4d spacefilling branes \mathcal{M} , associated to the internal boundary state of the instanton brane M of interest. The $\mathcal{M}\mathcal{M}$ leads to a 4d $N = 1$ $U(m)$ vector multiplet and a number $L_{\mathcal{M}\mathcal{M}'}$ of adjoint chiral multiplets. The orientifold action maps it to the $\mathcal{M}'\mathcal{M}'$ sector, hence we may keep just the former and impose no projection. The $\mathcal{M}\mathcal{M}'$ sector is mapped to itself under the orientifold projection. Denoting by $\gamma_{\Omega, \mathcal{M}}$ the action on Chan-Paton indices, the $\mathcal{M}\mathcal{M}'$ modes split into sets $L_{\mathcal{M}\mathcal{M}'}^{\pm}, L_{\mathcal{M}'\mathcal{M}'}^{\pm}$, which suffer a projection

$$\lambda = \pm \gamma_{\Omega, \mathcal{M}} \lambda^T \gamma_{\Omega, \mathcal{M}}^{-1} \quad (5.8)$$

leading, for $\gamma_{\Omega, \mathcal{M}} = \mathbf{1}_m$, to $L_{\mathcal{M}\mathcal{M}'}^+, L_{\mathcal{M}\mathcal{M}'}^-$ chiral multiplets in the \boxed{f}, \boxed{f} , and $L_{\mathcal{M}'\mathcal{M}'}^+, L_{\mathcal{M}'\mathcal{M}'}^-$ chiral multiplets in the \boxed{f}, \boxed{f} . The net number of chiral multiplets in the \boxed{f}, \boxed{f} is $I_{\mathcal{M}\mathcal{M}'}^+ = L_{\mathcal{M}\mathcal{M}'}^+ - L_{\mathcal{M}'\mathcal{M}'}^+, I_{\mathcal{M}\mathcal{M}'}^- = L_{\mathcal{M}\mathcal{M}'}^- - L_{\mathcal{M}'\mathcal{M}'}^-$. And oppositely for $\gamma_{\Omega, \mathcal{M}} = \epsilon_m$.

Finally, the $\mathcal{M}a, \mathcal{M}a'$ and related sectors lead, after the orientifold projection, to $L_{\mathcal{M}a'}, L_{\mathcal{M}a}, L_{\mathcal{M}'a'}, L_{\mathcal{M}'a}$ chiral multiplets in the $(\square_{\mathcal{M}}, \bar{\square}_a), (\square_{\mathcal{M}}, \square_a), (\bar{\square}_{\mathcal{M}}, \bar{\square}_a), (\bar{\square}_{\mathcal{M}}, \square_a)$. In order to simplify notation, we replace $\mathcal{M} \rightarrow M$ in these expressions in our discussions of instanton zero modes.

Let us now consider m brane instantons M and compute their zero mode spectrum in terms of the above. In the MM (and its image $M'M'$) sector there are four scalar modes and four fermions in the adjoint of the $U(m)$ gauge symmetry group; these are related to the 4d vector multiplet in the $\mathcal{M}\mathcal{M}$ sector. In addition, there are $L_{MM'}$ sets of scalars and fermions in the adjoint, related to the $L_{\mathcal{M}\mathcal{M}'}$ non-universal chiral multiplets in the $\mathcal{M}\mathcal{M}$ sector. The MM' sector is mapped to itself, and one has to impose the orientifold projection (recalling that the matrix $\gamma_{\Omega,M}$ differs from $\gamma_{\Omega,\mathcal{M}}$). For $\gamma_{\Omega,\mathcal{M}} = \mathbf{1}$, hence $\gamma_{\Omega,M} = \epsilon$, we obtain L_{MM}^+ , L_{MM}^- chiral multiplets in the \square , \square , and $L_{M'M'}^+$, $L_{M'M'}^-$ chiral multiplets in the $\bar{\square}$, $\bar{\square}$. The net number of chiral multiplets in the \square , \square is $I_{MM'}^+ = L_{MM}^+ - L_{M'M'}^+$, $I_{MM'}^- = L_{MM}^- - L_{M'M'}^-$. And oppositely for $\gamma_{\Omega,\mathcal{M}} = \epsilon$ hence $\gamma_{\Omega,M} = \mathbf{1}$.

In the Ma , Ma' and related sectors, there are generically no bosonic zero modes, and there are $L_{\mathcal{M}a}$, $L_{\mathcal{M}'a'}$, $L_{\mathcal{M}a'}$, $L_{\mathcal{M}'a}$ chiral fermion zero modes in the (\square_M, \square_a) , $(\bar{\square}_M, \bar{\square}_a)$, $(\square_M, \bar{\square}_a)$, and $(\bar{\square}_M, \square_a)$ respectively. The net number of chiral fermion zero modes in the $(\square_M, \bar{\square}_a)$ and $(\bar{\square}_M, \square_a)$ is given by $I_{\mathcal{M}a} = L_{\mathcal{M}a'} - L_{\mathcal{M}'a'}$ and $I_{\mathcal{M}a'} = L_{\mathcal{M}a} - L_{\mathcal{M}'a}$. In order to simplify notation, we replace $\mathcal{M} \rightarrow M$ in these expressions in our discussions of instanton zero modes.

The results for orientifold projections for real branes are shown in table 4.

6 Search for M instantons

In this section we perform a search of models which admit an instanton inducing a right-handed neutrino Majorana mass operator. Namely, for each model with the chiral content of the SM in the classification described in Section 4.2, we first scan over boundary states, searching for all instantons with the required uncharged and charged fermion zero mode structure to yield neutrino masses. We then relax our criteria a bit and allow for instantons with correct charged zero mode structure but having extra non-chiral zero modes (both charged and uncharged). The idea is that these non-chiral zero modes could be lifted by diverse effects, as discussed.

It is important to recall that the cubic couplings between instanton zero modes and 4d chiral multiplets are difficult to compute in Gepner model orientifolds. Hence, we will simply assume that such couplings are non-zero if there is no symmetry forbidding them.

Proj.	Multiplet in \mathcal{M}	\mathcal{M} (before orient.)	\mathcal{M} (after orient.)	M (after orient.)
Any	$N = 1$ vect. mult.	$U(m) \times U(m)'$	$U(m)$	$U(m)$ $4 \text{ Ad}_f + 4 \text{ Ad}_b$
	$N = 1$ ch. mult.	$p_{\text{adj}} \text{ Ad} + p_{\text{adj}} \text{ Ad}'$	$p_{\text{adj}} \text{ Ad}$	$2p_{\text{adj}} (\text{ Ad}_f + \text{ Ad}_b)$
SO	$N = 1$ ch.mult.	$L_{\mathcal{M}\mathcal{M}}(\square_{\mathcal{M}}, \square_{\mathcal{M}'})$	$L_{\mathcal{M}\mathcal{M}}^+ \square_{\mathcal{M}} + L_{\mathcal{M}\mathcal{M}}^- \bar{\square}_{\mathcal{M}}$	$2L_{\mathcal{M}\mathcal{M}}^+ \bar{\square}_{b,f} + 2L_{\mathcal{M}\mathcal{M}}^- \square_{b,f}$
		$L_{\mathcal{M}'\mathcal{M}'}(\bar{\square}_{\mathcal{M}}, \bar{\square}_{\mathcal{M}'})$	$L_{\mathcal{M}'\mathcal{M}'}^+ \bar{\square}_{\mathcal{M}} + L_{\mathcal{M}'\mathcal{M}'}^- \square_{\mathcal{M}}$	$2L_{\mathcal{M}'\mathcal{M}'}^+ \bar{\square}_{b,f} + 2L_{\mathcal{M}'\mathcal{M}'}^- \square_{b,f}$
Sp	$N = 1$ ch.mult.	$L_{\mathcal{M}\mathcal{M}}(\square_{\mathcal{M}}, \square_{\mathcal{M}'})$	$L_{\mathcal{M}\mathcal{M}}^+ \bar{\square}_{\mathcal{M}} + L_{\mathcal{M}\mathcal{M}}^- \square_{\mathcal{M}}$	$2L_{\mathcal{M}\mathcal{M}}^+ \square_{b,f} + 2L_{\mathcal{M}\mathcal{M}}^- \bar{\square}_{b,f}$
		$L_{\mathcal{M}'\mathcal{M}'}(\bar{\square}_{\mathcal{M}}, \bar{\square}_{\mathcal{M}'})$	$L_{\mathcal{M}'\mathcal{M}'}^+ \bar{\square}_{\mathcal{M}} + L_{\mathcal{M}'\mathcal{M}'}^- \square_{\mathcal{M}}$	$L_{\mathcal{M}'\mathcal{M}'}^+ \square_{b,f} + L_{\mathcal{M}'\mathcal{M}'}^- \bar{\square}_{b,f}$
Any	$N = 1$ ch. mult.	$L_{\mathcal{M}a'}(\square_{\mathcal{M}}, \bar{\square}_a) +$ \dots \dots \dots	$L_{\mathcal{M}a'}(\square_{\mathcal{M}}, \bar{\square}_a)$ $L_{\mathcal{M}a}(\square_{\mathcal{M}}, \square_a)$ $L_{\mathcal{M}'a'}(\bar{\square}_{\mathcal{M}}, \bar{\square}_a)$ $L_{\mathcal{M}'a}(\bar{\square}_{\mathcal{M}}, \square_a)$	$L_{\mathcal{M}a'}(\square_{\mathcal{M}}, \bar{\square}_a)_f$ $L_{\mathcal{M}a}(\square_{\mathcal{M}}, \square_a)_f$ $L_{\mathcal{M}'a'}(\bar{\square}_{\mathcal{M}}, \bar{\square}_a)_f$ $L_{\mathcal{M}'a}(\bar{\square}_{\mathcal{M}}, \square_a)_f$
			net $I_{\mathcal{M}a}(\square_{\mathcal{M}}, \bar{\square}_a)$ net $I_{\mathcal{M}a'}(\square_{\mathcal{M}}, \square_a)$	net $I_{\mathcal{M}a}(\square_{\mathcal{M}}, \bar{\square}_a)_f$ net $I_{\mathcal{M}a'}(\square_{\mathcal{M}}, \square_a)_f$

Table 4: Orientifold projection for complex branes: Massless modes of the 4d space-filling branes \mathcal{M} (before and after the orientifold projection) and zero modes on the instanton branes M (denoted with sub-indices b, f for bosonic and fermionic modes)

6.1 The instanton scan

Our detailed strategy will become clear along the description of the results. Given a set of **a,b,c,d** standard model branes, we must look for additional boundary states M that satisfy the requirements of a $(B - L)$ -violating instanton. From the internal CFT point of view this is just another boundary state, differing from 4d spacefilling branes only in the fully localized 4d spacetime structure. The minimal requirement for such a boundary state is $B - L$ violation, which means explicitly

$$I_{\mathcal{M}\mathbf{a}} - I_{\mathcal{M}\mathbf{a}'} - I_{\mathcal{M}\mathbf{d}} + I_{\mathcal{M}\mathbf{d}'} \neq 0 \quad (6.1)$$

It is easy to see that the existence of such an instanton implies (and hence requires) the existence of a Stückelberg coupling making $B - L$ massive. To see this, consider adding to the Standard Model configuration a 4d spacefilling brane \mathcal{M} (in fact used in Section 5) associated to the boundary state M (RR tadpoles can be avoided by simultaneously including M antibranes, which will not change the argument). The new sector in the chiral spectrum charged under the branes \mathcal{M} can be obtained by reversing the argument in Section 5, and is controlled by the intersection numbers of M . From

the above condition it follows that the complete system has mixed $U(1)_{B-L} \times (G_{\mathcal{M}})^2$ anomalies, where $G_{\mathcal{M}}$ is the Chan-Paton-factor of brane \mathcal{M} . These anomalies are cancelled by a Green-Schwarz mechanism involving a $(B-L)$ -axion bilinear coupling, which ends up giving a mass to $B-L$ via the Stückelberg mechanism. This coupling is in fact not sensitive to the presence of the brane \mathcal{M} , hence it must have been present already in the initial model (without \mathcal{M}).

Hence the existence of a boundary label M that satisfies (6.1) implies that $B-L$ is massive. Unfortunately the converse is not true: even if $B-L$ has a Stückelberg mass, this still does not imply the existence of suitable instantons satisfying (6.1)¹⁶ Indeed, in several models we found not a single boundary state satisfying (6.1).

Note that, since hypercharge must be massless, one can use the reverse argument and obtain that

$$I_{M\mathbf{a}} - I_{M\mathbf{a}'} - I_{M\mathbf{c}} + I_{M\mathbf{c}'} - I_{M\mathbf{d}} + I_{M\mathbf{d}'} = 0 \quad (6.2)$$

in all models. We verified this for all models we considered as a check on the computations.

As already discussed in Section 4.2, in the search for SM constructions in Gepner orientifold, there are 391 models with massless hypercharge and massive $B-L$. In these models we found a total of 29680 instantons with $B-L$ violation, i.e. with intersection numbers satisfying (6.1). Of course, in order to serve our purpose of generating a Majorana mass superpotential, the instantons have to satisfy some more conditions. Let us consider them in order of importance, and start with the conditions on the net number of chiral fermion zero modes charged under the 4d observable sector. Clearly we need $I_{M\mathbf{a}} = I_{M\mathbf{a}'}$ and $I_{M\mathbf{b}} = I_{M\mathbf{b}'}$. The latter condition is automatically satisfied in this case, because the b -brane is real in all 391 models. The chiral conditions on the zero modes charged under the branes \mathbf{c} and \mathbf{d} are as in [3]¹⁷ and are given in equations (3.6), (3.7) (3.8) of the present paper. These are the instantons of most interest, and on which we mainly focus. However, as discussed in Section 3.4, other important B - and/or L - violating operators (such as the Weinberg operator or the LH operator) can

¹⁶From intuition in geometric compactifications, one expects that there may always exist a D-brane with the appropriate topological pairings, but there is no guarantee that there is a supersymmetric representative in that topological sector, and even less that it would have no additional fermion zero modes. Note also that even if such D-brane instantons exists, there is no guarantee that it will fall in the scan over RCFT boundary states.

¹⁷Note that there is a sign change in the contribution of the $U(1)_{\mathbf{d}}$ generator to Y in comparison to [3]

be generated by instantons with similar intersection numbers, up to a factor of 2 and a sign, see table 2. For this reason we also allow at this stage any instanton which has the correct number of charged zero modes to generate them. Imposing these conditions reduces the number of candidate instantons potentially contributing to neutrino masses in any of the models to 1315.

All instantons satisfying these requirements are summarized in the table 5. In columns 1,2 and 3 we list the tensor combination, MIPF and orientifold choice for which the model occurred. The latter two numbers codify simple current data that describe respectively a MIPF and an orientifold. MIPFs are in general defined by means of a subgroup \mathcal{H} of the simple current group \mathcal{G} , plus a certain matrix X of rational numbers [55]. Orientifolds are defined by a simple current and a set of signs [43]. In previous work [6] we have enumerated these quantities (up to permutation symmetries) and assigned integer labels to them for future reference. We only refer to these numbers here, but further details are available upon request. Usually for each MIPF and orientifold which contains the standard model there are several choices **a,b,c,d** for which it is obtained. For a given choice of tensor combination, MIPF and orientifold and SM branes there may be several instantons. For clarity we put all such instantons together in the information in table 5. In column 4 we indicate which type of instanton branes were found. Five types are distinguished: $O1$, $O2$, $S2$, $U1$ and $U2$, corresponding to $O(1)$, $O(2)$, $Sp(2)$, $U(1)$ and $U(2)$ Chan-Paton symmetry on the instanton volume. The number indicates the instanton brane multiplicity that gives the correct number of instanton charged zero modes from the **a**, **b**, **c**, **d** branes, to lead to right-handed neutrino Majorana masses. The number of zero modes is in general the product of the instanton brane multiplicity and ‘intersection number’ with the corresponding 4d spacefilling brane. As discussed in Section 5, for symplectic branes the smallest possible brane multiplicity is 2. As we discussed there, only $O1$ instantons may have the required universal minimal set of two zero modes in the uncharged sector. Still we look for all $O(1)$, $Sp(2)$ and $U(1)$ instantons which may yield a superpotential if the extra uncharged fermion zero modes. In this vein we also include a search for $O2$ and $U2$ instantons. Note also that such $O2$ or $U2$ instantons imply the existence of other instantons involving the same boundary state, but with multiplicity 1, which may lead to the R-parity violating operator LH . We will discuss the generation of R-parity violating operators at the end of this section. The third character (+ or -) in the instanton in table 5 is the sign of $I_{Mc'} - I_{Mc}$. For the instantons giving rise to right-handed neutrino Majorana masses this sign should be negative, whereas it should

be positive for instantons giving rise to the Weinberg operator (or the LH operator), see table 2.

The 1315 instantons are divided in the following way over the different types: 3 of types $O1+$ and $O1-$, 46 of type $U1+$, 24 of type $U1-$, 550 $S2+$, 627 $S2-$, 27 of types $U2+$ and $U2-$ and four of types $O2+$ and $O2-$. Notice that the vast majority (97.5%) of the instanton solutions are of type $S2+$ and $S2-$. This is encouraging given the nice properties of such instantons, concerning e.g. R-parity conservation. Note also that in almost all cases both $S-$ and $S+$ are simultaneously present,¹⁸ so both sources of physical neutrino Majorana masses (from the see-saw mechanism or the Weinberg operator) are present. The other instanton classes possibly generating right-handed neutrino masses are $O1-$ and $U1-$, which are much less abundant. There is just one orientifold with $O1-$ instantons, for which one can obtain cancellation of RR tadpoles, see below. On the other hand we have found no orientifold with $U1-$ instantons and cancellation of tadpoles, see below.

Table 5: *Summary of instanton branes.*

Tensor	MIPF	Orientifold	Instanton	Solution
(1,16,16,16)	12	0	$S2^+, S2^-$	Yes
(2,4,12,82)	19	0	$S2^-!$?
(2,4,12,82)	19	0	$U2^+, U2^-!$	No
(2,4,12,82)	19	0	$U1^+, U1^-$	No
(2,4,14,46)	10	0		
(2,4,14,46)	16	0		
(2,4,16,34)	15	0		
(2,4,16,34)	15	1		
(2,4,16,34)	16	0	$S2^+, S2^-$	Yes
(2,4,16,34)	16	1		
(2,4,16,34)	18	0	$S2^-$	Yes
(2,4,16,34)	18	0	$U1^+, U1^-, U2^+, U2^-$	No
(2,4,16,34)	49	0	$U2^+, S2^-!, U1^+$	Yes
Continued on next page				

¹⁸In some models contributing many instantons there is an *exact* symmetry between $S-$ and $S+$. This explains the *approximate* symmetry in the full set. In some cases this symmetry can be understood in terms of flipping the degeneracy labels of boundary states. We regard it as accidental, since it is not found in all models.

Table 5 – continued from previous page

Tensor	MIPF	Orientifold	Instanton	Solution
(2,4,16,34)	49	0	$U1^-$	No
(2,4,18,28)	17	0		
(2,4,22,22)	13	3	$S2^+, S2^-!$	Yes!
(2,4,22,22)	13	2	$S2^+, S2^-!$	Yes
(2,4,22,22)	13	1	$S2^+, S2^-$	No
(2,4,22,22)	13	0	$S2^+, S2^-$	Yes
(2,4,22,22)	31	1	$U1^+, U1^-$	No
(2,4,22,22)	20	0		
(2,4,22,22)	46	0		
(2,4,22,22)	49	1	$O2^+, O2^-, O1^+, O1^-$	Yes
(2,6,14,14)	1	1	$U1^+$	No
(2,6,14,14)	22	2		
(2,6,14,14)	60	2		
(2,6,14,14)	64	0		
(2,6,14,14)	65	0		
(2,6,10,22)	22	2		
(2,6,8,38)	16	0		
(2,8,8,18)	14	2	$S2^+, S2^-!$	Yes
(2,8,8,18)	14	0	$S2^+, S2^-!$	No
(2,10,10,10)	52	0	$U1^+, U1^-$	No
(4,6,6,10)	41	0		
(4,4,6,22)	43	0		
(6,6,6,6)	18	0		

Most models have a hidden sector containing extra boundary states beyond the SM ones. In the same spirit of imposing chiral conditions first, we should require that $I_{Mh} = I_{Mh'}$, where h is a hidden sector brane. This is to guarantee that the generated superpotential does not violate some hidden sector gauge symmetry which would require the presence of hidden sector fields along with the ν_R bilinear. The latter condition is not imposed on the previously known hidden sector (*i.e.* the one in [6, 7]), but instead a new search for tadpole solutions was performed, for each M , restricting the candidate hidden sector branes to those satisfying $I_{Mh} = I_{Mh'}$ (as discussed in

Section 5). This is because in general the known hidden sector in [6, 7] is just a sample out of a huge number of possibilities.

In column 5 we indicate for which instantons it was possible to satisfy the tadpole conditions with this additional constraint. With regard to observable-hidden matter we use the same condition as in [6], namely that it is allowed only if it is vector-like. Such a solution could be found for 879 of the 1315 instantons, with ten cases inconclusive (*i.e* it was computationally too difficult to decide if a solution does or does not exist). The latter are indicated with a question mark in column 5 (for most of the undecidable cases there is a tadpole solution for a different instanton with the same characteristics; for that reason just one question mark appears).

Independently of the RR tadpole condition (since there may be alternative sources for its cancellation, or hidden sectors which fall beyond the reach of RCFT), we can also consider the further constraint that the number of charged fermion zero modes is exactly right, not just in the chiral sense. This means $I_{M\mathbf{a}} = I_{M\mathbf{a}'} = I_{M\mathbf{b}} = I_{M\mathbf{b}'} = 0$, $I_{M\mathbf{c}} = 2, I_{M\mathbf{c}'} = 0$ and $I_{M\mathbf{d}} = -2, I_{M\mathbf{d}'} = 0$ or vice-versa. Furthermore we require that there are no adjoint or rank-2 tensor zero-modes (note that the latter could be chiral if the instanton brane is complex, and indeed they are in some of the 1315 cases). This reduces the 1315 instantons to 263. In column 4 we indicate those cases with an exclamation mark. It is noteworthy that the success rate for solving the tadpole conditions is highest for these instantons: 254 of the 263 allow a solution (with 3 undecided). If an exclamation mark appears in column 4, this only indicates that *some* of the instantons are free of the aforementioned zero modes, not that all of them are. But in all cases, if there are tadpole solutions, they exist in particular for the configurations with an exclamation mark. Finally we may impose the condition that I_{Mh} and $I_{Mh'}$ are separately zero. This is indicated with an exclamation mark in column 5. This turns out to be very restrictive. The only cases where this happens have no hidden sector at all.

It is worth remarking that the only instantons having exactly the correct set of charged zero modes and cancelling tadpoles are of $S2^\pm$ type. Also those instantons are the only cases marked with an exclamation mark in column 4 and 5. These examples, which will be discussed below in some detail, also have just the minimal set of fermion zero modes, except for the universal sector (which for $Sp(2)$ instantons contains two extra triplets).

The main conclusion about this scan is that we did not find any instantons with exactly the zero mode fermions to generate the neutrino mass superpotential. However

we have found a number of examples which come very close to that, with exactly the required charged zero modes and a very reduced set of extra uncharged zero modes from the universal sector. These extra zero modes are non-chiral and hence one expects that e.g. RR/NS fluxes or other effects may easily lift them, as we discussed in section 2. Concerning $O(1)$ instantons, which have just the two required fermion zero modes in the universal sector, we have found one example, with the appropriate net structure of charged zero modes. However, it has plenty of other extra zero modes. We discuss examples of $O(1)$ and $Sp(2)$ instantons in the following subsections.

6.2 An O1 example

Let us first discuss the case of $O(1)$ instantons. In principle they would be the more attractive since they have no undesirable universal zero modes at all. Unfortunately this type of instanton is rare within the set we scanned, and we found just one example with a solution to the tadpole equations without any unwanted chiral zero-modes. The instanton however has a very large number of uncharged and charged vector-like zero modes.

The standard model brane configuration occurs for tensor product $(2, 4, 22, 22)$, MIPF 49, orientifold 1, boundaries $(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}) = (487, 1365, 576, 486)$. As usual we only provide this information in order to locate this model in the database. Further details are available on request.

The bi-fundamental fermion spectrum of this model in the $(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d})$ sector is fairly close to the MSSM: there is an extra up-quark mirror pair, two mirror pairs of leptons with down quark charges and one with up-quark charges, plus two extra right-handed neutrinos (*i.e.* a total of five right-handed neutrinos). There are three MSSM Higgs pairs. The tensor spectrum is far less appealing, in particular for brane \mathbf{c} : this has 25 adjoints and 7 vector-like pairs of anti-symmetric tensors.

As we said, there is just one instanton brane of type $O1-$. It has exactly the right number of zero-modes with brane d , but five superfluous pairs of vector-like zero-modes with brane c , plus one vector-like pair with brane a . In addition there are four symmetric tensor zero-modes on the instanton brane (which of course are vector-like, since it is a real brane): the parameter p_+ in table 3 is equal to 2.

The tadpole solution that is (chirally speaking) compatible with this instanton has a large hidden sector: $O(1) \times O(2)^4 \times O(3) \times U(1)^2 \times Sp(2)^2 \times U(3)$ (there are other possibilities, but no simple ones). This hidden sector introduces more undesirable features: vector-like observable/hidden matter, vector-like instanton/hidden sector modes, plus

chiral and non-chiral matter within the hidden sector. Finally the coupling ratios are as follows: $\alpha_3/\alpha_2 = .54$, $\sin^2\theta_w = .094$, and the instanton coupling is 3.4 times weaker than the QCD coupling ($\alpha_3/\alpha_{\text{Instanton}} = 3.4$).

Despite these unappealing features this model does demonstrate the existence of this kind of solution.

6.3 The S2 models

As we have mentioned, these are the examples which come closer to the minimal set of fermion zero modes. As we see in Table 5, all such instantons satisfying the criteria on the zero mode structure (except for the extra universal zero modes) appear for models based on the same CFT orientifold. It is the one obtained from the (2, 4, 22, 22) Gepner model with MIPF 13 and orientifold 3 in the table. The model is obtained as follows.

6.3.1 The closed string sector

We start with the tensor product (2,4,22,22). This yields a CFT with 12060 primary fields, 48 of which are simple currents, forming a discrete group $\mathcal{G} = \mathbf{Z}_{12} \times \mathbf{Z}_2 \times \mathbf{Z}_2$. After taking into account the permutation symmetry of the last two factors, we find that this tensor product has 54 symmetric MIPFs, and we choose one of them to build the model of interest. For convenience we specify all quantities in terms of a standard minimal model notation, but also in terms of the labelling of the computer program “kac” that generates the spectrum. This particular MIPF is nr. 13. To build it we choose a subgroup of \mathcal{G} , which is isomorphic to $\mathcal{H} = \mathbf{Z}_{12} \times \mathbf{Z}_2$. The generator of the \mathbf{Z}_{12} factor is primary field nr. 1, $(0, 0, 0, \{24, -24, 0\}, \{24, 20, 0\})$, and the \mathbf{Z}_2 factor is generated by primary field nr. 24, $(0, 0, 0, 0, \{24, 20, 2\})$. The representations are specified on a basis (NSR, $k = 2, k = 4, k = 22, k = 22$), *i.e.* the boundary conditions of the NSR-fermions and the four minimal models in the tensor product. Here 0 indicates the CFT vacuum, and for all other states we use the familiar (l, q, s) notation for the $N = 2$ minimal models. The first generator has conformal weight $h = \frac{11}{12}$ and has ground state dimension 1. The second has weight $h = \frac{1}{2}$ and has ground state dimension 2: the ground state contains both $(0, 0, 0, 0, \{24, 20, 2\})$ and $(0, 0, 0, \{24, 20, 2\}, 0)$. The matrix X defining the MIPF according to the prescription given in [53][54][55] is

$$\begin{pmatrix} \frac{1}{12} & 0 \\ 0 & \frac{1}{2} \end{pmatrix} \tag{6.3}$$

This simple current modification is applied to the charge conjugation invariant of the tensor product. This defines a MIPF that corresponds to an automorphism of the fusion rules, and that pairs all the primaries in the CFT off-diagonally. The number of Ishibashi states, and hence the number of boundary states is 1080. The MIPF is invariant under exchange of the two $k = 22$ factors: this maps current 24 to itself, and current 1 to current 11, which is also in \mathcal{H} . Hence this symmetry of the tensor product maps \mathcal{H} into itself, and it also preserves the matrix X .

To define an orientifold, we must specify a ‘‘Klein bottle current’’ plus two signs defined on the basis of the simple current group. For the current K we use the generator of the second \mathbf{Z}_2 in \mathcal{G} , primary field nr. 12. This is the representation $(0, 0, \{4, -4, 0\}, \{(24, 16, 2)\}, \{(24, -12, 2)\})$ which is degenerate with nine other states, all of dimension 1 and conformal weight 7. The crosscap signs are chosen, on the aforementioned basis of \mathcal{H} as $(+, -)$. This results in a crosscap coefficient of 0.0464731, and it is orientifold nr. 3 of a total of 8. The orientifold is also invariant under permutation of the identical factors.

The closed string spectrum contains 14 vector multiplets and 60 chiral multiplets.

6.3.2 The standard model branes

To build a standard model configuration we have to specify the boundary state labels. It turns out that we have four choices for label **a** and **b**, one for **c** and two for **d**. This leads to a total of 32 possibilities. Among these 32 there are 22 have distinct spectra (distinguished by the number of vector-like states), but for all 32 choices one obtains the same set of dilaton couplings. It seems plausible that these choices simply correspond to putting the **a**, **b** and **d** branes in slightly different positions, so that we move the configuration in brane moduli space. The choices are as follows (these are boundary labels assigned by the computer program, and can be decomposed in terms of minimal model representations; this will be explained in table 6 below)

$$\begin{aligned}
 \mathbf{a} : & \quad 10, 22, 130, 142 \\
 \mathbf{b} : & \quad 210, 282, 290, 291 \\
 \mathbf{c} : & \quad 629 \\
 \mathbf{d} : & \quad 712, 797
 \end{aligned}$$

There are additional possibilities, but they do not give rise to additional distinct spectra.

Table 6: *Branes appearing in standard model configurations*

Label	Orbit/Deg.	Reps	Weight	Dimension
10	240	$(0, 0, 0, 0, \{10, 0, 0\})$	5/4	1
130	2760	$(0, 0, 0, \{10, 0, 0\}, 0)$	5/4	1
22	[528,0]	$(0, 0, 0, \{1, -1, 0\}, \{11, 1, 0\})$	3/2	1
		$(0, 0, 0, \{1, 1, 0\}, \{11, -1, 0\})$	3/2	1
142	[3048,0]	$(0, 0, 0, \{11, -1, 0\}, \{1, 1, 0\})$	3/2	1
		$(0, 0, 0, \{11, 1, 0\}, \{1, -1, 0\})$	3/2	1
210	4248	$(0, 0, \{3, 3, 0\}, \{3, -3, 0\}, \{9, -9, 0\})$	1/2	1
282	5760	$(0, 0, \{3, 3, 0\}\{9, -9, 0\}\{3, -3, 0\})$	1/2	1
290	[5952,0]	$(0, 0, \{1, 1, 0\}\{9, 7, 0\}\{11, -11, 0\})$	5/6	1
291	[5952,24]	$(0, 0, \{1, 1, 0\}\{9, 7, 0\}\{11, -11, 0\})$	5/6	1
629	[9348,30]	$(0, (1, -1, 0), 0, \{9, 9, 0\}\{5, -3, 0\})$	7/12	1
712	[9852,0]	$(0, \{1, 1, 0\}\{3, -3, 0\}\{1, 1, 0\}\{5, 5, 0\})$	1/2	2
		$(0, \{1, 1, 0\}\{1, -1, 0\}\{1, 1, 0\}\{5, -3, 0\})$	1/2	2
797	[10356,30]	$(0, \{1, 1, 0\}\{3, -3, 0\}\{5, 5, 0\}\{1, 1, 0\})$	1/2	2
		$(0, \{1, 1, 0\}\{1, -1, 0\}\{5, -3, 0\}\{1, 1, 0\})$	1/2	2

The second column gives the boundary labels in terms of a primary field label and a degeneracy label (boundaries not indicated by square brackets are not degenerate). The labels appearing in columns 1 and 2 are assigned by the computer program, and are listed here only for the purpose of reproducing the results using that program. In column 2, the boundary labels are expressed in terms of primary field labels, as in formula (A.4). If a single number appears, this is a representative of an \mathcal{H} -orbit corresponding to the boundary. If square brackets are used, this means that the \mathcal{H} -orbit has fixed points, and that it corresponds to more than one boundary label. The second entry in the square brackets is the degeneracy label, and refers to a character of the ‘‘Central Stabilizer’’ defined in [43]; the details of the definition and the labelling will not be important here. In this case the first entry within the square brackets refers to an orbit representative.

These orbit representatives can also be expressed in a standard form for minimal model tensor products. This is done in column 3. This is basically the same expansion shown in (A.4), except that the degeneracy label Ψ_I turns out to be trivial in all cases,

both for the standard model and for the instanton branes shown below (although the theory does contain primaries with non-trivial Ψ 's). In columns 4 and 5 we specify the weight and ground state dimension of the corresponding highest weight representation. These data are not directly relevant for the boundary state, but helps in identifying it.

Since boundaries are specified by orbit representatives, it is not straightforward to compare them, since the standard choice (the one listed in column 2) is arbitrary. For this reason we have used another representative in columns 3, 4 and 5, selected by an objective criterion: we choose the one of minimal dimension and minimal conformal weight (in that order). If there is more than one representative satisfying these criteria we list all.

6.3.3 The open string spectrum

In Table 7 we summarize the spectra of the 32 models. The first four columns list the **a,b,c,d** brane labels. The last eight columns specify the total number of multiplets of types Q (quark doublet), U (up quark singlet), D (down quark singlet), L (lepton doublet), E (charged lepton singlet), N (neutrino singlet), Y (lepto-quark) and H (Higgs). The numbers given are for the total number of lefthanded fermions in the representation, plus their complex conjugates. So for example a 7 in column ‘‘Q’’ means that there are 5 quark doublets in the usual representation $(3, 2, \frac{1}{6})$, plus two in the complex conjugate representation $(3^*, 2, -\frac{1}{6})$.

This yields the required three families of quark doublets, plus two mirror pairs. Hence the smallest number that can occur in the six columns QUDLEN is three, if there are no mirrors (note that cubic anomaly cancellation requires three right-handed neutrinos in this class of models). The lepto-quarks Y are all in the same representation as the down-quarks (D), or the conjugate thereof, and they occur only as vector-like mirror pairs. They differ from D-type mirror quarks because they carry lepton number, because they come from open strings ending on the **d**-brane instead of the **c**-brane. In general, there can also exist U-type lepto-quarks, but in these models they do not occur. Finally the numbers 10, 18 and 26 in column ‘H’ mean that there are 5, 9 or 13 MSSM Higgs pairs $H + \bar{H}$. It is worth noticing that right-handed quarks U, D and neutrinos $N = \nu_R$ do not have vectorlike copies. On the other hand right-handed leptons E always have one and the left-handed fields Q, L may have up to 3 vector-like copies.

Table 7: *Spectrum all 32 configurations.*

$U(3)$	$Sp(2)$	$U(1)$	$U(1)$	Q	U	D	L	E	N	Y	H
10	210	629	712	7	3	3	9	5	3	6	10
22	210	629	712	7	3	3	9	5	3	6	10
130	210	629	712	3	3	3	9	5	3	2	10
142	210	629	712	3	3	3	9	5	3	2	10
10	282	629	712	3	3	3	5	5	3	6	26
22	282	629	712	3	3	3	5	5	3	6	26
130	282	629	712	7	3	3	5	5	3	2	26
142	282	629	712	7	3	3	5	5	3	2	26
10	290	629	712	3	3	3	3	5	3	6	18
22	290	629	712	3	3	3	3	5	3	6	18
130	290	629	712	3	3	3	3	5	3	2	18
142	290	629	712	3	3	3	3	5	3	2	18
10	291	629	712	3	3	3	3	5	3	6	18
22	291	629	712	5	3	3	3	5	3	6	18
130	291	629	712	3	3	3	3	5	3	2	18
142	291	629	712	3	3	3	3	5	3	2	18
10	210	629	797	7	3	3	5	5	3	2	10
22	210	629	797	7	3	3	5	5	3	2	10
130	210	629	797	3	3	3	5	5	3	6	10
142	210	629	797	3	3	3	5	5	3	6	10
10	282	629	797	3	3	3	9	5	3	2	26
22	282	629	797	3	3	3	9	5	3	2	26
130	282	629	797	7	3	3	9	5	3	6	26
142	282	629	797	7	3	3	9	5	3	6	26
10	290	629	797	3	3	3	3	5	3	2	18
22	290	629	797	3	3	3	3	5	3	2	18
130	290	629	797	3	3	3	3	5	3	6	18
142	290	629	797	3	3	3	3	5	3	6	18
10	291	629	797	3	3	3	3	5	3	2	18
22	291	629	797	5	3	3	3	5	3	2	18
130	291	629	797	3	3	3	3	5	3	6	18

Continued on next page

Table 7 – continued from previous page

$U(3)$	$Sp(2)$	$U(1)$	$U(1)$	Q	U	D	L	E	N	Y	H
142	291	629	797	3	3	3	3	5	3	6	18

In the following table we list the multiplicities L_{aa} and $L_{aa'}$ of the branes that occur in these models, leading to vector-like sets of adjoints and rank-2 tensors. Since brane **b** is symplectic, the number of adjoints is equal to the number of symmetric tensors.

Table 8: *4d matter from the aa and aa' sectors.*

Boundary	Adjoints	Anti-symm.	Symm.
a (10)	2	2	6
a (22)	2	2	2
a (130)	2	2	6
a (142)	2	2	2
b (210)	-	14	10
b (282)	-	14	10
b (290)	-	14	6
b (291)	-	14	6
c (629)	9	-	14
d (712)	3	-	6
d (797)	3	-	6

It should be emphasized that CFT constructions generically correspond to particular points in moduli space of CY orientifolds. Due to this, they usually have an ‘enhanced’ massless particle content with extra vector-like matter and closed string gauge interactions. Thus one would expect that many of the massless vector-like chiral fields present in this class of models could gain masses while moving to a nearby point in moduli space.

6.3.4 The instantons

Each of these 32 Standard Model compactifications admits 8 instantons. The instanton labels are identical for all the 32 models. They are listed in Table 9. The first five columns use the same notation as for the standard model boundary labels. In column 6 we list the numerical value of the dilaton coupling to the instanton brane. This quantity is proportional to $\frac{1}{g^2}$. It is instructive to compare these couplings to the gauge couplings, in order to gain intuition on the suppression factor for our instantons. In these models the $U(3)$ dilaton couplings are 0.00622, so that the instantons are more strongly coupled than QCD¹⁹. On the other hand in this particular model the ratio α_3/α_2 at the string scale is 3.23 (the value of $\sin^2\theta_w$ at the string scale is 0.527). All of these couplings are subject to renormalization group running, and there are plenty of vector-like states to contribute to this, if one assumes that they acquire masses at a sufficiently low scale. One should perform a detailed renormalization group analysis to check whether one may obtain consistency with the gauge couplings measured at low-energies. Let us emphasize however that one expects that moving in moduli space many of these vector-like states will gain masses and also the values of the different gauge couplings will also generically vary.

Since the value of the Type II dilaton is a free parameter at this level, one can get the appropriate (intermediate) mass scale for the right-handed neutrino Majorana masses by choosing an appropriate value for the dilaton. In this context, it is satisfactory to verify that the instanton couplings are unrelated to the gauge couplings, as expected since they do not correspond to gauge instantons [3], and are in fact less suppressed than the latter.

Note that the 8 instantons fall into two distinct classes (evidently not related by any discrete symmetry, since the conformal weight on the boundary orbit is distinct, and the coupling is different as well). Within each class, the orbits of the four instanton boundaries appear to be related by the \mathbf{Z}_2 symmetries of interchange of the last two tensor factors, and simultaneous inversion of the charge q of the minimal model. However, one has to be very careful in reading off symmetries directly from the labels in columns 3 of Tables (6) and (9) for a number of reasons. First of all the entries in column 3 are representatives of boundary orbits, and these representatives themselves are merely representatives of extension orbits. Secondly the action of any discrete

¹⁹ Note that the Type II dilaton in this compactifications is an arbitrary parameter which can always be chosen so that we consistently work at weak coupling. It is the relative value of gauge couplings which we are comparing here.

Table 9: *Instantons for all 32 configurations*

Lbl.	Orbit/Deg.	Reps	Weight	Dim.	coupling
414	[8064,0]	$(0, \{1, 1, 0\}, 0, \{22, -22, 0\}, \{20, 16, 0\})$	5/2	1	0.0016993
417	[8076,30]	$(0, \{1, -1, 0\}, 0, \{22, 22, 0\}, \{20, -16, 0\})$	5/2	1	0.0016993
456	[8316,0]	$(0, \{1, 1, 0\}, 0, \{20, 16, 0\}, \{22, -22, 0\})$	5/2	1	0.0016993
459	[8328,30]	$(0, \{1, -1, 0\}, 0, \{20, -16, 0\}, \{22, 22, 0\})$	5/2	1	0.0016993
418	[8088,0]	$(0, \{1, 1, 0\}, 0, \{22, -22, 0\}, \{18, 16, 0\})$	5/3	1	0.0027033
420	[8100,0]	$(0, \{1, -1, 0\}, 0, \{22, 22, 0\}, \{18, -16, 0\})$	5/3	1	0.0027033
502	[8592,0]	$(0, \{1, 1, 0\}, 0, \{18, 16, 0\}, \{22, -22, 0\})$	5/3	1	0.0027033
505	[8604,30]	$(0, \{1, -1, 0\}, 0, \{18, -16, 0\}, \{22, 22, 0\})$	5/3	1	0.0027033

symmetry on the degeneracy labels can be non-trivial. In appendix B we discuss these symmetries in more detail.

6.4 Other examples

The $Sp(2)$ instanton examples just discussed are the ones which get closer to the required minimal set of fermion zero modes. Under slightly weaker conditions, we find many more solutions. In all these cases some additional mechanism beyond exact RCFT will be needed to lift some undesirable zero modes.

The simplest such case is the following. The tensor product is $(2, 8, 8, 18)$, MIPF nr. 14, orientifold 2 (the precise spectra may be found using this information in the database www.nikhef.nl/~t58/filtersols.php). There are three distinct brane configurations for which almost perfect instantons exist, namely $(\mathbf{a}, \mathbf{b}, \mathbf{c}, \mathbf{d}) = (64, 562, 389, 67)$ and $(64, 577, 389, 67)$ and $(65, 560, 189, 66)$. Each has six instantons, three of type $S2+$ and three of type $S2-$. As in the foregoing example, the six instantons are identical for the three standard model configurations. In this example, they have three different dilaton coupling strengths: .00254, .00665 and .0108 (each value occurs once for $S2+$ and once for $S2-$). By comparison, the $U(3)$ -brane dilaton coupling strength is 0.0119338, so that the instanton brane coupling is quite a bit stronger than the QCD coupling. This is again an interesting point if we want that ν_R masses are not too much

suppressed. Furthermore in this example there are three distinct instanton couplings, so that one may expect three non-zero eigenvalues (with a hierarchy) in the mass matrix. As in the previous examples there is not gauge coupling unification, one rather has $\alpha_3/\alpha_2 = .4813$ and $\sin^2(\theta_w) = .183$ at the string scale. Again a full renormalization group analysis should be performed in order to check consistency with the measured low-energy gauge coupling values.

These models all have a hidden sector consisting of a single $Sp(2)$ factor. They have respectively 3, 1 and 3 susy Higgs pairs, and a spectrum of bi-fundamentals that is closer to that of the standard model than the previously discussed $Sp(2)$ examples: quarks and leptons do not have vector-like copies (there are only some vector-like leptoquarks), and even one of the three models have the minimal set of Higgs fields of the MSSM. The rest of the spectrum is purely vector-like, and contains a number of rank-2 tensors, including eight or six adjoints of $U(3)$. Furthermore there is vector-like observable-hidden matter. The only undesirable instanton zero-mode is a single bi-fundamental between the hidden sector $Sp(2)$ brane and the instanton brane. Still, these SM brane configurations without the hidden sector, provide interesting and very simple local models of D-brane sectors admitting instantons generating neutrino masses (with the additional ingredients required to eliminate the extra universal triplets of fermion zero modes).

6.5 R-parity violation

We now turn to the generation of other possible superpotentials violating $B - L$. An instanton violates R-parity if the amount of $B - L$ violation,

$$I_{M\mathbf{a}} - I_{M\mathbf{a}'} - I_{M\mathbf{d}} + I_{M\mathbf{d}'} \quad (6.4)$$

is odd. Examples of instantons with that property were found in the following tensor product/MIPF/orientifold combinations: $[(1, 16, 16, 16), 12, 0]$, $[(2, 4, 16, 34), 49, 0]$, $[(2, 4, 12, 82), 19, 0]$, $[(2, 4, 22, 22), 49, 0]$ and $[(2, 4, 16, 34), 18, 0]$. Note that all cases for which $O2$ or $U2$ instantons were found necessarily have R-parity violating instantons as well: the corresponding $O1$ and $U1$ instantons have $I_{M\mathbf{d}}$ or $I_{M\mathbf{d}'}$ equal to ± 1 , whereas the intersection with the \mathbf{a} is non-chiral. In principle, there are many more ways to obtain R-parity violating instantons (either due to non-vanishing contributions to $I_{M\mathbf{a}} - I_{M\mathbf{a}'}$ or higher values of $I_{M\mathbf{d}} - I_{M\mathbf{d}'}$), and indeed, many such instantons turn out to exist. But the number of tensor product/MIPF/orientifold combinations where they occur hardly increases: only in the case $[(1, 16, 16, 16), 12, 0]$ we found R-parity

violating instantons, but no $U1$ or $O1$ instantons. This suggests that in the other cases R-parity is a true symmetry of the model. Unfortunately we have no way of rigorously ruling out any other non-perturbative effects, but at least the set we can examine respects R-parity. This includes in particular the models without hidden sector (found for $[(2, 4, 22, 22), 13, 3]$) discussed above.

The following table list the total number of instantons with the chiral intersections listed in table 2. The total number of instantons (boundaries violating the sum rule, as defined in (6.1)) is 29680, for all standard model configurations combined. The last four columns indicate how many unitary instantons satisfy the sum rule exactly as listed in table (6.1), how many satisfy it with $I_{M\mathbf{x}} \leftrightarrow -I_{M\mathbf{x}'}$ (the column U'), and how many O-type and S-type instantons there are. Here 'S' refers to boundaries with a symplectic Chan-Paton group if the boundary is used as an instanton brane. All intersection numbers for type S have been multiplied by 2 before comparing with table 2. For real branes, the relevant quantities used in the comparison are $I_{M\mathbf{a}} - I_{M\mathbf{a}'}$, $I_{M\mathbf{c}} - I_{M\mathbf{c}'}$ and $I_{M\mathbf{d}} - I_{M\mathbf{d}'}$, while $I_{M\mathbf{b}} = 0$. There are fewer unitary instantons possibly generating Majorana masses then the numbers mentioned above because the conditions we use here are stricter: we require here that $I_{M\mathbf{x}}$ and $I_{M\mathbf{x}'}$ match exactly, not just their difference. Note however that this still allows additional vector-like zero-modes. If we only wish to consider cases without *any* spurious zero-modes, we may limit ourselves to the O-type instantons in the last column. There are very few to inspect, and all of them turn out to have a few non-universal zero modes.

$D = 4$ Operator	U	U'	S	O
$\nu_R \nu_R$	1	2	627	3
$LHLH$	0	5	550	3
LH	3	3	0	4
QDL	8	4	0	4
UDD	0	0	0	4
LLE	8	4	0	4
$QQQL$	0	4	0	3892
$UUDE$	4	0	0	3880

Table 10: Number of instantons in our search which may induce neutrino masses (first 2 rows), R-parity violation (next 4 rows) or proton decay operators (last 2 rows).

The last two cases are $B - L$ preserving dimension five operators, and obviously do not come from the set of 29680 $B - L$ violating instantons. They were searched for

separately, but the search was limited to the same 391 models we used in the rest of the paper. Obviously, one could equally well look for such instantons in the full database, since their existence does not require a massive $B - L$.

It is interesting to note that in the classes of MSSM-like models discussed earlier in this section with the closest to minimal zero mode structure, there are no instantons at all generating either R-parity violating or the $B - L$ dim=5 operators in the table. This makes them particularly attractive.

Note that all numbers in table 10 refer to the occurrence of instantons in the set of 391 tadpole-free models with massive B-L, but without checking the presence of zero-modes between the hidden sector and the instanton. It makes little sense to use the hidden sector in the database for such a check, since this is just one sample from a (usually) large number of possibilities. A meaningful question would be: can one find a hidden sector that has no zero-modes with the instanton. We have done such a search for the $B - L$ violating instantons (see the exclamation marks in the last column of table (5)), but not for the $B - L$ preserving instantons.

7 Conclusions and outlook

In this paper we have presented a systematic search for MSSM-like Type II Gepner orientifold models allowing for boundary states associated to instantons giving rise to neutrino Majorana masses. This search is very well motivated since neutrino masses are not easily accommodated in the semi-realistic compactifications constructed up to now. String instanton induced Majorana masses provides a novel and promising way to understand the origin of neutrino masses in the string theory context.

The string instantons under discussion are not gauge instantons. Thus, for example, they not only break $B + L$ symmetry (like 't Hooft instantons do) but also $B - L$, allowing for Majorana neutrino mass generation. The obtained mass terms are of order $M_s \exp(-V/g^2)$ but this suppression is unrelated to the exponential suppression of e.g. electroweak instantons and may be mild. In fact we find in our most interesting examples that the instanton action is typically substantially smaller than that of QCD or electroweak instantons, and hence these effects are much less suppressed than those coming from gauge theory instantons.

To perform our instanton search we have analyzed the structure of the zero modes that these instantons must have in order to induce the required superpotential. This analysis goes beyond the particular context of Gepner orientifolds and has general

validity for Type II CY orientifolds. We have found that instantons with $O(1)$ CP symmetry have the required universal sector of just two fermionic zero modes for the superpotential to be generated. Instantons with $Sp(2)$ and $U(1)$ CP symmetries have extra unwanted universal fermionic zero modes, which however may be lifted in a variety of ways in more general setups, as we discuss in the text. In fact we find in our search that around 98 % of the instantons with the correct structure of charged zero modes have $Sp(2)$ CP symmetry. Indeed, from a number of viewpoints the $Sp(2)$ instantons are specially interesting. The instantons we find with the simplest structure of fermionic zero modes are $Sp(2)$ instantons which are also the ones which are present more frequently in the MSSM-like class of Gepner constructions considered. They have also some interesting features from the phenomenological point of view. Indeed, due to the non-Abelian structure of the CP symmetry, the structure in flavor space of the neutrino Majorana masses factorizes. This makes that, irrespective of what particular compactification is considered, $Sp(2)$ instantons may easily lead to a hierarchical structure of neutrino masses. It would be important to further study the possible phenomenological applications of the present neutrino mass generating mechanism.

String instanton effects can also give rise to other B- or L-violating operators. Of particular interest is the dimension 5 Weinberg operator giving direct Majorana masses to the left-handed neutrinos. We find that in the most interesting cases, different instantons giving rise to the Weinberg operator and to ν_R Majorana masses are both simultaneously present. Which effect is the dominant one in the generation of the physical light neutrino masses depends on the values of the instanton actions and amplitudes as well as on the value of the string scale. Instantons may also generate $\text{dim} < 5$ operators violating R-parity. We find however that instantons inducing such operators are extremely rare, and in fact are completely absent in the Gepner models with the simplest $Sp(2)$ instantons inducing neutrino masses.

There are many avenues yet to be explored. It would be important to understand better the possible sources (moving in moduli space, addition of RR/NS backgrounds etc.) of uplifting for the extra uncharged fermionic zero modes in the most favoured $Sp(2)$ instantons. A second important question is that we have concentrated on checking the existence of instanton zero modes appropriate to generate neutrino masses; one should further check that the required couplings among the fermionic zero modes and the relevant 4d superfields (i.e. ν_R or $L\bar{H}$) are indeed present in each particular case. This is in principle possible in models with a known CFT description but could be difficult in practice for the Gepner models here described.

Instantons can also generate other superpotentials with interesting physical applications. One important example is the generation of a Higgs bilinear (i.e. a μ -term) in MSSM-like models [4, 3]. Thus, e.g., one could perform a systematic search for instantons (boundary states) generating a μ -term in the class of CFT Gepner orientifolds considered in the present article. Other possible application is the search for instantons inducing superpotential couplings involving only closed string moduli. The latter may be useful for the moduli-fixing problem, or for non-perturbative corrections to perturbatively allowed couplings [56].

Finally, it would be important to search for analogous instanton effects inducing neutrino masses in other string constructions (heterotic, M-theory etc.). A necessary condition is that the anomaly free $U(1)_{B-L}$ gauge boson should become massive due to a Stückelberg term.

The importance of neutrino masses in physics beyond the Standard Model is unquestionable. We have shown that string theory instantons provide an elegant and simple mechanism to implement them in semi-realistic MSSM-like string vacua, and a powerful constraint in model building. In our opinion, the conditions of the existence of appropriate instantons to generate neutrino masses should be an important guide in a search for a string description of the Standard Model.

Acknowledgements

We thank M. Bertolini, R. Blumenhagen, S. Franco, M. Frau, S. Kachru, E. Kiritsis, A. Lerda, D. Lüst, F. Marchesano, T. Weigand for useful discussions. A.M.U. thanks M. González for encouragement and support. The research of A.N. Schellekens was funded in part by program FP 57 of the Foundation for Fundamental Research of Matter (FOM), and Research Project FPA2005-05046 of de Ministerio de Educacion y Ciencia, Spain. The research by L.E. Ibáñez and A.M. Uranga has been supported by the European Commission under RTN European Programs MRTN-CT-2004-503369, MRTN-CT-2004-005105, by the CICYT (Spain), and the Comunidad de Madrid under project HEPHACOS P-ESP-00346.

Appendix

A CFT Notation

Here we summarize the labelling conventions for various CFT quantities. Further details and explanations can be found in [43].

It is important to keep in mind that there are four steps in the construction, each involving choices of some quantities. The steps are

- A CFT tensor product
- An extension of the chiral algebra of this tensor product
- The choice of a MIPF
- The choice of an orientifold

The second and third step are easily confused. A MIPF can itself be of extension type (although it can also be of automorphism or mixed type), meaning that it implies an extension of the chiral algebra. The crucial difference between step two and three in that case is that in step 2 all fields that are non-local with respect to the extension are projected out, and the symmetry of the extension is imposed on all states of the CFT, *i.e* in particular on all boundary states. The extension in step three acts as a bulk invariant, but the boundary states are not required to respect the symmetry implied by the extension.

Primary fields of $N = 2$ minimal models are labeled in the usual way by three integers (l, q, s) . In addition to these minimal models, one building block of our CFT's is of course a set of NSR fermions in four dimensions. They can be represented by the four conjugacy classes $(0), (v), (s), (c)$ analogous to those of a root lattice of type D.

Primary fields in a tensor product of M factors are therefore labelled as

$$I = ((x), (l_1, q_1, s_1), \dots, (l_M, q_M, s_M)) \tag{A.1}$$

where $x = 0, v, s$ or c .

This tensor product is extended by the alignment currents and the spin-1 field corresponding to the space-time supersymmetry generator. This organizes the tensor product fields into orbits, which can be labelled by one of the elements of the orbit. We always choose the field of minimal conformal weight (or one of them, in case there

are more) as the orbit representative labelling the orbit. The supersymmetry generator may have fixed points, leading to orbits appearing more than once as primary fields of the extended theory. In those cases we need an additional degeneracy label to distinguish them. It is convenient to choose for this label a character of the discrete group that is causing the degeneracy, the “untwisted stabilizer”, which depends on I . Denoting this character as Ψ_I we get then the following set of labels for the primaries of the extended CFT

$$i = [I, \Psi_I] \tag{A.2}$$

where I has the form (A.1). If there are no degeneracies we will leave out the square brackets and the Ψ_I .

In boundary CFT’s two new labels appear: the labels of Ishibashi-states that propagate in the transverse channel of the annulus, and the boundary labels. In the simplest, “Cardy” case both sets of labels are in one-to-one correspondence with the extended CFT labels i . But if we consider non-trivial MIPFs Z_{ij} both sets of labels are different. The Ishibashi states are in one-to-one correspondence with the fields i with $Z_{ii^c} \neq 0$. Degeneracies can occur here if $Z_{ii^c} \neq 0$. This requires the introduction of a degeneracy label. Such degeneracies may occur if the stabilizer of i (the set of simple currents that fix i) is non-trivial. It is convenient to use elements J of the stabilizer as degeneracy labels, so that the Ishibashi labels get the following form

$$m = (i, J) , \tag{A.3}$$

where i is an extended CFT label, as defined above (to be precise, in some cases a non-trivial degeneracy label is introduced even if $Z_{ii^c} = 1$. The details will not matter here).

Boundary states correspond to orbits of the simple current group \mathcal{H} that defines a MIPF. To label such orbits we choose a representative. There is no obvious canonical representative (one could use one of minimal conformal weight, but the conformal weight of orbit members of a boundary state does not play any rôle in the formalism, unlike the conformal weight of a primary). So in this case we just make an arbitrary choice. Once again there can be degeneracies. In this case they are due to a subgroup of the stabilizer called the “Central Stabilizer”. It is convenient to label the boundary states by an orbit representative i and a character ψ_i of the central stabilizer. If we expand the boundary state label in all of its components we get

$$a = [i, \psi_i] = [[I, \Psi_I], \psi_i] = [((x), (l_1, q_1, s_1), \dots, (l_M, q_M, s_M)), \Psi_I], \psi_i] . \tag{A.4}$$

Note that i is just a representative of a boundary orbit, and that I is just a representative of an orbit of the extension of the CFT.

B Instanton boundary symmetries

In the hidden-sector free example discussed in some detail in section 6 we have encountered $Sp(2)$ -type instantons, the most common kind in our scan. This particular model is the one that comes closest to the required zero mode count, although the only superfluous zero modes are rather awkward. Let us assume that the effect of these superfluous universal zero-modes instantons can be avoided. Then there is still another problem we have to face, namely that the two zero modes α^i and γ^i are related by an $Sp(2)$ transformation of the label i . Then we need at least three independent instantons (with unrelated couplings) to generate three non-zero neutrino masses, as discussed in Section 3.3. Since the technology to compute the couplings is not yet available, we cannot be completely sure that the relevant couplings are distinct, or indeed that they are non-vanishing, but at least we can inspect if there are obvious symmetries relating them.

The unextended tensor product $(2, 4, 22, 22)$ has 64 discrete symmetries: five separate charge conjugations of the factors (including the NSR space-time factor) and the interchange of the two identical $k = 22$ minimal models. To get space-time supersymmetry this tensor product is extended with the product of the simple current Ramond ground states of each factor. These Ramond ground states are not invariant under charge conjugation. Therefore this extension breaks the discrete symmetries to $\mathbf{Z}_2 \times \mathbf{Z}_2$, the combined charge conjugation of all five factors and the permutation of the two identical factors. The combined charge conjugation also acts non-trivially on the Ramond ground states in each factor, but the result is the charge conjugate of the space-time supersymmetry generator, which is in the chiral algebra. The combined conjugation is in fact the charge conjugation symmetry of the extended CFT. It turns out that only a \mathbf{Z}_2 subgroup of $\mathbf{Z}_2 \times \mathbf{Z}_2$ acts non-trivially on the simple currents of the extended CFT: the permutation of the $k = 22$ factors acts in the same way as charge conjugation. The action of these symmetries on the complete set of primary fields is more complicated. It is easy to see that the permutation acts differently than charge conjugation. In general the primary fields of the extended CFT are labelled as $i = [((x), (i_1), \dots, (i_4)), \Psi]$ (see Appendix A). The action of the permutation is to interchange i_3 and i_4 , but in cases with a non-trivial degeneracy it is not *a priori*

clear which of the degenerate states is the image of the map. This can be resolved by examining the fusion rules, which should be invariant under the permutation:

$$N_{ijk} = N_{\pi(i)\pi(j)\pi(k)} , \quad (\text{B.1})$$

where N is a fusion coefficient and π a permutation or other automorphism. In general there may be more than one way to resolve these ambiguities, resulting in additional automorphism of the CFT. The standard example of this situation is the extension of the affine algebra A_1 level 4 by the simple current. The resulting CFT has an outer automorphism, non-trivial charge conjugation, that has no counterpart in A_1 level 4.

As mentioned above, the \mathbf{Z}_2 permutation symmetry is respected by the MIPF and the orientifold, and since charge conjugation acts in the same way on the simple currents as the permutation, charge conjugation is respected as well.

In this way we end up with (at least) a surviving $\mathbf{Z}_2 \times \mathbf{Z}_2$ discrete symmetry acting on the boundary labels, or a larger discrete symmetry if that symmetry is extended by the action on the degeneracy labels of the extension. The foregoing story repeats itself for the action on the boundary labels. The boundary labels are given in terms of the CFT labels plus a second degeneracy label, the one indicated by the second entry in the square brackets in column 2 of tables (6) and (9). Once again one has to determine not only how a symmetry acts on the first entry (this is just the action of the symmetry in the extended CFT, respecting its fusion rules), but also how it acts on the degeneracy labels. In this case the precise action can be determined from the invariance of the annulus coefficients

$$A_{ab}^i = A_{\hat{\pi}(a)\hat{\pi}(b)}^{\pi(i)} , \quad (\text{B.2})$$

where π is the action on the primaries of the extended CFT (as determined above) and $\hat{\pi}$ is the action on the boundary labels induced by π .

Since the orientifold choice is non-trivial, boundary charge conjugation does not coincide with CFT charge conjugation. Indeed, the eight instanton boundary states are invariant under boundary charge conjugation (which they must be in order to produce a “real” $Sp(2)$ -type instanton). However, just as permutations, CFT charge conjugation may induce a non-trivial discrete symmetry on the boundary states.

In addition to these “outer automorphisms” there is the notion of boundary simple currents, introduced in the appendix of [6]. These may be thought of as remnants of the original simple currents, and imply relations between annulus amplitudes of the form

$$A_{ab'}^i = A_{J_a(Jb)'}^i \quad (\text{B.3})$$

All of the aforementioned symmetries might relate instanton couplings, and hence threaten their numerical independence. However, in order to do that they have to be symmetries of the full standard model/instanton configuration, not just relate some of the eight instantons to each other. It is easy to see that the permutation of the $k = 22$ factors changes the standard model brane configuration. Consider brane c : it turns out that under permutation boundary state 629 is mapped to boundary state 544 or 545 (depending on the action on the degeneracy label), which in any case is distinct. Hence even if the instanton boundaries 414 and 456 resp. 418 and 502 are mapped to each other by boundary permutation, at the same time the standard model configuration is mapped to a distinct one.

This means that we may expect at least four distinct couplings, which should be sufficient. It is of course possible to work out the discrete symmetries exactly, but in view of this argument this would not yield any additional insight.

We do know the exact boundary orbits. The orbit of instanton label 414 is (414, 415, 416, 417), so that instantons 414 and 417 are related. But the orbit of brane c under the same action is (629, 628, 626, 627). Hence the action that relates 414 and 417 maps 629 to 627. In fact all four standard model boundaries **a,b,c,d** are mapped to different ones. This implies that instantons 414 and 417 may produce different couplings as well, so that all eight instantons may contribute in a different way.

These distinctions concern the disk correlators $d_a^{(r)}$ in (3.12). The factors $\exp(-\text{Re } U_r)$ will be related by discrete symmetries, and it seems reasonable to expect them to be identical for instantons 414, 417, 456 and 459, which is indeed correct. However there is no reason to expect the other four instantons to have the same suppression factor, and indeed they do not.

Note that these symmetries imply the existence of a much larger set of standard model configurations than the 32 discussed here. However, as mentioned before, the 32 models considered here display all possible distinct spectra.

References

- [1] P. Minkowski, Phys.Lett.**B67**,421(1977); M. GellMann, P. Ramond and R. Slansky, in *Supergravity*, p.315, ed. P. Nieuwenhuizen and D. Friedman, (North Holland, Amsterdam, 1979); T. Yanagida, in *Proc. of the Workshop on Unified Theories and the Baryon Number of the Universe*, ed. O. Sawada and A. Sugamoto,

- (KEK, Japan, 1979); R.N. Mohapatra and G. Senjanovic, Phys.Rev.Lett.**44**, 912 (1980).
- [2] J. Giedt, G. L. Kane, P. Langacker and B. D. Nelson, Phys. Rev. D **71** (2005) 115013 [arXiv:hep-th/0502032].
- [3] L. E. Ibanez and A. M. Uranga, “Neutrino Majorana masses from string theory instanton effects,” arXiv:hep-th/0609213.
- [4] R. Blumenhagen, M. Cvetič and T. Weigand, “Spacetime instanton corrections in 4D string vacua - the seesaw mechanism for D-brane models,” arXiv:hep-th/0609191.
- [5] L. E. Ibáñez, F. Marchesano and R. Rabadan, JHEP **0111** (2001) 002 [arXiv:hep-th/0105155].
- [6] T. P. T. Dijkstra, L. R. Huiszoon and A. N. Schellekens, Nucl. Phys. B **710** (2005) 3, hep-th/0411129]
- [7] P. Anastasopoulos, T. P. T. Dijkstra, E. Kiritsis and A. N. Schellekens, “Orientifolds, hypercharge embeddings and the standard model,” hep-th/0605226, to appear in Nucl. Phys. B.
- [8] M. Cvetič, R. Richter, T. Weigand, “Computation of D-brane instanton induced superpotential couplings: Majorana masses from string theory”, arXiv:th/0703028.
- [9] R. Blumenhagen, L. Goerlich, B. Kors and D. Lust, JHEP **0010** (2000) 006 [arXiv:hep-th/0007024].
- [10] G. Aldazabal, S. Franco, L. E. Ibáñez, R. Rabadan and A. M. Uranga, J. Math. Phys. **42** (2001) 3103 [arXiv:hep-th/0011073]; JHEP **0102** (2001) 047 [arXiv:hep-ph/0011132].
- [11] A. M. Uranga, Class. Quant. Grav. **20** (2003) S373 [arXiv:hep-th/0301032]; R. Blumenhagen, M. Cvetič, P. Langacker and G. Shiu, Ann. Rev. Nucl. Part. Sci. **55** (2005) 71 [arXiv:hep-th/0502005]
- [12] C. Bachas, [arXiv:hep-th/9503030].

- [13] C. Angelantonj, I. Antoniadis, E. Dudas and A. Sagnotti, Phys. Lett. B **489** (2000) 223 [arXiv:hep-th/0007090].
- [14] E. Witten, Nucl. Phys. B **474** (1996) 343 [arXiv:hep-th/9604030].
- [15] S. Weinberg, “Varieties Of Baryon And Lepton Nonconservation,” Phys. Rev. D **22**, 1694 (1980).
- [16] R. Argurio, M. Bertolini, G. Ferretti, A. Lerda, C. Petersson, arXiv:hep-th/0704.0262
- [17] M. Bianchi, F. Fucito, J.F. Morales, arXiv:hep-th/07040784
- [18] A. Sagnotti, Phys. Lett. B **294** (1992) 196 [arXiv:hep-th/9210127].
- [19] L. E. Ibanez, R. Rabadan and A. M. Uranga, Nucl. Phys. B **542** (1999) 112 [arXiv:hep-th/9808139].
- [20] I. Antoniadis, E. Kiritsis and J. Rizos, Nucl. Phys. B **637** (2002) 92 [arXiv:hep-th/0204153].
- [21] G. 't Hooft, Phys. Rev. D **14** (1976) 3432 [Erratum-ibid. D **18** (1978) 2199].
- [22] D. Baumann, A. Dymarsky, I. R. Klebanov, J. Maldacena, L. McAllister and A. Murugan, JHEP **0611** (2006) 031 [arXiv:hep-th/0607050].
- [23] R. Blumenhagen, M. Cvetič and T. Weigand, [arXiv:hep-th/0609191].
- [24] M. Haack, D. Krefl, D. Lust, A. Van Proeyen and M. Zagermann, JHEP **0701** (2007) 078 [arXiv:hep-th/0609211].
- [25] B. Florea, S. Kachru, J. McGreevy and N. Saulina, arXiv:hep-th/0610003.
- [26] M. Bianchi, E. Kiritsis, “Non-perturbative and flux superpotentials for type I strings on the $Z(3)$ orbifold”, arXiv:hep-th/0702015.
- [27] O. J. Ganor, Nucl. Phys. B **499** (1997) 55 [arXiv:hep-th/9612077].
- [28] M. Billo, M. Frau, I. Pesando, F. Fucito, A. Lerda, A. Liccardo, “Classical gauge instantons from open strings”, JHEP **0302** (2003) 045. arXiv:hep-th/0211250.

- [29] L. Gorlich, S. Kachru, P. K. Tripathy and S. P. Trivedi, JHEP **0412** (2004) 074, [arXiv:hep-th/0407130];
P. G. Camara, L. E. Ibanez and A. M. Uranga, Nucl. Phys. B **708** (2005) 268, [arXiv:hep-th/0408036]; J. Park, arXiv:hep-th/0507091; R. Kallosh, A. K. Kashani-Poor and A. Tomasiello, JHEP **0506** (2005) 069, [arXiv:hep-th/0503138]. E. Bergshoeff, R. Kallosh, A. K. Kashani-Poor, D. Sorokin and A. Tomasiello, JHEP **0510** (2005) 102, [arXiv:hep-th/0507069].
- [30] K. Becker, M. Becker, C. Vafa, J. Walcher, “Moduli stabilization in non-geometric backgrounds”, arXiv:hep-th/0611001.
- [31] A. Sen, “F-theory and orientifolds”, Nucl. Phys. B **475** (1996) 562, arXiv:hep-th/9605150.
- [32] M. Bershadsky, A. Johansen, T. Pantev, V. Sadov and C. Vafa, Nucl. Phys. B **505** (1997) 153 [arXiv:hep-th/9612052].
- [33] B. S. Acharya, arXiv:hep-th/0011089.
- [34] N. Akerblom, R. Blumenhagen, D. Lust, E. Plauschinn and M. Schmidt-Sommerfeld, arXiv:hep-th/0612132.
- [35] C. Angelantonj, M. Bianchi, G. Pradisi, A. Sagnotti and Y. S. Stanev, “*Comments on Gepner models and type I vacua in string theory,*” Phys. Lett. B **387** (1996) 743 [ArXiv:hep-th/9607229].
- [36] R. Blumenhagen and A. Wisskirchen, “*Spectra of 4D, N = 1 type I string vacua on non-toroidal CY threefolds,*” Phys. Lett. B **438**, 52 (1998) [ArXiv:hep-th/9806131].
- [37] G. Aldazabal, E. C. Andres, M. Leston and C. Nunez, “*Type IIB orientifolds on Gepner points,*” JHEP **0309**, 067 (2003) [ArXiv:hep-th/0307183].
- [38] I. Brunner, K. Hori, K. Hosomichi and J. Walcher, “*Orientifolds of Gepner models,*” [ArXiv:hep-th/0401137].
- [39] R. Blumenhagen and T. Weigand, “*Chiral supersymmetric Gepner model orientifolds,*” JHEP **0402** (2004) 041 [ArXiv:hep-th/0401148].
- [40] T. P. T. Dijkstra, L. R. Huiszoon and A. N. Schellekens, Phys. Lett. B **609** (2005) 408 [arXiv:hep-th/0403196].

- [41] G. Aldazabal, E. C. Andres and J. E. Juknevich, “*Particle models from orientifolds at Gepner-orbifold points,*” JHEP **0405**, 054 (2004) [ArXiv:hep-th/0403262].
- [42] G. Aldazabal, E. Andres and J. E. Juknevich, JHEP **0607** (2006) 039 [arXiv:hep-th/0603217].
- [43] J. Fuchs, L. R. Huiszoon, A. N. Schellekens, C. Schweigert and J. Walcher, “*Boundaries, crosscaps and simple currents,*” Phys. Lett. B **495** (2000) 427 [ArXiv:hep-th/0007174].
- [44] G. Pradisi, A. Sagnotti, and Ya.S. Stanev, *The open descendants of non-diagonal $SU(2)$ Wess-Zumino-Witten models,* Phys. Lett. B 356 (1995) 230
- [45] G. Pradisi, A. Sagnotti, and Ya.S. Stanev, *Completeness conditions for boundary operators in 2D Conformal Field Theory,* Phys. Lett. B 381 (1996) 97
- [46] L. R. Huiszoon, A. N. Schellekens and N. Sousa, Phys. Lett. B **470**, 95 (1999) [arXiv:hep-th/9909114].
- [47] L. R. Huiszoon and A. N. Schellekens, Nucl. Phys. B **584**, 705 (2000) [arXiv:hep-th/0004100].
- [48] A. M. Uranga, “*D-brane probes, RR tadpole cancellation and K-theory charge,*” Nucl. Phys. B **598** (2001) 225 [ArXiv:hep-th/0011048].
- [49] B. Gato-Rivera and A. N. Schellekens, “*Remarks on global anomalies in RCFT orientifolds,*” Phys. Lett. B **632** (2006) 728 [ArXiv:hep-th/0510074].
- [50] E. Gimon, J. Polchinski, “Consistency conditions for orientifolds and D-manifolds”, Phys. Rev. D**54** (1996) 1667, arXiv:hep-th/9601038.
- [51] G. Pradisi and A. Sagnotti, Phys. Lett. B **216** (1989) 59.
- [52] E. Witten, Nucl. Phys. B **460** (1996) 541 [arXiv:hep-th/9511030].
- [53] B. Gato-Rivera and A. N. Schellekens, Commun. Math. Phys. **145**, 85 (1992).
- [54] B. Gato-Rivera and A. N. Schellekens, Nucl. Phys. B **353**, 519 (1991).
- [55] M. Kreuzer and A. N. Schellekens, Nucl. Phys. B **411**, 97 (1994) [arXiv:hep-th/9306145].
- [56] S.A. Abel, M.D. Goodsell, arXiv:hep-th/0612110.