

Insulin-releasing and insulin-like activity of the traditional anti-diabetic plant *Coriandrum sativum* (coriander)

Alison M. Gray* and Peter R. Flatt

School of Biomedical Sciences, University of Ulster, Coleraine BT52 1SA, UK

(Received 3 April 1998 – Revised 5 October 1998 – Accepted 8 October 1998)

Coriandrum sativum (coriander) has been documented as a traditional treatment of diabetes. In the present study, coriander incorporated into the diet (62.5 g/kg) and drinking water (2.5 g/l, prepared by 15 min decoction) reduced hyperglycaemia of streptozotocin-diabetic mice. An aqueous extract of coriander (1 mg/ml) increased 2-deoxyglucose transport (1.6-fold), glucose oxidation (1.4-fold) and incorporation of glucose into glycogen (1.7-fold) of isolated murine abdominal muscle comparable with 10^{-8} M-insulin. In acute 20 min tests, 0.25–10 mg/ml aqueous extract of coriander evoked a stepwise 1.3–5.7-fold stimulation of insulin secretion from a clonal B-cell line. This effect was abolished by 0.5 mM-diazoxide and prior exposure to extract did not alter subsequent stimulation of insulin secretion by 10 mM-L-alanine, thereby negating an effect due to detrimental cell damage. The effect of extract was potentiated by 16.7 mM-glucose and 10 mM-L-alanine but not by 1 mM-3-isobutyl-1-methylxanthine. Insulin secretion by hyperpolarized B-cells (16.7 mM-glucose, 25 mM-KCl) was further enhanced by the presence of extract. Activity of the extract was found to be heat stable, acetone soluble and unaltered by overnight exposure to acid (0.1 M-HCl) or dialysis to remove components with molecular mass < 2000 Da. Activity was reduced by overnight exposure to alkali (0.1 M-NaOH). Sequential extraction with solvents revealed insulin-releasing activity in hexane and water fractions indicating a possible cumulative effect of more than one extract constituent. These results demonstrate the presence of antihyperglycaemic, insulin-releasing and insulin-like activity in *Coriandrum sativum*.

Coriander: Insulin: Diabetes mellitus

Before the discovery of insulin in the early 1920s and the later development of oral hypoglycaemic agents, the major form of treatment of diabetes mellitus involved starvation, dietary manipulation and the use of plant therapies (Bailey & Flatt, 1990). More than 400 plants worldwide have been documented as beneficial in the treatment of diabetes (Bailey & Day, 1989; Swanston-Flatt *et al.* 1991; Gray & Flatt, 1997a). However, most of these await proper scientific or medical evaluation, and the World Health Organization (1980) has recommended accordingly that assessment of traditional plant treatments for diabetes merits further investigation.

Coriandrum sativum (coriander) has been reported to have a number of possible medicinal attributes including antispasmodic, carminative and stomachic properties (Lust, 1986). Additionally, coriander has been advocated as an anti-diabetic remedy (Farnsworth & Segelman, 1971; Lewis & Elvin-Lewis, 1977). More recent studies have confirmed the antihyperglycaemic effect of coriander in streptozotocin-diabetic mice (Swanston-Flatt *et al.* 1990), suggesting

that further studies are warranted on the antihyperglycaemic actions of this plant. Anti-diabetic agents can exert beneficial effects in the diabetic environment by improving and/or mimicking insulin action and/or by enhancing insulin secretion (Gray & Flatt, 1997b). To understand better the mechanisms by which coriander ameliorates hyperglycaemia, the present study investigated *in vitro* actions of aqueous extracts of coriander on glucose metabolism by isolated murine abdominal muscle and on insulin secretion by a clonal B-cell line (BRIN-BD11).

Materials and methods

Plant material

Dried coriander seeds were obtained from a commercial source in Birmingham, West Midlands, UK. Seeds were homogenized to a fine powder and stored at room temperature ($20 \pm 2^\circ$) in opaque screw-top jars. Powdered coriander was used for incorporation into test animal diet. An aqueous

Abbreviations: KRB, Krebs-Ringer bicarbonate buffer; STZ, streptozotocin.

* **Corresponding author:** Dr Alison Gray, present address: Young Hearts Project, Level 3 McKinney House, Musgrave Park Hospital, Belfast BT9 7JB, UK, fax +44 (0) 1232 382008, email youngheartsproject@hpsppop.n-i.nhs.uk

extract of coriander was prepared by a method of decoction as described previously (Gray & Flatt, 1998a). In brief, 1 g powdered material was placed in 40 ml cold (distilled) water, brought to the boil, then removed from the heat source and allowed to infuse for 15 min. This suspension was filtered (Whatman no. 1) and the volume readjusted to 40 ml with distilled water. For *in vivo* studies 10 ml portions of extract were stored at -20° until use when they were diluted tenfold with tap water (2.5 g/l). For *in vitro* studies 1 ml portions of extract were brought to dryness under vacuum (Savant speedvac; Savant Instrumentation Incorp., Framingdale, NY, USA), stored at -20° and reconstituted on the day of use with incubation buffer. Incubations within a single experiment using isolated muscle or BRIN-BD11 cells were always conducted using the same batch of extract. This allowed for any variation in potency of different batches of extract.

Animal studies

Heterozygous lean (*ob/+*) male mice derived from a colony maintained at Aston University, Birmingham, UK (Flatt & Bailey, 1981) were used at 21–24 weeks of age. Groups of two or three mice were housed in an air-conditioned room at $22 \pm 2^{\circ}$ with a lighting schedule of 12 h light (08.00–20.00 hours) and 12 h dark. Animals had free access to a standard pellet diet (Mouse Breeding Diet, Pillsbury Ltd, Birmingham, W. Midlands, UK) and tap water. The overall nutrient composition of the diet was (g/kg): carbohydrate 362, protein 209, fat 44 and fibre 385 with a metabolizable energy content of 11.8 MJ/kg. The experimental procedure for *in vivo* studies was similar to that previously described (Gray & Flatt, 1997b). For animal diets, coriander was incorporated into powdered mouse diet and mixed thoroughly, then distilled water was added and the mixture was combined to a stiff paste. The diet was then pelleted and placed at 45° until dry. Control diet was prepared by the same method to ensure there were no end differences in vitamin and mineral content as a result of the drying process. Coriander was incorporated into the diet (62.5 g/kg) and drinking water (2.5 g/l) of a group of seven mice 5 d before and after intraperitoneal administration of streptozotocin (STZ; Sigma Chemical Co., Poole, Dorset, UK) at 200 mg/kg body weight in 0.1 M-sodium citrate buffer (pH 4.5). Daily measures of body weight, food intake and fluid intake were made. Non-fasting blood samples obtained from the cut tail-tip of conscious mice were collected at the same time (09.00–10.00 hours) for plasma glucose analysis (Stevens, 1971). Groups of six normal mice and six STZ-treated mice with free access to unsupplemented diet and drinking water were used as controls.

Glucose transport and glucose metabolism *in vitro*

Recently-weaned non-fasting male mice (3–5 weeks) were killed by cervical dislocation and pieces of abdominal muscle (approximately 10–20 mg) were prepared. In order to replicate by mouse, pieces of muscle from each mouse were designated to each of the treatment groups. Glucose uptake was determined as described previously (Gray &

Flatt, 1998b). In brief, muscle pieces were incubated at 30° for 30 min in Krebs-Ringer bicarbonate buffer (KRB) supplemented with 2 mM-sodium pyruvate, 3.70 kBq/ml 2-deoxy-D-[1- 3 H]glucose, 0.37 kBq/ml L-[1- 14 C]glucose in the presence and absence of 10^{-8} M-human insulin and 1 mg/ml extract of coriander (2×2 factorial design). Oxidative glucose metabolism to CO_2 and incorporation of glucose into glycogen were determined as described previously (Gray & Flatt, 1998b). In brief, muscles were incubated at 37° for 60 min in KRB supplemented with 10 mM-glucose, 18.50 kBq/ml D-[U- 14 C]glucose in the presence and absence of 10^{-8} M-human insulin and 1 mg/ml extract of coriander (2×2 factorial design). Following incubation, CO_2 was captured onto a NaOH-saturated filter paper and muscles were removed for glycogen analysis (Gray & Flatt, 1998b).

Insulin secretion *in vitro*

BRIN-BD11 cells, produced by electrofusion of immortal RINm5F cell with New England Deaconess Hospital rat pancreatic B-cell, were used to evaluate insulin secretion (McClenaghan *et al.* 1996; Gray & Flatt, 1997b). In brief, wells containing 0.2×10^6 insulin-secreting cells were washed with KRB containing 1.1 mM-glucose and preincubated for 40 min at 37° (Gray & Flatt, 1997b). Unless otherwise stated, cells were then incubated for 20 min with KRB at 1.1 mM-glucose in the absence and presence of plant extract, diazoxide (an established opener of K^+ -ATP channels) and other test agents. Following incubation, portions were removed from each well and stored at -20° for insulin assay (Flatt & Bailey, 1981). Modified neutral red assay (Hunt *et al.* 1987) confirmed that 0.25–10 mg/ml coriander extract did not influence the viability of BRIN-BD11 cells during the test period.

To assess the importance of heat during extract preparation, aqueous extracts of coriander were prepared by the normal method of decoction (normal extract) or by cold infusion (cold extract; plant material placed in cold water, allowed to stand for 15 min, then filtered as before). Modified aqueous extract was freshly reconstituted in KRB and the effect on insulin secretion evaluated at a concentration equivalent to 1 mg/ml compared with normal extract (produced by 15 min decoction, as described previously). To further evaluate the nature of the insulin-releasing component(s), the normal aqueous extract of coriander prepared by decoction was subjected to heat, overnight dialysis, acid-alkali or acetone treatment. Heat: aqueous extract was boiled for 1 h immediately after preparation. Dialysis: aqueous extract was dialysed overnight (Spectra/Por molecular mass cut-off 2000 Da; Spectrum, Los Angeles, CA, USA) against deionized water (Milli-Q, Millipore Corp., Milford, MA, USA) at 4° . Acid-alkali treatment: portions of aqueous extract were added to 5 M-HCl or 5 M-NaOH to produce 0.1 M-HCl or 0.1 M-NaOH, allowed to stand at room temperature overnight, then neutralized. Acetone treatment: 1 ml aqueous extract (1 mg/ml) was added to 10 ml ice-cold acetone, allowed to stand for 30 min (on ice) and centrifuged (1000 rev./min; 5 min) to obtain acetone-soluble and acetone-insoluble fractions. Portions of untreated extract and modified aqueous extracts were dried under vacuum. All modified

aqueous extracts were freshly reconstituted in KRB and effects on insulin secretion at a concentration equivalent to 1 mg/ml were compared with that of untreated extract.

In another series of experiments ground coriander seeds were subjected to sequential extraction by increasingly polar solvents. Plant material (0.25 g) was placed in 5 ml hexane, agitated for 15 min and centrifuged (1200 rev./min; 5 min). The precipitate was dried under vacuum, extracted with a further 5 ml hexane and centrifuged as before. The extraction supernatant fractions were pooled, filtered (Whatman no. 1) and the volume adjusted to 10 ml hexane. The extraction precipitate (dried under vacuum) was subsequently extracted (as described earlier) with two 5 ml volumes of ethyl acetate, then methanol and finally with water. All extract fractions were brought to dryness under vacuum and freshly reconstituted in KRB for investigation of effects on insulin secretion at a concentration equivalent to 1 mg/ml compared with untreated extract.

Statistical analyses

Data were evaluated using Student's unpaired *t* test, one-way ANOVA or two-way ANOVA where appropriate. Groups were considered to be significantly different if $P < 0.05$. When a significant *F* value was obtained for ANOVA the differences between all pairs were tested using the Student-Newman-Keuls multiple comparisons test. If standard deviations were significantly different (Bartlett's test for homogeneity of variances) data were transformed ($\log_{10}[x]$).

Results

Studies in vivo

Compared with normal mice, STZ-treated mice showed significant ($P < 0.05$) weight loss, polydipsia and hyperglycaemia (Table 1). Administration of coriander in the diet and drinking water significantly decreased the hyperglycaemia by study day 20 (Table 1). Polydipsia increased in coriander supplemented *v.* unsupplemented STZ-treated mice (Table 1) and may indicate the presence of diuretic components in coriander seed. Plasma glucose concentrations of STZ-treated mice receiving coriander were comparable to those of normal mice at day 20 (Table 1). Throughout the study levels of food intake by treatment groups did not differ.

Glucose transport and glucose metabolism in vitro

Aqueous extract of coriander (1 mg/ml) increased glucose uptake (1.6-fold), glucose oxidation (1.4-fold), and glyco-genesis (1.7-fold) during incubations without insulin but did not significantly alter the stimulatory effect of 10^{-8} M-insulin (Table 2).

Insulin secretion in vitro

Aqueous extract of coriander (0.25–10 mg/ml) had a dose-dependent 1.3–5.7-fold stimulatory effect on insulin secretion by BRIN-BD11 cells at 1.1 mM-glucose (Fig. 1). It was confirmed that these concentrations of extract did not influence the viability of BRIN-BD11 cells during the test period as evaluated by modified neutral red assay (Hunt *et*

Table 1. Effects of coriander, administered in the diet (62.5 g/kg) and drinking water (2.5 g/l) on body weight, fluid intake, food intake and plasma glucose concentrations of streptozotocin (STZ)-treated mice†

(Mean values with their standard errors for six or seven mice)

	Normal mice		STZ mice		STZ mice + coriander	
	Mean	SEM	Mean	SEM	Mean	SEM
Body weight (g)						
Study day 0	45.5	2.90	42.3	0.94	43.9	2.44
Study day 12	47.2	1.12	37.9**	1.89	38.7*	1.57
Study day 20	48.1	1.19	37.7**	2.38	39.2**	1.59
Fluid intake (ml/d)						
Study day 0	4.5	0.20	4.3	0.24	5.0	0.61
Study day 12	5.5	0.38	5.5	0.26	7.0	0.48
Study day 20	5.2	0.31	10.0***	0.50	12.6***††	0.46
Food intake (g/d)						
Study day 0	4.4	0.49	4.4	0.43	4.4	0.49
Study day 12	4.1	0.19	3.5	0.20	4.1	0.19
Study day 20	5.2	0.40	4.9	0.40	5.0	0.40
Plasma glucose (mmol/l)						
Study day 12	6.6	0.45	13.5**	1.68	10.3*	1.20
Study day 20	9.9	0.74	22.3**	2.57	15.0†	2.20

Mean values were significantly different from those for normal mice: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Mean values were significantly different from those for STZ-treated mice receiving normal diet: † $P < 0.05$, †† $P < 0.01$.

‡ For details of diets and procedures, see pp. 203–205. STZ was administered 4 d after the introduction of coriander (day 0).

Table 2. Glucose uptake, oxidation and incorporation into glycogen by isolated mouse abdomen muscle incubated with aqueous extract of coriander in the presence or absence of 10^{-8} M-insulin
(Mean values with their standard errors for six or eleven samples)

	Control		Insulin (10^{-8} M)		Extract (1 mg/ml)		Insulin + extract	
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM
Glucose uptake (dpm/mg per h) (<i>n</i> 6)	244	37	439*	51	398*	44	422*	66
Glucose oxidation (nmol/mg per h) (<i>n</i> 11)	0.44	0.05	0.70*	0.04	0.63***	0.05	0.76***	0.05
Incorporation of glucose into glycogen (nmol/mg per h) (<i>n</i> 11)	0.23	0.03	0.47*	0.09	0.39*	0.06	0.42**	0.06

Mean values were significantly different from those for control incubations: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

al. 1987). The presence of 0.5 mM-diazoxide inhibited the stimulatory effect of the extract (Fig. 2(a)), indicating that the enhancement of insulin release was not a mere consequence of cellular damage. Consistent with this view, prior exposure of BRIN-BD11 cells to extract for 20 min did not alter the subsequent insulin secretory response to 10 mM-L-alanine (Fig. 2(b)). The insulin-releasing effect of 1 mg/ml extract was markedly potentiated by the presence of high (16.7 mM) glucose (Fig. 3(a)) and by the presence of 10 mM-L-alanine (Fig. 3(b)). However, the action of extract (1 mg/ml) was not potentiated by 1 mM-3-isobutyl-1-methyl-xanthine, which increases cyclic AMP in insulin-secreting cells (Sharp, 1979) (Fig. 3(c)). In contrast, aqueous extract of coriander further enhanced the insulin release by depolarized BRIN-BD11 cells (16.7 mM-glucose, 25 mM-KCl) (Fig. 3(d)).

Temperature of extraction did not alter the effect of the extract in enhancing insulin secretion (results not shown). Prolonged exposure to heat, overnight dialysis or exposure to acid did not significantly alter the insulin-releasing effect

of the plant extract (Table 3). Prolonged exposure to an alkaline environment reduced the insulin-enhancing effect (Table 3). The insulin-releasing activity was completely retained in the acetone-soluble fraction (Table 3). Both the hexane and water fractions of coriander (produced by sequential extraction) exerted an insulin-enhancing effect comparable with normal aqueous extract of coriander (Fig. 4). Although less potent than the normal aqueous extract, the ethyl acetate and methanol fractions of coriander (produced by sequential extraction) exerted insulin-releasing effects (Fig. 4).

Discussion

Early experiments involving administration of coriander fruit as a decoction did not reveal effects on fasting blood




Fig. 1. Effects of aqueous extract of coriander on insulin secretion by BRIN-BD11 cells. Values are means for groups of four to six observations with their standard errors indicated by vertical bars. Mean values were significantly different from those for control incubations: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.




Fig. 2. (a) Insulin secretion by BRIN-BD11 cells in response to 1.1 mM-glucose (□, control) and 1.1 mM-glucose plus aqueous extract of coriander (0.5 mg/ml) in the absence (□) or presence (■) of 0.5 mM-diazoxide. (b) Insulin secretion by BRIN-BD11 cells in response to 10 mM-L-alanine after 20 min prior exposure to 1.1 mM-glucose (□, control) or coriander extract (0.5 mg/ml) (■). Values are means for groups of six observations with their standard errors indicated by vertical bars. Mean values were significantly different from those for control incubations: * $P < 0.05$, *** $P < 0.001$. Mean values were significantly different from those for incubation with extract: †† $P < 0.01$.




Fig. 3. Insulin secretion by BRIN-BD11 cells incubated in the presence or absence of aqueous coriander extract (1 mg/ml) with various other substances. (a) (□), 1.1 mM-glucose (control) and (▨) 16.7 mM-glucose; (b) (□), 1.1 mM-glucose (control) and (▨), 10 mM-L-alanine; (c) (□), 1.1 mM-glucose (control) and (▨), 1 mM-3-isobutyl-1-methylxanthine (IBMX); (d) (□), 16.7 mM-glucose (control) and (▨), 25 mM-KCl. Values are means for groups of six observations with their standard errors indicated by vertical bars. Two-way ANOVA revealed: (a) extract effect ($P < 0.001$), glucose effect ($P < 0.001$) and extract \times glucose interaction ($P < 0.004$); (b) extract effect ($P < 0.001$), L-alanine effect ($P < 0.001$) and extract \times L-alanine interaction ($P = 0.021$); (c) extract effect ($P < 0.001$) and IBMX effect ($P < 0.001$); (d) extract effect ($P < 0.001$), KCl effect ($P < 0.001$) and extract \times KCl interaction ($P = 0.02$).

sugar levels of normal and alloxan diabetic rats, but demonstrated alleviation of adrenaline-induced hyperglycaemia (Sharaf *et al.* 1963). Subsequent studies involving longer-term administration of coriander seed in the diet showed that the plant purported as a traditional treatment for diabetes indeed decreased the hyperglycaemia of STZ-diabetic mice (Swanston-Flatt *et al.* 1990). The present study not only confirms this antihyperglycaemic action but indicates the presence of natural products in coriander which exhibit insulin-releasing and insulin-like actions. The possibility also exists that dietary administration of coriander affords protection against STZ-induced pancreatic B-cell destruction, but this requires further evaluation.

The possibility that part of the glucose lowering action of coriander may be due to an effect on glucose handling by skeletal muscle was evaluated *in vitro* using an insulin-sensitive mouse skeletal muscle preparation (Gray & Flatt,

1998b). These experiments revealed that an aqueous extract of coriander enhanced glucose transport, glucose oxidation and glycogenesis to an extent comparable with 10^{-8} M-insulin. The combined actions of coriander and insulin on muscle were not additive, suggesting that the active constituent(s) of coriander extract acts through a pathway similar in part to that utilized by insulin. Interestingly, the action of coriander differs significantly from that of the established antihyperglycaemic drug, metformin, which exerts effects on glucose transport via insulin-mediated peripheral glucose uptake (Bailey & Puah, 1986; Prager *et al.* 1986).

Experiments using clonal pancreatic B-cells showed that coriander extract stimulated insulin secretion in a dose-dependent fashion over the concentration range 0.25–10 mg/ml. The maximal fivefold insulin response can be compared with an approximate doubling of insulin release in this cell line induced by micromolar concentrations of




Fig. 4. Insulin secretion by BRIN-BD11 cells in response to extracts of coriander. Coriander was subjected to sequential extraction by solvents with increasing polarity; all extracts were tested at concentrations equivalent to 1 mg/ml. Values are means for groups of six observations with their standard errors represented by vertical bars. Mean values were significantly different from those for incubations without extract: ** $P < 0.01$, *** $P < 0.001$. Mean values were significantly different from those for incubations with normal extract (produced by 15 min decoction): ††† $P < 0.001$.

Table 3. Effect of heat, dialysis, acid-alkali treatment and acetone treatment on ability of aqueous extract of coriander to enhance insulin secretion by BRIN-BD11 cells‡

(Mean values with their standard errors for six observations)

Test	Insulin secretion (ng/10 ⁶ cells per 20 min)	
	Mean	SEM
Control (without extract)	1.57	0.06
Untreated extract	2.87***	0.16
Boiled extract	2.78**	0.38
Dialysed extract	2.43***	0.16
Acid-exposed extract	2.36**	0.23
Alkali-exposed extract	2.08*††	0.16
Acetone-insoluble extract	1.44†††	0.08
Acetone-soluble extract	2.76***	0.16

Mean values were significantly different from that for the control incubation: * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Mean values were significantly different from that for the untreated extract (produced by 15 min decoction): †† $P < 0.01$, ††† $P < 0.001$.

‡ Normal extract was subject to heat, overnight dialysis, acid-alkali or acetone treatment. For details of procedures, see pp. 203–205.

sulfonylureas, tolbutamide and glibenclamide (McClenaghan *et al.* 1998; McClenaghan & Flatt, 1998). The stimulatory action of coriander extract was also glucose-dependent and potentiated by the established amino acid stimulator L-alanine. However, insulin release was not enhanced by the phosphodiesterase inhibitor 3-isobutyl-1-methylxanthine which increases cyclic AMP and enhances hormone-mediated insulin release (Sharp, 1979). Importantly, prior exposure to 0.5 mg/ml extract did not diminish the subsequent insulin secretory response to L-alanine, thereby negating a possible deleterious effect of extract on cell viability in the insulin secretion experiments. The non-toxic nature of coriander extract was also confirmed by toxicity testing using the modified neutral red assay.

Studies performed to evaluate further the possible mechanisms underlying the insulin-releasing action of coriander extract confirmed an action similar to that of sulfonylureas. These agents act by binding to sulfonylurea receptors, resulting in closure of plasma membrane K⁺-ATP channels, membrane depolarization, opening of voltage-dependent Ca channels and elevation of intracellular Ca²⁺ (Rorsman, 1997). Diazoxide, by holding open K⁺-ATP channels, inhibits the stimulatory action of sulfonylureas (Dunne *et al.* 1994). In the present study, diazoxide also inhibited the insulin-releasing action of coriander. Like sulfonylureas (Flatt *et al.* 1994; Eliasson *et al.* 1996) coriander also stimulated insulin release from chemically depolarized cells incubated with 25 mM-KCl. This suggests that constituents of coriander extract have K⁺-ATP channel and distal actions similar to sulfonylureas. However, the greater potency of coriander extract compared with maximally effective concentrations of sulfonylureas suggests possible additional actions on B-cell nutrient metabolism or second-messenger pathways.

Preliminary investigations into the nature of the insulin-releasing agent(s) in coriander indicate heat stability, acetone solubility and resistance to acidic or alkaline environments. Extract activity was not due to smaller molecules or ions as suggested by retention of potency following dialysis to remove components of low molecular mass (<2000 Da). Studies using sequential solvent extractions of coriander seeds point towards cumulative insulin-releasing effects of more than one active constituent of coriander. The involvement of these or other extract components in mediating the effects of coriander on glucose uptake and metabolism clearly merit further investigation.

In conclusion, the present study has shown that the antihyperglycaemic action of coriander is associated with stimulation of insulin secretion and enhancement of glucose uptake and metabolism by muscle, reflecting the effects of more than one active constituent. *Coriandrum sativum* therefore represents a possible antihyperglycaemic dietary adjunct and potential source of a new orally active agent(s) for diabetes therapy.

Acknowledgements

These studies were supported by the Department of Health and Social Services, Northern Ireland and NI Developmental Research funding.

References

- Bailey CJ & Day C (1989) Traditional plant medicines as treatments for diabetes. *Diabetes Care* **12**, 553–564.
- Bailey CJ & Flatt PR (editors) (1990) *New Antidiabetic Drugs*, pp. 1–295. London: Smith-Gordon.
- Bailey CJ & Puaah JA (1986) Effect of metformin on glucose metabolism in mouse soleus muscle. *Diabète et Metabolisme* **12**, 212–218.
- Dunne MJ, Harding EA, Jaggar JH, Ayton BJ & Squires PE (1994) Endogenous and chemical activators of ATP-regulated potassium channels in insulin-secreting cells: possible mechanisms and physiological significance. In *Frontiers of Insulin Secretion and Pancreatic B-Cell Research*, pp. 153–159 [PR Flatt and S Lenzen, editors]. London: Smith-Gordon.
- Eliasson L, Renström E, Åmmälä C, Berggren P-O, Bertorello AM, Bokvist K, Chibalin A, Deeney JT, Flatt PR, Gabel J, Gromada J, Larsson O, Lindström P, Rhodes CJ & Rorsman P (1996) PKC-Dependent stimulation of exocytosis by sulfonylureas in pancreatic B-cells. *Science* **271**, 813–815.
- Farnsworth NR & Segelman AB (1971) Hypoglycaemic plants. *Tile and Till* **57**, 52–55.
- Flatt PR & Bailey CJ (1981) Abnormal plasma glucose and insulin responses in heterozygous lean (*ob/+*) mice. *Diabetologia* **20**, 573–577.
- Flatt PR, Shibir O, Szczowka J & Berggren P-O (1994) New perspectives on the actions of sulphonylureas and hyperglycaemic sulphonamides on the pancreatic B-cell. *Diabète et Metabolisme* **20**, 157–162.
- Gray AM & Flatt PR (1997a) Nature's own pharmacy: the diabetes perspective. *Proceedings of the Nutrition Society* **56**, 507–517.
- Gray AM & Flatt PR (1997b) Pancreatic and extra-pancreatic effects of the traditional anti-diabetic plant, *Medicago sativa* (lucerne). *British Journal of Nutrition* **78**, 325–334.
- Gray AM & Flatt PR (1998a) Actions of the traditional anti-diabetic plant, *Agrimony eupatoria* (agrimony): effects on hyperglycaemia, cellular glucose metabolism and insulin secretion. *British Journal of Nutrition* **80**, 109–114.
- Gray AM & Flatt PR (1998b) Insulin-releasing and insulin-like activity of *Agaricus campestris* (mushroom). *Journal of Endocrinology* **157**, 259–266.
- Hunt SM, Chrzanowska C, Barnett CR, Brand HN & Fawell JK (1987) A comparison of *in vitro* cytotoxicity assays and their application to water samples. *Alternatives to Laboratory Animals* **15**, 20–29.
- Lewis WH & Elvin-Lewis MPF (1977) *Medical Botany: Plants Affecting Man's Health*. New York, NY: Wiley.
- Lust J (1986) *The Herb Book*. London: Bantam Books.
- McClenaghan NH, Barnet CR, Ah-Sing E, Abdelwahab YHA, O'Harte FPM, Yoon T-W, Swanston-Flatt SK & Flatt PR (1996) Characterization of a novel glucose-responsive insulin-secreting cell line, BRIN-BD11, produced by electrofusion. *Diabetes* **45**, 1132–1140.
- McClenaghan NH & Flatt PR (1998) Engineering cultured insulin-secreting pancreatic B-cell lines. *Journal of Molecular Medicine* (In the Press).
- McClenaghan NH, Flatt PR & Bailey CJ (1998) Insulin-releasing action of the novel antidiabetic agent BTS 67 582. *British Journal of Pharmacology* **123**, 400–404.
- Prager R, Scherthaser G & Graf H (1986) Effect of metformin on peripheral insulin sensitivity in non-insulin dependent diabetes mellitus. *Diabète et Metabolisme* **12**, 346–350.
- Rorsman P (1997) The pancreatic B-cell as a fuel sensor: an electrophysiologist's view point. *Diabetologia* **40**, 487–495.
- Sharaf AA, Hussein AM & Mansour MY (1963) Studies on the antidiabetic effects of some plants. *Planta Medica* **11**, 159–168.
- Sharp GWG (1979) The adenylate cyclase–cyclic AMP system in islets of Langerhans and its role in the control of insulin release. *Diabetologia* **16**, 287–297.
- Stevens JF (1971) Determination of glucose by automatic analyser. *Clinica Chimica Acta* **32**, 199–201.
- Swanston-Flatt SK, Day C, Bailey CJ & Flatt PR (1990) Traditional plant treatments for diabetes: studies in normal and streptozotocin diabetic mice. *Diabetologia* **33**, 462–464.
- Swanston-Flatt SK, Flatt PR, Day C & Bailey CJ (1991) Traditional dietary adjuncts for the treatment of diabetes mellitus. *Proceedings of the Nutrition Society* **50**, 641–651.
- World Health Organization (1980) *World Health Organization Expert Committee on Diabetes Mellitus, Second Report. Technical Report Series* no. 646, p. 66. Geneva: WHO.

A leading
international
journal
publishing
authoritative,
timely and
in-depth
reviews...

Nutrition Research Reviews

General Editor
JM Forbes (UK)

Editors
JDJ Millward (UK)
HJ Powers (UK)
DI Thurnham (UK)
P Yaqoob (UK)

International Editors
J Black (Australia)
L Hambræus (Sweden)
J Okumura (Japan)
NW Solomons (Guatemala)
CM Weaver (USA)

 **CABI Publishing**
A division of CAB INTERNATIONAL

publishes *Nutrition Research Reviews*
on behalf of The Nutrition Society

Contributions by international experts:

- advance new concepts
- encourage fresh thinking
- highlight timely reports on important techniques and methods in experimental work
- provide thorough overviews on specific subject areas
- promote scientific communications on new ideas

Reviews of large-scale, international projects and surveys such as the EURESTA project feature prominently in *Nutrition Research Reviews*.

Leading nutritionists look back on a lifetime of involvement in important developments in nutrition research.

Forthcoming papers include:

- Homocysteine as a risk factor for cardiovascular and related disease: nutritional implications
- Irritable bowel syndrome: the interactions of diet and stress
- Perspectives on ruminant nutrition and metabolism
- Lipids and infant formulas

1999, Volume 12
Bi-annual publication (*June and December*)
ISSN: 0954 4224
£100.00 (US\$180.00 Americas only)

To order your 1999 subscription to this important journal contact:

CABI Publishing
CAB International
Wallingford, Oxon, OX10 8DE, UK
Tel: +44 (0) 1491 832111
Fax: +44 (0) 1491 829292
Email: publishing@cabi.org

CABI Publishing
CAB International
10 East 40th Street, Suite 3203,
New York, NY 10016, USA
Tel: +1 (212) 481 7018
Toll-free: 1 800 528 4841
Fax: +1 (212) 686 7993
Email: cabi-nao@cabi.org

Available on the Internet in 1999
<http://nutrition.cabweb.org>