

SHORT REPORT

Intellectual disability without epilepsy associated with *STXBP1* disruption

Fadi F Hamdan^{1,2}, Julie Gauthier^{1,3}, Sylvia Dobrzaniecka^{1,3}, Anne Lortie², Laurent Mottron^{1,4}, Michel Vanasse², Guy D'Anjou², Jean Claude Lacaille⁵, Guy A Rouleau^{1,3} and Jacques L Michaud^{*,1,2}

STXBP1 (Munc18-1) is a component of the machinery involved in the fusion of secretory vesicles to the presynaptic membrane for the release of neurotransmitters. *De novo* missense mutations in *STXBP1* were recently reported in patients with Ohtahara syndrome, a form of encephalopathy with severe early-onset epilepsy. In addition, sequencing of the coding region of *STXBP1* in 95 patients with non-syndromic intellectual disability (NSID) revealed *de novo* truncating mutations in two patients who also showed severe non-specific epilepsy, suggesting that *STXBP1* disruption has the potential of causing a wide spectrum of epileptic disorders in association with intellectual disability. Here, we report on the mutational screening of *STXBP1* in a different series of 50 patients with NSID and the identification of a novel *de novo* truncating mutation (c.1206delT/ p.Y402X) in a male with NSID, but surprisingly with no history of epilepsy. This is the first report of a patient with a truncating mutation in *STXBP1* that does not show epilepsy, thus, expanding the clinical spectrum associated with *STXBP1* disruption.

European Journal of Human Genetics (2011) 19, 607–609; doi:10.1038/ejhg.2010.183; published online 2 March 2011

Keywords: *STXBP1*; epilepsy; intellectual deficiency; mental retardation

INTRODUCTION

A growing body of work suggests that disruption of synapse development or function explains a large fraction of patients with intellectual disability (ID; also referred as mental retardation).^{1,2} One of the key events of synaptic transmission is the fusion of secretory vesicles to the presynaptic membrane leading to the release of the neurotransmitters into the synaptic cleft. This process is mediated by a complex composed of SNARE proteins and of Munc18-1/*STXBP1*.^{3–5} Consistent with its important role for neurotransmission, recent studies showed that mutations in *STXBP1* cause severe ID and epilepsy.^{6,7} Using array genomic hybridization, Saitsu *et al*⁶ identified a *de novo* 2 Mb deletion encompassing *STXBP1* in a patient with Ohtahara syndrome, an infantile form of epileptic encephalopathy characterized by burst suppressions and severe ID. Analysis of candidate genes mapped to the deletion revealed missense mutations in *STXBP1* in four other patients with Ohtahara syndrome. These mutations were found to arise *de novo* in at least three of these cases and to disrupt *STXBP1* biochemical properties.⁶ Subsequently, our group reported on the identification of *de novo* truncating mutations in *STXBP1* in two patients with severe non-syndromic ID (NSID) and intractable partial complex epilepsy.⁷ In order to better understand the contribution of *STXBP1* mutations to NSID, we studied 50 additional patients and identified a novel *de novo* *STXBP1* truncating mutation in a patient with NSID, but surprisingly with no history of epilepsy. This observation indicates that *STXBP1* mutations are associated with a

wide spectrum of phenotypes, from NSID with or without epilepsy to syndromic forms of epilepsy.

METHODS

NSID subjects

We recruited 50 sporadic cases (24 males and 26 females) with unexplained NSID, the majority of which were of French Canadian origin. We selected sporadic cases to increase the likelihood of finding *de novo* mutation. In the context of this study, and of our previous one,⁷ NSID was defined using the following criteria: (1) diagnosis of ID established on a clinical basis using standardized developmental or IQ tests; (2) absence of specific dysmorphic features, as assessed by an experienced clinical geneticist; (3) birth weight and postnatal growth within normal limits; (4) normal head circumference at birth; (5) absence of risk factors such as neonatal asphyxia, prematurity or exposure to teratogenic drugs and (6) negative standard investigations, including comparative genomic hybridization studies, molecular testing for the common expansion mutation in *FMRI* and brain CT-scan or MRI. A subset of these NSID patients ($n=11$) displayed epilepsy. Blood samples were collected from all members of the NSID cohort, as well as from their parents after approval by the Sainte Justine Hospital Research Center Medical Ethics Committee for genetic studies. Written consent forms were obtained from all guardians (parents) of the patients participating in this study.

STXBP1 sequencing

Genomic DNA was extracted from blood samples using the Puregene DNA kit (Gentra System; Qiagen, Mississauga, ON, Canada). Paternity and maternity of each NSID patient were confirmed using six informative unlinked microsatellite

¹The Centre of Excellence in Neuromics of Université de Montréal and Synapse to Disease (S2D) group, Montréal, Quebec, Canada; ²Research Center CHU Sainte-Justine, Montréal, Quebec, Canada; ³CHUM Research Center and Department of Medicine, Université de Montréal, Montreal, Canada; ⁴Centre de Recherche Fernand-Séguin, Hôpital Rivière-des-Prairies, Montréal, Quebec, Canada; ⁵Le Groupe de Recherche sur le Système Nerveux Central, Department of Physiology, Université de Montréal, Montreal, Quebec, Canada

*Correspondence: Dr JL Michaud, Research Center, CHU Sainte-Justine, 3175 Côte Sainte-Catherine, Montreal, Quebec, Canada H3T 1C5. Tel: +1 514 345 4931, ext: 6900; Fax: +1 514 345 4766; E-mail: jacques.michaud@recherche-ste-justine.qc.ca

Received 1 June 2010; revised 6 September 2010; accepted 7 October 2010; published online 2 March 2011

markers. PCR amplification and sequencing of all of the *STXBPI* exons (19 in total) and intronic splice junctions based on the structure of the longest isoform (Refseq no. NM_001032221) were performed as previously described.⁷

RESULTS

We sequenced all *STXBPI* exons and splice junction boundaries from the genomic DNA of 50 patients with NSID. We identified a heterozygous 1-bp deletion that creates a premature stop codon at amino acid position 402 (c.1206delT/p.Y402X) in one of the patients. This mutation was not identified in the DNA of both parents, indicating that it originated *de novo* (Figure 1). It lies in domain-3 of STXBPI, in close proximity to a previously reported *de novo* truncating mutation (p.R388X) identified by our group in a patient with NSID and severe epilepsy (Figure 1).⁷ Domain-3 together with domain-1 of STXBPI form the central cavity providing surfaces for Syntaxin-1 binding, an essential step for SNARE complex assembly and subsequent neurotransmitter release.^{3,8} No other amino acid altering or splicing mutations were identified in *STXBPI* in the remaining patients. We previously sequenced *STXBPI* in healthy French Canadian controls ($n=190$) and did not identify any amino acid changing or splicing mutations.⁷

The patient with the c.1206delT (referred to as patient 3) is a man, aged 21 years, born to non-consanguineous French Canadian parents with no family history of developmental disabilities (Table 1). He was delivered at term after an unremarkable pregnancy. Immediate neonatal course was uneventful. His psychomotor development was characterized by global delay. He first walked at 2 years of age. He could only say 20 words at 4 years of age. Cognitive evaluation with the Leiter cognitive test was not possible at that age because he did not understand the tasks. Another attempt to evaluate his cognitive profile with the Leiter and the WPPSI-R tests also failed at 5 years and

9 months for the same reason. Currently, he can use short and incomplete sentences to communicate his needs. He can evoke events of the past, but he cannot have a conversation. He can dress by himself, use tools and play simple video games but he cannot draw, cut his meals or tie his shoes. He is treated with methylphenidate for attention disorder without hyperactivity or impulsivity. There is no history of seizures. Physical examination did not reveal any specific dysmorphic features. Neurological exam showed the presence of diffuse tremor of the extremities that was increased upon voluntary movements, likely contributing to his poor fine motor skills. His tonus is normal. There was no dysmetria or ataxia. He walks with an increased polygon of sustentation without retropulsions. Electroencephalography performed at 21 years of age did not reveal any abnormality except for intermittent slow dysfunction in the left temporal area. Brain CT scan performed at 4 years of age was normal.

DISCUSSION

We report here, a novel *de novo* truncating mutation, c.1206delT/p.402X, in a NSID patient with severe cognitive deficit, but with no history of epilepsy. This mutation is likely to be pathogenic as it is predicted to truncate STXBPI towards the middle of its sequence, in the same functional domain as p.R388X, which was recently reported in a patient with severe NSID and non-syndromic epilepsy.⁷ This conclusion is further supported by the observations that truncation of the *Caenorhabditis elegans* orthologue of STXBPI downstream of position p.Y402 results in defects in synaptic vesicle docking⁹ and that *Stxbp1* haploinsufficiency causes impaired neurotransmission in mice.¹⁰

All seven known patients with pathogenic point mutations in *STXBPI* show severe ID. Four of them, all carrying missense




Figure 1 Pathogenic *STXBPI* mutations. (a) Positions of the reported pathogenic *STXBPI* mutations. Missense mutations reported to cause Ohatahira syndrome⁶ are indicated on top of the *STXBPI* sequence, while splicing and truncating mutations reported in NSID with nonspecific epilepsy⁷ (c.169+1G>A and p.R388X) or without epilepsy (current study, p.Y402X, highlighted) are shown below the *STXBPI* sequence. The positions of the various *STXBPI* protein domains are depicted based on the rat *STXBPI* crystal structure.⁸ (b) Chromatograms of the *STXBPI* *de novo* mutation identified in patient 3 and the corresponding representative sequence from parents of patient 3 (both the mother's and father's *STXBPI* sequences lack c.1206delT mutation). Wild-type (WT) and mutant (MT) *STXBPI* DNA sequences are shown along with their corresponding amino acids. Numbering of the coding nucleotides and amino acids are based on isoform a of *STXBPI* (Refseq NM_003165).

Table 1 Clinical profiles of patients with *de novo* truncating *STXBP1* mutations

Description	Patient 1 Hamdan <i>et al</i> (2009) ⁷	Patient 2 Hamdan <i>et al</i> (2009) ⁷	Patient 3 current study
<i>De novo</i> mutation	c.169+1G>A/p.157NfsX7	c.1162C>T/p.R388X	c.1206delT/p.Y402X
Age (years) when studied	27	15	21
Gender	Female	Female	Male
Ethnic origin	FC	FC	FC
ID	Severe	Severe	Severe
<i>Epilepsy</i>	+	+	–
Age at first seizures	6 weeks	2 years 9 months	NA
Initial type of seizures	Partial complex	Partial complex	NA
Initial electroencephalogram	Focalized	Focalized	Normal
<i>Neurological examination</i>			
Head circumference (centile)	56 cm (75th)	55.7 cm (50–75th)	58 cm (90–97th)
Hypotonia	+	+	–
Tremor	+	+	+
Hyperventilation	+	–	–
<i>Brain imaging</i>			
MRI	ND	Normal	ND
CT-scan	Normal	Normal	Normal

Abbreviations: FC, French Canadian; NA, not applicable; ND, not determined.

mutations, presented with Ohtahara syndrome and, with one exception, developed infantile spasms,⁶ whereas two other patients, bearing truncating mutations, show severe NSID with non-syndromic intractable partial complex epilepsy,⁷ indicating that mutations in *STXBP1* can result in different types of epileptic manifestations. The case described here shows that *STXBP1* disruption can also cause severe ID without epilepsy. Additional clinical clues suggesting *STXBP1* mutations include the presence of tremor, hypotonia and hyperventilation, which were observed in a subset of patients (Table 1).^{6,7}

Although the small number of patients identified at this point in time precludes drawing definitive correlations between genotypes and phenotypes, mutations leading to a decrease of *STXBP1* function appear to be associated with variable expressivity. First, different genetic lesions in *STXBP1* can result in the same phenotype. For instance, although missense mutations were exclusively identified in patients with Ohtahara syndrome, another patient with this condition was found to carry a *de novo* 2Mb deletion encompassing *STXBP1*.⁶ Second, similar genetic lesions in *STXBP1* can lead to different phenotypes, as illustrated by the variable presence of severe epilepsy in our patients with truncating mutations (Table 1).

By combining this study and our previous one, we have identified three *de novo* truncating *STXBP1* mutations in 145 sporadic NSID cases (~2%). This high rate of deleterious mutations suggests that analysis of *STXBP1* should be considered in sporadic cases of severe NSID, with or without epilepsy.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGEMENTS

We thank the patients and their parents for participating to this study. We are grateful for the dedicated work of members of the S2D team (CHUM Research

Center, Montreal), including management (Claude Marineau and Ronald G Lafrenière), bioinformatics (Edouard Henrion, Ousmane Diallo and Dan Spiegelman) and genetic screening divisions (Amelie Piton, Annie Raymond, Annie Levert, Pascale Thibodeau, Sandra Laurent and Karine Lachapelle). We are also thankful for the McGill University and Genome Québec Innovation Centre Sequencing (Pierre Lepage, Sébastien Brunet and Hao Fan Yam) and Bioinformatics (Louis Létourneau and Louis Dumond Joseph) groups. Supported by grants from the Canadian Institute of Health Research (CIHR; to JLM, GAR and JCL), Réseau de Génétique Médicale Appliquée/Fonds de la Recherche en Santé (FRSQ; to JLM), Genome Canada and Genome Quebec and co-funding by Université de Montréal for the S2D project (to GAR). JLM is a recipient of a Clinical Investigatorship Award of the CIHR (Institute of Genetics) and of a Senior Scientist Award from the FRSQ.

- Chelly J, Khelifaoui M, Francis F, Cherif B, Bienvu T: Genetics and pathophysiology of mental retardation. *Eur J Hum Genet* 2006; **14**: 701–713.
- Humeau Y, Gambino F, Chelly J, Vitale N: X-linked mental retardation: focus on synaptic function and plasticity. *J Neurochem* 2009; **109**: 1–14.
- Burkhardt P, Hattendorf DA, Weis WI, Fasshauer D: Munc18a controls SNARE assembly through its interaction with the syntaxin N-peptide. *Embo J* 2008; **27**: 923–933.
- Fisher RJ, Pevsner J, Burgoyne RD: Control of fusion pore dynamics during exocytosis by Munc18. *Science* 2001; **291**: 875–878.
- Gerber SH, Rah JC, Min SW *et al*: Conformational Switch of Syntaxin-1 Controls Synaptic Vesicle Fusion. *Science* 2008; **321**: 1507–1510.
- Saito H, Kato M, Mizuguchi T *et al*: *De novo* mutations in the gene encoding STXBP1 (MUNC18-1) cause early infantile epileptic encephalopathy. *Nat Genet* 2008; **40**: 782–788.
- Hamdan FF, Piton A, Gauthier J *et al*: *De novo* STXBP1 mutations in mental retardation and nonsyndromic epilepsy. *Ann Neurol* 2009; **65**: 748–753.
- Misura KM, Scheller RH, Weis WI: Three-dimensional structure of the neuronal-Sec1-syntaxin 1a complex. *Nature* 2000; **404**: 355–362.
- Weimer RM, Richmond JE, Davis WS, Hadwiger G, Nonet ML, Jorgensen EM: Defects in synaptic vesicle docking in unc-18 mutants. *Nat Neurosci* 2003; **6**: 1023–1030.
- Toonen RF, Wierda K, Sons MS *et al*: Munc18-1 expression levels control synapse recovery by regulating readily releasable pool size. *Proc Natl Acad Sci USA* 2006; **103**: 18332–18337.