

Interaction of Pregnancy-Specific Glycoprotein 1 With Integrin A5 β 1 Is a Modulator of Extravillous Trophoblast Functions

RATTILA, Shemona, *et al.*

Abstract

Human pregnancy-specific glycoproteins (PSGs) serve immunomodulatory and pro-angiogenic functions during pregnancy and are mainly expressed by syncytiotrophoblast cells. While PSG mRNA expression in extravillous trophoblasts (EVTs) was reported, the proteins were not previously detected. By immunohistochemistry and immunoblotting, we show that PSGs are expressed by invasive EVT_s and co-localize with integrin α 5. In addition, we determined that native and recombinant PSG1, the most highly expressed member of the family, binds to α 5 β 1 and induces the formation of focal adhesion structures resulting in adhesion of primary EVT_s and EVT-like cell lines under 21% oxygen and 1% oxygen conditions. Furthermore, we found that PSG1 can simultaneously bind to heparan sulfate in the extracellular matrix and to α 5 β 1 on the cell membrane. Wound healing assays and single-cell movement tracking showed that immobilized PSG1 enhances EVT migration. Although PSG1 did not affect EVT invasion in the *in vitro* assays employed, we found that the serum PSG1 concentration is lower in African-American women diagnosed with early-onset and late-onset [...]

Reference

RATTILA, Shemona, *et al.* Interaction of Pregnancy-Specific Glycoprotein 1 With Integrin A5 β 1 Is a Modulator of Extravillous Trophoblast Functions. *Cells*, 2019, vol. 8, no. 11, p. 1369

DOI : 10.3390/cells8111369

PMID : 31683744

Available at:

<http://archive-ouverte.unige.ch/unige:126818>




Disclaimer: layout of this document may differ from the published version.



UNIVERSITÉ
DE GENÈVE

Article

Interaction of Pregnancy-Specific Glycoprotein 1 With Integrin $\alpha 5\beta 1$ Is a Modulator of Extravillous Trophoblast Functions

Shemona Rattila ¹, Caroline E. Dunk ², Michelle Im ³, Olga Grichenko ¹, Yan Zhou ⁴, Marie Cohen ⁵, Maria Yanez-Mo ⁶, Sandra M. Blois ^{7,8}, Kenneth M. Yamada ⁹, Offer Erez ^{10,11,12}, Nardhy Gomez-Lopez ^{10,12,13}, Stephen J. Lye ², Boris Hinz ³, Roberto Romero ^{14,15,16,17,18} and Gabriela Dveksler ^{1,*}

- ¹ Department of Pathology, Uniformed Services University of Health Sciences, Bethesda, MD 20814, USA; shemona.rattila@usuhs.edu (S.R.); olgri4@gmail.com (O.G.)
- ² Lunenfeld Tanenbaum Research Institute, Sinai Health System, Toronto, ON M5T 3H7, Canada; dunk@lunenfeld.ca (C.E.D.); Lye@lunenfeld.ca (S.J.L.)
- ³ Laboratory of Tissue Repair and Regeneration, Faculty of Dentistry, University of Toronto, Toronto, ON M5G 1G6, Canada; michelleim27@gmail.com (M.I.); boris.hinz@utoronto.ca (B.H.)
- ⁴ Department of Obstetrics, Gynecology and Reproductive Sciences, Center for Reproductive Sciences, University of California San Francisco, San Francisco, CA 94143, USA; Yan.Zhou@ucsf.edu
- ⁵ Department of Pediatrics, Gynecology and Obstetrics, University of Geneva, 1206 Geneva, Switzerland; Marie.Cohen@hcuge.ch
- ⁶ Department of Molecular Biology, Universidad Autónoma de Madrid (UAM), 28049 Madrid, Spain; maria.yanez@cbm.csic.es
- ⁷ Experimental and Clinical Research Center, a Cooperation between the Max Delbrück Center for Molecular Medicine in the Helmholtz Association, and the Charité-Universitätsmedizin Berlin, AG GlycoImmunology, 13125 Berlin, Germany; sandra.blois@gmail.com
- ⁸ Berlin Institute of Health, Institute for Medical Immunology, Charité-Universitätsmedizin Berlin, corporate member of Freie Universität Berlin, Humboldt-Universität zu Berlin, 13353 Berlin, Germany
- ⁹ Cell Biology Section, National Institute of Dental and Craniofacial Research, National Institute of Health, Bethesda, MD 20892, USA; kyamada@dir.nidcr.nih.gov
- ¹⁰ Perinatology Research Branch, Division of Obstetrics and Maternal-Fetal Medicine, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health, U. S. Department of Health and Human Services, Bethesda, MD 20892, and Detroit, MI 48201, USA; offererez@gmail.com (O.E.); ngomezlo@med.wayne.edu (N.G.-L.)
- ¹¹ Maternity Department "D," Division of Obstetrics and Gynecology, Soroka University Medical Center, School of Medicine, Faculty of Health Sciences, Ben Gurion University of the Negev, Beer-Sheva 8410501, Israel
- ¹² Department of Obstetrics and Gynecology, Wayne State University School of Medicine, Detroit, MI 48201, USA
- ¹³ Department of Biochemistry, Microbiology and Immunology, Wayne State University School of Medicine, Detroit, MI 48201, USA
- ¹⁴ Department of Obstetrics and Gynecology, University of Michigan Health System, Ann Arbor, MI 48109, USA; prbchiefstaff@med.wayne.edu
- ¹⁵ Department of Epidemiology and Biostatistics, Michigan State University, East Lansing, MI 48824, USA
- ¹⁶ Center for Molecular Medicine and Genetics, Wayne State University, Detroit, MI 48201, USA
- ¹⁷ Detroit Medical Center, Detroit, MI 48201, USA
- ¹⁸ Department of Obstetrics and Gynecology, Florida International University, Miami, FL 33199, USA
- * Correspondence: gabriela.dveksler@usuhs.edu

Received: 4 October 2019; Accepted: 25 October 2019; Published: 31 October 2019



Abstract: Human pregnancy-specific glycoproteins (PSGs) serve immunomodulatory and pro-angiogenic functions during pregnancy and are mainly expressed by syncytiotrophoblast cells. While PSG mRNA expression in extravillous trophoblasts (EVTs) was reported, the proteins were

not previously detected. By immunohistochemistry and immunoblotting, we show that PSGs are expressed by invasive EVT_s and co-localize with integrin $\alpha 5\beta 1$. In addition, we determined that native and recombinant PSG1, the most highly expressed member of the family, binds to $\alpha 5\beta 1$ and induces the formation of focal adhesion structures resulting in adhesion of primary EVT_s and EVT-like cell lines under 21% oxygen and 1% oxygen conditions. Furthermore, we found that PSG1 can simultaneously bind to heparan sulfate in the extracellular matrix and to $\alpha 5\beta 1$ on the cell membrane. Wound healing assays and single-cell movement tracking showed that immobilized PSG1 enhances EVT migration. Although PSG1 did not affect EVT invasion in the in vitro assays employed, we found that the serum PSG1 concentration is lower in African-American women diagnosed with early-onset and late-onset preeclampsia, a pregnancy pathology characterized by shallow trophoblast invasion, than in their respective healthy controls only when the fetus was a male; therefore, the reduced expression of this molecule should be considered in the context of preeclampsia as a potential therapy.

Keywords: pregnancy-specific glycoproteins; extravillous trophoblasts; integrin $\alpha 5\beta 1$; adhesion; migration; preeclampsia

1. Introduction

Pregnancy-specific glycoproteins (PSGs), also known as Schwangerschafts Protein 1 (SP1) and pregnancy-specific beta 1 glycoproteins, are secreted placental proteins found in the maternal circulation [1–4]. PSGs belong to the carcinoembryonic antigen cell adhesion molecules (CEACAM) family but while CEACAMs are mostly membrane-associated proteins, PSGs are secreted [5]. All species that express PSGs have multiple PSG-encoding genes [6]. In mice and rats, 17 genes (*Psg16–Psg32*) and eight genes (*Psg36–Psg43*) have been identified, respectively [7]. In humans, there are 11 *PSG* genes; while one of them has been reported to be a pseudogene, the expression of 10 PSGs designated as PSG1–PSG9 and PSG11 is predicted, although specific antibodies for each member of the family are not available [8]. The PSG1 messenger RNA (mRNA) is abundantly expressed throughout pregnancy and the protein concentration of PSGs reaches its maximum level in maternal plasma at term [9,10]. PSGs are found only in species with hemochorial placentation in which maternal blood comes into direct contact with fetal cells, posing a risk of rejection by the maternal immune system [11]. Interestingly, our group and others have shown that human PSGs and murine PSG23 have immune-regulatory activity consistent with the hypothesis that these proteins may participate in tolerance to the fetal semi-allograft [12–16]. In addition, we have reported that PSG1 has pro-angiogenic activity as it induces endothelial tubulogenesis [17,18]. Furthermore, human PSG1 and PSG9 and mouse PSG23 were shown to have anti-thrombotic activity [9].

PSGs are expressed predominantly, but not exclusively in trophoblasts as low levels of expression were detected in a healthy colon and in the squamous epithelium of the esophagus [19,20]. PSG expression has also been reported in tumors of trophoblastic and non-trophoblastic origin [21–23]. Similar to the presence of human PSGs in non-placental tissues, mouse PSG18 is expressed in the follicle-associated epithelium of Peyer's patches potentially playing a role in the interplay between epithelial cells and immune cells in mucosa-associated lymphoid tissue [24].

Trophoblasts are a specialized cell population in the placenta serving various functions ranging from attachment, migration and invasion to vascular remodeling [25]. In the human placenta, cytotrophoblasts (CTBs) proliferate and differentiate into spatially distinct populations [26]. In the floating villi, fusion of CTBs generates multinucleated syncytiotrophoblast (STB) [27–29]. STB produces pregnancy hormones, transport nutrients and oxygen from the mother to the fetus and remove fetal waste products [27,28]. In the anchoring villi that physically anchor the placenta to the uterine wall, differentiation of CTBs starts with the formation of trophoblast cell-columns, in which the proximal cell-column trophoblasts are highly proliferative, and the distal cell-column trophoblasts

are non-proliferative, migratory and eventually differentiate into invasive extravillous trophoblasts (EVTs) [28]. EVT migrate towards and invade into the maternal decidua to transform the uterine spiral arteries of the fetal-maternal interface [30,31]. In human pregnancy, exclusive expression of PSGs by STB was reported more than two decades ago [32]. More recent studies have indicated the presence of PSG mRNA in EVT [33,34]. Therefore, we first examined whether PSGs are expressed in EVT using two PSG-specific antibodies and investigated the interaction of PSG1 with EVT.

Some PSG1 ligands have been identified; human PSG1 and mouse PSG17, PSG22 and PSG23 bind to heparan sulfate proteoglycans (HSPGs) [17,35,36]. The interaction of PSG1 with HSPGs was shown to be required for the ability of PSG1 to induce endothelial tube formation [17]. Besides binding to HSPGs, PSG1, PSG9 and PSG23 bind to integrin α IIb β 3 and thereby inhibit fibrinogen binding to platelets [9]. As differentiation into EVT is accompanied by a sequential alteration of integrin expression referred to as “integrin switching”, which is regulated in a spatial and temporal manner, we next investigated the interaction of PSG1 with integrins. Proliferative CTBs anchored to the basement membrane express integrin α 6 β 4 [37]. Their differentiation into EVT near the distal cell column is accompanied by down-regulation of integrin α 6 β 4 and up-regulation of integrin α 5 β 1 expression, associating with acquisition of a migratory cell phenotype [31,38]. The cells that invade the uterine wall express integrin α 1 β 1 along with α 5 β 1 and loose α 6 β 4 expression [37,39,40]. In this study we show via numerous functional assays that immobilized PSG1 induces adhesion of primary EVT and EVT-like cell lines in an integrin α 5 β 1-dependent manner and that PSG1 directly interacts with this integrin. Furthermore, EVT seeded on PSG1 have increased migration when compared to cells seeded on a control protein as determined in wound healing assays and single-cell movement tracking experiments.

The spatial regulation of integrin expression is altered in various placental disorders involving either excessive invasion, such as in placenta accrete, or insufficient invasion, such as in pre-eclampsia (PE), suggesting that alteration of PSG levels may in part be connected to these defects [40–43]. In this study using a specific ELISA, we report reduced serum concentrations of PSG1 in African American women diagnosed with early-onset and late-onset PE but only when they carried a male fetus. Altogether our results indicate that PSGs play important roles in several processes required for normal placentation.

2. Materials and Methods

2.1. Protein Production and Purification

PSG1-Fc, PSG1-His, PSG1N-Fc and the proteins used interchangeably as negative controls, Fc and CEACAM9-Fc, were generated from the supernatant of stably transfected CHO-K1 single-cell clones established in our laboratory and grown in hollow fiber cartridge bioreactors (FiberCell Systems, Frederick, MD, USA), as previously described [44]. Plasmids encoding the complementary DNAs (cDNAs) of the single-domain proteins (PSG1A2-Fc and PSG1B2-Fc) were generated as previously reported [44]. For the generation of the PSG1A2 and PSG1B2-Fc single-domain proteins, the plasmids were transfected into ExpiCHO cells (Thermo Fisher Scientific, Waltham, MA, USA) following the manufacturer’s recommendations and the supernatants were collected 5–6 days post-transfection based on cell viability. PSG1 from the serum of pregnant women and PSG1-His generated from the stable cell line in the bioreactor were purified using an anti-PSG1 mAb#4 column as described previously [16]. The identity of native PSG1 was confirmed by matrix-assisted laser desorption/ionization-time of flight (MALDI-TOF) mass spectrometry (MS). The anti-PSG1 mAb#4, which reacts with the N-terminal domain of PSG1, was obtained from Dr. S. Jonjic (University of Rijeka, Rijeka, Croatia). For the purification of Fc-tagged proteins, the supernatants were applied to protein A columns (GE Healthcare, Chicago, IL, USA) and the proteins were eluted with 0.1 M glycine buffer pH 2.7, followed by immediate neutralization with 1 M Tris-HCl pH 8 (Thermo Fisher Scientific). The eluted fractions were pooled, concentrated, and buffer-exchanged with phosphate-buffered saline (PBS) using an Amicon Ultra-10K centrifugal filter unit (MilliporeSigma, Burlington, MA, USA). For quantitation, the purified proteins

were separated on NuPAGE 4%–12% Bis-Tris gels (Life Technologies, Carlsbad, CA, USA) at different dilutions alongside known concentrations of bovine serum albumin (Thermo Fisher Scientific) used as standards. The proteins on the gel were stained with GelCode Blue (Thermo Fisher Scientific) and were quantified by densitometry. The identity of the proteins were confirmed by Western blot with specific Abs.

2.2. Cell Lines

The human first trimester trophoblast cell line HTR8/SVneo was provided by Dr. Caroline Dunk (Mount Sinai Hospital, Toronto, Ontario, Canada) and Dr. Charles Graham (Queen's University, Kingston, Ontario, Canada), and was cultured in RPMI 1640 (Corning, Corning, NY, USA). The human first trimester trophoblast cell line Swan71 was provided by Dr. Gil Mor (Yale University School of Medicine, New Haven, Connecticut, USA) and CHO-K1 cells were obtained from American Type Culture Collection (Manassas, Virginia, USA), and were cultured in Dulbecco's Modified Eagle Medium (DMEM) (Lonza, Basel, Switzerland). The cell line CHO-B2 (integrin $\alpha 5$ deficient) was a gift from Dr. Rudy L. Juliano (University of North Carolina, Chapel Hill, North Carolina, USA) and was cultured in Minimum Essential Medium (MEM) alpha modified with ribo- and deoxyribonucleosides (Thermo Fisher Scientific). All cell lines were maintained by using the media indicated above, supplemented with 10% fetal bovine serum (FBS) (Innovative Research Inc., Novi, MI, USA), 100 $\mu\text{g}/\text{mL}$ penicillin/streptomycin (Corning) and 100 $\mu\text{g}/\text{mL}$ normocin (InvivoGen, San Diego, CA, USA) in a 37 °C humidified incubator with 5% CO_2 .

2.3. Placental Tissue Collection and Primary Cells Isolation

Placental and decidual tissues were obtained from women carrying a normal pregnancy after legal termination during the first- or second-trimester of pregnancy. All patients who donated a placenta provided their informed written consent prior to their inclusion in the study in accordance with the Declaration of Helsinki. Approvals on the use of human subject materials were obtained from the following institutions and their respective committees on human research: the Research Ethics Board (REB # 12-0007E) of Mount Sinai Hospital (Toronto, Ontario, Canada), the local ethics committee of the Maternity and Pediatrics Department (project code # Gyn 02-007) of the Geneva University Hospital (Switzerland) and the University of California San Francisco (California, USA). Fresh tissue specimens were collected in ice-cold PBS and were washed several times in sterile Hanks balanced salt solution (HBSS). Placentas were microdissected to collect EVT columns at the tips of the villi. Approximately 30 EVT columns per placenta were digested with 10 mL of an enzyme cocktail (0.125% trypsin, 100 mM HEPES, 40 mM MgSO_4 , 0.15 mg/mL DNase I and 100 $\mu\text{g}/\text{mL}$ normocin in HBSS) for 15 min in a shaking water bath at 37 °C. After neutralization with FBS, the cells were resuspended in DMEM and the suspension was filtered through 70 μm mesh and counted. The cells were confirmed as >95% EVTs based on the expression of cytokeratin 8 (CK8), and human leukocyte antigen-G (HLA-G) and the absence of vimentin and then used for single-cell tracking experiments shown in Figure 1D–F. The EVTs utilized for the experiments shown in Figure 4B were purified on Percoll gradients as previously described [45].




Figure 1. Pregnancy-specific glycoproteins (PSGs) localize to extravillous trophoblasts (EVTs) in first-trimester placenta and interstitial EVTs in the decidua. Representative photographs of dual fluorescent immunohistochemistry localizing cytokeratin 8 (red) and PSGs (green with both BAP3 and #4 mAbs) are shown in both individual and dual channels in sections of first-trimester placenta and decidua basalis. Serial sections stained with a mAb against HLA-G are shown in (D), (I) and (M) (red) to confirm localization in the EVTs. (A–E) 5-week old placenta. A: CK8, B: PSG (BAP3), C: CK8 and PSG (BAP3), D: HLA-G and E: guinea pig and mouse IgG control. (F–I) 7-week old placenta. F: CK8, G: PSG (mAb#4) with the insert showing the STB layer, H: CK8 and PSG (mAb#4) and I: HLA-G. (J–M) 6.5-week old decidua. J: CK8, K: PSG (mAb#4), L: CK8 and PSG (mAb#4) and M: HLA-G. Scale bars = 25 μm for (A–C), (E–H) and (J–K), and 45 μm for (D), (I) and (M).

CTBs were isolated from the chorionic villi of second-trimester placentas and purified on Percoll gradients according to previously described protocols [46,47]. After initial purification of CTBs, the remaining leukocytes were removed with anti-CD45-coupled magnetic beads. Purified CTBs were cultured in serum-free DMEM, with 2% Nutridoma (Boehringer Mannheim Biochemicals, Indianapolis, Indiana, USA) on a Matrigel-coated surface for 14–20 h for differentiation into EVTs.

2.4. Fluorescence Immunohistochemistry for Analysis of PSGs Expression and Localization with Integrin $\alpha 5$ in First-Trimester Placentas

Tissues collected from the placenta at 5–8 weeks of gestation were fixed in paraformaldehyde (PFA) and processed to 5 μm thick paraffin sections on Superfrost Plus glass slides (VWR, Mississauga, Ontario, Canada). Tissue sections were deparaffinized using xylene and rehydrated with a descending concentration gradient of ethanol. Heat-induced antigen retrieval with sodium citrate buffer (pH 6) was performed to avoid epitope masking. Autofluorescence was minimized with the treatment of 0.1% Sudan Black in 70% ethanol for 1 min. Blocking with a cocktail of 10% normal goat serum, 2% human serum, 2% donkey serum and 2% rabbit serum for 1 h was done to prevent non-specific binding. Tissue sections were incubated overnight at 4 $^{\circ}\text{C}$ with the following primary Abs: 1:250

dilution of anti-PSG mouse mAb-BAP3 (cat#GM-0507; Aldevron, Fargo, North Dakota, USA) or anti-PSG mAb #4, which react with the B2 and N-domain of PSG1 respectively, and anti-cytokeratin 8 guinea pig polyclonal (cat# ab194130; Abcam, Cambridge, UK) at a 1:100 dilution. These Abs were applied at appropriate combination for dual immunofluorescence images shown in Figure 1; anti-PSG mouse mAb-BAP3 and anti-integrin $\alpha 5$ rabbit mAb (cat# ab112183; Abcam) at 1:500 dilution were applied in combination for dual immunofluorescence images shown in Figure A1. Serial sections were incubated with anti-HLA-G (cat#11-499-c100; Exbio, Vestec, Czechia) to identify the EVT in the placental sections for Figure 1. As controls, mouse (cat#X0931, Agilent Technologies, Santa Clara, CA, USA), guinea pig (cat# NBP197036; Novus Biologicals, Centennial, CO, USA) and rabbit IgGs (cat#ab171870, Abcam) instead of specific primary Abs were applied at the same concentrations and combinations as the specific primary Abs. Secondary Abs used for Figure 1 were anti-mouse Alexa Fluor-488 (cat#A-28175; Thermo Fisher Scientific) at 1:200 dilution and anti-guinea pig biotin (cat#106-066-003; Jackson ImmunoResearch Laboratories Inc., West Grove, PA, USA) at 1:1000 dilution and amplification was carried out using Streptavidin Alexa Fluor-546 (Thermo Fisher Scientific). Secondary Abs used for Figure A1 were anti-mouse Alexa Fluor-546 (at 1:300) (cat#A-11030; Thermo Fisher Scientific) in combination with anti-rabbit biotin (at 1:1000) (cat#ab6720; Abcam) and amplification was carried out using Streptavidin Alexa Fluor-488 at 1:1000 dilution (cat#S-32354; Thermo Fisher Scientific). Slides were washed in PBS and mounted in aqueous immune-mount (Thermo Fisher Scientific). Cells were counterstained with the nuclear dye DAPI (5 $\mu\text{g}/\text{mL}$). Images were captured using the Quorum Wave FX spinning disc confocal system comprising a Leica DMI 6000B microscope (Leica Microsystems, Wetzlar, Germany) with a Yokogawa Spinning Head and Image EM Hamamatsu EMCCD camera and Velocity imaging software version 6.30 (Quorum Technologies Inc, Puslinch, Ontario, Canada). To ensure fair comparison across gestation, exposure time and laser intensity was kept constant across all images.

2.5. Immunoblot Analysis of PSG1 Expression in Second-Trimester Trophoblasts

Cell lysates were prepared from purified second-trimester CTBs following isolation (0 h) or cultured on Matrigel for different periods of time (14 and 20 h). Ten μg of cell lysate was loaded onto a 4%–12% SDS-PAGE gradient gel and transferred to a polyvinylidene fluoride (PVDF) membrane. Anti-PSG mAb-BAP3 (Aldevron, Fargo, North Dakota, USA) and anti-PSG mAb#4 were used at 1 $\mu\text{g}/\text{mL}$. The Anti- α tubulin mAb (mouse) was utilized to normalize protein loading (clone#B-5-1-2, cat#T5168; Sigma-Aldrich, St. Louis, MO). The signal was developed with horseradish peroxidase conjugated (HRP) anti-mouse secondary Abs (cat#715-0350150; Jackson ImmunoResearch Laboratories Inc.). Images were captured and analyzed with ImageJ software.

2.6. Cell Adhesion

Adhesion assays were performed in 96-well polystyrene high protein binding wells (Thermo Fisher Scientific). Wells were coated with 100 μL of the purified proteins diluted in PBS at the indicated concentrations and incubated overnight at 4 $^{\circ}\text{C}$. Wells were then incubated with 300 μL of 1% bovine serum albumin (BSA)/PBS for 1 h at room temperature to block non-specific binding. After blocking, EVT-like cell lines, removed from tissue culture flasks with Accutase (Innovative Cell Technologies, Inc., San Diego, California, USA), were seeded in their corresponding serum-free media supplemented with 1% BSA-10mM HEPES at a final concentration of 2×10^4 cells/well and incubated for 2 h at 37 $^{\circ}\text{C}$ with 5% CO_2 . When purified primary EVTs were applied, 1×10^5 cells/well were seeded. After incubation, wells were washed three times with PBS to remove unbound cells and the remaining cells were incubated for 2 h with CellTiter 96 Aqueous One Solution (MTS; Promega, Madison, WI, USA), following the manufacturer's recommendation, and absorbance at 490 nm was recorded. When indicated, cells were pre-incubated with 500 μM RGD-peptide (GRGDNP) or RGD-control peptide (GRADSP; Enzo Life Sciences, New York, NY, USA), or with echistatin (MilliporeSigma, Burlington, MA, USA) at 1 $\mu\text{g}/\text{mL}$ for 30 min at room temperature or with function-blocking mAbs: mouse anti-human integrin $\alpha 1\beta 1$

(clone#5E8D9, cat# 05-246), mouse anti-human integrin α V β 3 (clone#LM609, cat#MAB1976) and mouse anti-human integrin α V β 5 (clone#P1F6, cat#1961Z; Millipore Sigma); rat anti-human integrin α 5 mAb16 and rat anti-human integrin β 1 mAb13 [48,49] at the indicated concentrations. Mouse isotype control and rat isotype control Abs were purchased from R&D Systems and BD Biosciences, respectively. To determine the effect of heparin on PSG1-mediated cell adhesion, cells were incubated with 50 μ g/mL heparin (Millipore Sigma), which was maintained during cell seeding. Experiments in hypoxic conditions with primary EVT_s and EVT-like cell lines were performed using a hypoxic chamber in an incubator regulated to maintain 1% oxygen exposure. Hypoxic-like conditions for the EVT-like cell lines were chemically induced with 100 μ M CoCl₂ added for 4 h prior and kept for the duration of the experiment.

2.7. Integrin Binding ELISA

Wells of a 96-well plate were coated overnight at 4 °C with 100 μ L of protein diluted in PBS at the concentrations indicated. After washing with tris buffered saline-tween (TBST), the wells were blocked with 300 μ L/well blocking buffer (1% BSA in TBS) for 2 h at 37 °C. Recombinant human integrin α 5 β 1 (R&D Systems, Minneapolis, MN, USA) prepared in 1% BSA in 1 \times TBS with 1 mM MnCl₂ was added to the wells and incubated for 2 h at 37 °C. Following washes, 1 μ g/mL of biotin-conjugated anti-human integrin β 1 Ab (cat#BAF1778, R&D Systems) was added to the wells for 2 h at room temperature. The wells were washed again followed by the addition of Streptavidin-HRP and TMB substrate (R&D Systems). Reactions were stopped by addition of 2 N of sulfuric acid and the absorbance at 450 nm was recorded using a GloMax Discover multimode detection instrument (Promega, Madison, WI, USA). When indicated, the integrin was added to the wells together with either 5 μ M RGD peptide or the RGD-control peptide.

2.8. FAK Phosphorylation Analysis by Western Blot

Thirty-five mm poly-D-lysine dishes were coated with 30 μ g/mL PSG1-Fc or control protein overnight at 4 °C. Cells maintained overnight in media with 2% FBS, were detached with Accutase and were resuspended in 1% BSA containing serum-free media prior to the addition at 3.5×10^5 cells/dish and incubated 60 min at 37 °C. Dishes were placed on ice, washed with PBS and cells were lysed with 200 μ L LDS-sample buffer (Invitrogen, Carlsbad, CA, USA) supplemented with protease and phosphatase inhibitors (Millipore Sigma). Protein concentration was determined with the EZQ Protein Quantitation Kit (Invitrogen) and 50 μ g were loaded per lane and separated in a NuPAGE 3–8% tris-acetate gel (Thermo Fisher Scientific) before transferring to a PVDF membrane. The blot was probed sequentially with anti-FAK (cat#3285), anti-pFAK Y397 (clone#D20B1, cat#8556) and anti- α tubulin (clone#11H10, cat#2125; Cell Signaling Technology, Danvers, MA, USA) Abs, followed by HRP-conjugated bovine anti-rabbit IgG (cat#sc-2370, Santa Cruz Biotechnology, Inc., Dallas, TX, USA) and the West Pico Chemiluminescent Substrate (Thermo Scientific). Images were captured using Fujifilm laser imager (LAS 4000) (GE Healthcare, Chicago, IL, USA) and quantitated with the ImageJ software Version 1.52r (NIH, Bethesda, MD, USA).

2.9. Focal Adhesion Structure Analysis by Fluorescence Microscopy

HTR8/SVneo cells were seeded in serum free media onto PSG1-Fc (20 μ g/mL) or poly-L-lysine (0.01%)-coated coverslips and allowed to adhere for 2 h at 37 °C. After removal of the media, cells were fixed in 4% PFA for 5 min at room temperature (RT), washed twice with TBS, and permeabilized with 0.5% Triton X-100 for 5 min. Cells were incubated with following primary mouse mAbs overnight at 4 °C: anti-paxillin (cat#610051; BD Biosciences, San Jose, CA, USA) and anti-vinculin (cat#V9131, Millipore Sigma), followed by species-matching secondary Abs coupled to Alexa Fluor-488 (cat#A28175, Thermo Fisher Scientific) for 1 h at room temperature. Co-staining was performed with Alexa-Fluor-555-Phalloidin (cat#A34055; Thermo Fisher Scientific) and mounting with Prolong (Invitrogen, Carlsbad, CA, USA) containing DAPI. Images were obtained with a Leica LSM510

inverted confocal microscope and analyzed with the Leica confocal LAS software version 3.8.0 (Leica Microsystems, Wetzlar, Germany).

2.10. PSG1-ECM Binding Assay

MRC-5 fibroblast cells were seeded at 2×10^5 cells/dish onto 60-mm dishes and grown for 7 days for extracellular matrix (ECM) deposition. After washing extensively with PBS, 1.5 mL of decellularization solution (20 mM NH_4OH with 0.5% Triton X-100) was applied four times to remove the cells followed by a second round of extensive washing with PBS. The deposited ECM was incubated with or without the addition of heparin (50 $\mu\text{g}/\text{mL}$) for 3 h at 37 °C. PSG1-Fc (10 $\mu\text{g}/\text{mL}$) for 1 h at room temperature. The ECM was washed with PBS for removal of unbound proteins and then lysed with 40 μL 1% Triton X-100 solution supplemented with a protease inhibitor cocktail. Fifty μg of lysed ECM was loaded onto a 10% SDS-PAGE gel and transferred to a PVDF membrane. The following Abs were used for Western blot analysis: goat anti-human Fc at 1:1000 dilution (cat#H10500, Thermo Fisher Scientific), rabbit anti-fibronectin at 1:2000 dilution (cat#F3648, Millipore Sigma), mouse anti-GAPDH and appropriate secondary Abs (1:1000; LI-COR Biosciences, Lincoln, NE, USA).

2.11. Wound Healing Assay

Cell migration was measured in a wound healing assay using a 2-chamber culture-insert with a middle gap (Ibidi GmbH, Munich, Germany). The culture-inserts were placed in each well of a 24-well poly-D-lysine plate previously coated overnight with PSG1-His or control protein. Then, either 6.5×10^4 HTR8/SVneo or 6×10^4 Swan71 in 80 μL of serum-free media (SFM) with 1% BSA was seeded in each of the two chambers of the insert and incubated at 37 °C with 5% CO_2 for 3 h. Cells that did not attach to the plate were gently washed off and the inserts were removed 6–7 h post-cell seeding. The wells were washed with SFM and the inserts filled with 1.5 mL 1% BSA in SFM. Cell migration into the 500 μm gap was recorded by time-lapse microscopy (Leica AF6000 DMI6000B with built in camera DFC 365FX-566903911) for 20 h. Images were captured every 20 min. Movies were edited using the LAS X software version 3.3.2 (Leica Microsystems, Wetzlar, Germany), analyzed with the “Wound Healing ACAS Image Analysis” (MetaVi Labs Inc., Bottrop, Germany) and expressed as gap closure area (measurement of difference between gap area at $t = 0$ and $t = 20$ h) and gap closure speed (measurement of gap area covered per unit time). To mimic hypoxic conditions, cells were pre-treated for 4 h with 100 μM CoCl_2 in serum-containing media. The wound healing assays were carried out with 100 μM CoCl_2 in SFM with B-27 Supplement (Thermo Fisher Scientific) or using a hypoxic chamber in an incubator regulated to maintain 1% oxygen exposure. Results shown are mean \pm S.D. of seven fields of four wells from one representative of three independent experiments.

2.12. Tracking of Single Cell Migration

Wells of a 24-well poly-D-lysine plate were coated with 20 $\mu\text{g}/\text{mL}$ PSG1-His or control protein overnight at 4 °C. Wells were blocked with 1% BSA/PBS, and HTR8/SVneo and Swan71 cells were seeded in Opti-MEM (Thermo Fisher Scientific) at a density of 5000 cells/well and allowed to adhere overnight at 37 °C in a humidified 5% CO_2 incubator. For experiments with primary EVT_s, a 48-well plate was used and the cells were seeded with 2% FBS containing Opti-MEM at a density of 2500 cells/well. Imaging was performed using a Zeiss Axiovert (Carl Zeiss Microscopy, Jena, Germany) 135 microscope with a temperature-controlled chamber maintained at 37 °C with 5.0% CO_2 . Images were captured at 10 min intervals for a total of 100 frames and were analyzed with ImageJ using the plug-in MTrackJ, which in the ‘Measure’ option provided output as track length, velocity, and distance to previous point for the individual cells that were tracked. Results are shown as mean \pm S.D. of four experiments consisting of $n = 10$ for control protein, and $n = 22$ for PSG1-His in each experiment.

2.13. HIF1 α Stabilization Analysis by Western Blot

Swan71 and HTR8/SVneo cells were plated in standard growth medium at a density of 2.5×10^6 cells or 2.2×10^6 cells per 100-mm dish, respectively. The following day, cells were washed and treated with 100 μ M CoCl₂ (Millipore Sigma) in 0.1% BSA containing media for 4 h. Cells were lysed with 400 μ L ice-cold LDS sample buffer (Invitrogen) containing protease inhibitors (Millipore Sigma). The lysates were centrifuged at 15,000 rpm for 30 min and the protein concentration of the soluble material was determined with the EZQ Protein Quantitation Kit (Invitrogen). Fifty μ g of protein were loaded per lane and separated on a NuPAGE 4%–8% Bis-Tris gel (Invitrogen). The proteins were transferred to a PVDF membrane and, after blocking, incubated with mouse anti-HIF1 α (clone#241809, cat#MAB1536; R&D Systems) and rabbit anti- α -tubulin (clone#11H10, cat#2125; Cell Signaling), followed by the respective HRP-conjugated secondary antibodies. The signal was developed with the SuperSignal West Pico Chemiluminescence substrate (Thermo Fisher Scientific) and visualized in a Fujifilm laser imager (LAS 4000).

2.14. Flow Cytometry

Expression of integrin $\alpha 5\beta 1$ in HTR8/SVneo and Swan71 cells was studied by flow cytometry. For flow cytometry, cells were detached with Accutase, and 1×10^6 cells in 100 μ L of Fluorescens Activated Cell Sorting (FACS) analysis buffer (PBS with 2% BSA and 0.05% NaN₃) were incubated with 1 μ g of anti-human $\alpha 5\beta 1$ mAb (clone#HA5, cat#MAB1999; Millipore Sigma) or mouse isotype control (clone#11711, cat#MAB002; R&D systems) for 1 h on ice, followed by 0.25 μ g allophycocyanin (APC) conjugated anti-mouse Ig (BD Biosciences) for 30 min. Excess antibodies were removed by washing with the FACS buffer between steps. Cells were analyzed using the BD LSR II (BD Biosciences). A total of 50,000 events were collected for each treatment using the FACS Diva software (BD Biosciences), and the FlowJo software V10.0.8 (BD Biosciences, San Jose, CA, USA) was used for post-acquisition analysis.

2.15. Invasion Assays

Cell invasion was examined using a transwell invasion assay and a Biogel cell invasion assay. For the transwell invasion assay, 24-well format cell culture inserts (BD Biosciences) were used. Insert membranes were coated with Matrigel (Corning) or Geltrex (Thermo Fisher) diluted at 0.1 or 0.2 mg/mL with culture media. PSG1-His or control protein at 60 μ g/mL was added to the diluted matrix before coating. Coating was performed by incubating at 37 °C for 2 h followed by drying under laminar air flow for 5 h. After rehydrating the dried matrix-coated insert for 2 h with 100 μ L SFM, HTR8/SVneo and Swan71 cells in SFM were plated in the inserts at 8×10^4 /insert and 6×10^4 /inserts, respectively. Serum containing (10% FBS) media was added into the lower wells and plates were incubated at 37 °C for 24 h. Non-invaded cells from the upper surface of the insert were removed by scrubbing with cotton swabs and the cells on the under surface were fixed with 5% glutaraldehyde (Electron Microscopy Sciences) for 10 min, followed by staining with 2% crystal violet (Sigma Aldrich) for 5 min. For each experiment, cells in 10 randomly chosen fields of each filter were counted manually and the experiment was repeated three times. The collagen I coated, 96-well cell migration Biogel assay plates (Enzo Life Sciences) were used for cell invasion assay according to the manufacturer's instruction with some modifications. HTR8/SVneo and Swan71 cell suspensions prepared with 2% serum containing media were plated at 2.5×10^4 /well and 2.2×10^4 /well, respectively. Plates were incubated at 37 °C for 2 h in which the cell-free detection zone in the center of the wells became available after the Biogel dissolved. After gentle washes with SFM, 50 μ L of Geltrex at 0.1 or 0.2 mg/mL mixed with 60 μ g/mL PSG1-His or control protein was added onto the seeded cells in the wells and incubated for 2 h to allow for gel formation. Some wells were used as reference for pre-invasion conditions and fixed immediately without adding Geltrex. After adding 100 μ L of media with 1% FBS to each well with the gelled matrix, plates were further incubated for 24 h. After that time, cells were fixed with 5% glutaraldehyde for 10 min and stained with TRITC-phalloidin (Thermo Scientific) for 40 min followed

by DAPI (Thermo Scientific) for 5 min. Images were captured using LAX AF600 software compatible with a Leica AF600 fluorescence imaging system with a built in camera (DFC 365FX-566903911). All images were analyzed using ImageJ software Version 1.52r (NIH, Bethesda, MD, USA) and the number of cells invaded into the detection zones was determined by comparison to the reference wells.

2.16. Determination of PSG1 Concentration in Serum of Pregnant Women

Plasma samples were collected at the time of a prenatal visit and all patients provided written informed consent prior to sample collection. Women were enrolled as participants of a study conducted at the Center for Advanced Obstetrical Care and Research of the Perinatology Research Branch, NICHD/NIH/DHHS, the Detroit Medical Center and Wayne State University (Detroit, Michigan, USA). Women with multiple gestations, severe chronic maternal morbidity (i.e., renal insufficiency, congestive heart disease and/or chronic respiratory insufficiency), acute maternal morbidity (i.e., asthma exacerbation requiring systemic steroids and/or active hepatitis) or fetal chromosomal abnormalities and congenital anomalies were excluded from the study. PE was defined as new-onset hypertension that developed after 20 weeks of gestation (systolic or diastolic blood pressure ≥ 140 mm Hg and/or ≥ 90 mm Hg, respectively, measured on at least two occasions, 4 hours to 1 week apart) and proteinuria (≥ 300 mg in a 24 h urine collection, or two random urine specimens obtained 4 h to 1 week apart containing $\geq 1+$ by dipstick or one dipstick demonstrating $\geq 2+$ protein) [50]. Early-onset PE was defined as PE diagnosed and delivery ≤ 34 weeks of gestation, and late-onset PE was defined as PE delivered and delivery ≥ 34 weeks of gestation [51].

We determined the concentration of PSG1 with the PSG1 Quantikine ELISA kit following the manufacturer's recommendations (R&D Systems). Serum samples were tested at 1:100 dilution for the early-onset control and early-onset PE groups (obtained at weeks 20.0–33.3 of gestation) and at 1:200 dilution for the late-onset PE and control (obtained between 37 to 41.7 weeks of gestation). For the few samples in which the values of PSG1 obtained were lower than the lowest standard, samples were diluted 1:50 or 1:13 and retested. All samples, obtained from the study population of African American women, were analyzed in triplicate. Data were analyzed using one-way analysis of variance (ANOVA) followed by pairwise comparisons using Tukey's adjustment for multiple comparisons. Two-way ANOVA, followed by pairwise tests of simple main effects, was used for comparisons stratified by fetal gender.

3. Results

3.1. Expression of PSGs in EVT_s

While PSG mRNA expression for different members of the family has been reported in EVT_s, previous studies using immunohistochemistry identified the STBs of the placenta as the sole site for PSG expression [32,33]. Therefore, we performed dual immunofluorescence staining of tissues obtained from the first-trimester placental villi and decidua basalis to examine PSGs expression employing more sensitive techniques by using two anti-PSG mAbs. EVT_s were identified in the 5-week and 7-week old placentas (1A, 1F), and 6.5-week old decidua (1J) with anti-CK-8, and HLA-G antibodies (1D, 1I, 1M). Expression of PSGs was examined with either anti-PSG BAP3 (1B) or anti-PSG mAb#4 (1G, 1K). Co-staining of 5-week and 7-week-old placentas with anti-PSG and anti-CK8 Abs shows that PSGs are localized to the EVT cell edges. Stronger staining was seen in the more distal fully invasive EVT_s (Figure 1C,H). As shown in panel 1L, PSG expression is also observed in the EVT_s actively invading the decidua and localizing to points possibly associated with focal adhesions (arrows). Only minimal background was observed with control IgGs (1E). In addition, we observed substantial co-localization of PSGs and integrin $\alpha 5$ in the first-trimester EVT_s (Figure A1a). PSG expression was also detected in the EVT_s that were differentiated in vitro from second-trimester CTBs (Figure A1b) and in second-trimester EVT_s as examined by immunohistochemistry (data not shown). Altogether, these results indicate that besides the high expression by STBs, PSGs are expressed by EVT_s in the cell

column where they colocalize with integrin $\alpha 5$, and by EVT s invading the decidua. In addition, PSG s are expressed by EVT s differentiated *in vitro* from CTBs.

3.2. Immobilized PSG1 Induces Adhesion of EVT s by Binding to Integrin $\alpha 5\beta 1$

To investigate whether PSG1 interacts with EVT s and affects their adhesive capacity, Swan71 cells were seeded onto non-tissue culture wells that had been pre-coated with either PSG1-Fc or the Fc-tag as control protein. As shown in the pictures taken 2 h post-seeding, cells adhered to the PSG1-coated wells (Figure 2A, top), but not to the control protein-coated wells (Figure 2A, bottom). Similar results were obtained with the HTR8/SVneo cell line (data not shown). We also determined that this PSG1-mediated effect was observed with PSG1 with a His tag and was dose-dependent with a maximal response observed at a PSG1 concentration of 60 $\mu\text{g}/\text{mL}$. (Figure 2B). The PSG1-mediated cell adhesion was inhibited when the cells were pre-treated with the chelating agent EDTA or with the disintegrin echistatin, indicating a requirement for a divalent-cation dependent receptor and a potential involvement of integrins (data not shown). To confirm the requirement of an integrin receptor for the adhesion of cells to PSG1, the cells were incubated with an RGD or control peptide prior to seeding onto the PSG1-coated wells. As shown in Figure 2C, pre-incubation with the RGD peptide prevented adhesion of the cells to PSG1 while the control peptide did not. Studies were then undertaken to define which of the integrins on EVT s were involved in this interaction. EVT s have been reported to express several integrins including $\alpha 1\beta 1$, $\alpha 4\beta 1$, $\alpha V\beta 3$, $\alpha V\beta 5$ and $\alpha 5\beta 1$ [52]. We found that anti- $\alpha V\beta 3$, anti- $\alpha V\beta 5$ and anti- $\alpha 1\beta 1$ neutralizing Abs did not inhibit PSG1-mediated cell adhesion (Figure 2D). On the other hand, the adhesion of Swan71 (Figure 2F, left) and HTR8/SVneo (Figure A1c) cells to immobilized PSG1 were significantly reduced in the presence of neutralizing Abs to the integrin $\alpha 5$ and integrin $\beta 1$ subunits, in ambient oxygen conditions (21%), CoCl_2 -induced hypoxia-like conditions (Figure A1d) and under low oxygen (1%) conditions (data not shown). Hypoxia more closely reflects the conditions during the first-trimester of pregnancy and has been shown to modulate integrin expression [53–56]. Importantly, expression of $\alpha 5\beta 1$ on the membrane of these cells was confirmed by flow cytometry (Figure A1e).

Given the limited availability of native PSG1 protein, we performed cell adhesion experiments with PSG1 purified from the serum of pregnant women only in 21% oxygen conditions and demonstrated that the adhesion of HTR8/SVneo cells to native PSG1 was reduced by 45% and 90% in the presence of neutralizing antibodies to integrin $\alpha 5$ and integrin $\beta 1$, respectively (Figure 2E). To determine whether the interaction between PSG1 and $\alpha 5\beta 1$ was also observed in primary cells, we performed adhesion experiments with primary EVT s maintained in 21% and 1% oxygen. Blocking of integrin $\alpha 5$ and integrin $\beta 1$ in primary EVT s resulted in 70%–90% reduction of the adhesion mediated by PSG1 (Figure 2F, right). Moreover, we observed that PSG1 induced adhesion of CHO-K1 cells, which express integrin $\alpha 5$, but did not induce adhesion of CHO-B2 cells, which lack expression of this integrin [57] (Figure 2G). Thus, these results indicate that immobilized native and recombinant PSG1 induce cell adhesion in an integrin $\alpha 5\beta 1$ -dependent manner in primary and EVT-like cell lines under 21% oxygen and 1% oxygen conditions. To determine whether PSG1 interacts directly with integrin $\alpha 5\beta 1$, we performed an ELISA with purified proteins. Figure 2H–J shows that binding of native and recombinant PSG1 to $\alpha 5\beta 1$ is direct and dose-dependent and can be specifically blocked by the RGD-peptide. Altogether, these results indicate that PSG1 is a new ligand for integrin $\alpha 5\beta 1$.




Figure 2. PSG1-mediated adhesion of the EVTs is dependent on a direct interaction with integrin $\alpha 5\beta 1$. Micrographs of Swan71 cells on wells coated with 30 $\mu\text{g}/\text{mL}$ PSG1-Fc for 1.5 h in serum-free media (A, top) or a protein consisting of just the Fc-tag used as control (B, bottom). Images were taken at 40 \times magnification. (B) HTR8/SVneo and Swan71 cells were seeded in wells coated with PSG1-His or control protein at various concentrations. Cells were incubated for 2 h at 37 $^{\circ}\text{C}$ and the wells were washed to remove non-adherent cells. Cells remaining in the wells were incubated with cell titer aqueous solution and cell adhesion was quantified as described in Materials and Methods. The adhesion of Swan71 cells to wells coated with 60 $\mu\text{g}/\text{mL}$ PSG1-His is considered as 100%. (C) HTR8/SVneo and Swan71 cells, pre-incubated with 500 μM RGD peptide or control (ctrl) peptide for 30 min at room temperature were seeded in wells coated with 30 $\mu\text{g}/\text{mL}$ PSG1-His or control protein. The adhesion of control peptide-treated cells to PSG1-His is considered as 100%. (D) HTR8/SVneo cells were pre-incubated with the indicated mAbs for 30 min at RT, after which they were seeded in wells coated with 30 $\mu\text{g}/\text{mL}$ PSG1-Fc or control protein. (E) HTR8/SVneo cells were pre-incubated with the indicated mAbs for 30 min at RT and seeded on wells coated with 30 $\mu\text{g}/\text{mL}$ of native PSG1 (nPSG1) or control protein. (F) Swan71 cells and purified primary EVTs, pre-incubated with the indicated mAbs were seeded on wells coated with 30 $\mu\text{g}/\text{mL}$ PSG1-His or control protein and adhesion assays were performed in 21% oxygen and CoCl_2 -induced hypoxic-like condition or in 1% oxygen conditions as described in Materials and Methods. The adhesion of control Ab-treated cells to PSG1 is considered as 100% in (D), (E) and (F). (G) CHO-K1 (integrin $\alpha 5$ -expressing) or CHO-B2 (integrin $\alpha 5$ -deficient) cells were seeded on wells coated with 30 $\mu\text{g}/\text{mL}$ PSG1-Fc or control protein. The adhesion of CHO-K1 cells to PSG1-Fc is considered as 100%. (H) nPSG1 or protein control at 20 $\mu\text{g}/\text{mL}$ or bovine fibronectin (FN) at 2 $\mu\text{g}/\text{mL}$ was coated on wells. After blocking, 1 $\mu\text{g}/\text{mL}$ of $\alpha 5\beta 1$ in 1 \times TBS/1mM MnCl_2 was added and binding of the integrin was detected with biotin-labeled anti- $\beta 1$ mAb followed by Streptavidin-HRP. I. Wells were coated with PSG1-Fc or protein control (20 $\mu\text{g}/\text{mL}$). After blocking, 1 $\mu\text{g}/\text{mL}$ of $\alpha 5\beta 1$ in 1 \times TBS/1 mM MnCl_2 was added in combination with 0.5 μM RGD or control peptides and binding of the integrin was detected as indicated in (H). The binding of the integrin to the control protein is considered as 1 in (H) and (I).

(J) Wells were coated with 20 µg/mL PSG1-Fc or CEACAM9-Fc. After blocking, $\alpha 5\beta 1$ in 1× TBS/1 mM $MnCl_2$ was added at various concentrations and binding of the integrin was detected as described above. The graph shows values for PSG1 obtained after subtracting the average of the control values. (K). HTR8/SVneo cells were seeded on wells coated with the indicated proteins (20 µg/mL) and adhesion experiment was carried out as described in Figure 2. Cell adhesion to PSG1-His is considered as 100%. Results shown are mean ± S.D. of triplicates from one representative of three independent experiments. *p* values were obtained by a one-way ANOVA followed by Sidak's multiple comparison tests for (H), (I) and (K), by a two-way ANOVA followed by Tukey's multiple comparison tests for (B), (C), (E) and (F), or followed by Sidak's multiple comparison tests for (G) (**** *p* < 0.0001).

3.3. More Than One Domain of PSG1 Mediates EVT Adhesion

PSG1 is comprised of four Ig-like domains designated as N, A1, A2 and B2 and a short cytoplasmic tail of varying length depending on the splice variant [6]. The PSG1-Fc utilized in our initial studies is a natural splice variant that lacks the A1 domain; therefore, we concluded that this domain is not essential for the ability of PSG1 to bind to $\alpha 5\beta 1$ and induce cell adhesion [6]. We generated recombinant single-domain proteins with an Fc tag and tested them for their ability to induce EVT adhesion. We found that the single-domain proteins composed of either the N- or the B2- domain induced adhesion of HTR8/SVneo while the A2 domain did not (Figure 2K). Similar results were obtained with the Swan71 cell line (data not shown).

3.4. PSG1 Induces Focal Adhesion (FA) Structures and Focal Adhesion Kinase (FAK) Phosphorylation

Ligand binding to integrins initiates the recruitment of adaptor proteins and signaling molecules, followed by cytoskeleton reorganization leading to the assembly of focal adhesion (FA) structures [58]. When EVTs were seeded on PSG1-coated covered slips, we observed the assembly of FA structures as determined by the detection of the adaptor proteins vinculin and paxillin associated with F-actin in HTR8/SVneo cells (Figure 3A). In addition, HTR8/SVneo adhesion to PSG1-coated wells induced phosphorylation of the Tyr-397 of FAK, which was not observed in the poly-D-lysine-coated wells (Figure 3B). Similarly, phosphorylation of FAK at Tyr-397 was detected in Swan71 cells when adhered to PSG1-coated wells but not in the cells adhered to poly-D-lysine-coated wells (data not shown).

3.5. PSG1 Binds to HSPGs in the ECM and can Concurrently Bind to Integrin $\alpha 5\beta 1$ on the Cell Surface

We observed PSG1-mediated cell adhesion only when PSG1 was immobilized on a solid surface but not when the protein was added to the media (data not shown). Previous results from our group showed that PSG1 binds to the ECM and to HSPGs [13,17]. To confirm these results and to determine whether the interaction of PSG1 with the ECM is mediated by binding to HSPGs, we tested whether PSG1-Fc could bind to the ECM deposited by a fibroblast cell line in the presence or absence of excess heparin, a highly sulfated form of HSPGs [59,60]. MRC-5 fibroblasts were cultured for 7 days for ECM deposition after which the cells were removed as confirmed by the absence of reactivity with the anti-GAPDH Ab (Figure 3C). We observed that binding of PSG1-Fc to the ECM was inhibited in the presence of heparin (Figure 3C). We also examined whether binding of PSG1 to heparin could interfere with PSG1's ability to induce adhesion of EVTs. We observed that the presence of added heparin did not interfere with PSG1-Fc-mediated adhesion of HTR8/SVneo cells (Figure 3D). These results strongly suggest that PSG1 can concurrently bind to HSPGs in the ECM and to integrin $\alpha 5\beta 1$ on the cell surface.




Figure 3. Binding of PSG1 leads to focal adhesion formation and PSG1 can concurrently bind to heparan sulfate proteoglycans (HSPGs) affecting EVT migration. **(A)** Representative immunofluorescence images of HTR8/SVneo cells on PSG1-Fc coated cover slips co-stained with Abs to vinculin, paxillin, and F-actin. DAPI was used to stain the nuclei. Arrowheads show the presence of paxillin (left) and vinculin (right) in the FA structures. Scale bars = 25 μm . **(B)** Representative immunoblot (left) of lysates of HTR8/SVneo cells seeded on poly-D-lysine or PSG1-Fc-coated wells with the corresponding densitometric analysis (right) normalized to total FAK are shown. **(C)** Fibroblast-deposited ECM was decellularized and PSG1-Fc (10 $\mu\text{g}/\text{mL}$) was added in the presence or absence of heparin for 1 h at RT. After several washes, the ECM was lysed and 5 μg was loaded on an SDS-PAGE gel and probed with Abs to human Fc (for PSG1-Fc detection), FN (for ECM detection) and GAPDH (to demonstrate successful cell removal; left). Densitometric analyses of the results are presented in arbitrary units (a.u.; right). **(D)** HTR8/SVneo cells were seeded in wells coated with PSG1-Fc or protein control (30 $\mu\text{g}/\text{mL}$) in the presence or absence of 50 $\mu\text{g}/\text{mL}$ heparin. The cell adhesion experiment was carried out as described in Figure 2. The adhesion of untreated cells to PSG1-Fc is considered as 100%. **(E)** PSG1-His or protein control were immobilized on poly-D-lysine-coated wells. Swan71 cells were seeded inside the 2-chamber inserts placed on the protein-coated wells and grown until confluent. Inserts were removed (0 h) to create a cell-free gap area. Cell migration was recorded periodically for 20 h and representative images are shown. **(F)** Quantitative analyses of a representative wound healing assay performed in 21% O_2 and in CoCl_2 -induced hypoxia-like conditions with Swan71 cells seeded on PSG1-His or control protein-coated wells. Gap closure speed and gap covered area were calculated as described in Materials and Methods. The average of the replicates obtained with the cells seeded on the control protein-coated wells in 21% O_2 is considered as 1. Results shown are mean \pm S.D. of triplicates in **(B–D)**. **(G)** Quantitative analyses of migration of single cells (primary EVTs, HTR8/SVneo and Swan71) seeded on PSG1-His or control protein-coated wells measured as track length in μm (**G**, left), velocity in $\mu\text{m}/\text{sec}$ (**G**, middle) and distance to previous point (D2P) in μm (**G**, right). *p* values were obtained by Student's *t*-test for **B** (** *p* < 0.003), by a one-way ANOVA followed by Sidak's multiple comparison tests for **C** (** *p* < 0.002) and by a two-way ANOVA followed by Sidak's multiple comparison tests for **(F–G)** (**** *p* < 0.0001, *** *p* < 0.0002, ** *p* < 0.002 and * *p* < 0.03).

3.6. PSG1 Increases EVT Migration

Integrin $\alpha 5\beta 1$ expression on EVTs is required for their migratory activity [61,62]. Therefore, we explored whether PSG1 can modulate migration of EVTs. We performed wound healing assays and found that Swan71 cells on the PSG1-His-coated wells migrated faster and covered the wound area almost completely in 20 h in comparison to the cells on the control protein-coated wells (Figure 3E). Cells on the PSG1-coated wells showed a faster gap closure speed and covered a larger area, which was 2–2.5 times over that of cells on the control protein-coated wells both in an oxygen concentration of 21% and in the presence of CoCl_2 (Figure 3F). We observed similar results when the wound healing assay was performed with HTR8/SVneo cells in 21% or 1% oxygen (data not shown). The observed increase in cell migration was not the result of increased proliferation or viability induced by PSG1 as HTR8/SVneo and Swan71 did not proliferate differently or differ in viability on PSG1 when compared to poly-D-lysine coated wells as determined with the MTS assay (data not shown). We also studied cell migration at the single cell level, which allowed us to investigate the effect of PSG1 on the migration of purified first-trimester EVTs (Figure 3G). We observed higher migration of primary EVTs, HTR8/SVneo and Swan71 on the PSG1-coated wells when compared to the control protein-coated wells as determined by measurements of track length, velocity and distance from previous point (D2P) of the tracked cells (Figure 3G). The average track length obtained for primary EVTs, HTR8/SVneo and Swan71 cells on the PSG1-coated wells were 908–1292 μm , which were more than twice of that of the cells on the control protein-coated wells (394–498 μm ; 3G, left). The velocities of single cells on the PSG1-coated wells were 0.019–0.033 $\mu\text{m}/\text{sec}$, which were also more than two times higher than the values obtained for the cells on the control protein-coated wells (0.0083–0.01 $\mu\text{m}/\text{s}$; Figure 3G, middle). Similarly, D2P values of single cells on the PSG1-coated wells (19.6–39.4 μm) were almost three times higher than the values obtained for the cells on the wells coated with control protein (8.9–12.4 μm ; Figure 3G, right). Altogether, the results indicate that PSG1 induces increased migration of EVTs.

3.7. The PSG1 Concentration Is Significantly Reduced in Pregnant Women Diagnosed with Early-Onset and Late-Onset PE

We explored whether PSG1 can modulate the invasive capacity of HTR8/SVneo and Swan71 cells on Matrigel in transwell and Biogel cell invasion assays. We were unable to detect any significant differences in the invasive capacity of these cells in the presence of PSG1 (data not shown). Invasion of EVTs into the endometrium is an essential process in placenta formation, and shallow invasion of the decidua and incomplete transformation of the spiral arteries by EVTs is believed to be an initiating cause of PE [28,63–66]. Although we did not observe any direct effect of PSG1 on EVT invasion, these in vitro studies have limitations [67]. Previous reports suggest that lower than normal levels of PSGs are found in some placental pathologies, including PE [68]; therefore, we investigated the PSG1 levels in PE with a validated specific PSG1 ELISA. We measured PSG1 in serum samples obtained from African American women diagnosed with early-onset and late-onset PE and the respective gestational age-matched controls with no pregnancy complications. We found that the concentration of PSG1 was significantly lower in women diagnosed with early-onset and late-onset PE compared to their respective controls (Figure 4A). The data was then analyzed taking into consideration the gender of the fetus. Interestingly, we found no statistical difference in the PSG1 concentration between early-onset or late-onset PE and the respective controls when the fetuses were female. On the other hand, when the fetuses were male, there was a significant difference for early-onset ($p = 0.002$) and late-onset ($p = 0.001$) PE and their respective gestational age-matched controls (Figure 4B).




Figure 4. Serum PSG1 concentration is lower in women diagnosed with early-onset and late-onset pre-eclampsia (PE) carrying a male fetus compared to gestational age-matched controls. (A) The PSG1 concentration was determined in triplicate in each serum sample using the Quantikine PSG1 ELISA kit. (B) PSG1 concentrations shown in A were analyzed considering the gender of the baby. *p* values were obtained by one-way ANOVA followed by pairwise comparisons using Tukey's adjustment for A and by a two-way ANOVA followed by pairwise comparison using Holm-Sidak's adjustment for B.

4. Discussion

PSGs were previously shown to be expressed exclusively by STBs by immunohistochemistry; however, some recent studies have reported expression of PSG mRNAs in EVT [32–34,69]. Therefore, to determine whether PSGs are present in EVTs at the protein level, we utilized two specific anti-PSG mAbs that do not react with any member of the CEACAM family, previously shown to be expressed in the placenta (data not shown) [70,71]. At present, there are no available Abs to a specific PSG family member that could be employed for immunohistochemistry. Characterization of the two mAbs utilized in this study showed that mAb#4 binds to the N-terminal domain of PSG1, PSG6, PSG7 and PSG8, and BAP3 binds to the B2 domain of PSG1, PSG3, PSG4, PSG6, PSG7 and PSG8 (data not shown). We observed positive reactivity with both anti-PSG mAbs in EVTs and STB of 5-week and 7-week-old placentas, with more intense staining in STBs as anticipated based on previously published RNA-seq data [34]. Analysis of staining in EVTs showed stronger staining in the more distal areas, suggesting increased expression of PSGs in the EVTs with an increased invasive cell phenotype. PSG expression in the EVTs invading into the decidua was mostly seen as localized puncta, possibly associated with FA structures. The discrepancy in our results from those of Zhou and co-workers, using the BAP3 mAb, is likely due to the difference in the tissue processing; as for these studies, we utilized antigen retrieval [32]. In addition, we showed co-localized expression of PSGs and integrin $\alpha 5$ in the first-trimester EVTs. EVTs differentiated from second-trimester CTBs also showed expression of PSGs as determined by the immunoblot analysis [46,47]. Although our study reports for the first time the presence of human PSGs in invasive EVTs, expression of mouse PSGs has been reported in trophoblast giant cells, which are similar to human EVTs, as well as in spongiotrophoblasts [35,72,73].

Here we show that immobilized PSG1 induces adhesion of two EVT-like cell lines, HTR8/SVneo and Swan71, and primary EVTs. Inhibition of PSG1-mediated adhesion by EDTA, an RGD-containing peptide, or the disintegrin echistatin pointed to the involvement of an integrin receptor [74,75]. Among the RGD-binding integrins, $\alpha V\beta 3$, $\alpha V\beta 5$ and $\alpha 5\beta 1$ are expressed on trophoblast cells at various stages of differentiation [52]. Integrin switching from $\alpha 6\beta 4$ in villous CTB, to $\alpha 5\beta 1$ in cell-column trophoblasts,

and to $\alpha 1\beta 1$ in mature invasive EVT_s has been reported in normal placental development [37]. Using function-blocking Abs to different integrins, we determined that PSG1-mediated adhesion was dependent on integrin $\alpha 5\beta 1$, a major fibronectin receptor [76]. The first-trimester placenta grows in a low oxygen or environment required for successful placenta development [77,78]. Since the differentiation of EVT_s occurs most extensively in the first-trimester, the developing EVT_s in the growing cell-columns of anchoring villi arise within an oxygen gradient with a shift from proliferative to non-proliferative phenotypes and changes in integrin expression, including that of integrin $\alpha 5$ [26]. Integrin $\alpha 5\beta 1$ has been shown to be up-regulated in low oxygen conditions via HIF1 α stabilization, which is only expressed in EVT_s of the early placenta [28,56]. We observed PSG1-mediated adhesion of EVT_s in ambient (21%) and in low (1%) oxygen conditions. In both cases, cell adhesion was dependent on the interaction of PSG1 with $\alpha 5\beta 1$. The requirement for $\alpha 5\beta 1$ integrin expression was further confirmed using a CHO cell line lacking integrin $\alpha 5$ as these cells did not attach to PSG1-coated wells while wild type cells did. Furthermore, we determined that the interaction of $\alpha 5\beta 1$ and native or recombinant PSG1 is direct by ELISA using purified proteins.

An interaction of PSGs with a different integrin, platelet $\alpha IIb\beta 3$, has been previously reported by Shanley and co-workers [9]. They observed that PSG1, PSG9 and mouse PSG23 bind to $\alpha IIb\beta 3$ and inhibit the interaction of platelets with fibrinogen. Similar to their observation, we show that more than one domain of the protein mediates the interaction with $\alpha 5\beta 1$ integrin as we observed that both the N and B2 domains of PSG1 individually induce EVT adhesion.

PSG1 induced adhesion of EVT_s only when immobilized on a well; therefore, as PSG1 was previously shown to bind to HSPGs, we hypothesized that PSG1 may bind simultaneously to the ECM and to $\alpha 5\beta 1$ on the cell membrane [17]. We observed that PSG1 binds to the ECM deposited by fibroblast and that this binding can be inhibited in the presence of excess heparin. On the other hand, PSG1-induced cell adhesion was not inhibited by heparin indicating that cells can potentially bind to PSG1 anchored to the decidual ECM via its interaction with HSPGs, facilitating the attachment of EVT_s to the decidua.

Clustering of ligand-engaged integrins leads to FA assembly that involves the recruitment of signaling molecules similar to FAK and cytoskeletal adaptor proteins, such as vinculin and paxillin, to the integrin cytoplasmic tails [79]. FAK, a mediator of integrin-mediated signaling pathways, has been shown to regulate anchorage-dependent behaviors such as cell proliferation, migration and anoikis suppression [79]. The observed localization of vinculin and paxillin only in the EVT_s bound to PSG1 and not to poly-D-lysine is consistent with the PSG1-induced FAK phosphorylation [80]. Integrin $\alpha 5\beta 1$ -dependent anchorage of trophoblast to the uterine ECM and FAK signaling are well documented as important regulators of EVT migration during early placental development [31,80]. Integrin $\alpha 5\beta 1$ expression in EVT_s has been shown to be essential for their migration and IGF-stimulated enhancement in EVT migration is $\alpha 5\beta 1$ dependent [31,61,62]. In accordance with these previous findings, we show that adhesion to PSG1 increased the migratory capacity of primary EVT_s and EVT-like cell lines.

The influence of integrin $\alpha 5$ on EVT invasion is controversial with some groups proposing that binding to fibronectin via this integrin has invasion-restraining and others proposing an invasion-promoting effect [37,81]. In our studies using two different invasion assays, we did not observe differences in invasion between cells incubated with PSG1 and control protein. While the interaction of $\alpha 5\beta 1$ with fibronectin likely differs from the interaction of $\alpha 5\beta 1$ with PSG1, the inherent limitations of the in vitro assays employed does not properly address the complexity of the in vivo EVT invasion process including the composition of the ECM and the presence of PSG1 in the matrix rather than in the solution.

Several studies conducted to investigate the correlation of serum PSG levels with various pregnancy complications reported conflicting results [68,82]. However, inherent limitations of the assays used to measure PSG concentration as well as significant inter- and intra- patient variability may explain some of these discrepancies [83–85]. Earlier studies reporting on the association between the PSG level and pregnancy complications did not take into consideration the difference in incidence of PE across

different ethnicities and have generally involved small numbers of patients insufficient to produce reproducible and generalizable conclusions [86–88]. However, some recent studies identified some risk factors that are differentially associated with certain forms of PE, such as African American race for early-onset PE [89]. Recent reports have revisited the association of PE with PSG levels. A proteomic study identified some PSGs as potential predictive markers of early-onset PE [90]. Another study using global RNA profiling has reported down-regulated PSG expression in patients with severe PE [33]. Down-regulation of mRNA coding for some PSG family members and of PSG-derived peptides, had yet to be validated at the protein level [33,90,91]. We measured the concentration of PSG1, the most highly expressed member of the family based on mRNA expression data, to determine whether prior observations that PSGs are found at lower concentrations in pregnant women suffering from PE could be replicated with an ELISA using a validated anti-PSG1 antibody pair [9]. We observed significant differences in PSG1 concentration in African American women diagnosed with early-onset and late-onset PE compared to their gestational age-matched controls. It is likely that a major fraction of PSG1 measured in maternal serum is derived from the multinucleated STB layer of the placenta. Therefore, both the STBs and EVT_s can contribute to the circulating levels of PSG1 in the mother. Abnormally low levels of PSG1 in women with preeclampsia may reflect dysfunctional or stressed STBs, which may in turn contribute to the pathogenesis of this placental syndrome [92]. In contrast, in this study we show that staining of PSG in EVT is located at the membrane where its functional interaction with integrin $\alpha 5$ and focal adhesions mediate migration. Due to the limited availability of placental bed biopsies from preeclamptic women we were unable to assess levels of PSG1 in this EVT context in this study. Whether PSGs also affect the initial penetration of the uterine epithelium by blastocytic STBs that occurs in the earliest stages of implantation requires further investigation. Fetal gender-specific differences in gene expression during pregnancy have been reported for cytokines, pro-angiogenic factors, and some galectins [93,94]. In addition, carrying of a male fetus has been identified as an increased risk factor for adverse pregnancy outcomes [95,96]. Therefore, we also analyzed the expression of PSG1 considering a fetal gender. Interestingly, our results indicate that samples obtained from women carrying a male fetus are the major contributors to the observed differences in PSG1 concentration. At present, the mechanisms that regulate the expression of PSGs in the placenta are not completely understood, and whether fetal sex influences their expression has never been investigated. Additional studies using well-validated reagents, which do not cross-react with members of the CEACAM family and can detect only PSG family members, are required to determine whether the concentration of PSGs as a whole are diminished in PE and are also lower than normal in women of different ethnicities.

Author Contributions: C.E.D., S.M.B., K.M.Y., and G.D. conceptualized and designed the research. S.R., C.E.D., M.I., O.G., Y.Z., M.C., and M.Y.-M. performed the research; S.R., C.E.D., M.I., Y.Z., M.Y.-M., K.M.Y., O.E., N.G.-L., B.H., S.J.L., R.R., and G.D. contributed in data interpretation. C.E.D., M.C., K.M.Y., O.E., N.G.-L., and R.R. contributed with essential reagents, primary cells and human serum samples. All authors revised the manuscript; S.R. and G.D. wrote the manuscript and S.M.B. helped in generating figures.

Funding: The research was supported by a grant from the Collaborative Health Initiative Research Program (CHIRP#64532) to GD. The opinions expressed here are those of the authors and should not be construed as official or reflecting the views of the Uniformed Services University of the Health Sciences or the Department of Defense. CD and SJL were funded by FDN-143262. SMB was supported by the Deutsche Forschungsgemeinschaft (DFG) through the Heisenberg Program (BL1115/3-1) and BL1115/4-1. The research of BH was funded through a Foundation Grant by the Canadian Institutes of Health Research (CIHR, #375597). This research was supported, in part, by the Perinatology Research Branch, Division of Obstetrics and Maternal-Fetal Medicine, Division of Intramural Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development, National Institutes of Health (NIH), U.S. Department of Health and Human Services (NICHD/NIH/DHHS); and, in part, with Federal funds from NICHD/NIH/DHHS under Contract No. HHSN275201300006C. Dr. Romero has contributed to this work as part of his official duties as an employee of the United States Federal Government. In addition, this work was supported in part by the Intramural Research Program of the NIH, NIDCR. N.G.-L. is supported by the Wayne State University Perinatal Research Initiative. No conflicts of interests are declared by the authors.

Acknowledgments: The authors thank the donors and the Research Centre for Women’s and Infants’ Health BioBank Program of the CIHR Group in Development and Fetal Health (CIHR MGC-13299), the Samuel Lunenfeld Research Institute, and the MSH/UHN Dept. of Obstetrics & Gynecology for the human specimens used in this study. We would also like to thank Cara Olsen for assistance with the statistical analysis and James Warren for his technical contribution with protein purification.

Conflicts of Interest: The authors report no financial or other conflict of interest relevant to this study. The funders had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

Appendix A




Figure A1. Co-localization of PSGs with integrin $\alpha 5$, PSG1 expression in EVTs differentiated in vitro, PSG1-mediated adhesion of HTR8/SVneo cells, HIF1 α stabilization by CoCl_2 and integrin $\alpha 5\beta 1$ expression in EVT-like cell lines. (a) Representative photographs of dual fluorescent immunohistochemistry localizing PSGs (red) and integrin $\alpha 5$ (green) are shown in both individual and dual channels in sections of first-trimester placenta. Scale bars = 45 μm . ctrl = control Ab. (b) Immunoblot of cytotrophoblast (CTB) lysates with two different anti-PSG mAbs shows no PSG expression in undifferentiated (0 h) second-trimester CTBs but high expression in the EVTs that were differentiated in vitro (14 h and 20 h) from second-trimester CTBs. (c) HTR8/SVneo cells that were pre-incubated with the indicated Abs were seeded on wells coated with 30 $\mu\text{g}/\text{mL}$ PSG1-His or control protein. When indicated, cells were pre-treated 4 h with 100 μM CoCl_2 and the treatment was maintained during the experiment. Cell adhesion experiments were carried out as described in Figure 2. (d) Representative image of Western blot showing HIF1 α stabilization in HTR8/SVneo and Swan71 cells treated with 100 μM CoCl_2 for 4 h. (e) HTR8/SVneo and Swan71 were incubated with anti-integrin $\alpha 5\beta 1$ or isotype control mAb followed by APC-conjugated anti-mouse Ig. MFI = mean fluorescence intensity.

References

- Zhou, G.Q.; Hammarstrom, S. Pregnancy-specific glycoprotein (PSG) in baboon (*Papio hamadryas*): Family size, domain structure, and prediction of a functional region in primate PSGs. *Biol. Reprod.* **2001**, *64*, 90–99. [[CrossRef](#)]
- Kammerer, R.; Mansfeld, M.; Hanske, J.; Missbach, S.; He, X.; Kollner, B.; Mouchantat, S.; Zimmermann, W. Recent expansion and adaptive evolution of the carcinoembryonic antigen family in bats of the Yangochiroptera subgroup. *BMC Genom.* **2017**, *18*, 717. [[CrossRef](#)]

3. Sorensen, S. Pregnancy-“specific” beta 1-glycoprotein (SP1): Purification, characterization, quantification and clinical application in malignancies (a review). *Tumour Biol.* **1984**, *5*, 275–302.
4. Bohn, H. Detection and characterization of pregnancy proteins in the human placenta and their quantitative immunochemical determination in sera from pregnant women. *Arch. Gynakol.* **1971**, *210*, 440–457. [[CrossRef](#)]
5. Kammerer, R.; Zimmermann, W. Coevolution of activating and inhibitory receptors within mammalian carcinoembryonic antigen families. *BMC Biol.* **2010**, *8*, 12. [[CrossRef](#)]
6. Moore, T.; Dveksler, G.S. Pregnancy-specific glycoproteins: Complex gene families regulating maternal-fetal interactions. *Int. J. Dev. Biol.* **2014**, *58*, 273–280. [[CrossRef](#)]
7. McLellan, A.S.; Fischer, B.; Dveksler, G.; Hori, T.; Wynne, F.; Ball, M.; Okumura, K.; Moore, T.; Zimmermann, W. Structure and evolution of the mouse pregnancy-specific glycoprotein (Psg) gene locus. *BMC Genom.* **2005**, *6*, 4. [[CrossRef](#)]
8. Teglund, S.; Olsen, A.; Khan, W.N.; Frangsmyr, L.; Hammarstrom, S. The pregnancy-specific glycoprotein (PSG) gene cluster on human chromosome 19: Fine structure of the 11 PSG genes and identification of 6 new genes forming a third subgroup within the carcinoembryonic antigen (CEA) family. *Genomics* **1994**, *23*, 669–684. [[CrossRef](#)]
9. Shanley, D.K.; Kiely, P.A.; Golla, K.; Allen, S.; Martin, K.; O’Riordan, R.T.; Ball, M.; Aplin, J.D.; Singer, B.B.; Caplice, N.; et al. Pregnancy-specific glycoproteins bind integrin alphaIIb beta3 and inhibit the platelet-fibrinogen interaction. *PLoS ONE* **2013**, *8*, e57491. [[CrossRef](#)]
10. Wurz, H.; Geiger, W.; Kunzig, H.J.; Jabs-Lehmann, A.; Bohn, H.; Luben, G. Radioimmunoassay of SP1 (pregnancy-specific beta1-glycoprotein) in maternal blood and in amniotic fluid normal and pathologic pregnancies. *J. Perinat. Med.* **1981**, *9*, 67–78. [[CrossRef](#)]
11. McLellan, A.S.; Zimmermann, W.; Moore, T. Conservation of pregnancy-specific glycoprotein (PSG) N domains following independent expansions of the gene families in rodents and primates. *BMC Evol. Biol.* **2005**, *5*, 39. [[CrossRef](#)]
12. Jones, K.; Ballesteros, A.; Mentink-Kane, M.; Warren, J.; Rattila, S.; Malech, H.; Kang, E.; Dveksler, G. PSG9 stimulates increase in FoxP3+ regulatory T-Cells through the TGF-beta1 pathway. *PLoS ONE* **2016**, *11*, e0158050. [[CrossRef](#)]
13. Warren, J.; Im, M.; Ballesteros, A.; Ha, C.; Moore, T.; Lambert, F.; Lucas, S.; Hinz, B.; Dveksler, G. Activation of latent transforming growth factor-beta1, a conserved function for pregnancy-specific beta 1-glycoproteins. *Mol. Hum. Reprod.* **2018**, *24*, 602–612. [[CrossRef](#)]
14. Snyder, S.K.; Wessner, D.H.; Wessells, J.L.; Waterhouse, R.M.; Wahl, L.M.; Zimmermann, W.; Dveksler, G.S. Pregnancy-Specific glycoproteins function as immunomodulators by inducing secretion of IL-10, IL-6 and TGF-beta1 by human monocytes. *Am. J. Reprod. Immunol.* **2001**, *45*, 205–216. [[CrossRef](#)]
15. Motran, C.C.; Diaz, F.L.; Montes, C.L.; Bocco, J.L.; Gruppi, A. In vivo expression of recombinant pregnancy-specific glycoprotein 1a induces alternative activation of monocytes and enhances Th2-type immune response. *Eur. J. Immunol.* **2003**, *33*, 3007–3016. [[CrossRef](#)]
16. Blois, S.M.; Sulkowski, G.; Tirado-Gonzalez, I.; Warren, J.; Freitag, N.; Klapp, B.F.; Rifkin, D.; Fuss, I.; Strober, W.; Dveksler, G.S. Pregnancy-specific glycoprotein 1 (PSG1) activates TGF-beta and prevents dextran sodium sulfate (DSS)-induced colitis in mice. *Mucosal Immunol.* **2014**, *7*, 348–358. [[CrossRef](#)]
17. Lisboa, F.A.; Warren, J.; Sulkowski, G.; Aparicio, M.; David, G.; Zudaire, E.; Dveksler, G.S. Pregnancy-specific glycoprotein 1 induces endothelial tubulogenesis through interaction with cell surface proteoglycans. *J. Biol. Chem.* **2011**, *286*, 7577–7586. [[CrossRef](#)]
18. Ha, C.T.; Waterhouse, R.; Warren, J.; Zimmermann, W.; Dveksler, G.S. N-Glycosylation is required for binding of murine pregnancy-specific glycoproteins 17 and 19 to the receptor CD9. *Am. J. Reprod. Immunol.* **2008**, *59*, 251–258. [[CrossRef](#)]
19. Houston, A.; Williams, J.M.; Rovis, T.L.; Shanley, D.K.; O’Riordan, R.T.; Kiely, P.A.; Ball, M.; Barry, O.P.; Kelly, J.; Fanning, A.; et al. Pregnancy-Specific glycoprotein expression in normal gastrointestinal tract and in tumors detected with novel monoclonal antibodies. *MAbs* **2016**, *8*, 491–500. [[CrossRef](#)]
20. Shupert, W.L.; Chan, W.Y. Pregnancy specific beta 1-glycoprotein in human intestine. *Mol. Cell Biochem.* **1993**, *120*, 159–170. [[CrossRef](#)]

21. Lee, J.N.; Salem, H.T.; Al-Ani, A.T.; Chard, T.; Huang, S.C.; Ouyang, P.C.; Wei, P.Y.; Seppala, M. Circulating concentrations of specific placental proteins (human chorionic gonadotropin, pregnancy-specific beta-1 glycoprotein, and placental protein 5) in untreated gestational trophoblastic tumors. *Am. J. Obstet. Gynecol.* **1981**, *139*, 702–704. [[CrossRef](#)]
22. Campo, E.; Algaba, F.; Palacin, A.; Germa, R.; Sole-Balcells, F.J.; Cardesa, A. Placental proteins in high-grade urothelial neoplasms. An immunohistochemical study of human chorionic gonadotropin, human placental lactogen, and pregnancy-specific beta-1-glycoprotein. *Cancer* **1989**, *63*, 2497–2504. [[CrossRef](#)]
23. Salahshor, S.; Goncalves, J.; Chetty, R.; Gallinger, S.; Woodgett, J.R. Differential gene expression profile reveals deregulation of pregnancy specific beta1 glycoprotein 9 early during colorectal carcinogenesis. *BMC Cancer* **2005**, *5*, 66. [[CrossRef](#)]
24. Kawano, K.; Ebisawa, M.; Hase, K.; Fukuda, S.; Hijikata, A.; Kawano, S.; Date, Y.; Tsuneda, S.; Itoh, K.; Ohno, H. Psg18 is specifically expressed in follicle-associated epithelium. *Cell Struct. Funct.* **2007**, *32*, 115–126. [[CrossRef](#)]
25. Maltepe, E.; Fisher, S.J. Placenta: The forgotten organ. *Annu. Rev. Cell Dev. Biol.* **2015**, *31*, 523–552. [[CrossRef](#)]
26. Chang, C.W.; Wakeland, A.K.; Parast, M.M. Trophoblast lineage specification, differentiation and their regulation by oxygen tension. *J. Endocrinol.* **2018**, *236*, R43–R56. [[CrossRef](#)]
27. Tarrade, A.; Lai Kuen, R.; Malassiné, A.; Tricottet, V.; Blain, P.; Vidaud, M.; Evain-Brion, D. Characterization of human villous and extravillous trophoblasts isolated from first trimester placenta. *Lab. Investig.* **2001**, *81*. [[CrossRef](#)]
28. Knofler, M.; Haider, S.; Saleh, L.; Pollheimer, J.; Gamage, T.; James, J. Human placenta and trophoblast development: Key molecular mechanisms and model systems. *Cell Mol. Life Sci.* **2019**. [[CrossRef](#)]
29. Pollheimer, J.; Vondra, S.; Baltayeva, J.; Beristain, A.G.; Knöfler, M. Regulation of placental extravillous trophoblasts by the maternal uterine environment. *Front Immunol.* **2018**, *9*. [[CrossRef](#)]
30. Pujal, J.M.; Roura, S.; Munoz-Marmol, A.M.; Mate, J.L.; Bayes-Genis, A. Fetal-maternal interface: A chronicle of allogeneic coexistence. *Chimerism* **2012**, *3*, 18–20. [[CrossRef](#)]
31. Gleeson, L.M.; Chakraborty, C.; McKinnon, T.; Lala, P.K. Insulin-Like growth factor-binding protein 1 stimulates human trophoblast migration by signaling through alpha 5 beta 1 integrin via mitogen-activated protein Kinase pathway. *J. Clin. Endocrinol. Metab.* **2001**, *86*, 2484–2493. [[CrossRef](#)] [[PubMed](#)]
32. Zhou, G.Q.; Baranov, V.; Zimmermann, W.; Grunert, F.; Erhard, B.; Mincheva-Nilsson, L.; Hammarstrom, S.; Thompson, J. Highly specific monoclonal antibody demonstrates that pregnancy-specific glycoprotein (PSG) is limited to syncytiotrophoblast in human early and term placenta. *Placenta* **1997**, *18*, 491–501. [[CrossRef](#)]
33. Gormley, M.; Ona, K.; Kapidzic, M.; Garrido-Gomez, T.; Zdravkovic, T.; Fisher, S.J. Preeclampsia: Novel insights from global RNA profiling of trophoblast subpopulations. *Am. J. Obstet. Gynecol.* **2017**, *217*. [[CrossRef](#)] [[PubMed](#)]
34. Okae, H.; Toh, H.; Sato, T.; Hiura, H.; Takahashi, S.; Shirane, K.; Kabayama, Y.; Suyama, M.; Sasaki, H.; Arima, T. Derivation of human trophoblast stem cells. *Cell Stem Cell* **2018**, *22*, 50–63. [[CrossRef](#)] [[PubMed](#)]
35. Blois, S.; Tirado-Gonzalez, I.; Wu, J.; Barrientos, G.; Johnson, B.; Warren, J.; Freitag, N.; Klapp, B.; Irmak, S.; Ergun, S.; et al. Early expression of pregnancy-specific glycoprotein 22 (PSG22) by trophoblast cells modulates angiogenesis in mice. *Biol. Reprod.* **2012**, *86*, 1–9. [[CrossRef](#)]
36. Sulkowski, G.N.; Warren, J.; Ha, C.T.; Dveksler, G.S. Characterization of receptors for murine pregnancy specific glycoproteins 17 and 23. *Placenta* **2011**, *32*, 603–610. [[CrossRef](#)]
37. Damsky, C.H.; Librach, C.; Lim, K.H.; Fitzgerald, M.L.; McMaster, M.T.; Janatpour, M.; Zhou, Y.; Logan, S.K.; Fisher, S.J. Integrin switching regulates normal trophoblast invasion. *Development* **1994**, *120*, 3657–3666.
38. Korhonen, M.; Ylänne, J.; Laitinen, L.; Cooper, H.M.; Quaranta, V.; Virtanen, I. Distribution of the alpha 1-alpha 6 integrin subunits in human developing and term placenta. *Lab. Investig.* **1991**, *65*, 347–356.
39. Aplin, J.D. Expression of integrin alpha 6 beta 4 in human trophoblast and its loss from extravillous cells. *Placenta* **1993**, *14*, 203–215. [[CrossRef](#)]
40. Silva, J.F.; Serakides, R. Intrauterine trophoblast migration: A comparative view of humans and rodents. *Cell Adh. Migr.* **2016**, *10*, 88–110. [[CrossRef](#)]
41. Bronson, R.A.; Fusi, F.M. Integrins and human reproduction. *Mol. Hum. Reprod.* **1996**, *2*, 153–168. [[CrossRef](#)] [[PubMed](#)]

42. Bilban, M.; Haslinger, P.; Prast, J.; Klinglmüller, F.; Woelfel, T.; Haider, S.; Sachs, A.; Otterbein, L.E.; Desoye, G.; Hiden, U.; et al. Identification of novel trophoblast invasion-related genes: Heme oxygenase-1 controls motility via peroxisome proliferator-activated receptor gamma. *Endocrinology* **2009**, *150*, 1000–1013. [[CrossRef](#)] [[PubMed](#)]
43. Zhou, Y.; Damsky, C.H.; Chiu, K.; Roberts, J.M.; Fisher, S.J. Preeclampsia is associated with abnormal expression of adhesion molecules by invasive cytotrophoblasts. *J. Clin. Investig.* **1993**, *91*, 950–960. [[CrossRef](#)] [[PubMed](#)]
44. Ballesteros, A.; Mentink-Kane, M.M.; Warren, J.; Kaplan, G.G.; Dveksler, G.S. Induction and activation of latent transforming growth factor-beta1 are carried out by two distinct domains of pregnancy-specific glycoprotein 1 (PSG1). *J. Biol. Chem.* **2015**, *290*, 4422–4431. [[CrossRef](#)]
45. Cohen, M.; Ribaux, P.; Epiney, M.; Irion, O. Role of prostate apoptosis response 4 in translocation of GRP78 from the endoplasmic reticulum to the cell surface of trophoblastic cells. *PLoS ONE* **2013**, *8*, e80231. [[CrossRef](#)]
46. Fisher, S.J.; Cui, T.Y.; Zhang, L.; Hartman, L.; Grahl, K.; Zhang, G.Y.; Tarpey, J.; Damsky, C.H. Adhesive and degradative properties of human placental cytotrophoblast cells in vitro. *J. Cell Biol.* **1989**, *109*, 891–902. [[CrossRef](#)]
47. Zhou, Y.; Fisher, S.J.; Janatpour, M.; Genbacev, O.; Dejana, E.; Wheelock, M.; Damsky, C.H. Human cytotrophoblasts adopt a vascular phenotype as they differentiate. A strategy for successful endovascular invasion? *J. Clin. Investig.* **1997**, *99*, 2139–2151. [[CrossRef](#)]
48. Akiyama, S.K.; Yamada, S.S.; Chen, W.T.; Yamada, K.M. Analysis of fibronectin receptor function with monoclonal antibodies: Roles in cell adhesion, migration, matrix assembly, and cytoskeletal organization. *J. Cell Biol.* **1989**, *109*, 863–875. [[CrossRef](#)]
49. Yamada, K.M.; Kennedy, D.W.; Yamada, S.S.; Gralnick, H.; Chen, W.T.; Akiyama, S.K. Monoclonal antibody and synthetic peptide inhibitors of human tumor cell migration. *Cancer Res.* **1990**, *50*, 4485–4496.
50. ACOG Committee on Obstetric Practice. ACOG practice bulletin. Diagnosis and management of preeclampsia and eclampsia. *Obstet. Gynecol.* **2002**, *99*, 159–167.
51. Von Dadelszen, P.; Magee, L.A.; Roberts, J.M. Subclassification of preeclampsia. *Hypertens. Pregnancy* **2003**, *22*, 143–148. [[CrossRef](#)] [[PubMed](#)]
52. Harris, L.K.; Jones, C.J.; Aplin, J.D. Adhesion molecules in human trophoblast—A review. II. extravillous trophoblast. *Placenta* **2009**, *30*, 299–304. [[CrossRef](#)] [[PubMed](#)]
53. Jauniaux, E.; Watson, A.L.; Hempstock, J.; Bao, Y.P.; Skepper, J.N.; Burton, G.J. Onset of maternal arterial blood flow and placental oxidative stress. A possible factor in human early pregnancy failure. *Am. J. Pathol.* **2000**, *157*, 2111–2122. [[CrossRef](#)]
54. Rodesch, F.; Simon, P.; Donner, C.; Jauniaux, E. Oxygen measurements in endometrial and trophoblastic tissues during early pregnancy. *Obstet. Gynecol.* **1992**, *80*, 283–285. [[PubMed](#)]
55. Yamanaka-Tatematsu, M.; Nakashima, A.; Fujita, N.; Shima, T.; Yoshimori, T.; Saito, S. Autophagy induced by HIF1alpha overexpression supports trophoblast invasion by supplying cellular energy. *PLoS ONE* **2013**, *8*, e76605. [[CrossRef](#)] [[PubMed](#)]
56. Arimoto-Ishida, E.; Sakata, M.; Sawada, K.; Nakayama, M.; Nishimoto, F.; Mabuchi, S.; Takeda, T.; Yamamoto, T.; Isobe, A.; Okamoto, Y.; et al. Up-Regulation of alpha5-integrin by E-cadherin loss in hypoxia and its key role in the migration of extravillous trophoblast cells during early implantation. *Endocrinology* **2009**, *150*, 4306–4315. [[CrossRef](#)] [[PubMed](#)]
57. Wu, C.; Bauer, J.S.; Juliano, R.L.; McDonald, J.A. The alpha 5 beta 1 integrin fibronectin receptor, but not the alpha 5 cytoplasmic domain, functions in an early and essential step in fibronectin matrix assembly. *J. Biol. Chem.* **1993**, *268*, 21883–21888.
58. Orecchia, A.; Mettouchi, A.; Uva, P.; Simon, G.C.; Arcelli, D.; Avitabile, S.; Ragone, G.; Meneguzzi, G.; Pfenninger, K.H.; Zambruno, G.; et al. Endothelial cell adhesion to soluble vascular endothelial growth factor receptor-1 triggers a cell dynamic and angiogenic phenotype. *FASEB J.* **2014**, *28*, 692–704. [[CrossRef](#)]
59. Chen, C.P.; Liu, S.H.; Lee, M.Y.; Chen, Y.Y. Heparan sulfate proteoglycans in the basement membranes of the human placenta and decidua. *Placenta* **2008**, *29*, 309–316. [[CrossRef](#)]
60. Sarrazin, S.; Lamanna, W.C.; Esko, J.D. Heparan sulfate proteoglycans. *Cold Spring Harb. Perspect. Biol.* **2011**, *3*. [[CrossRef](#)]
61. Irving, J.A.; Lala, P.K. Functional role of cell surface integrins on human trophoblast cell migration: Regulation by TGF-beta, IGF-II, and IGFBP-1. *Exp. Cell Res.* **1995**, *217*, 419–427. [[CrossRef](#)] [[PubMed](#)]

62. Kabir-Salmani, M. The role of 51-integrin in the IGF-I-induced migration of extravillous trophoblast cells during the process of implantation. *Mol. Hum. Reprod.* **2004**, *10*, 91–97. [[CrossRef](#)] [[PubMed](#)]
63. Dunk, C.E.; Pappas, J.J.; Lye, P.; Kibschull, M.; Javam, M.; Bloise, E.; Lye, S.J.; Szyf, M.; Matthews, S.G. P-Glycoprotein (P-gp)/ABCB1 plays a functional role in extravillous trophoblast (EVT) invasion and is decreased in the pre-eclamptic placenta. *J. Cell Mol. Med.* **2018**, *22*, 5378–5393. [[CrossRef](#)] [[PubMed](#)]
64. Zhou, Y.; Damsky, C.H.; Fisher, S.J. Preeclampsia is associated with failure of human cytotrophoblasts to mimic a vascular adhesion phenotype. One cause of defective endovascular invasion in this syndrome? *J. Clin. Investig.* **1997**, *99*, 2152–2164. [[CrossRef](#)]
65. Aplin, J.D.; Beristain, A.; DaSilva-Arnold, S.; Dunk, C.; Duzyj, C.; Golos, T.G.; Kemmerling, U.; Knofler, M.; Mitchell, M.D.; Olson, D.M.; et al. IFPA meeting 2016 workshop report III: Decidua-trophoblast interactions; trophoblast implantation and invasion; immunology at the maternal-fetal interface; placental inflammation. *Placenta* **2017**. [[CrossRef](#)]
66. Khong, T.Y.; De Wolf, F.; Robertson, W.B.; Brosens, I. Inadequate maternal vascular response to placentation in pregnancies complicated by pre-eclampsia and by small-for-gestational age infants. *Br. J. Obstet. Gynaecol.* **1986**, *93*, 1049–1059. [[CrossRef](#)]
67. Kramer, N.; Walzl, A.; Unger, C.; Rosner, M.; Krupitza, G.; Hengstschlager, M.; Dolznig, H. In vitro cell migration and invasion assays. *Mutat. Res.* **2013**, *752*, 10–24. [[CrossRef](#)]
68. Pihl, K.; Larsen, T.; Laursen, I.; Krebs, L.; Christiansen, M. First trimester maternal serum pregnancy-specific beta-1-glycoprotein (SP1) as a marker of adverse pregnancy outcome. *Prenat. Diagn.* **2009**, *29*, 1256–1261. [[CrossRef](#)]
69. Camolotto, S.; Racca, A.; Rena, V.; Nores, R.; Patrino, L.C.; Genti-Raimondi, S.; Panzetta-Dutari, G.M. Expression and transcriptional regulation of individual pregnancy-specific glycoprotein genes in differentiating trophoblast cells. *Placenta* **2010**, *31*, 312–319. [[CrossRef](#)]
70. Bamberger, A.M.; Sudahl, S.; Loning, T.; Wagener, C.; Bamberger, C.M.; Drakakis, P.; Coutifaris, C.; Makrigiannakis, A. The adhesion molecule CEACAM1 (CD66a, C-CAM, BGP) is specifically expressed by the extravillous intermediate trophoblast. *Am. J. Pathol.* **2000**, *156*, 1165–1170. [[CrossRef](#)]
71. Shao, L.; Jacobs, A.R.; Johnson, V.V.; Mayer, L. Activation of CD8+ regulatory T cells by human placental trophoblasts. *J. Immunol.* **2005**, *174*, 7539–7547. [[CrossRef](#)] [[PubMed](#)]
72. Kromer, B.; Finkenzeller, D.; Wessels, J.; Dveksler, G.; Thompson, J.; Zimmermann, W. Coordinate expression of splice variants of the murine pregnancy-specific glycoprotein (PSG) gene family during placental development. *Eur. J. Biochem.* **1996**, *242*, 280–287. [[CrossRef](#)] [[PubMed](#)]
73. Wynne, F.; Ball, M.; McLellan, A.S.; Dockery, P.; Zimmermann, W.; Moore, T. Mouse pregnancy-specific glycoproteins: Tissue-specific expression and evidence of association with maternal vasculature. *Reproduction* **2006**, *131*, 721–732. [[CrossRef](#)] [[PubMed](#)]
74. Carlson, T.R.; Feng, Y.; Maisonpierre, P.C.; Mrksich, M.; Morla, A.O. Direct cell adhesion to the angiopoietins mediated by integrins. *J. Biol. Chem.* **2001**, *276*, 26516–26525. [[CrossRef](#)]
75. Wang, Z.; Collighan, R.J.; Gross, S.R.; Danen, E.H.; Orend, G.; Telci, D.; Griffin, M. RGD-independent cell adhesion via a tissue transglutaminase-fibronectin matrix promotes fibronectin fibril deposition and requires syndecan-4/2 alpha5beta1 integrin co-signaling. *J. Biol. Chem.* **2010**, *285*, 40212–40229. [[CrossRef](#)]
76. Pankov, R.; Yamada, K.M. Fibronectin at a glance. *J. Cell Sci.* **2002**, *115*, 3861–3863. [[CrossRef](#)]
77. Huppertz, B.; Gauster, M.; Orendi, K.; Konig, J.; Moser, G. Oxygen as modulator of trophoblast invasion. *J. Anat.* **2009**, *215*, 14–20. [[CrossRef](#)]
78. Jauniaux, E.; Watson, A.; Burton, G. Evaluation of respiratory gases and acid-base gradients in human fetal fluids and uteroplacental tissue between 7 and 16 weeks' gestation. *Am. J. Obstet. Gynecol.* **2001**, *184*, 998–1003. [[CrossRef](#)]
79. Rainero, E.; Norman, J.C. Endosomal integrin signals for survival. *Nat. Cell Biol.* **2015**, *17*, 1373–1375. [[CrossRef](#)]
80. MacPhee, D.J.; Mostachfi, H.; Han, R.; Lye, S.J.; Post, M.; Caniggia, I. Focal adhesion kinase is a key mediator of human trophoblast development. *Lab. Investig.* **2001**, *81*, 1469–1483. [[CrossRef](#)]
81. Zeng, B.X.; Fujiwara, H.; Sato, Y.; Nishioka, Y.; Yamada, S.; Yoshioka, S.; Ueda, M.; Higuchi, T.; Fujii, S. Integrin alpha5 is involved in fibronectin-induced human extravillous trophoblast invasion. *J. Reprod. Immunol.* **2007**, *73*, 1–10. [[CrossRef](#)] [[PubMed](#)]

82. Bersinger, N.A.; Odegard, R.A. Second- and third-trimester serum levels of placental proteins in preeclampsia and small-for-gestational age pregnancies. *Acta Obstet. Gynecol. Scand.* **2004**, *83*, 37–45. [[CrossRef](#)] [[PubMed](#)]
83. Amsalem, H.; Gaiger, A.; Mizrahi, S.; Yagel, S.; Rachmilewitz, J. Characterization of a lymphocyte subset displaying a unique regulatory activity in human decidua. *Int. Immunol.* **2008**, *20*, 1147–1154. [[CrossRef](#)] [[PubMed](#)]
84. Salem, H.T.; Lee, J.N.; Seppala, M.; Vaara, L.; Aula, P.; Al-Ani, A.T.; Chard, T. Measurement of placental protein 5, placental lactogen and pregnancy-specific beta 1 glycoprotein in mid-trimester as a predictor of outcome of pregnancy. *Br. J. Obstet. Gynaecol.* **1981**, *88*, 371–374. [[CrossRef](#)] [[PubMed](#)]
85. Towler, C.M.; Horne, C.H.; Jandial, V.; Campbell, D.M.; MacGillivray, I. Plasma levels of pregnancy-specific beta 1-glycoprotein in complicated pregnancies. *Br. J. Obstet. Gynaecol.* **1977**, *84*, 258–263. [[CrossRef](#)] [[PubMed](#)]
86. Irwin, D.E.; Savitz, D.A.; Hertz-Picciotto, I.; St Andre, K.A. The risk of pregnancy-induced hypertension: Black and white differences in a military population. *Am. J. Public Health* **1994**, *84*, 1508–1510. [[CrossRef](#)]
87. Deepthi, G.; Chaithri, P.K.; Latha, P.; Usha Rani, V.; Rahman, P.F.; Jahan, P. TGFB1 functional gene polymorphisms (C-509T and T869C) in the maternal susceptibility to pre-eclampsia in South Indian Women. *Scand. J. Immunol.* **2015**, *82*, 390–397. [[CrossRef](#)]
88. Tanaka, M.; Jaamaa, G.; Kaiser, M.; Hills, E.; Soim, A.; Zhu, M.; Shcherbatykh, I.Y.; Samelson, R.; Bell, E.; Zdeb, M.; et al. Racial disparity in hypertensive disorders of pregnancy in New York State: A 10-year longitudinal population-based study. *Am. J. Public Health* **2007**, *97*, 163–170. [[CrossRef](#)]
89. Lisonkova, S.; Joseph, K.S. Incidence of preeclampsia: Risk factors and outcomes associated with early-versus late-onset disease. *Am. J. Obstet. Gynecol.* **2013**, *209*. [[CrossRef](#)]
90. Blankley, R.T.; Fisher, C.; Westwood, M.; North, R.; Baker, P.N.; Walker, M.J.; Williamson, A.; Whetton, A.D.; Lin, W.; McCowan, L.; et al. A label-free SRM workflow identifies a subset of pregnancy specific glycoproteins as potential predictive markers of early-onset pre-eclampsia. *Mol. Cell Proteom.* **2013**. [[CrossRef](#)]
91. Sulem, P.; Helgason, H.; Oddson, A.; Stefansson, H.; Gudjonsson, S.A.; Zink, F.; Hjartarson, E.; Sigurdsson, G.T.; Jonasdottir, A.; Jonasdottir, A.; et al. Identification of a large set of rare complete human knockouts. *Nat. Genet.* **2015**, *47*, 448–452. [[CrossRef](#)] [[PubMed](#)]
92. Redman, C.W.; Staff, A.C. Preeclampsia, biomarkers, syncytiotrophoblast stress, and placental capacity. *Am. J. Obstet. Gynecol.* **2015**, *213*. [[CrossRef](#)] [[PubMed](#)]
93. Hutter, S.; Knabl, J.; Andergassen, U.; Mayr, D.; Hofmann, S.; Kuhn, C.; Mahner, S.; Arck, P.; Jeschke, U. Fetal gender specific expression of tandem-repeat galectins in placental tissue from normally progressed human pregnancies and intrauterine growth restriction (IUGR). *Placenta* **2015**, *36*, 1352–1361. [[CrossRef](#)] [[PubMed](#)]
94. Blois, S.M.; Dveksler, G.; Vasta, G.R.; Freitag, N.; Blanchard, V.; Barrientos, G. Pregnancy galectinology: Insights into a complex network of glycan binding proteins. *Front. Immunol.* **2019**, *10*, 1166. [[CrossRef](#)] [[PubMed](#)]
95. Di Renzo, G.C.; Rosati, A.; Sarti, R.D.; Cruciani, L.; Cutuli, A.M. Does fetal sex affect pregnancy outcome? *Gend. Med.* **2007**, *4*, 19–30. [[CrossRef](#)]
96. Thagard, A.S.; Slack, J.L.; Estrada, S.M.; Kazanjian, A.A.; Chan, S.; Burd, I.; Napolitano, P.G.; Ieronimakis, N. Long-term impact of intrauterine neuroinflammation and treatment with magnesium sulphate and betamethasone: Sex-specific differences in a preterm labor murine model. *Sci. Rep.* **2017**, *7*, 17883. [[CrossRef](#)]

