

1 **Interactive effects of fearfulness and geographical location on**
2 **bird population trends**

3

4 **Mario Díaz^a, José Javier Cuervo^b, Tomas Grim^c, Einar Flensted-Jensen^d,**
5 **Juan Diego Ibáñez-Álamo^e, Jukka Jokimäki^f, Gábor Markó^{g,h,i}, Piotr**
6 **Tryjanowski^j and Anders Pape Møller^k**

7

8 ^aDepartment of Biogeography and Global Change, Museo Nacional de Ciencias Naturales
9 (BGC-MNCN-CSIC), c/Serrano 115bis, E-28006 Madrid, Spain, ^bDepartment of
10 Evolutionary Ecology, Museo Nacional de Ciencias Naturales, CSIC, E-28006 Madrid,
11 Spain, ^cPalacky University, Department of Zoology and Laboratory of Ornithology, CZ-
12 77146 Olomouc, Czech Republic, ^dCypresvej 1, DK-9700 Brønderslev, Denmark,
13 ^eDepartamento de Zoología, Facultad de Ciencias, Universidad de Granada, E-18071
14 Granada, Spain, ^fNature Inventory and EIA-services, Arctic Centre, University of Lapland, P.
15 O. Box 122, FI-96101 Rovaniemi, Finland, ^gEcology Research Group, Hungarian Academy
16 of Sciences, Hungarian Natural History Museum, Eötvös Loránd University, Pázmány Péter
17 sétány 1/c, H-1117 Budapest, Hungary, ^hEötvös Loránd University, Behavioral Ecology
18 Group, Department of Systematics, Zoology and Ecology, Pázmány Péter sétány 1/c, H-1117
19 Budapest, Hungary, ⁱDepartment of Plant Pathology, Corvinus University of Budapest,
20 Ménesi út 44, H-1118 Budapest, Hungary, ^jInstitute of Zoology, Poznań University of Life
21 Sciences, Wojska Polskiego 71C, PL-60-625 Poznań, Poland, ^kLaboratoire d'Ecologie,
22 Systématique et Evolution, CNRS UMR 8079, Université Paris-Sud, Bâtiment 362, F-91405
23 Orsay Cedex, France

24

25

26

27 Address correspondence to M. Díaz. E-mail Mario.Diaz@ccma.csic.es

28

29

30

31

32 Short title: Population trends and fearfulness

33 **ABSTRACT**

34 Animal populations are currently under pressure from multiple factors that
35 include human land use and climate change. They may compensate for such
36 effects by reducing, either by habituation or by natural selection, the distance at
37 which they flee from humans (i.e., flight initiation distance, FID), and this
38 adaptation may improve their population trends. We analyzed population trends
39 of common breeding birds in relation to FID and geographical location (latitude,
40 longitude, and marginality of the breeding distribution) across European
41 countries from Finland in the north to Spain in the south, while also considering
42 other potential predictors of trends like farmland habitat, migration, body size
43 and brain size. We found evidence of farmland, migratory and smaller-sized
44 species showing stronger population declines. In contrast, there was no
45 significant effect of relative brain size on population trends. We did not find
46 evidence for main effects of FID and geographical location on trends after
47 accounting for confounding and interactive effects; instead, FID and location
48 interacted to generate complex spatial patterns of population trends. Trends were
49 more positive for fearful populations northwards, westwards and (marginally)
50 towards the centre of distribution areas, and more negative for fearless
51 populations toward the south, east, and the margins of distribution ranges. These
52 findings suggest that it is important to consider differences in population trends
53 among countries, but also interaction effects among factors, because such
54 interactions can enhance or compensate for negative effects of other factors on
55 population trends.

56

57

58 **Key-words:** breeding birds, flight initiation distance, latitude, longitude,
59 marginality of distribution.

60

61

62 **INTRODUCTION**

63 Human disturbance of wild organisms is a common cause of concern in a world
64 with a rapidly increasing human population (Wong and Candolin 2012; Ehrlich
65 and Ehrlich 2013). Such effects of disturbance include release of stress
66 hormones (Wingfield and Ramenofsky 1999), increased metabolic rate
67 (Belanger and Bédard 1990), reduction in foraging activity (Madsen 1998a;
68 1998b), displacement from preferred foraging and roosting sites and changes in
69 diurnal rhythms (Madsen and Fox 1995) and non-lethal effects of predation
70 (Abrams 1991). These factors on their own and in combination may have effects
71 on the condition of animals and hence on their reproduction and survival
72 prospects. A common behavioral measure of proneness to disturbance by
73 humans and animals alike is the flight initiation distance (FID): The distance at
74 which an animal takes flight when approached by a potential predator (Cooper
75 and Blumstein 2015). Because all animals continuously have to weigh the risk
76 of falling prey to a predator by fleeing too late when approached against the
77 benefits of staying put and hence continuing to feed and/or rest, FID constitutes
78 an instantaneous measure of this individual trade-off. Cooke (1980) noticed that
79 urban birds had much shorter flight distances than rural populations of the same
80 species, and that this difference depended on body size, the difference being
81 larger in small species with high metabolism. This change in behavior between
82 urban and rural habitats allowed birds to coexist with humans even at high
83 human population densities, which are a cause of frequent disturbance. Parallel
84 latitudinal trends in FID and raptor abundance in paired urban and rural sites
85 suggest that birds, besides responding to human presence, also adjust their
86 behavior in response to natural levels of disturbance by predators (see Díaz et al.
87 2013 and references therein).

88 It has been noticed that human disturbance at seabird colonies linked to
89 escape behavior and FID could result in altered habitat use and reduced
90 reproductive performance (Burger 1981; Burger and Gochfeld 1981). Therefore,
91 FID can be a useful tool in conservation including assessment of levels of
92 disturbance and susceptibility to disturbance (Madsen 1995; 1998a; 1998b;

93 Weston et al. 2012). The population consequences of FIDs can be investigated
94 by relating population trends to FID (Møller 2008). We should expect species
95 with long FIDs for their body size to show declining population trends because
96 such species should be more prone to get disturbed by humans. Among 56
97 species of birds, FID accounted for 33% of the variance in population trends in
98 Denmark, with effect sizes ranging from 0.36 to 0.58 in different analyses.
99 Therefore, species with long FIDs for their body size had declining populations
100 while species with short FIDs had increasing populations even when controlling
101 statistically for potentially confounding effects. However, a study on population
102 trends in the UK in relation to predictors that included FID recorded in Denmark
103 did not find significant relationship between FID and population trends (Thaxter
104 et al. 2010). This raises the question whether population trends and FID should
105 originate from the same geographic location to make analyses meaningful.

106 Many national and international monitoring programs tally population
107 trends of organisms as diverse as birds, mammals, butterflies and bumblebees.
108 In particular, birds have been popular targets for monitoring since the 1960's in
109 many countries in Europe, and population trends based on European continent-
110 wide monitoring have been published since 1980 (European Bird Census
111 Council, <http://www.ebcc.info/index.php?ID=509>). According to these data,
112 while many species have increased in distribution and abundance, a majority, at
113 least in specific habitats such as farmland, have shown a clear decline. Although
114 humans either directly or indirectly play a major role in determining long-term
115 population trends of birds in Europe (Reif 2013), the underlying mechanisms
116 remain poorly understood. In addition, population trends vary across the
117 distribution range of species. Cuervo and Møller (2013) found stronger increases
118 in northern populations and greater fluctuations in marginal populations,
119 somewhat expected from influences of global warming on climatic niches
120 (Hampe and Petit 2005), and Donald, Green and Heath (2001) and Reif et al.
121 (2011) showed longitudinally varying trends. Reif et al. (2011) also showed an
122 interesting difference in the effect of relative brain size on trends at both sides of

123 the iron curtain, consistent with the differences in land-use intensity across
124 Europe. These intriguing and varying patterns, and the need to optimize
125 conservation priorities, mean that there are good reasons to investigate patterns
126 of population trends at different spatial scales in an attempt to elucidate the
127 underlying mechanisms, including the potential effects of FID.

128 The objectives of this study were to test whether population trends were
129 related to FID, and whether these influences varied across the European
130 continent. If spatial changes in FID could partially compensate for the main
131 effects of factors of global change on trends, we predicted significant
132 interactions between FID and latitude, longitude and marginality on trends. We
133 also tested whether previously established predictors of population trends such
134 as farmland habitat, migration distance, body mass or brain mass affected the
135 relationship between population trend and FID. Overall, elucidating
136 geographical variation in the relationships between trends and recent responses
137 of organisms to changes in the level of human activities will help us to
138 understand our impact on wild populations of animals and eventually to reduce
139 such impacts.

140

141 **METHODS**

142 We recorded FID for a total of 159 species during the breeding seasons 2009-
143 2010 at nine locations from eight countries along a wide latitudinal gradient
144 across Europe, from Finland in the north to Spain in the south, by using a
145 standard procedure developed by Blumstein (2006). These data are reported in
146 Díaz et al. (2013). In brief, we walked at ordinary walking speed towards a bird
147 recording the distance from the bird when we started walking, the distance at
148 which the birds initiated escape, and the bird's height in the vegetation. This
149 information was used to estimate FID. In order to account for the height at
150 which individuals were perched, FID was calculated as the Euclidean distance
151 between the approaching human and the focal bird (which equals the square-root
152 of the sum of the squared flight distance and the squared height in the

153 vegetation). Observers wore neutrally colored clothes and behaved as normal
154 pedestrians. FID was measured by a number of trained observers and therefore
155 data were pooled for analysis. We used the FID estimates for rural populations
156 in each location, which consisted of paired rural and urban sites (Díaz et al.
157 2013), because the population size estimates used to assess trends for each
158 country are mostly based on data coming from non-urban populations (Cuervo
159 and Møller 2013). Data for the two Spanish sites were averaged to obtain a
160 single country-level estimate.

161 Population trends for breeding birds in all European countries for which
162 we had information on FID (Finland, Norway, Denmark, Poland, Czech
163 Republic, Hungary, France and Spain) were obtained from Cuervo and Møller
164 (2013). Available population size estimates for each bird species and country
165 were regressed on years, and the slope of this regression was used as a proxy for
166 population trend. We used time series of 7-27 years gathered until 2004-2008
167 depending on countries and species (see Cuervo and Møller 2013 for details and
168 a full discussion of the quality of trend estimates).

169 Latitude and longitude for each country were estimated as the coordinates
170 of the mid-point between the northernmost and the southernmost, and between
171 the easternmost and the westernmost, mainland points of every country,
172 excluding islands except for Denmark. Latitude and longitude for each country
173 were considered the latitude and longitude for all bird populations in that
174 particular country regardless of the actual distribution of every species within
175 the country. Marginality of each bird population was estimated by comparing
176 two distances (in degrees): L is the distance between the population (i.e., the
177 country) latitude and the northernmost or the southernmost (the one that resulted
178 in a shorter distance) limits of the breeding distribution range of the species. L
179 was set to zero in the few cases in which the country latitude index was more
180 southern than the southernmost limit of the species range or more northern than
181 the northernmost limit of the species range. C is the distance between the
182 population latitude and the latitude of the mid-point between the northernmost

183 and the southernmost limits of the breeding distribution range of the species.
184 Marginality was computed as $\log_{10}(C+1) - \log_{10}(L+1)$, with positive values
185 representing marginal populations (the distance to the range centre was larger
186 than the distance to the nearest limit) and negative values central populations
187 (the distance to the range centre was smaller than the distance to the nearest
188 limit). These values were transformed by adding the absolute value of the most
189 negative number and dividing by the largest value resulting from the previous
190 addition, to ensure that marginality estimates ranged from 0 (central population)
191 to 1 (marginal population; see Cuervo and Møller 2013 for details).

192 Bird population trends have previously been shown to be systematically
193 affected by body size, migration distance, farmland habitat and relative brain
194 size (reviews in Møller 2008; Møller, Rubolini and Lehikoinen 2008; Reif
195 2013). We extracted information on mean body mass of adult birds of each
196 species from Cramp and Perrins (1977-1994). Migration distances (mean of the
197 northernmost and the southernmost latitudes of the breeding distribution range
198 minus the corresponding mean for the wintering distribution range) were taken
199 from Møller, Rubolini and Lehikoinen (2008). Farmland habitat was coded as 1
200 (species depending on arable and/or mixed farmland) or 0 (species depending on
201 other habitat types) following Appendix 2 in Tucker and Evans (1997). Relative
202 brain sizes were the residuals of a log-log phylogenetically corrected regression
203 of brain mass on body mass based on a sample of 567 bird species (Møller,
204 2008); brain mass data were obtained from Garamszegi, Møller and Erritzøe
205 (2002), Iwaniuk and Nelson (2002), Galván and Møller (2011) and Møller and
206 Erritzøe (2014).

207 We \log_{10} -transformed FID, population trend and migration distance
208 before analyses. Within-species repeatability of FID and trends across Europe
209 was computed following Lessells and Boag (1987), and differences between
210 them and the null hypothesis of zero repeatability were tested following Becker
211 (1984). Significant repeatabilities imply statistical dependence of estimates for
212 the same species in different countries, a fact that will bias results based on

213 phylogenetically-structured databases (Garamszegi and Møller 2010). As
214 species occupy a variable number of study locations and countries (Díaz et al.
215 2013; Cuervo and Møller 2013), geographical trends could be partly due to
216 phylogenetic effects. To control for such relationships we used phylogenetic
217 generalized least square regression (PGLS) models implemented in R (Díaz et
218 al. 2013). After estimating the phylogenetic scaling parameter lambda (λ), we
219 calculated the phylogenetically corrected partial correlations between the
220 variables of interest. Different populations of the same species were considered
221 as polytomies with a constant small genetic distance of 10^{-10} between them. We
222 used the R script and the edited phylogeny supplied as Supplementary Files S1
223 and S2 in Díaz et al. (2013), but using the function *pglm3.3.r* instead of the
224 *pglm3.1.r* to fit type III (orthogonal) models. We used the phylogeny reported in
225 Thuiller et al. (2011). The dependent variable was the population trend,
226 confounding variables were farmland habitat, migration distance, body size and
227 relative brain size, and predictors FID, latitude, longitude, marginality, and the
228 first-order interaction between FID and geographical variables. Predictors were
229 computed from the corresponding input variables (\log_{10} FID and geographical
230 variables) by standardizing them (i.e., by subtracting sample means and dividing
231 by standard deviations), in order to allow direct comparison of effect sizes
232 (Pearson's product-moment correlation coefficients computed from P values of
233 t-tests according to Lipsey and Wilson 2001) and to make main effects
234 biologically interpretable even when involved in interactions (Schielzeth 2010).

235

236 **RESULTS**

237 We collected data on mean FID and on recent population trends from 338
238 populations of 129 bird species. Data on farmland habitat, body size and
239 migration distance were available for all of them, while there were no data on
240 brain size for 9 species (Appendix A). Both FID and trends were significantly
241 repeatable within species ($F_{1, 209} = 3.08$, $P < 0.001$ and $F_{1, 209} = 1.45$, $P = 0.009$,
242 respectively). FID was significantly more repeatable than population trends ($r =$

243 0.45 ± 0.04 (SD) vs. $r = 0.15 \pm 0.05$; $t_{338} = 4.0$, $P < 0.001$; Becker 1984); in
244 other words, geographical variation within species was larger for population
245 trends than for mean fearfulness as reflected by FID.

246 Log-transformed population trends were significantly related to \log_{10} FID
247 ($F_{1,337} = 7.96$, $P = 0.005$, $r^2 = 0.02$), but not to latitude ($F_{1,337} = 0.00$, $P = 0.967$,
248 $r^2 = 0.00$), longitude ($F_{1,337} = 0.40$, $P = 0.530$, $r^2 = 0.00$) or marginality ($F_{1,337} =$
249 0.62 , $P = 0.432$, $r^2 = 0.00$) when predictor effects were analyzed one by one.
250 The relationship with FID vanished, however, after correcting for significant
251 effects of farmland habitat, migration distance and body mass (effect sizes for
252 these three confounding variables ranged from 0.14 to 0.16), while also
253 accounting for phylogenetic effects (Table 1). Trends were more negative for
254 farmland birds, long-distance migrants and smaller species (Table 1). Relative
255 brain size showed no significant effects on population trends, which did not
256 show significant geographical trends either (Table 1). However, FID showed
257 significant interactive effects with latitude and longitude, and marginally-
258 significant interactive effects with marginality, with effect sizes ranging from
259 0.10 to 0.13 (Table 1, Fig. 1). FID-trend relationships were more positive
260 northwards, westwards and (marginally) towards the centre of distribution areas
261 (Table 1, Fig. 1). These interactions implied that trends were more negative for
262 fearless populations toward the south, east, and the margins of distribution
263 ranges.

264

265 **DISCUSSION**

266 Many different factors have been proposed to account for population trends of
267 birds (reviewed in Reif 2013). These variables range from migration and the
268 perils of living under different climate regimes (Hjort and Lindholm 1978;
269 Baillie and Peach 1992; Sanderson et al. 2006; Reif 2013), relative brain mass
270 that facilitates the ability to cope with changing environments (Shultz et al.
271 2005; Møller, Rubolini and Lehikoinen 2008; Reif et al. 2011), thermal range
272 and hence the ability to cope with changing climatic conditions (Jiguet et al.

273 2007, 2010), the number of broods with species producing more broods doing
274 better (Julliard, Jiguet and Couvet 2004), and body mass with large sized species
275 with smaller total populations having negative population trends (Bennett and
276 Owens 2002).

277 Geographical variation in trends within breeding ranges of species are
278 also expected due to geographical changes in the suitability of environmental
279 conditions (the niche variation hypothesis; Brown 1984), in the intensity of
280 global change drivers (Hampe and Petit 2005; Reif et al. 2011; Tryjanowski et
281 al. 2011) or in both (Díaz et al. 1998). It has been suggested that population
282 responses of birds to environmental gradients may be highly species-specific,
283 even precluding broad generalizations (Taper, Böhning-Gaese and Brown
284 1995); however, Cuervo and Møller (2013) have recently shown that changes in
285 population size of breeding birds in Europe are the strongest at the margins of
286 the breeding distribution, but are particularly negative at the southern-most
287 range margins, where increasing temperatures may render environmental
288 conditions for maintenance of viable populations the most difficult. Climate
289 change has affected the distribution of many species, and range margins have on
290 average moved pole-wards (Chen et al. 2011), and recent work has shown fine-
291 grained effects of climate change on local population trends (Jiguet et al. 2010).
292 Longitudinal variation due to differences in land-use intensity between Western
293 and Eastern Europe have also been documented, especially for farmland birds
294 (Donald, Green and Heath 2001). However, we did not find evidence for direct
295 effects of these variables after accounting for effects of third variables and their
296 interactions. Reif et al. (2011) suggested that longitudinal effects of the iron
297 curtain dividing industrialized Western Europe from more extensive land use in
298 Eastern Europe interacted with relative brain mass to account for spatial
299 heterogeneity in population trends. Here we found no evidence of an effect of
300 relative brain mass on population trends contrary to previous reports (Shultz et
301 al. 2005; Thaxter et al. 2010; Reif et al. 2011). We hypothesize that these
302 differences among studies may arise from the inclusion of different predictors

303 and their interactions, but also from inclusion of multiple countries that differ in
304 significant predictors of population trends. Studies such as this, encompassing
305 the widest ranges of variation of relevant variables available even at the expense
306 of lower precision within ranges, are thus essential to detect non-linear and
307 interactive relationships of geographically-varying conditions on local
308 abundance and trends (e.g. Jiguet et al. 2010; Concepción et al. 2012).

309 Bird species breeding on farmland displayed the steepest declines. This
310 is probably a consequence of agriculture having become ever more
311 industrialized and intensified and thereby disproportionately negatively affecting
312 farmland specialists (Fuller et al. 1995; Chamberlain et al. 2000; Møller,
313 Rubolini and Lehikoinen 2008; Reif 2013). Here we found evidence consistent
314 with this general trend, with farmland species showing more negative population
315 trends than non-farmland birds. Migration has been predicted to affect
316 population trends because migrants are affected negatively by land-use and
317 climate change in their breeding range, during migration and in their winter
318 quarters (Hjort and Lindholm 1978; Baillie and Peach 1992; Sanderson et al.
319 2006; Møller, Rubolini and Lehikoinen 2008; Reif 2013). Here we found a
320 negative effect of migration distance on population trends, when accounting for
321 the effects of the remaining variables.

322 We hypothesized that population trends would be negatively related to
323 FID, as reported by Møller (2008) for European birds. Most recent work
324 indicates that FID can be considered a general measure of the willingness of
325 animals to be involved in risky activities such as foraging and courtship under
326 perceived risky conditions (reviewed in Cooper and Blumstein 2014, 2015).
327 Such willingness to take risks would depend on levels of risk (abundance and
328 identity of predators and other sources of risk, such as humans), but also on
329 potential fitness benefits (ie. it will be worth taking more risks if the expected
330 fitness consequence of the reward is larger, as under food shortage or time-
331 limited conditions), after accounting for species- and population-specific
332 proneness to risk-taking associated with phylogeny, urban habitat or life-history

333 traits such as body size or migratory behavior (Díaz et al. 2013). We found an
334 overall main effect of FID in this study, which however vanished when
335 considering interactive effects with geographical location. This fact suggests that
336 the observed geographical variation in trends would in fact be the net result of
337 complex interactions between spatial variations in many factors proposed to
338 drive population trends (Reif 2013), as well as on the varying effects of risk-
339 taking behaviors on trends. Our results showed that fearfulness of bird
340 populations (i.e., long FIDs) enhanced population trends where such trends were
341 already less negative, as in northern European populations (Cuervo and Møller
342 2013), or where land use intensity is higher, as in western European countries
343 (Tryjanowski et al. 2011), but these relationships reversed at more stressful
344 extremes of spatial gradients, such as southern and marginal locations. We
345 interpret these interactions as implying that we cannot assess predictors by
346 considering solely their main effects. We are unaware of any previous studies
347 investigating such interaction effects as predictors of population trends.

348 In conclusion, we have analyzed for the first time how geographical
349 patterns of population trends of birds in Europe, as related to natural and man-
350 made geographical variation in environmental factors such as climate, predation
351 risk and land use, interact with a measure of the tolerance of birds to human
352 disturbance. Overall we found that proneness to risk-taking as estimated by
353 short FIDs enhanced population resilience to disturbance in a changing world, as
354 more tolerant individuals will reduce the costs associated with escape behaviors
355 (Cooper and Blumstein 2014). In contrast, bird species and populations less
356 tolerant of frequent disturbance, by humans or wild and domestic predators,
357 would perform worse, especially at the southern- and eastern-most edges of
358 breeding distributions. Further studies including fine-grained estimates of FID,
359 trends and secondary influences on them (eg. Jiguet et al. 2010) carried out over
360 wide geographical gradients would be needed to ascertain whether these patterns
361 were due to geographical variations in risks, fitness benefits of risk-taking, or
362 both.

363

364

365 ACKNOWLEDGMENTS

366 Comments by two anonymous referees were very helpful during revision. We
367 thank Martin Flade, Magne Husby, Jaroslav Krogulec, Robert Kwak, Jiří Reif,
368 Norbert Teufelbauer, Tibor Szép and Dansk Ornithologisk Forening for
369 information on population size estimates. JJC was supported by the Spanish
370 National Research Council (grant EST001196). This paper is a contribution by
371 MD to the project RiskDisp (CGL2009-08430) and to the thematic networks
372 GlobiMed (www.globimed.net) and REMEDINAL III.

373

374

375

376

377 APPENDIX A. SUPPLEMENTARY MATERIAL

378 Information on bird species, country, population trend, mean FID (m), migration
379 distance (°latitude), body mass (g), latitude and longitude of the
380 population/country, marginality of the population within the species breeding
381 range, relative brain size and farmland habitat. Relative brain size is residuals
382 from a log-log phylogenetically corrected regression of brain mass on body
383 mass. See Material and methods for sources and details. Nomenclature and basic
384 phylogeny follows Thuiller et al. (2011).

385

386 REFERENCES

- 387 Abrams PA. 1991. Strengths of indirect effects generated by optimal foraging.
388 *Oikos* 62:167-176.
- 389 Baillie SR, Peach WJ. 1992. Population limitation in Palaearctic-African
390 migrant passerines. *Ibis* 134:120–132.

- 391 Becker WA. 1984. A manual of quantitative genetics. Washington: Pullman
392 Academic Enterprises.
- 393 Belanger L, Bédard J. 1990. Energetic cost of man-induced disturbance to
394 staging snow geese. *J. Wildl. Manage.* 54:36-41.
- 395 Bennett PM, Owens IPF. 2002. Evolutionary ecology of birds. Oxford: Oxford
396 Univ. Press.
- 397 Blumstein DT. 2006. Developing an evolutionary ecology of fear: How life
398 history and natural history traits affect disturbance tolerance in birds.
399 *Anim. Behav.* 71:389–399.
- 400 Brown JH. 1984. On the relationship between abundance and distribution of
401 species. *Am. Nat.* 124:255-279.
- 402 Burger J, Gochfeld M. 1981. Discrimination of the threat of direct versus
403 tangential approach to the nest by incubating herring and great black-
404 backed gulls. *J. Comp. Physiol. Psychol.* 95:676–684.
- 405 Burger J. 1981. The effect of human activity on birds at a coastal bay. *Biol.*
406 *Conserv.* 21:231–241.
- 407 Chamberlain DE, Fuller RJ, Bunce RGH, Duckworth JC, Shrubbs M. 2000.
408 Changes in the abundance of farmland birds in relation to the timing of
409 agricultural intensification in England and Wales. *J. Appl. Ecol.* 37:771–
410 788.
- 411 Chen IC, Hill JK, Ohlemüller R, Roy DB, Thomas CD. 2011. Rapid range
412 shifts of species associated with high levels of climate warming. *Science*
413 333:1024-1026.
- 414 Concepción ED, Díaz M, Kleijn D, Báldi A, Batáry P, Clough Y, Gabriel D,
415 Herzog F, Holzschuh A, Knop E, Marshall JP, Tschamntke T, Verhulst J
416 2012. Interactive effects of landscape context constrains the effectiveness
417 of local agri-environmental management. *J. Appl. Ecol.* 49:695-705.
- 418 Cooke AS. 1980. Observations on how close certain passerine species will
419 tolerate an approaching human in rural and suburban areas. *Biol. Conserv.*
420 18:85-88.

- 421 Cooper Jr WE, Blumstein DT. 2014. Novel effects of monitoring predators on
422 costs of fleeing and not fleeing explain flushing early in economic escape
423 theory. *Behav. Ecol.* 25:44-52.
- 424 Cooper Jr WE, Blumstein DT. 2015. Escaping from predators: An integrative
425 view of escape decisions and refuge use. Cambridge: Cambridge Univ.
426 Press.
- 427 Cramp S, Perrins CM (eds) 1977-1994. The birds of the Western Palearctic.
428 Oxford: Oxford Univ. Press.
- 429 Cuervo JJ, Møller AP. 2013. Temporal variation in population size of European
430 bird species: Effects of latitude and marginality of distribution. *PLoS One*
431 8:e77654.
- 432 Díaz M, Carbonell R, Santos T, Tellería JL. 1998. Breeding bird communities in
433 pine plantations of the Spanish plateaux: Biogeography, landscape and
434 vegetation effects. *J. Appl. Ecol.* 35:562-574.
- 435 Díaz M, Møller AP, Flensted-Jensen E, Grim T, Ibáñez-Álamo JD, Jokimäki J,
436 Markó G, Tryjanowski P. 2013. The geography of fear: A latitudinal
437 gradient in anti-predator escape distances of birds across Europe. *PloS One*
438 8:e64634.
- 439 Donald PF, Green RE, Heath MF. 2001. Agricultural intensification and the
440 collapse of Europe's farmland bird populations. *Proc. R. Soc. B* 268:25–29.
- 441 Ehrlich PR, Ehrlich AH. 2013. Can a collapse of global civilization be avoided?
442 *Proc. R. Soc. B* 280:20122845.
- 443 Forstmeier W, Schielzeth H. 2011. Cryptic multiple hypotheses testing in linear
444 models: overestimated effect sizes and the winner's curse. *Behav. Ecol.*
445 *Sociobiol.* 65:47-55.
- 446 Fuller R J, Gregory RD, Gibbons DW, Marchant JH, Wilson JD, Baillie SR,
447 Carter N. 1995. Population declines and range contractions among lowland
448 farmland birds in Britain. *Conserv. Biol.* 9:1425-1441.
- 449 Galván I, Møller AP. 2011. Brain size and the expression of pheomelanin-based
450 colour in birds. *J. Evol. Biol.* 24:999-1006.

- 451 Garamszegi LZ, Møller AP. 2010. Effects of sample size and intraspecific
452 variation in phylogenetic comparative studies: a meta-analytic review. *Biol.*
453 *Rev.* 85:797–805.
- 454 Garamszegi LZ, Møller AP, Erritzøe J. 2002. Coevolving avian eye size and
455 brain size in relation to prey capture and nocturnality. *Proc. R. Soc. B*
456 269:961–967.
- 457 Hampe A, Petit RJ. 2005. Conserving biodiversity under climate change: the
458 rear edge matters. *Ecol. Lett.* 8:461-467.
- 459 Hjort C, Lindholm CG. 1978. Annual bird ringing totals and population
460 fluctuations. *Oikos* 30:387–392.
- 461 Iwaniuk AN, Nelson JE. 2002. Can endocranial volume be used as an estimate
462 of brain size in birds? *Can. J. Zool.* 80:16-23.
- 463 Jiguet F, Devictor V, Ottvall R, Van Turnhout C, Van der Jeugd H, Lindström
464 Å. 2010. Bird population trends are linearly affected by climate change
465 along species thermal ranges. *Proc. R. Soc. B* 277:3601–3608.
- 466 Jiguet F, Gadot AS, Julliard R, Newson SE, Couvet D. 2007. Climate envelope,
467 life history traits and the resilience of birds facing global change. *Global*
468 *Change Biol.* 13:1672–1684.
- 469 Julliard R, Jiguet F, Couvet D. 2004. Common birds facing global changes:
470 What makes a species at risk? *Global Change Biol.* 10:148 –154.
- 471 Lessells CM, Boag PT. 1987. Unrepeatable repeatabilities: a common
472 mistake. *Auk* 104:116-121.
- 473 Lipsey MW, Wilson DB. 2001. Practical meta- analysis. *Applied Social*
474 *Research Methods Series* 39. London: Sage.
475 [www.campbellcollaboration.org/escalc/html/EffectSizeCalculator-](http://www.campbellcollaboration.org/escalc/html/EffectSizeCalculator-Home.php)
476 [Home.php](http://www.campbellcollaboration.org/escalc/html/EffectSizeCalculator-Home.php)
- 477 Madsen J. 1995. Impacts of disturbance on migratory waterfowl. *Ibis* 137:S67–
478 S74.

- 479 Madsen J. 1998a. Experimental refuges for migratory waterfowl in Danish
480 wetlands. I. Baseline assessment of disturbance effects of recreational
481 activities. *J. Appl. Ecol.* 35:386–397.
- 482 Madsen J. 1998b. Experimental refuges for migratory waterfowl in Danish
483 wetlands. II. Tests of hunting disturbance effects. *J. Appl. Ecol.* 35:398–
484 417.
- 485 Madsen J, Fox AD. 1995. Impacts of hunting disturbance on waterbirds: A
486 review. *Wildl. Biol.* 1:193-207.
- 487 Møller AP. 2008. Flight distance and population trends in European breeding
488 birds. *Behav. Ecol.* 19:1095–1102.
- 489 Møller AP, Erritzøe J. 2014. Predator–prey interactions, flight initiation distance
490 and brain size. *J. Evol. Biol.* 27:34-42.
- 491 Møller AP, Rubolini D, Lehikoinen E. 2008. Populations of migratory bird
492 species that did not show a phenological response to climate change are
493 declining. *Proc. Nat. Acad. Sci. USA* 105:16195–16200.
- 494 Reif J. 2013. Long-term trends in bird populations: a review of patterns and
495 potential drivers in North America and Europe. *Acta Ornithol.* 48:1-16.
- 496 Reif J, Böhning-Gaese K, Flade M, Schwarz J, Schwager M. 2011. Bird
497 population trends across iron curtain, brain matters. *Biol. Conserv.*
498 144:2524–2533.
- 499 Sanderson FJ, Donald PF, Pain DJ, Burfield IJ, van Bommel FPJ. 2006. Long-
500 term population declines in Afro-Palearctic migrant birds. *Biol. Conserv.*
501 131:93–105.
- 502 Schielzeth H. 2010. Simple means to improve the interpretability of regression
503 coefficients. *Met. Ecol. Evol.* 1:103–113.
- 504 Shultz S, Bradbury RB, Evans KL, Gregory RD, Blackburn TM. 2005. Brain
505 size and resource specialization predict long-term population trends in
506 British birds. *Proc. R. Soc. B* 272:2305–2311.
- 507 Taper ML, Böhning-Gaese K, Brown JH. 1995. Individualistic responses of bird
508 species to environmental change. *Oecologia* 101:478-486.

- 509 Thaxter CB, Joys AC, Gregory RD, Baillie SR, Noble DG. 2010. Hypotheses to
510 explain patterns of population change among breeding bird species in
511 England. *Biol. Conserv.* 143 2006-2019.
- 512 Thuiller W, Lavergne S, Roquet C, Boulangeat I, Lafourcade B, Araújo MB
513 2011. Consequences of climate change on the tree of life in Europe. *Nature*
514 470:531–534.
- 515 Tryjanowski P, Hartel T, Báldi A, Szymański P, Tobolka M, Herzon I, Goławski
516 A, Konvička M, Hromada M, Jerzak L, Kujawa K, Lenda M, Orłowski M,
517 Panek M, Skórka P, Sparks TH, Tworek S, Wuczyński A, Żmihorski M.
518 2011. Conservation of farmland birds faces different challenges in Western
519 and Central-Eastern Europe. *Acta Ornithol.* 46:1–12.
- 520 Tucker GM, Evans MI (comp) 1997. Habitats for birds in Europe. A
521 conservation strategy for the wider environment. Cambridge: BirdLife
522 International.
- 523 Weston MA, McLeod EM, Blumstein DT, Guay PJ. 2012. A review of flight-
524 initiation distances and their application to managing disturbance to
525 Australian birds. *Emu* 112:269-286.
- 526 Wingfield JC, Ramenofsky M. 1999. Hormones and the behavioral ecology of
527 stress. In: *Stress Physiology in Animals*. Balm PHM, editor. Sheffield:
528 Sheffield Academic Press. pp. 1-51.
- 529 Wong B, Candolin U. 2012. Behavioural responses to a changing world:
530 Mechanisms and consequences. Oxford: Oxford Univ. Press.
531
532
533
534
535
536

537 Fig. 1. Latitudinal and longitudinal variation in standardized regression
538 coefficients ($\beta \pm SE$) between population trends of European birds (residuals
539 from a multiple regression between log-transformed trends, migration distance,
540 body mass, relative brain mass, farmland habitat, marginality, and either
541 longitude or latitude) and fearfulness (flight initiation distance, FID, log-
542 transformed). Lines are best-fit regressions.

543

544

545

546

547

548

549

550

551

552


553

554

555

556

557



558 Table 1. Relationships between population trends of European birds (response
 559 variable) and geographical location (latitude, longitude and marginality) and
 560 fearfulness (flight initiation distance, FID), after accounting for effects of
 561 farmland habitat, migration distance, body mass and relative brain size on trends
 562 and correcting for the effect of the phylogenetic structure of the data set, that
 563 was, however, not significant ($\lambda = 0.000$, $\chi^2 = -0.012$, $P = 1.000$). The full model
 564 (no removal of non-significant terms was done, as recommended by Forstmeier
 565 and Schielzeth 2011) had the statistics $F = 4.73$, d.f. = 12, 329, adjusted $r^2 =$
 566 0.11, $P < 0.0001$. Effect sizes are Pearson's product-moment correlation
 567 coefficients.

568

569

Source	estimate (SE)	<i>t</i>	<i>P</i>	Effect size
Farmland	-0.008(0.003)	-2.50	0.013	0.14
Migration distance	-0.006(0.002)	-2.71	0.007	0.15
Body mass	0.018(0.006)	2.93	0.004	0.16
Relative brain size	-0.014(0.010)	-1.46	0.147	0.08
FID	0.000(0.002)	0.16	0.876	0.01
Latitude	-0.002(0.002)	-1.07	0.286	0.06
Longitude	0.003(0.002)	1.50	0.134	0.08
Marginality	0.001(0.002)	0.38	0.706	0.02
FID x Latitude	0.005(0.002)	2.29	0.023	0.13
FID x Longitude	-0.004(0.002)	-2.27	0.024	0.12
FID x Marginality	-0.003(0.002)	-1.76	0.079	0.10

570

571

572

573 **APPENDIX A. SUPPLEMENTARY MATERIAL**

574 Information on bird species, country, population trend, mean FID (m), migration
 575 distance (°latitude), body mass (g), latitude and longitude of the
 576 population/country, marginality of the population within the species breeding
 577 range, relative brain size and farmland habitat. Relative brain size is residuals
 578 from a log-log phylogenetically corrected regression of brain mass on body
 579 mass. See Material and methods for sources and details. Nomenclature and basic
 580 phylogeny follows Thuiller et al. (2011).

581

Species	Country	Popul. trend	FID (m)	Migration distance (° latitude)	Body mass (g)	Latitude	Longitude	Marginality	Rel. brain size	Farmland
<i>Accipiter nisus</i>	Denmark	0.002	36.14	12.79	204.0	56.16	10.38	0.585	0.464	0
<i>Acrocephalus palustris</i>	Czech Rep.	-0.010	14.14	66.84	12.0	49.80	15.48	0.599	-0.281	0
<i>Acrocephalus palustris</i>	Denmark	0.012	8.76	66.84	12.0	56.16	10.38	0.683	-0.281	0
<i>Acrocephalus schoenobaenus</i>	Denmark	-0.003	6.93	62.10	11.9	56.16	10.38	0.595	-0.357	0
<i>Acrocephalus scirpaceus</i>	Denmark	-0.009	6.59	44.60	11.8	56.16	10.38	0.659	-0.330	0
<i>Aegithalos caudatus</i>	France	0.000	5.27	0.00	8.8	46.71	1.72	0.489	-0.385	0
<i>Aegithalos caudatus</i>	Spain	-0.010	10.84	0.00	8.8	39.90	-2.99	0.500	-0.385	0
<i>Aegithalos caudatus</i>	Hungary	0.155	6.36	0.00	8.8	47.16	19.51	0.489	-0.385	0
<i>Alauda arvensis</i>	Denmark	-0.008	31.42	13.02	36.4	56.16	10.38	0.581	-0.033	1
<i>Alauda arvensis</i>	Poland	0.017	45.44	13.02	36.4	51.92	19.13	0.533	-0.033	1
<i>Alcedo atthis</i>	France	-0.042	8.54	0.00	32.4	46.71	1.72	0.429	-0.099	0
<i>Alectoris rufa</i>	Spain	0.004	37.08	0.00	477.5	39.90	-2.99	0.732	0.303	1
<i>Anas platyrhynchos</i>	Denmark	0.048	28.76	8.13	1119.0	56.16	10.38	0.579	0.720	0
<i>Anas platyrhynchos</i>	France	0.012	4.81	8.13	1119.0	46.71	1.72	0.533	0.720	0
<i>Anas platyrhynchos</i>	Norway	0.123	11.11	8.13	1119.0	64.56	18.01	0.709	0.720	0
<i>Anas platyrhynchos</i>	Poland	0.014	88.00	8.13	1119.0	51.92	19.13	0.531	0.720	0
<i>Anser anser</i>	Denmark	0.107	180.00	12.28	3464.5	56.16	10.38	0.578	1.080	0
<i>Anthus pratensis</i>	Denmark	-0.027	13.22	15.64	19.3	56.16	10.38	0.595	-0.312	0
<i>Anthus pratensis</i>	Finland	0.002	28.31	15.64	19.3	64.95	26.07	0.730	-0.312	0
<i>Anthus pratensis</i>	Norway	0.001	5.82	15.64	19.3	64.56	18.01	0.722	-0.312	0
<i>Anthus pratensis</i>	Poland	-0.046	32.00	15.64	19.3	51.92	19.13	0.619	-0.312	0
<i>Anthus spinoletta</i>	France	-0.037	7.50	21.27	21.5	46.71	1.72	0.549	-0.237	0
<i>Anthus trivialis</i>	Denmark	-0.013	10.90	47.07	23.4	56.16	10.38	0.587	-0.210	0
<i>Apus apus</i>	Denmark	0.006	38.10	59.39	39.7	56.16	10.38	0.585	-0.205	0
<i>Ardea cinerea</i>	Denmark	0.195	62.08	1.50	1433.0	56.16	10.38	0.330	0.903	0
<i>Ardea cinerea</i>	France	0.049	26.00	1.50	1433.0	46.71	1.72	0.289	0.903	0
<i>Ardea cinerea</i>	Norway	0.207	50.00	1.50	1433.0	64.56	18.01	0.596	0.903	0
<i>Athene noctua</i>	Spain	-0.008	25.73	0.00	168.0	39.90	-2.99	0.439	0.579	1
<i>Aythya fuligula</i>	Denmark	0.015	10.68	17.08	656.5	56.16	10.38	0.603	0.651	0
<i>Buteo buteo</i>	Czech Rep.	0.012	55.26	29.57	806.5	49.80	15.48	0.502	0.896	0
<i>Buteo buteo</i>	Denmark	0.033	60.01	29.57	806.5	56.16	10.38	0.591	0.896	0
<i>Carduelis cannabina</i>	Czech Rep.	-0.011	14.93	4.11	19.0	49.80	15.48	0.547	-0.187	0
<i>Carduelis cannabina</i>	Denmark	-0.011	10.80	4.11	19.0	56.16	10.38	0.635	-0.187	0
<i>Carduelis cannabina</i>	Poland	-0.062	12.89	4.11	19.0	51.92	19.13	0.574	-0.187	0
<i>Carduelis cannabina</i>	Spain	-0.009	18.51	4.11	19.0	39.90	-2.99	0.532	-0.187	0
<i>Carduelis carduelis</i>	Czech Rep.	-0.009	14.95	1.16	15.6	49.80	15.48	0.564	-0.240	0
<i>Carduelis carduelis</i>	Denmark	0.169	10.77	1.16	15.6	56.16	10.38	0.667	-0.240	0
<i>Carduelis carduelis</i>	France	-0.002	11.00	1.16	15.6	46.71	1.72	0.524	-0.240	0

<i>Carduelis carduelis</i>	Spain	-0.022	11.52	1.16	15.6	39.90	-2.99	0.490	-0.240	0
<i>Carduelis carduelis</i>	Hungary	0.032	8.99	1.16	15.6	47.16	19.51	0.530	-0.240	0
<i>Carduelis chloris</i>	Czech Rep.	-0.021	12.98	1.34	27.7	49.80	15.48	0.507	-0.057	0
<i>Carduelis chloris</i>	Denmark	0.036	6.53	1.34	27.7	56.16	10.38	0.581	-0.057	0
<i>Carduelis chloris</i>	Finland	0.253	15.22	1.34	27.7	64.95	26.07	0.730	-0.057	0
<i>Carduelis chloris</i>	France	-0.020	7.65	1.34	27.7	46.71	1.72	0.502	-0.057	0
<i>Carduelis chloris</i>	Spain	0.033	15.92	1.34	27.7	39.90	-2.99	0.518	-0.057	0
<i>Carduelis chloris</i>	Hungary	0.050	18.43	1.34	27.7	47.16	19.51	0.501	-0.057	0
<i>Carduelis chloris</i>	Norway	0.022	5.88	1.34	27.7	64.56	18.01	0.720	-0.057	0
<i>Carduelis chloris</i>	Poland	0.009	15.11	1.34	27.7	51.92	19.13	0.531	-0.057	0
<i>Carduelis flammea</i>	Denmark	-0.001	4.50	9.46	13.1	56.16	10.38	0.622	-0.263	0
<i>Carduelis flammea</i>	Norway	-0.027	12.00	9.46	13.1	64.56	18.01	0.653	-0.263	0
<i>Carduelis spinus</i>	Finland	0.015	10.11	6.83	13.8	64.95	26.07	0.740	-0.272	0
<i>Carpodacus erythrinus</i>	Finland	-0.014	9.44	0.00	13.8	64.95	26.07	0.804	--	0
<i>Certhia brachydactyla</i>	France	0.031	7.12	0.00	9.2	46.71	1.72	0.635	-0.338	0
<i>Certhia brachydactyla</i>	Spain	0.024	11.15	0.00	9.2	39.90	-2.99	0.603	-0.338	0
<i>Certhia familiaris</i>	Denmark	0.015	4.47	0.00	9.2	56.16	10.38	0.582	-0.297	0
<i>Cettia cetti</i>	Spain	0.000	26.84	2.11	14.1	39.90	-2.99	0.564	--	0
<i>Cisticola juncidis</i>	Spain	-0.011	31.89	0.00	8.5	39.90	-2.99	0.337	--	0
<i>Clamator glandarius</i>	Spain	0.185	55.60	4.58	153.5	39.90	-2.99	0.344	0.205	0
<i>Coccothraustes coccothraustes</i>	Czech Rep.	-0.020	22.17	5.07	54.7	49.80	15.48	0.591	0.222	0
<i>Coccothraustes coccothraustes</i>	France	0.038	5.10	5.07	54.7	46.71	1.72	0.551	0.222	0
<i>Coccothraustes coccothraustes</i>	Hungary	0.038	24.00	5.07	54.7	47.16	19.51	0.556	0.222	0
<i>Columba livia</i>	France	0.024	8.00	0.00	261.0	46.71	1.72	0.403	0.303	1
<i>Columba livia</i>	Spain	0.007	30.87	0.00	261.0	39.90	-2.99	0.291	0.303	1
<i>Columba livia</i>	Hungary	0.350	6.00	0.00	261.0	47.16	19.51	0.410	0.303	1
<i>Columba oenas</i>	France	-0.012	17.75	3.45	494.5	46.71	1.72	0.546	0.333	1
<i>Columba palumbus</i>	Czech Rep.	0.028	27.26	2.03	494.5	49.80	15.48	0.521	0.365	1
<i>Columba palumbus</i>	Denmark	0.018	28.17	2.03	494.5	56.16	10.38	0.606	0.365	1
<i>Columba palumbus</i>	Finland	0.024	30.00	2.03	494.5	64.95	26.07	0.812	0.365	1
<i>Columba palumbus</i>	France	0.052	14.43	2.03	494.5	46.71	1.72	0.485	0.365	1
<i>Columba palumbus</i>	Spain	0.011	45.40	2.03	494.5	39.90	-2.99	0.461	0.365	1
<i>Columba palumbus</i>	Hungary	0.008	21.47	2.03	494.5	47.16	19.51	0.490	0.365	1
<i>Columba palumbus</i>	Norway	0.016	7.00	2.03	494.5	64.56	18.01	0.796	0.365	1
<i>Columba palumbus</i>	Poland	0.018	60.41	2.03	494.5	51.92	19.13	0.548	0.365	1
<i>Corvus corax</i>	Denmark	0.293	78.06	0.00	1200.6	56.16	10.38	0.448	1.189	0
<i>Corvus cornix</i>	Czech Rep.	-0.019	13.89	5.71	544.5	49.80	15.48	0.528	0.944	0
<i>Corvus cornix</i>	Denmark	0.011	41.15	5.71	544.5	56.16	10.38	0.579	0.944	0
<i>Corvus cornix</i>	Finland	-0.013	31.69	5.71	544.5	64.95	26.07	0.717	0.944	0
<i>Corvus cornix</i>	Hungary	0.113	24.09	5.71	544.5	47.16	19.51	0.536	0.944	0
<i>Corvus cornix</i>	Norway	0.001	17.30	5.71	544.5	64.56	18.01	0.708	0.944	0
<i>Corvus corone</i>	France	0.001	20.41	5.71	544.5	46.71	1.72	0.537	0.944	0
<i>Corvus frugilegus</i>	Denmark	0.031	46.53	2.32	453.5	56.16	10.38	0.645	0.904	1
<i>Corvus monedula</i>	Denmark	0.000	32.11	0.29	249.0	56.16	10.38	0.635	0.660	1
<i>Corvus monedula</i>	Finland	0.168	60.00	0.29	249.0	64.95	26.07	0.872	0.660	1
<i>Corvus monedula</i>	Spain	-0.030	46.86	0.29	249.0	39.90	-2.99	0.573	0.660	1
<i>Corvus monedula</i>	Poland	0.019	18.23	0.29	249.0	51.92	19.13	0.576	0.660	1
<i>Cuculus canorus</i>	Denmark	-0.010	21.72	49.38	120.5	56.16	10.38	0.562	0.164	0
<i>Cyanopica cyanus</i>	Spain	0.060	56.86	0.00	71.0	39.90	-2.99	0.579	0.314	0
<i>Delichon urbica</i>	Spain	0.026	35.51	44.25	19.6	39.90	-2.99	0.378	-0.262	0
<i>Dendrocopos major</i>	Czech Rep.	0.018	18.56	0.00	89.7	49.80	15.48	0.498	0.405	0
<i>Dendrocopos major</i>	Denmark	0.005	14.20	0.00	89.7	56.16	10.38	0.581	0.405	0
<i>Dendrocopos major</i>	France	0.033	14.00	0.00	89.7	46.71	1.72	0.460	0.405	0
<i>Dendrocopos major</i>	Spain	0.069	58.14	0.00	89.7	39.90	-2.99	0.411	0.405	0
<i>Dendrocopos major</i>	Hungary	-0.014	32.98	0.00	89.7	47.16	19.51	0.466	0.405	0
<i>Dendrocopos major</i>	Norway	0.097	9.85	0.00	89.7	64.56	18.01	0.745	0.405	0
<i>Dendrocopos syriacus</i>	Hungary	0.023	13.99	0.00	76.8	47.16	19.51	0.726	--	0
<i>Dryocopus martius</i>	France	0.069	38.71	0.00	273.0	46.71	1.72	0.517	0.870	0

<i>Dryocopus martius</i>	Poland	0.025	50.16	0.00	273.0	51.92	19.13	0.542	0.870	0
<i>Egretta garzetta</i>	France	0.041	24.50	7.53	532.5	46.71	1.72	0.400	--	0
<i>Emberiza cirulus</i>	Spain	-0.008	15.00	0.48	23.8	39.90	-2.99	0.583	-0.135	1
<i>Emberiza citrinella</i>	Czech Rep.	-0.014	15.54	4.72	26.8	49.80	15.48	0.611	-0.106	1
<i>Emberiza citrinella</i>	Denmark	-0.016	9.79	4.72	26.8	56.16	10.38	0.600	-0.106	1
<i>Emberiza citrinella</i>	Finland	-0.002	10.43	4.72	26.8	64.95	26.07	0.743	-0.106	1
<i>Emberiza citrinella</i>	Norway	-0.023	8.20	4.72	26.8	64.56	18.01	0.734	-0.106	1
<i>Emberiza citrinella</i>	Poland	-0.016	3.08	4.72	26.8	51.92	19.13	0.597	-0.106	1
<i>Emberiza schoeniclus</i>	Czech Rep.	0.001	28.07	10.52	18.8	49.80	15.48	0.510	-0.167	0
<i>Emberiza schoeniclus</i>	Denmark	-0.008	9.17	10.52	18.8	56.16	10.38	0.575	-0.167	0
<i>Emberiza schoeniclus</i>	Norway	-0.022	8.00	10.52	18.8	64.56	18.01	0.705	-0.167	0
<i>Emberiza schoeniclus</i>	Poland	0.026	36.51	10.52	18.8	51.92	19.13	0.527	-0.167	0
<i>Erithacus rubecula</i>	Czech Rep.	-0.001	16.71	5.00	16.4	49.80	15.48	0.510	-0.196	0
<i>Erithacus rubecula</i>	Denmark	0.006	8.91	5.00	16.4	56.16	10.38	0.585	-0.196	0
<i>Erithacus rubecula</i>	France	0.023	5.19	5.00	16.4	46.71	1.72	0.503	-0.196	0
<i>Erithacus rubecula</i>	Norway	-0.001	5.47	5.00	16.4	64.56	18.01	0.727	-0.196	0
<i>Erithacus rubecula</i>	Spain	0.015	2.24	5.00	16.4	39.90	-2.99	0.519	-0.196	0
<i>Falco tinnunculus</i>	Czech Rep.	-0.003	50.58	5.60	174.5	49.80	15.48	0.228	0.570	1
<i>Falco tinnunculus</i>	Denmark	0.019	28.35	5.60	174.5	56.16	10.38	0.294	0.570	1
<i>Falco tinnunculus</i>	France	-0.040	6.32	5.60	174.5	46.71	1.72	0.290	0.570	1
<i>Falco tinnunculus</i>	Spain	-0.012	115.49	5.60	174.5	39.90	-2.99	0.350	0.570	1
<i>Falco tinnunculus</i>	Hungary	-0.007	25.00	5.60	174.5	47.16	19.51	0.284	0.570	1
<i>Ficedula hypoleuca</i>	Denmark	-0.034	5.39	43.00	14.4	56.16	10.38	0.583	-0.364	0
<i>Ficedula hypoleuca</i>	Finland	0.013	6.59	43.00	14.4	64.95	26.07	0.726	-0.364	0
<i>Ficedula hypoleuca</i>	Norway	-0.019	7.07	43.00	14.4	64.56	18.01	0.718	-0.364	0
<i>Fringilla coelebs</i>	Czech Rep.	-0.010	13.62	5.54	24.2	49.80	15.48	0.499	-0.126	0
<i>Fringilla coelebs</i>	Denmark	0.011	9.39	5.54	24.2	56.16	10.38	0.571	-0.126	0
<i>Fringilla coelebs</i>	Finland	-0.002	8.50	5.54	24.2	64.95	26.07	0.710	-0.126	0
<i>Fringilla coelebs</i>	France	-0.003	6.55	5.54	24.2	46.71	1.72	0.500	-0.126	0
<i>Fringilla coelebs</i>	Spain	0.047	17.26	5.54	24.2	39.90	-2.99	0.515	-0.126	0
<i>Fringilla coelebs</i>	Norway	0.008	7.30	5.54	24.2	64.56	18.01	0.701	-0.126	0
<i>Fringilla coelebs</i>	Poland	0.002	8.76	5.54	24.2	51.92	19.13	0.522	-0.126	0
<i>Fulica atra</i>	Denmark	0.010	19.87	3.32	732.5	56.16	10.38	0.158	0.484	0
<i>Galerida cristata</i>	Spain	-0.013	36.40	0.00	44.7	39.90	-2.99	0.417	0.033	1
<i>Gallinago gallinago</i>	Denmark	-0.028	25.83	7.05	106.5	56.16	10.38	0.279	0.117	0
<i>Gallinago gallinago</i>	Finland	-0.009	54.15	7.05	106.5	64.95	26.07	0.542	0.117	0
<i>Gallinula chloropus</i>	Denmark	-0.015	20.00	0.00	348.5	56.16	10.38	0.400	0.296	0
<i>Gallinula chloropus</i>	France	0.018	10.32	0.00	348.5	46.71	1.72	0.263	0.296	0
<i>Garrulus glandarius</i>	Czech Rep.	0.041	39.15	0.00	161.7	49.80	15.48	0.484	0.605	0
<i>Garrulus glandarius</i>	Denmark	0.001	20.95	0.00	161.7	56.16	10.38	0.564	0.605	0
<i>Garrulus glandarius</i>	France	0.030	11.16	0.00	161.7	46.71	1.72	0.446	0.605	0
<i>Garrulus glandarius</i>	Hungary	0.011	27.87	0.00	161.7	47.16	19.51	0.452	0.605	0
<i>Grus grus</i>	Poland	0.051	100.00	39.10	4541.5	51.92	19.13	0.572	1.246	1
<i>Haematopus ostralegus</i>	Denmark	-0.009	40.01	21.39	531.0	56.16	10.38	0.587	0.583	0
<i>Haematopus ostralegus</i>	Norway	-0.010	19.00	21.39	531.0	64.56	18.01	0.714	0.583	0
<i>Hippolais icterina</i>	Denmark	-0.034	7.76	71.34	13.3	56.16	10.38	0.601	-0.293	0
<i>Hippolais polyglotta</i>	Spain	0.028	16.57	30.63	11.5	39.90	-2.99	0.619	--	0
<i>Hirundo rustica</i>	Czech Rep.	-0.007	15.63	42.34	19.1	49.80	15.48	0.482	-0.269	1
<i>Hirundo rustica</i>	Denmark	-0.008	10.15	42.34	19.1	56.16	10.38	0.561	-0.269	1
<i>Hirundo rustica</i>	Finland	-0.013	11.18	42.34	19.1	64.95	26.07	0.712	-0.269	1
<i>Hirundo rustica</i>	Poland	0.007	10.77	42.34	19.1	51.92	19.13	0.507	-0.269	1
<i>Lanius collurio</i>	Czech Rep.	0.046	21.31	64.73	30.7	49.80	15.48	0.554	0.008	1
<i>Lanius collurio</i>	Denmark	-0.011	17.39	64.73	30.7	56.16	10.38	0.639	0.008	1
<i>Lanius collurio</i>	Hungary	0.017	11.91	64.73	30.7	47.16	19.51	0.559	0.008	1
<i>Lanius collurio</i>	Poland	-0.003	35.01	64.73	30.7	51.92	19.13	0.580	0.008	1
<i>Lanius excubitor</i>	Poland	0.072	22.36	4.87	66.9	51.92	19.13	0.555	0.199	1
<i>Lanius excubitor</i>	Spain	-0.044	31.05	4.87	66.9	39.90	-2.99	0.669	0.199	1
<i>Lanius senator</i>	Spain	-0.012	36.12	28.45	36.0	39.90	-2.99	0.575	--	1

<i>Larus argentatus</i>	Denmark	0.019	50.09	22.14	895.0	56.16	10.38	0.673	0.774	0
<i>Larus argentatus</i>	France	0.050	15.10	22.14	895.0	46.71	1.72	0.952	0.774	0
<i>Larus canus</i>	Denmark	-0.009	59.94	13.76	386.5	56.16	10.38	0.671	0.602	1
<i>Larus canus</i>	Norway	-0.024	14.00	13.76	386.5	64.56	18.01	0.729	0.602	1
<i>Larus fuscus</i>	France	0.170	22.00	34.35	817.5	46.71	1.72	0.646	0.726	0
<i>Larus marinus</i>	Denmark	0.220	57.52	7.28	1599.5	56.16	10.38	0.528	0.837	0
<i>Larus ridibundus</i>	Denmark	-0.033	37.75	23.50	280.5	56.16	10.38	0.624	0.453	0
<i>Locustella naevia</i>	Czech Rep.	-0.010	7.22	29.75	12.7	49.80	15.48	0.597	-0.222	1
<i>Loxia curvirostra</i>	Denmark	-0.040	4.74	0.00	40.6	56.16	10.38	0.571	0.173	0
<i>Luscinia luscinia</i>	Denmark	-0.016	15.89	63.10	25.0	56.16	10.38	0.670	-0.125	0
<i>Luscinia megarhynchos</i>	Czech Rep.	0.099	15.52	32.20	20.2	49.80	15.48	0.722	-0.179	0
<i>Luscinia megarhynchos</i>	Spain	0.018	25.87	32.20	20.2	39.90	-2.99	0.560	-0.179	0
<i>Luscinia megarhynchos</i>	Hungary	0.023	10.53	32.20	20.2	47.16	19.51	0.659	-0.179	0
<i>Luscinia megarhynchos</i>	Poland	-0.011	46.01	32.20	20.2	51.92	19.13	0.794	-0.179	0
<i>Merops apiaster</i>	Spain	0.008	94.95	19.86	55.1	39.90	-2.99	0.335	-0.098	1
<i>Miliaria calandra</i>	Denmark	-0.004	10.29	6.40	47.7	56.16	10.38	0.782	0.074	1
<i>Miliaria calandra</i>	Spain	-0.008	24.18	6.40	47.7	39.90	-2.99	0.576	0.074	1
<i>Miliaria calandra</i>	Poland	0.064	31.64	6.40	47.7	51.92	19.13	0.671	0.074	1
<i>Motacilla alba</i>	Czech Rep.	-0.014	18.03	18.13	20.8	49.80	15.48	0.480	-0.250	0
<i>Motacilla alba</i>	Denmark	0.049	11.62	18.13	20.8	56.16	10.38	0.556	-0.250	0
<i>Motacilla alba</i>	Finland	0.000	7.57	18.13	20.8	64.95	26.07	0.699	-0.250	0
<i>Motacilla alba</i>	France	0.002	6.33	18.13	20.8	46.71	1.72	0.445	-0.250	0
<i>Motacilla alba</i>	Spain	-0.019	24.36	18.13	20.8	39.90	-2.99	0.421	-0.250	0
<i>Motacilla alba</i>	Norway	0.026	7.15	18.13	20.8	64.56	18.01	0.690	-0.250	0
<i>Motacilla alba</i>	Poland	0.005	25.56	18.13	20.8	51.92	19.13	0.505	-0.250	0
<i>Motacilla cinerea</i>	Czech Rep.	-0.017	8.00	24.68	17.4	49.80	15.48	0.507	-0.279	0
<i>Motacilla cinerea</i>	Denmark	-0.027	6.87	24.68	17.4	56.16	10.38	0.562	-0.279	0
<i>Motacilla cinerea</i>	France	-0.015	9.07	24.68	17.4	46.71	1.72	0.503	-0.279	0
<i>Motacilla flava</i>	Denmark	-0.027	11.20	40.99	17.5	56.16	10.38	0.562	-0.371	1
<i>Motacilla flava</i>	Finland	-0.036	11.84	40.99	17.5	64.95	26.07	0.709	-0.371	1
<i>Motacilla flava</i>	Poland	-0.030	5.39	40.99	17.5	51.92	19.13	0.509	-0.371	1
<i>Muscicapa striata</i>	Czech Rep.	0.057	15.56	64.40	15.5	49.80	15.48	0.516	-0.292	0
<i>Muscicapa striata</i>	Denmark	0.010	10.08	64.40	15.5	56.16	10.38	0.584	-0.292	0
<i>Muscicapa striata</i>	Finland	-0.004	9.06	64.40	15.5	64.95	26.07	0.729	-0.292	0
<i>Muscicapa striata</i>	Spain	-0.025	21.98	64.40	15.5	39.90	-2.99	0.551	-0.292	0
<i>Numenius arquata</i>	Finland	-0.008	42.07	44.67	725.0	64.95	26.07	0.744	0.595	0
<i>Oriolus oriolus</i>	Spain	0.056	30.34	44.77	68.5	39.90	-2.99	0.485	0.117	0
<i>Oriolus oriolus</i>	Hungary	0.003	24.12	44.77	68.5	47.16	19.51	0.536	0.117	0
<i>Parus ater</i>	Czech Rep.	0.017	12.08	0.00	9.3	49.80	15.48	0.520	-0.264	0
<i>Parus ater</i>	Denmark	-0.009	5.41	0.00	9.3	56.16	10.38	0.605	-0.264	0
<i>Parus ater</i>	France	-0.049	4.00	0.00	9.3	46.71	1.72	0.483	-0.264	0
<i>Parus ater</i>	Spain	0.019	18.49	0.00	9.3	39.90	-2.99	0.456	-0.264	0
<i>Parus caeruleus</i>	Czech Rep.	0.002	10.08	0.00	11.8	49.80	15.48	0.540	-0.200	0
<i>Parus caeruleus</i>	Denmark	0.007	6.61	0.00	11.8	56.16	10.38	0.622	-0.200	0
<i>Parus caeruleus</i>	Finland	0.258	8.00	0.00	11.8	64.95	26.07	0.825	-0.200	0
<i>Parus caeruleus</i>	France	0.033	5.17	0.00	11.8	46.71	1.72	0.537	-0.200	0
<i>Parus caeruleus</i>	Spain	0.027	9.93	0.00	11.8	39.90	-2.99	0.570	-0.200	0
<i>Parus caeruleus</i>	Hungary	0.025	5.94	0.00	11.8	47.16	19.51	0.535	-0.200	0
<i>Parus caeruleus</i>	Norway	0.012	6.09	0.00	11.8	64.56	18.01	0.808	-0.200	0
<i>Parus cristatus</i>	Denmark	-0.013	7.20	0.00	11.2	56.16	10.38	0.636	-0.189	0
<i>Parus cristatus</i>	France	0.005	4.81	0.00	11.2	46.71	1.72	0.603	-0.189	0
<i>Parus major</i>	Czech Rep.	-0.002	10.47	0.00	18.5	49.80	15.48	0.370	-0.073	0
<i>Parus major</i>	Denmark	-0.009	5.46	0.00	18.5	56.16	10.38	0.479	-0.073	0
<i>Parus major</i>	Finland	0.021	8.61	0.00	18.5	64.95	26.07	0.649	-0.073	0
<i>Parus major</i>	France	0.017	4.60	0.00	18.5	46.71	1.72	0.307	-0.073	0
<i>Parus major</i>	Spain	0.013	11.92	0.00	18.5	39.90	-2.99	0.000	-0.073	0
<i>Parus major</i>	Hungary	0.027	7.65	0.00	18.5	47.16	19.51	0.317	-0.073	0
<i>Parus major</i>	Norway	0.003	5.74	0.00	18.5	64.56	18.01	0.639	-0.073	0

<i>Parus major</i>	Poland	-0.009	16.49	0.00	18.5	51.92	19.13	0.408	-0.073	0
<i>Parus montanus</i>	Norway	-0.004	5.61	0.00	11.7	64.56	18.01	0.716	-0.102	0
<i>Parus palustris</i>	Czech Rep.	-0.022	10.60	0.00	11.9	49.80	15.48	0.542	-0.151	0
<i>Parus palustris</i>	Denmark	-0.013	4.03	0.00	11.9	56.16	10.38	0.638	-0.151	0
<i>Parus palustris</i>	France	-0.008	6.60	0.00	11.9	46.71	1.72	0.503	-0.151	0
<i>Passer domesticus</i>	Czech Rep.	-0.026	15.28	0.00	30.4	49.80	15.48	0.435	-0.041	0
<i>Passer domesticus</i>	Denmark	-0.013	4.72	0.00	30.4	56.16	10.38	0.521	-0.041	0
<i>Passer domesticus</i>	Finland	-0.038	16.67	0.00	30.4	64.95	26.07	0.671	-0.041	0
<i>Passer domesticus</i>	France	0.004	4.83	0.00	30.4	46.71	1.72	0.392	-0.041	0
<i>Passer domesticus</i>	Spain	-0.004	19.76	0.00	30.4	39.90	-2.99	0.277	-0.041	0
<i>Passer domesticus</i>	Hungary	-0.004	7.34	0.00	30.4	47.16	19.51	0.398	-0.041	0
<i>Passer domesticus</i>	Norway	-0.008	4.24	0.00	30.4	64.56	18.01	0.662	-0.041	0
<i>Passer domesticus</i>	Poland	-0.027	11.05	0.00	30.4	51.92	19.13	0.463	-0.041	0
<i>Passer montanus</i>	Czech Rep.	-0.009	16.67	3.62	21.7	49.80	15.48	0.378	-0.123	1
<i>Passer montanus</i>	Denmark	0.036	6.35	3.62	21.7	56.16	10.38	0.487	-0.123	1
<i>Passer montanus</i>	Spain	-0.027	13.79	3.62	21.7	39.90	-2.99	0.041	-0.123	1
<i>Passer montanus</i>	Hungary	0.073	8.08	3.62	21.7	47.16	19.51	0.325	-0.123	1
<i>Passer montanus</i>	Poland	-0.051	14.45	3.62	21.7	51.92	19.13	0.415	-0.123	1
<i>Perdix perdix</i>	Denmark	-0.012	24.92	0.00	382.0	56.16	10.38	0.655	0.259	1
<i>Phoenicurus ochruros</i>	Czech Rep.	0.009	15.89	15.83	16.0	49.80	15.48	0.612	-0.254	0
<i>Phoenicurus ochruros</i>	France	0.006	4.12	15.83	16.0	46.71	1.72	0.567	-0.254	0
<i>Phoenicurus ochruros</i>	Hungary	0.061	15.10	15.83	16.0	47.16	19.51	0.573	-0.254	0
<i>Phoenicurus ochruros</i>	Poland	-0.011	22.40	15.83	16.0	51.92	19.13	0.648	-0.254	0
<i>Phoenicurus phoenicurus</i>	Czech Rep.	0.061	19.10	33.93	15.9	49.80	15.48	0.505	-0.321	0
<i>Phoenicurus phoenicurus</i>	Denmark	0.019	12.05	33.93	15.9	56.16	10.38	0.579	-0.321	0
<i>Phoenicurus phoenicurus</i>	Finland	0.020	20.02	33.93	15.9	64.95	26.07	0.725	-0.321	0
<i>Phylloscopus collybita</i>	Czech Rep.	0.010	11.03	22.55	7.7	49.80	15.48	0.507	-0.455	0
<i>Phylloscopus collybita</i>	Denmark	0.168	7.88	22.55	7.7	56.16	10.38	0.581	-0.455	0
<i>Phylloscopus collybita</i>	France	-0.018	5.25	22.55	7.7	46.71	1.72	0.500	-0.455	0
<i>Phylloscopus trochilus</i>	Denmark	-0.020	6.06	68.09	9.4	56.16	10.38	0.617	-0.507	0
<i>Phylloscopus trochilus</i>	Hungary	-0.034	2.00	68.09	9.4	47.16	19.51	0.725	-0.507	0
<i>Phylloscopus trochilus</i>	Norway	-0.011	5.05	68.09	9.4	64.56	18.01	0.725	-0.507	0
<i>Pica pica</i>	Denmark	0.008	37.59	0.00	228.0	56.16	10.38	0.551	0.736	0
<i>Pica pica</i>	Finland	0.002	29.00	0.00	228.0	64.95	26.07	0.694	0.736	0
<i>Pica pica</i>	France	-0.112	15.12	0.00	228.0	46.71	1.72	0.438	0.736	0
<i>Pica pica</i>	Spain	-0.008	38.13	0.00	228.0	39.90	-2.99	0.399	0.736	0
<i>Pica pica</i>	Hungary	0.016	21.46	0.00	228.0	47.16	19.51	0.443	0.736	0
<i>Pica pica</i>	Norway	0.000	13.98	0.00	228.0	64.56	18.01	0.685	0.736	0
<i>Pica pica</i>	Poland	0.022	26.66	0.00	228.0	51.92	19.13	0.499	0.736	0
<i>Picus viridis</i>	France	0.026	17.82	0.00	193.5	46.71	1.72	0.587	0.636	0
<i>Picus viridis</i>	Spain	-0.006	50.24	0.00	193.5	39.90	-2.99	0.657	0.636	0
<i>Pluvialis apricaria</i>	Finland	-0.010	18.00	17.97	175.5	64.95	26.07	0.741	0.314	0
<i>Prunella modularis</i>	Czech Rep.	-0.011	13.56	9.23	19.0	49.80	15.48	0.585	-0.163	0
<i>Prunella modularis</i>	Denmark	-0.018	7.43	9.23	19.0	56.16	10.38	0.594	-0.163	0
<i>Prunella modularis</i>	France	-0.006	4.61	9.23	19.0	46.71	1.72	0.606	-0.163	0
<i>Pyrrhula pyrrhula</i>	Denmark	0.010	7.65	0.00	31.1	56.16	10.38	0.592	-0.058	0
<i>Pyrrhula pyrrhula</i>	Finland	0.061	5.00	0.00	31.1	64.95	26.07	0.738	-0.058	0
<i>Pyrrhula pyrrhula</i>	Norway	-0.042	16.00	0.00	31.1	64.56	18.01	0.729	-0.058	0
<i>Regulus regulus</i>	Denmark	-0.008	5.41	0.00	5.8	56.16	10.38	0.578	-0.446	0
<i>Regulus regulus</i>	France	-0.003	3.24	0.00	5.8	46.71	1.72	0.484	-0.446	0
<i>Regulus regulus</i>	Norway	-0.007	5.11	0.00	5.8	64.56	18.01	0.718	-0.446	0
<i>Riparia riparia</i>	Denmark	-0.020	23.10	42.73	13.2	56.16	10.38	0.572	-0.466	0
<i>Saxicola rubetra</i>	Denmark	-0.040	15.43	34.84	16.6	56.16	10.38	0.602	-0.222	1
<i>Saxicola rubetra</i>	Finland	-0.018	23.09	34.84	16.6	64.95	26.07	0.751	-0.222	1
<i>Saxicola torquata</i>	Hungary	-0.006	15.98	3.98	14.9	47.16	19.51	0.280	--	1
<i>Scolopax rusticola</i>	Finland	0.020	9.00	14.32	309.5	64.95	26.07	0.732	0.388	0
<i>Serinus serinus</i>	Czech Rep.	-0.039	13.94	4.91	12.0	49.80	15.48	0.621	-0.371	0
<i>Serinus serinus</i>	France	-0.033	6.06	4.91	12.0	46.71	1.72	0.574	-0.371	0

<i>Serinus serinus</i>	Spain	-0.025	21.67	4.91	12.0	39.90	-2.99	0.551	-0.371	0
<i>Serinus serinus</i>	Hungary	-0.059	14.14	4.91	12.0	47.16	19.51	0.581	-0.371	0
<i>Serinus serinus</i>	Poland	-0.044	24.04	4.91	12.0	51.92	19.13	0.658	-0.371	0
<i>Sitta europaea</i>	Czech Rep.	0.014	13.39	0.00	23.9	49.80	15.48	0.492	0.017	0
<i>Sitta europaea</i>	France	-0.038	4.92	0.00	23.9	46.71	1.72	0.450	0.017	0
<i>Sitta europaea</i>	Norway	0.025	5.41	0.00	23.9	64.56	18.01	0.788	0.017	0
<i>Streptopelia decaocto</i>	Czech Rep.	0.026	22.42	0.00	201.5	49.80	15.48	0.460	0.157	0
<i>Streptopelia decaocto</i>	Denmark	0.000	26.43	0.00	201.5	56.16	10.38	0.547	0.157	0
<i>Streptopelia decaocto</i>	France	0.077	11.67	0.00	201.5	46.71	1.72	0.418	0.157	0
<i>Streptopelia decaocto</i>	Spain	0.538	23.61	0.00	201.5	39.90	-2.99	0.315	0.157	0
<i>Streptopelia decaocto</i>	Hungary	0.044	10.79	0.00	201.5	47.16	19.51	0.424	0.157	0
<i>Streptopelia decaocto</i>	Poland	0.005	13.31	0.00	201.5	51.92	19.13	0.488	0.157	0
<i>Streptopelia turtur</i>	Czech Rep.	-0.022	27.99	25.90	136.5	49.80	15.48	0.550	0.102	1
<i>Streptopelia turtur</i>	France	-0.006	7.07	25.90	136.5	46.71	1.72	0.509	0.102	1
<i>Streptopelia turtur</i>	Spain	-0.004	97.53	25.90	136.5	39.90	-2.99	0.440	0.102	1
<i>Sturnus unicolor</i>	Spain	0.018	42.70	0.00	90.6	39.90	-2.99	0.728	0.262	1
<i>Sturnus vulgaris</i>	Czech Rep.	0.030	36.44	2.63	80.5	49.80	15.48	0.493	0.235	1
<i>Sturnus vulgaris</i>	Denmark	-0.015	14.31	2.63	80.5	56.16	10.38	0.567	0.235	1
<i>Sturnus vulgaris</i>	France	-0.005	9.75	2.63	80.5	46.71	1.72	0.477	0.235	1
<i>Sturnus vulgaris</i>	Hungary	0.042	13.46	2.63	80.5	47.16	19.51	0.477	0.235	1
<i>Sturnus vulgaris</i>	Norway	-0.016	8.83	2.63	80.5	64.56	18.01	0.697	0.235	1
<i>Sturnus vulgaris</i>	Poland	0.049	31.28	2.63	80.5	51.92	19.13	0.517	0.235	1
<i>Sylvia atricapilla</i>	Czech Rep.	0.074	13.56	19.63	18.9	49.80	15.48	0.476	-0.207	0
<i>Sylvia atricapilla</i>	Denmark	0.044	8.48	19.63	18.9	56.16	10.38	0.557	-0.207	0
<i>Sylvia atricapilla</i>	France	0.013	4.83	19.63	18.9	46.71	1.72	0.438	-0.207	0
<i>Sylvia atricapilla</i>	Spain	0.044	11.47	19.63	18.9	39.90	-2.99	0.373	-0.207	0
<i>Sylvia atricapilla</i>	Hungary	0.056	10.81	19.63	18.9	47.16	19.51	0.443	-0.207	0
<i>Sylvia atricapilla</i>	Norway	0.119	3.16	19.63	18.9	64.56	18.01	0.704	-0.207	0
<i>Sylvia atricapilla</i>	Poland	0.031	11.32	19.63	18.9	51.92	19.13	0.502	-0.207	0
<i>Sylvia borin</i>	Czech Rep.	-0.012	19.10	63.25	19.1	49.80	15.48	0.584	-0.220	0
<i>Sylvia borin</i>	Denmark	-0.014	6.78	63.25	19.1	56.16	10.38	0.598	-0.220	0
<i>Sylvia cantillans</i>	Spain	0.028	5.22	20.82	8.1	39.90	-2.99	0.668	--	0
<i>Sylvia communis</i>	Czech Rep.	0.000	14.70	53.05	14.5	49.80	15.48	0.527	-0.281	1
<i>Sylvia communis</i>	Denmark	0.004	7.54	53.05	14.5	56.16	10.38	0.598	-0.281	1
<i>Sylvia communis</i>	Spain	-0.021	22.94	53.05	14.5	39.90	-2.99	0.570	-0.281	1
<i>Sylvia curruca</i>	Czech Rep.	0.041	17.26	27.79	12.4	49.80	15.48	0.514	-0.296	0
<i>Sylvia curruca</i>	Denmark	-0.022	5.72	27.79	12.4	56.16	10.38	0.590	-0.296	0
<i>Sylvia curruca</i>	Hungary	-0.007	8.55	27.79	12.4	47.16	19.51	0.509	-0.296	0
<i>Sylvia melanocephala</i>	Spain	-0.002	23.22	3.61	13.5	39.90	-2.99	0.660	-0.269	0
<i>Tadorna tadorna</i>	Denmark	0.013	37.71	9.35	1152.0	56.16	10.38	0.638	0.684	0
<i>Tringa hypoleucos</i>	France	-0.021	20.00	44.39	47.8	46.71	1.72	0.527	-0.108	0
<i>Tringa totanus</i>	Denmark	-0.030	29.71	35.28	112.0	56.16	10.38	0.576	0.149	0
<i>Troglodytes troglodytes</i>	Czech Rep.	0.011	10.55	1.34	8.9	49.80	15.48	0.497	-0.312	0
<i>Troglodytes troglodytes</i>	Denmark	0.029	7.51	1.34	8.9	56.16	10.38	0.577	-0.312	0
<i>Troglodytes troglodytes</i>	France	0.012	4.90	1.34	8.9	46.71	1.72	0.461	-0.312	0
<i>Troglodytes troglodytes</i>	Norway	-0.002	3.00	1.34	8.9	64.56	18.01	0.731	-0.312	0
<i>Turdus iliacus</i>	Finland	-0.002	14.05	10.77	62.9	64.95	26.07	0.741	0.133	0
<i>Turdus iliacus</i>	Norway	0.026	4.00	10.77	62.9	64.56	18.01	0.732	0.133	0
<i>Turdus merula</i>	Czech Rep.	0.016	21.51	3.98	95.9	49.80	15.48	0.438	0.284	0
<i>Turdus merula</i>	Denmark	0.015	12.22	3.98	95.9	56.16	10.38	0.524	0.284	0
<i>Turdus merula</i>	France	0.016	8.02	3.98	95.9	46.71	1.72	0.396	0.284	0
<i>Turdus merula</i>	Spain	0.007	21.98	3.98	95.9	39.90	-2.99	0.284	0.284	0
<i>Turdus merula</i>	Hungary	0.012	9.08	3.98	95.9	47.16	19.51	0.402	0.284	0
<i>Turdus merula</i>	Norway	0.015	9.65	3.98	95.9	64.56	18.01	0.665	0.284	0
<i>Turdus merula</i>	Poland	-0.018	22.59	3.98	95.9	51.92	19.13	0.466	0.284	0
<i>Turdus philomelos</i>	Czech Rep.	-0.001	29.03	14.65	70.5	49.80	15.48	0.576	0.196	0
<i>Turdus philomelos</i>	Denmark	-0.004	15.37	14.65	70.5	56.16	10.38	0.596	0.196	0
<i>Turdus philomelos</i>	France	0.034	8.10	14.65	70.5	46.71	1.72	0.595	0.196	0

<i>Turdus pilaris</i>	Czech Rep.	-0.003	29.31	10.77	92.1	49.80	15.48	0.709	0.261	1
<i>Turdus pilaris</i>	Denmark	-0.001	18.14	10.77	92.1	56.16	10.38	0.634	0.261	1
<i>Turdus pilaris</i>	Finland	0.037	20.95	10.77	92.1	64.95	26.07	0.735	0.261	1
<i>Turdus pilaris</i>	Norway	-0.015	8.56	10.77	92.1	64.56	18.01	0.727	0.261	1
<i>Turdus pilaris</i>	Poland	-0.038	5.50	10.77	92.1	51.92	19.13	0.677	0.261	1
<i>Turdus viscivorus</i>	Czech Rep.	0.024	49.90	4.36	117.8	49.80	15.48	0.526	0.313	0
<i>Turdus viscivorus</i>	Denmark	0.018	23.86	4.36	117.8	56.16	10.38	0.602	0.313	0
<i>Turdus viscivorus</i>	France	-0.011	27.00	4.36	117.8	46.71	1.72	0.534	0.313	0
<i>Upupa epops</i>	Spain	0.001	40.43	9.59	67.1	39.90	-2.99	0.339	0.102	0
<i>Vanellus vanellus</i>	Denmark	-0.011	33.99	12.08	218.5	56.16	10.38	0.598	0.326	1

582

583 **REFERENCE**

584 Thuiller W, Lavergne S, Roquet C, Boulangeat I, Lafourcade B, Araújo MB

585 2011. Consequences of climate change on the tree of life in Europe. *Nature*

586 470:531–534.

587