

Interferon Tau in the Ovine Uterus

Gwonhwa Song*, Jae Yong Han*, Thomas E. Spencer** and Fuller W. Bazer*, **

WCU Biomodulation Major, Department of Agricultural Biotechnology, and Research Institute for Agriculture and Life Sciences, Seoul National University, Korea*, Department of Animal Science, Center for Animal Biotechnology and Genomics, Texas A&M University, 442 Kleberg Center, 2471 TAMU, College Station, Texas 77843-2471, USA **

ABSTRACT

The peri-implantation period in mammals is critical with respect to survival of the conceptus (embryo/fetus and associated extraembryonic membranes) and establishment of pregnancy. During this period of pregnancy, reciprocal communication between ovary, conceptus, and endometrium is required for successful implantation and placentation. At this time, interferon tau (IFNT) is synthesized and secreted by the mononuclear trophoblastic cells of the conceptus between days 10 and 21~25. The actions of IFNT to signal pregnancy recognition and induce or increase expression of IFNT-stimulated genes (ISGs), such as *ISG15* and *OAS*, are mediated by the Type I IFN signal transduction pathway. This article reviews the history, signaling pathways of IFNT and the uterine expression of several IFNT-stimulated genes during the peri-implantation period. Collectively, these newly identified genes are believed to be critical to unraveling the mechanism(s) of reciprocal fetal-maternal interactions required for successful implantation and pregnancy.

(**Key words** : Interferon tau, Progesterone, Sheep, Uterus, Implantation)

INTRODUCTION

In eutherian mammals, including sheep, implantation of the blastocyst is a most important developmental event associated with viviparity (Spencer and Bazer, 2002; Spencer et al., 2004). During the peri-implantation period in the ovine uterus, the spherical blastocyst elongates to a tubular and then a filamentous form, and develops into a conceptus. At this time, interferon tau (IFNT) is synthesized and secreted by the mononuclear trophoblastic cells of the conceptus between days 10 and 21~25 (maximally on days 14 to 16) (Ashworth and Bazer, 1989; Farin et al., 1989; Bazer, 1992; Roberts et al., 1999). In the ovine uterus, IFNT acts directly on the endometrial luminal epithelium (LE) and superficial ductal glandular epithelium (sGE) to suppress transcription of estrogen receptor alpha (*ESR1*) and oxytocin receptor (*OXTR*) genes (Spencer and Bazer, 1996; Fleming et al., 2001), thereby preventing production of luteolytic pulses of prostaglandin F_{2α} (PGF) (Fig. 1).

During the estrous cycle, *ESR1* expression increases and progesterone receptor (*PGR*) expression decreases on days 11 to 13, and subsequently estrogen (E₂) induces *OXTR* expression on days 13 to 14 (Wathes and Hamon, 1993;

Spencer and Bazer, 1995). Thus, oxytocin from the posterior pituitary and/or corpus luteum (CL) can then induce release of luteolytic pulses of PGF on days 15 and 16 (Hooper et al., 1986). During early pregnancy, IFNT produced by the elongating ovine conceptus suppresses *ESR1* expression which then prevents *ESR1*-induced *OXTR* expression (Spencer and Bazer, 1996; Stevenson et al., 1994; Lamming et al., 1995; Spencer et al., 1995). Collectively, these indicate that the antiluteolytic actions of IFNT are to prevent increases in epithelial expression of E₂-responsive *ESR1*, *PGR*, and *OXTR* gene by directly inhibiting transcription of the *ESR1* gene and maintaining secretion of progesterone (P₄) by the CL (Fleming et al., 2001; Fleming et al., 2006; Spencer et al., 2004).

In the ovine uterus, establishment and maintenance of pregnancy requires reciprocal communication between the ovary, conceptus, and endometrium by means of endocrine and paracrine signals during implantation and synepitheliochorial placentation (Spencer and Bazer, 2002). P₄, the hormone of pregnancy, plays an important role in the establishment and maintenance of a uterine environment that supports conceptus development. Endometrial gland secretions, including growth factors, cytokines, and ions, are predominantly regulated by P₄ (Spencer et al., 1999) and are required for peri-implantation

Corresponding author : Dr. Gwonhwa Song, Seoul National University, 599 Gwanak-ro, Gwanak-gu, Seoul 151-921, Korea. Tel: +82-2-880-4849; FAX: +82-2-874-4811, E-mail: ghsong@snu.ac.kr




Fig. 1. Schematic illustration of the current working hypothesis on hormonal regulation of the endometrial antiluteolytic mechanism and cross-talk between the conceptus and the maternal endometrium. During the estrous cycle, *ESR1* expression increases and *PGR* expression decreases and then E2 induces *OXTR* expression, thereby allowing oxytocin from the posterior pituitary and/or CL to induce release of luteolytic pulses of *PGF*. In contrast, during early pregnancy, secreted IFNT from fully elongated conceptus silences *ESR1* expression which prevents E2-induced *OXTR* expression. Legend: IFNT, interferon tau; IFNAR, Type I IFN receptor; *PGR*, progesterone receptor; *ESR1*, estrogen receptor alpha; *OXTR*, oxytocin receptor; *IRF2*, interferon regulatory factor 2; LE, luminal epithelium; sGE, superficial ductal glandular epithelium. (Adapted from Spencer *et al.*, 2007.)

conceptus survival, elongation, and development (Gray *et al.*, 2001; Gray *et al.*, 2000). P4 acts via its cognate receptor, *PGR*. In the ovine endometrium, *PGR* are expressed in epithelia and stroma and allow P4 to directly regulate a variety of genes in the uterus. However, *PGR* expression is down-regulated by continuous exposure to P4 in ovine endometrial LE and GE after days 11 and 13 of pregnancy, respectively (Spencer and Bazer, 1995). The paradigm of loss of *PGR* in endometrial epithelia immediately before implantation is common to sheep (Spencer and Bazer, 2002; Spencer and Bazer, 1995), cattle (Kimmins and MacLaren, 2001), and pigs (Geisert *et al.*, 1994), and other mammals studied to date including humans and mice (see review by (Spencer *et al.*, 2004)).

During the peri-implantation period, uterine epithelial cell functions might be regulated by interactions between

reprogrammed epithelial cells following down-regulation of *PGR* and specific factors produced by *PGR*-positive stromal cells in response to P4, and/or products of the conceptus such as IFNT, placental lactogen, and placental growth hormone (see review by Spencer *et al.*, 2004). A large number of genes are induced by IFNT throughout the width of the uterine wall. These IFNT-stimulated genes (ISGs) are proposed to have biological roles in pregnancy recognition and uterine receptivity (Spencer and Bazer, 2002). In addition, induction of an antiviral state in the endometrium during early pregnancy may be beneficial by inhibiting sexually transmitted viruses as well as modulating local immune cells to promote tolerance of the allogeneic conceptus and stimulating production of cytokines beneficial for conceptus survival and growth (Hansen, 1995; Tekin and Hansen, 2002; Croy *et al.*, 2003).

Collectively, knowledge of the complex, precisely orchestrated interaction between P4 and IFNT during the implantation period should provide new insights to improve fertility in humans and domestic animals, and provide key knowledge for interpreting cross-talk mechanisms between maternal endometrium and conceptus.

OVERVIEW OF IFNT IN SHEEP

1. History of IFNT

The developing conceptus must signal its presence to the mother in order to ensure successful establishment and maintenance of pregnancy, a process termed maternal recognition of pregnancy (Short, 1969). In the 1960s, Moor and Rowson reported extension of the inter-estrous interval following a transfer of day 13 sheep blastocysts into recipient ewes on day 12 of the estrous cycle (Moor and Rowson, 1964), and that removal of blastocysts after day 13 significantly extended CL life-span (Moor and Rowson, 1966). In addition, infusion of the homogenates of the sheep conceptus collected between days 14 and 15 into the uterine lumen of recipient ewes (on or before day 12 of their cycle) extend CL life-span and the inter-estrous interval of cyclic ewes. However, infusion of pig conceptus homogenates had no effects on estrous cycle length in ewes (Rowson and Moor, 1967). The transfer of trophoblastic vesicles from blastocysts collected between days 11 and 13, without the embryonic disc, to recipient ewes on day 12 of the estrous cycles maintained CL function (Heyman *et al.*, 1984). The first report of secretion of low molecular weight acidic proteins by day 16 ovine conceptuses was by

Wilson *et al.* (1979). Later, in 1982, Godkin and colleagues characterized secretion of the low molecular weight acid protein by cultured ovine conceptuses collected between days 13 and 21 of pregnancy and termed it protein X (Godkin *et al.*, 1982). In a later study, protein X from ovine trophoderm was termed ovine trophoblast protein 1 (oTP-1) (Godkin *et al.*, 1984). Subsequently, native purified or recombinant oTP-1 was shown to extend the inter-estrous interval of ewes and to attenuate oxytocin-induced PGF release in sheep (Fincher *et al.*, 1986; Vallet *et al.*, 1988; Ott *et al.*, 1993).

oTP-1 has been found to have an amino acid sequence homologous to that of bovine IFN alpha (Imakawa *et al.*, 1987) and also to possess the antiviral and antiproliferative properties (Pontzer *et al.*, 1988; Pontzer *et al.*, 1991; Roberts *et al.*, 1989). Therefore, oTP-1 has been renamed IFNT and designated as a member of the Type I IFN family (Roberts *et al.*, 1992; Bazer *et al.*, 1996) by the International Interferon Society. In the conceptus, *IFNT* mRNA increases from day 12 to 14 and then declines to day 22 and is localized to mononuclear trophodermal cells (Farin *et al.*, 1989; Hansen *et al.*, 1988). Like other Type I IFN family members such as IFN alpha, -beta, -delta, -epsilon, -kappa and -omega, IFNT possesses potent antiviral (Pontzer *et al.*, 1988; Pontzer *et al.*, 1991; Roberts *et al.*, 1989), antiproliferative (Roberts *et al.*, 1989; Fillion *et al.*, 1991), and immunomodulatory biological activities and effects (Fillion *et al.*, 1991; Tennakoon *et al.*, 2001). IFNT is most closely related to IFN omega encoding the 172 amino acid sequence (Ealy *et al.*, 1998). Even though IFNT shares a high-degree of DNA and amino acid sequence identity in ruminants (sheep, cattle, goats) (Imakawa *et al.*, 1987; Roberts *et al.*, 1992), the precise biochemical structure of IFNT is different among species because of different post-translational modifications. Ovine IFNT is not glycosylated, bovine IFNT is glycosylated, and caprine IFNT is found in both glycosylated and non-glycosylated forms (Roberts *et al.*, 1992; Nephew *et al.*, 1993; Baumbach *et al.*, 1990).

2. Type I IFN Signal Transduction Pathway

The actions of IFNT to signal pregnancy recognition and induce or increase expression of IFNT-stimulated genes (ISGs) are mediated by the Type I IFN signal transduction pathway (Fig. 2). Type I IFNs bind to a common Type I IFN receptor (*IFNAR*), a heterodimer consisting of two subunits, *IFNAR1* and *IFNAR2*, associated with janus kinase 1 (*JAK1*) and

tyrosine kinase 2 (*TYK2*) (Novick *et al.*, 1994; Domanski *et al.*, 1995). The receptor is present in all endometrial cell types, but is highest in endometrial LE (Rosenfeld *et al.*, 2002). *IFNAR* classically activate the JAK/STAT (signal transducers and activators of transcription) signaling pathway (Darnell, 1997;




Fig. 2. Schematic illustration of the current working hypothesis on IFNT signaling in the ovine endometrial stroma and glandular epithelium. IFNT binds to a common Type I IFN receptor, *IFNAR1* and *IFNAR2* containing tyrosine kinase such as *JAK1* and *TYK2*, and activates the JAK/STAT signaling pathway. Phosphorylated *STAT1* binds the phosphorylated *STAT2* to form a heterodimer and translocates to the nucleus after forming a heterotrimeric transcriptional complex by binding with *ISGF3G*, collectively termed *ISGF3*. In addition to *STAT1/2* heterodimerization, Type I IFN also induces formation of phosphorylated *STAT1* homodimers, termed *GAF*. In the nucleus, *ISGF3* binds to the IFN-stimulated response element (*ISRE*) in promoter regions of *ISGs* and activates their transcription in cooperation with several coactivators. Similarly, *GAF* enters the nucleus, binds to *GAS* elements, and stimulates the transcription of *ISGs*. Legend: IFNT, interferon tau; *IFNAR*, Type I IFN receptor; *JAK1*, janus kinase 1; *TYK2*, tyrosine kinase 2; *STAT*, signal transducers and activators of transcription; *GAF*, gamma IFN activation factor; *GAS*, GAF activation sequence; *IRF1*, interferon regulatory factor 1; *ISGF3G*, IFN-stimulated transcription factor 3, gamma 48-kDa; *ISGF3*, IFN-stimulated transcription factor 3; *ISRE*, IFN-stimulated response element; *ISG*, IFNT-stimulated gene; *ISG15*, IFNT-stimulated gene 15; *B2M*, beta-2-microglobulin; *MHC*, major histocompatibility complex; *OAS*, oligoadenylate synthetase; *CXCL10*, chemokine (C-X-C motif) ligand 10; *IFI56*, interferon-induced protein 56; *GBP2*, guanylate binding protein 2, interferon-inducible. (Adapted from Spencer *et al.*, 2007.)

Darnell et al., 1994; Stark et al., 1998). Upon cognate ligand binding, IFNAR1 and IFNAR2 heterodimerize, change their conformation, and activate TYK2 and JAK1 by tyrosine phosphorylation (Gauzzi et al., 1996; Muller et al., 1993). The activated TYK2 phosphorylates STAT2 through its SH2 (src homologous 2) domain and then recruits signal transducers and activators of transcription-1 (STAT1) (Colamonici et al., 1994; Yan et al., 1996). Phosphorylated STAT1 binds phosphorylated STAT2 to form a heterodimer. The STAT1-STAT2 dimer is subsequently released from the receptor and binds ISGF3G (IFN-stimulated transcription factor 3, gamma 48 kDa) to form a heterotrimeric transcriptional complex, collectively termed ISGF3, which translocates to the nucleus (Silvennoinen et al., 1993; Shuai et al., 1994). In addition to STAT1/2 heterodimerization, Type I IFN also induces formation of phosphorylated STAT1 homodimers, termed GAF (gamma IFN activation factor) (Haque and Williams, 1994). In the nucleus, ISGF3 binds to an IFN-stimulated response element (ISRE) in promoter regions of ISGs to activate transcription in cooperation with several coactivators, such as the cAMP response element binding protein (CREB)-binding protein (CBP)/p300 (Bhattacharya, et al., 1996). Similarly, GAF enters the nucleus, binds to GAS (GAF activation sequence) elements to stimulate transcription of ISGs (Pine et al., 1994).

3. IFNT-Stimulated Genes (ISGs)

Most IFNT-stimulated genes (ISGs) are expressed by endometrial stroma and middle to deep GE of the ovine uterus (Choi et al., 2001; Johnson et al., 1999; Johnson et al., 2001). These ISGs include *STAT1* and *STAT2* (Johnson et al., 1999; Stewart et al., 2001), *IRF1* (Johnson et al., 1999; Stewart et al., 2001; Spencer et al., 1998), *ISG15* (Johnson et al., 1999; Johnson et al., 2000; Stewart et al., 2001), *Mx* (Ott et al., 1998), 2',5'-oligoadenylate synthetase (*OAS*) (Mirando et al., 1991; Johnson et al., 2001), major histocompatibility complex (*MHC*) class I (Choi et al., 2003), and beta-2-microglobulin (*B2M*) (Choi et al., 2003; Vallet et al., 1991).

IRF2, a known transcriptional repressor of Type I ISGs in the ovine uterus is constitutively expressed in the endometrial LE and sGE, increases during early pregnancy, and prevents induction or increases in transcription of ISGs by IFNT (Fig. 3) (Choi et al., 2002; Kim et al., 2003). *WNT7A* (Kim et al., 2003) and *LGALS15* (also known as galectin-15) (Gray et al., 2004) are the only genes known to be induced in LE and sGE by IFNT utilizing an unknown non-classical signaling




Fig. 3. Schematic illustration of the current working hypothesis on IFNT-signaling in the ovine luminal and superficial ductal glandular epithelium. IRF-2, a known transcriptional repressor of Type I ISGs in the ovine uterus constitutively expressed in the endometrial LE and sGE, increases during early pregnancy, and prevents induction or increases in transcription of ISGs by IFNT. At present, *LGALS15* and *WNT7A* are the only genes known to be induced in LE and sGE by IFNT via an unknown non-classical signaling pathway that does not involve the classical STAT transcription factors. Legend: IFNT, interferon tau; IFNAR, Type I IFN receptor; JAK1, janus kinase 1; TYK2, tyrosine kinase 2; GAS, GAF activation sequence; IRF, interferon regulatory factor; ISG, IFNT-stimulated gene; *LGALS15*, galectin 15; *WNT7A*, wingless-type MMTV integration site family, member 7A; LE, luminal epithelium; sGE, superficial ductal glandular epithelium. (Adapted from Spencer et al., 2007).

pathway that is independent of the classical STAT transcription factors.

4. Representative STAT1-Dependent ISGs

(1) Ubiquitin Cross-Reactive Protein / IFN-Stimulated Gene 15/17.

In humans, IFN-stimulated gene 15 (*ISG15*) encoding a 15-kDa protein, which has been identified in tumor and lymphoblastoid cells, and is induced by Type I IFNs (IFN alpha and beta) to a greater extent than by Type II IFN (IFN gamma) (Farrell et al., 1979; Korant et al., 1984). However, *ISG15* was renamed ubiquitin cross-reactive protein (*UCRP*) because its sequence is highly homologous to a tandem diubiquitin repeat, and antibodies raised to *ISG15* cross-react with ubiquitin (Haas et al., 1987). In bovine

endometrium, a 17 kDa precursor form of UCRP was detected as a 16 kDa form that might have undergone proteolytic cleavage in the endometrium (Johnson et al., 1999; Austin et al., 1996). During early pregnancy, in sheep, *ISG15* mRNA abundance increases only in stroma and GE from days 11 to 15 and then declines thereafter (Johnson et al., 1999). This period is coincident with peak production of IFNT by the ovine conceptus and *ISG15* expression increases in immortalized ovine LE, GE, and stromal cells treated with IFNT (Johnson et al., 1999).

(2) 2',5'-Oligoadenylate Synthetase.

2',5'-oligoadenylate synthetase (*OAS*) is induced by Type I and -II IFNs and polymerizes ATP into 2'-5' linked oligomers in order to bind and activate RNase L which can destroy intracellular viral RNAs. Further, *OAS* is involved in antiviral activity, cell growth, differentiation, and apoptosis (Samuel, 1991; Lengyel, 1993; Salzberg et al., 1997). In the ovine uterus, *OAS* is expressed only in the stroma and deep GE in response to IFNT and P4 during early pregnancy (Johnson et al., 1999; Mirando et al., 1991).

(3) RSAD2.

Radical S-adenosyl methionine domain containing 2 (*RSAD2*), also known as viperin, is a cytoplasmic antiviral protein that consists of 361 amino acids, and is encoded for by a gene which contains putative IRF binding sites in the promoter region (Chin and Cresswell, 2001; Sun and Nie, 2004). In humans, stable expression of *RSAD2* in fibroblasts inhibits human cytomegalovirus infection (Chin and Cresswell, 2001). *RSAD2* is also a potential antiviral effector expressed in patients with atherosclerosis (Olofsson et al., 2005) and chronic hepatitis C virus (Helbig et al., 2005). Chin *et al.* reported that *RSAD2* expression is greater in response to Type I than Type II IFN (IFN gamma) and that *RSAD2* may have an antiviral function (Chin and Cresswell, 2001).

(4) MDA5 (IFIH1).

Melanoma differentiation associated gene 5 (*MDA5* also known as *IFIH1*) is a dsRNA-dependent ATPase that responds predominantly to Type I IFNs and is known to be induced during differentiation, cancer reversion, and programmed cell death (Kang et al., 2002; Kang et al., 2004). The *IFIH1* gene contains both CARD and RNA helicase motifs and acts as a positive regulator to sense intracellular viral infection and stimulate innate antiviral responses including the production of

Type I IFN (Kang et al., 2002; Yoneyama et al., 2005). The V proteins of a wide variety of paramyxoviruses bind IFIH1 and inhibit its ability to activate the IFNB promoter (Andrejeva et al., 2004). Further, IFN beta promoter stimulator 1, which can induce Type I IFN and IFN-inducible genes through activation of IRF3, IRF7 and NF- κ B transcription factors, is known as an adaptor during IFIH1-mediated antiviral immune response (Kawai et al., 2005).

5. Representative STAT1-Independent ISGs

(1) Wingless-Type Mouse Mammary Tumor Virus Integration Site Family, Member 7A (*WNT7A*).

Most members of the WNT family are involved in embryonic cell growth, development, and differentiation and also in maternal-fetal interactions during implantation (Mohamed et al., 2005). In the ovine uterus, *wingless type mouse mammary tumor virus integration site family, member 7A* (*WNT7A*) was first identified and the gene is induced by IFNT during early pregnancy and expressed only in LE and sGE (Kim et al., 2003). Ovine endometrial *WNT7A* may activate the canonical WNT signaling pathway to stimulate proliferation and differentiation of conceptus trophoblasts may also regulate important genes for uterine receptivity for implantation and conceptus survival (Spencer et al., 2007).

(2) Galectin-15 (*LGALS15*).

Galectins are widely distributed in a variety of mammalian species, as well as non-mammalian species including birds, fish, and amphibians (Cooper and Barondes, 1999). They are members of a superfamily of binding lectins that bind β -galactoside via a CRD (carbohydrate recognition domain) (Barondes et al., 1994). In sheep, *LGALS15* was identified as the novel 14 kDa form of a P4-modulated protein associated with crystalline inclusion bodies in endometrial LE and conceptus trophoblasts (Kazemi et al., 1990). In the ovine uterus, *LGALS15* mRNA is expressed only in endometrial LE and sGE where it is induced by P4 and stimulated by IFNT. In addition, *LGALS15* protein has a nucleocytoplasmic distribution within the LE and sGE and is also concentrated near and on the apical surface (Gray et al., 2004). Therefore, *LGALS15* is secreted into the uterine lumen by the LE and sGE, where it may promote adhesion during implantation, as well as is phagocytosed by the trophoblasts and formed intracellular crystals (Gray et al., 2004; Gray et al., 2005).

(3) Cathepsin L (CTSL).

Cathepsins (*CTS*) are a family of lysosomal proteinases that are active in an acidic environment (Kirschke et al., 1998). They can degrade extracellular matrix (ECM) molecules, including collagens, laminin, fibronectin and proteoglycans and are also involved in catabolism of intracellular proteins and processing of pro-hormones. Available evidence supports the concept that a variety of proteases, as well as their specific inhibitors regulate trophoblast invasion in many species (e.g. mouse, rat, cat, pig, and human) during conceptus implantation (Afonso et al., 1997; Elangovan and Moulton, 1980; Li et al., 1992; Verhage et al., 1989; Geisert et al., 1997; Roberts et al., 1976; Jokimaa et al., 2001). CTSL is normally localized in lysosomes where it plays a major role in intracellular protein catabolism. In rodents, interactions between *Ctsb*, *Ctsl*, and *Cst3* (*Ctsb* and *Ctsl* inhibitors) are important for implantation and placentation, because inhibition of endometrial *Ctsb* and *Ctsl* results in abnormal embryonic development and uterine decidualization during the peri-implantation period (Afonso et al., 1997). In cats, CTSL is localized to the GE and can be detected in the uterine lumen where it is implicated in blastocyst invasion (Li et al., 1992). In pigs, *CTSL* is expressed in the endometrial GE and it is a P4-regulated component of the uterine lumen during implantation and placentation (Geisert et al., 1997).

(4) Cystatin C (CST3).

Cystatin C (*CST3*) is a secreted inhibitor of lysosomal cysteine proteases *CTSB* and *CTSL* (Abrahamson et al., 1986; Abrahamson et al., 1990; Hall et al., 1995; Grubb and Lofberg, 1982). In mice, *Ctsb* and *Ctsl* are necessary for normal embryonic development and uterine decidualization. The decidua coordinately expresses *Cst3* to control *Ctsb* and *Ctsl* actions within the implantation site (Afonso et al., 1997). A variety of proteases, as well as their inhibitors, regulate endometrial remodeling and trophoblast invasion in many species (e.g. mouse, rat, cat, sheep, pig, and human) during conceptus implantation and placentation.

DISCUSSION

What are the molecular mechanisms and signal transduction pathways activated by IFNT to regulate transcription of the novel epithelial genes, such as *WNT7A*, *LGALS15*, *CTSL*, and *CST3*, only in LE and sGE in the ovine uterus during the peri-implantation period? The current working hypothesis is

that IFNT utilizes STAT1-independent signaling pathway(s) to stimulate transcription of those genes in the LE and sGE (Fig. 4). In the ovine endometrial LE and sGE, the essential components of the JAK/STAT signal transduction, such as STAT1, -2, and ISGF3G, are not expressed, but IRF2, a potent transcriptional repressor of ISGs, was identified specifically in those cells, where it could repress or suppress the transcriptional activity of the promoter regions of ISGs that contain ISREs and IRF-Es (see bottom panel) (Choi et al., 2001; Stewart et al., 2001). Further, in our *in silico* study, the enhancer/promoter regions of bovine *WNT7A*, *LGALS15*, *CTSL*, and *CST3* genes had conserved transcription factor(s) binding sites for AP-1, CEBPB, CREB, ELK1, GATA, and LEF1/TCF7, but not for STATs or IRFs. Are there unknown non-classical JAK/STAT signaling pathways that are independent of STAT1? Recently, Platanius *et al.* reported that the generation of responses to Type I IFN requires the coordination and cooperation of multiple distinctive signaling cascades including the mitogen-activated protein (MAP) kinase p38 pathway and the phosphatidylinositol 3-kinase (PI3K) pathway (see review by Platanius, 2005). The p38 MAP kinase is phosphorylated and activated in several IFN-sensitive cell lines in response to Type I IFN such as IFN alpha and its inhibitor (SB203580) blocks IFN-inducible transcription (Platanius, 2005; Uddin et al., 2000). Inhibition of p38 MAP kinase has no effects on the phosphorylation of STAT1 or -2, and formation of the ISGF3 transcriptional complex (Uddin et al., 1999; Platanius et al., 2005). In addition, Type II IFN (IFN gamma) did not activate p38 MAP kinase in several cell lines (Katsoulidis et al., 2005; Uddin et al., 2000). Further, in the bovine uterus, IFNT activates the p38 MAP kinase pathway for induction of PTGS2 in myometrial cells (Doualla-Bell and Koromilas, 2001). These results indicate that p38 MAP kinase may play a role in Type I IFN-mediated signal transduction that is independent of STATs. Therefore, IFNT activation of p38 MAP kinase may be one signaling pathway whereby IFNT stimulates transcription of certain genes independent of STAT1 in the ovine uterus. Meanwhile, PI3K is activated in response to Type I or II IFNs. In the case of the Type I IFN signaling pathway, Type I IFNs activate the PI3K-signal pathway downstream of JAKs, in an insulin receptor substrate (IRS)-dependent but STAT-independent manner (Platanius, 2005; Kaur et al., 2005). The proposed model for the IFNT signal transduction cascade that is STAT1-independent in the ovine LE and sGE is illustrated in the upper panel of Fig. 4. IFNT-activated JAK1/TYK2 may




Fig. 4. A proposed model of IFNT signal transduction cascades that is independent of STAT1 in the ovine LE and sGE. IFNT-activated JAK1/TYK2 may regulate the phosphorylation of PI3K, resulting in the downstream activation of PDK1 and AKT. The activated AKT translocate into the nucleus and phosphorylate a variety of target proteins such as CBP/p300 or NF- κ B. Also, IFNT may activate MAPKKK or Raf which is activated by activated Ras. Activated MAPKKK and/or Raf subsequently regulate activation of downstream effectors including MAPKK, p38 MAPK, or MEK, ERK, respectively. In addition, the mTOR-p70S6K pathway which is activated by PI3K or AKT, may be involved in mRNA translation of ISGs by phosphorylated RPS6 and translational repressor 4EBP1. Legend: IFNT, interferon tau; IFNAR, Type I IFN receptor; JAK1, janus kinase 1; TYK2, tyrosine kinase 2; GAS, GAF activation sequence; IRF, interferon regulatory factor; ISG, IFNT-stimulated gene; LGALS15, galectin 15; WNT7A, wingless-type MMTV integration site family, member 7A; CTSL, cathepsin L; CST3, cystatin C; PI3K, phosphatidylinositol 3-kinase; PDK1, phosphoinositide-dependent protein kinase 1; AKT, proto oncogenic protein kinase Akt; CREB, cAMP-response element binding protein; mTOR, mammalian target of rapamycin; 4EBP1, EIF4-E-binding protein 1; E1F4E, eukaryotic translation-initiation factor 4 E; p70S, p70 ribosomal protein S6 kinase; RPS6, ribosomal protein S6; MAPK³, mitogen-activated protein kinase (MAPK) kinase kinase; MAPK², MAPK kinase; ERK, extracellular signal-regulated kinase; LE, luminal epithelium ; sGE, superficial ductal glandular epithelium.

regulate the phosphorylation of PI3K, resulting in the downstream activation of phosphoinositide-dependent protein kinase 1 (PDK1) and proto oncogenic protein kinase Akt (AKT). The activated AKT translocates into the nucleus and then phosphorylates a variety of target proteins such as CREB (cAMP-response element binding protein)-binding protein (CBP)/p300 or NF- κ B. Also, IFNT may activate MAPK kinase (MAPKKK) or Raf which is activated by activated Ras. Activated MAPKKK and/or Raf subsequently regulate activation of downstream effectors including MAPK kinase (MAPKK), p38 MAPK, MEK, or extracellular signal-regulated kinase (ERK). In addition, the mammalian target of rapamycin (mTOR)-p70 ribosomal protein S6 kinase (p70S6K) pathways which are activated by PI3K or AKT, may be involved in mRNA translation of ISGs by phosphorylated ribosomal protein S6 (RPS6) and translational repressor 4EBP1 (eukaryotic translation-initiation factor 4 E (EIF4-E)-binding protein 1). This hypothesis is supported by available results that IFNT and growth factors including insulin-like growth factor 2 stimulate PI3K-AKT and MAPK signal transduction cascades in ovine trophodermal and LE cells. Meanwhile, another possible scenario in the ovine uterus during the peri-implantation period is that IFNT may induce WNT7A using the canonical WNT signaling pathway between days 12 and 16 of pregnancy and then WNT7A acts in an autocrine or paracrine manner to stimulate the *LGALS15*, *CTSL*, and *CST3* genes in endometrial LE and sGE. Because *WNT7A* is the only gene truly induced by IFNT, its expression is not detected on day 12 of pregnancy, but is induced by IFNT between days 14 and 16 (Farrell et al., 1979). In fact, *LGALS15*, *CTSL*, and *CST3* genes are stimulated by IFNT between days 14 and 16 of pregnancy,

Interestingly, the ovine placenta expresses large numbers of aspartic proteinase inhibitor genes, termed pregnancy-associated glycoproteins, and the endometrial glands also express large amounts of serine protease inhibitors, termed serpins or uterine milk proteins, that could regulate the activity of endometrial CTS identified in these studies. Therefore, the molecular control of expression of CTS in the ovine endometrium may play an important role in establishing a regulatory network of multiple proteolytic enzymes responsible for ECM remodeling during implantation and placentation. Further, coordinated increases in *CTSL* and *CTSB* with *CST3* occur in endometrial LE and sGE as well as in conceptus trophoderm during early pregnancy. Thus, one biological role of *CST3* may be to inhibit the actions of cysteine

proteases produced by the conceptus and endometrial epithelia in order to limit the invasive activity of the trophoblast. These results support the general idea that proteases and their inhibitors expressed at the maternal-fetal interface are important for uterine receptivity, endometrial remodeling and conceptus implantation during pregnancy in mammals.

CONCLUSIONS

During the peri-implantation period in sheep, *CTSL* and *CST3* are novel P4-induced and IFNT-stimulated genes in endometrial LE and sGE. The majority of ISGs including *RSAD2* and *IFIH1* are expressed in endometrial stroma and middle to deep glands as well as immune cells in response to cell signaling involving the classical STAT1-dependent JAK/STAT signal transduction pathway without a requirement for P4 in the ovine uterus. It has been reported and hypothesized that Type I IFNs and many common ISGs are upregulated for the implanting conceptus in the endometrium during pregnancy in humans, rodents, and domestic animals. Recent evidence that ISGs are among the most upregulated genes in human decidualized stromal cells by trophoblast conditioned medium supports the hypothesis that a lack of ISG expression would compromise pregnancy. In contrast, IFNT induction of several non-classical ISGs, such as *LGALS15*, *WNT7A*, *CTSL*, and *CST3* in endometrial LE and sGE is dependent on P4, which is hypothesized to involve P4-induced down-regulation of PGR in those epithelia, as well as induction of an unknown STAT1-independent signaling pathway. Thus, knowledge of mechanisms whereby IFNT stimulates *CTSL* and *CST3* gene expression in endometrial LE and sGE may elucidate a non-classical signaling pathway for Type I IFNs.

ACKNOWLEDGEMENTS

The authors are appreciative of Drs. Robert C. Burghardt, and Greg A. Johnson for helpful discussions and Mr. Tae-Hyun Kim for illustration. This research was funded by National Institutes of Health Grants 5 R01 HD32534 and 5 P30 ES09106, USA and was also supported by the WCU program (R31-2008-000-10056-0) of the Ministry of Education, Science, and Technology, Korea.

REFERENCES

1. Spencer, T. E. and Bazer, F. W. 2002. Biology of progesterone

- action during pregnancy recognition and maintenance of pregnancy. *Front Biosci* 7:d1879-98.
2. Spencer, T. E., Johnson, G. A., Bazer, F. W. and Burghardt, R. C. 2004. Implantation mechanisms: insights from the sheep. *Reproduction* 128:657-68.
 3. Ashworth, C. J. and Bazer, F. W. 1989. Changes in ovine conceptus and endometrial function following asynchronous embryo transfer or administration of progesterone. *Biol Reprod* 40:425-33.
 4. Farin, C. E., Imakawa, K. and Roberts, R. M. 1989. *In situ* localization of mRNA for the interferon, ovine trophoblast protein-1, during early embryonic development of the sheep. *Mol Endocrinol* 3:1099-107.
 5. Bazer, F. W. 1992. Mediators of maternal recognition of pregnancy in mammals. *Proc Soc Exp Biol Med* 199:373-84.
 6. Roberts, R. M., Ealy, A. D., Alexenko, A. P., Han, C. S. and Ezashi, T. 1999. Trophoblast interferons. *Placenta* 20:259-64.
 7. Spencer, T. E. and Bazer, F. W. 1996. Ovine interferon tau suppresses transcription of the estrogen receptor and oxytocin receptor genes in the ovine endometrium. *Endocrinology* 137:1144-7.
 8. Fleming, J. A., Choi, Y., Johnson, G. A., Spencer, T. E. and Bazer, F. W. 2001. Cloning of the ovine estrogen receptor-alpha promoter and functional regulation by ovine interferon-tau. *Endocrinology* 142:2879-87.
 9. Wathes, D. C. and Hamon, M. 1993. Localization of oestradiol, progesterone and oxytocin receptors in the uterus during the oestrous cycle and early pregnancy of the ewe. *J Endocrinol* 138:479-92.
 10. Spencer, T. E. and Bazer, F. W. 1995. Temporal and spatial alterations in uterine estrogen receptor and progesterone receptor gene expression during the estrous cycle and early pregnancy in the ewe. *Biol Reprod* 53:1527-43.
 11. Hooper, S. B., Watkins, W. B. and Thorburn, G. D. 1986. Oxytocin, oxytocin-associated neurophysin, and prostaglandin F2 alpha concentrations in the utero-ovarian vein of pregnant and nonpregnant sheep. *Endocrinology* 119:2590-7.
 12. Stevenson, K. R., Riley, P. R., Stewart, H. J., Flint, A. P. and Wathes, D. C. 1994. Localization of oxytocin receptor mRNA in the ovine uterus during the oestrous cycle and early pregnancy. *J Mol Endocrinol* 12:93-105.
 13. Lamming, G. E., Wathes, D. C., Flint, A. P., Payne, J. H., Stevenson, K. R. and Vallet, J. L. 1995. Local action of trophoblast interferons in suppression of the development of oxytocin and oestradiol receptors in ovine endometrium. *J Reprod Fertil* 105:165-75.
 14. Spencer, T. E., Becker, W. C., George, P., Mirando, M. A., Ogle, T. F. and Bazer, F. W. 1995. Ovine interferon-tau inhibits estrogen receptor up-regulation and estrogen-induced luteolysis in cyclic ewes. *Endocrinology* 136:4932-44.
 15. Spencer, T. E., Becker, W. C., George, P., Mirando, M. A., Ogle, T. F. and Bazer, F. W. 1995. Ovine interferon-tau regulates expression of endometrial receptors for estrogen and oxytocin but not progesterone. *Biol Reprod* 53:732-45.
 16. Fleming, J. G., Spencer, T. E., Safe, S. H. and Bazer, F. W. 2006. Estrogen regulates transcription of the ovine oxytocin receptor gene through GC-rich SP1 promoter elements. *Endocrinology* 147:899-911.
 17. Spencer, T. E., Johnson, G. A., Burghardt, R. C. and Bazer, F. W. 2004. Progesterone and placental hormone actions on the uterus: insights from domestic animals. *Biol Reprod* 71: 2-10.
 18. Spencer, T. E., Burghardt, R. C., Johnson, G. A. and Bazer, F. W. 2004. Conceptus signals for establishment and maintenance of pregnancy. *Anim Reprod Sci* 82-83:537-50.
 19. Spencer, T. E., Gray, A., Johnson, G. A., Taylor, K. M. and Gertler, A. 1999. Effects of recombinant ovine interferon-tau, placental lactogen, and growth hormone on the ovine uterus. *Biol Reprod* 61:1409-18.
 20. Gray, C. A., Bartol, F. F., Tarleton, B. J., Wiley, A. A. and Johnson, G. A. 2001. Developmental biology of uterine glands. *Biol Reprod* 65:1311-23.
 21. Gray, C. A., Bartol, F. F., Taylor, K. M., Wiley, A. A. and Ramsey, W. S. 2000. Ovine uterine gland knock-out model: effects of gland ablation on the estrous cycle. *Biol Reprod* 62:448-56.
 22. Gray, C. A., Bazer, F. W. and Spencer, T. E. 2001. Effects of neonatal progestin exposure on female reproductive tract structure and function in the adult ewe. *Biol Reprod* 64:797-804.
 23. Gray, C. A., Taylor, K. M., Bazer, F. W. and Spencer, T. E. 2000. Mechanisms regulating norgestomet inhibition of endometrial gland morphogenesis in the neonatal ovine uterus. *Mol Reprod Dev* 57:67-78.
 24. Gray, C. A., Taylor, K. M., Ramsey, W. S., Hill, J. R. and Bazer, F. W. 2001. Endometrial glands are required for preimplantation conceptus elongation and survival. *Biol Reprod* 64:1608-13.
 25. Kimmins, S. and MacLaren, L. A. 2001. Oestrous cycle and pregnancy effects on the distribution of oestrogen and progesterone receptors in bovine endometrium. *Placenta* 22:

- 742-8.
26. Geisert, R. D., Pratt, T. N., Bazer, F. W., Mayes, J. S. and Watson, G. H. 1994. Immunocytochemical localization and changes in endometrial progesterin receptor protein during the porcine oestrous cycle and early pregnancy. *Reprod Fertil Dev* 6:749-60.
 27. Hansen, P. J. 1995. Interactions between the immune system and the ruminant conceptus. *J Reprod Fertil Suppl* 49:69-82.
 28. Tekin, S. and Hansen, P. J. 2002. Natural killer-like cells in the sheep: functional characterization and regulation by pregnancy-associated proteins. *Exp Biol Med (Maywood)* 227: 803-11.
 29. Croy, B. A., Esadeg, S., Chantakru, S., van den Heuvel, M., Paffaro, V. A. 2003. Update on pathways regulating the activation of uterine Natural Killer cells, their interactions with decidual spiral arteries and homing of their precursors to the uterus. *J Reprod Immunol* 59:175-91.
 30. Short, R. V. 1969. Maternal recognition of pregnancy. In: Wolstenholm GEW, O'Conner M, editors. *Foetal Anatomy*. Churchill, London: Wiley, pp. 2-26.
 31. Moor, R. M. and Rowson, L. E. 1964. Influence of the Embryo and Uterus on Luteal Function in the Sheep. *Nature* 201:522-3.
 32. Moor, R. M. and Rowson, L. E. 1966. The corpus luteum of the sheep: effect of the removal of embryos on luteal function. *J Endocrinol* 34:497-502.
 33. Rowson, L. E. and Moor, R. M. 1967. The influence of embryonic tissue homogenate infused into the uterus, on the life-span of the corpus luteum in the sheep. *J Reprod Fertil* 13:511-6.
 34. Heyman, Y., Camous, S., Fevre, J., Meziou, W., Martal, J. 1984. Maintenance of the corpus luteum after uterine transfer of trophoblastic vesicles to cyclic cows and ewes. *J Reprod Fertil* 70:533-40.
 35. Wilson, M. E., Lewis, G. S., Bazer, F. W. 1979. Proteins of ovine blastocyst origin. *Proc Soc Study Reprod, Quebec, Canada*; p. 101A.
 36. Godkin, J. D., Bazer, F. W., Moffatt, J., Sessions, F., Roberts, R. M. 1982. Purification and properties of a major, low molecular weight protein released by the trophoblast of sheep blastocysts at day 13-21. *J Reprod Fertil* 65:141-50.
 37. Godkin, J. D., Bazer, F. W., Roberts, R. M. 1984. Ovine trophoblast protein 1, an early secreted blastocyst protein, binds specifically to uterine endometrium and affects protein synthesis. *Endocrinology* 114:120-30.
 38. Fincher, K. B., Bazer, F. W., Hansen, P. J., Thatcher, W. W., Roberts, R. M. 1986. Proteins secreted by the sheep conceptus suppress induction of uterine prostaglandin F-2 alpha release by oestradiol and oxytocin. *J Reprod Fertil* 76:425-33.
 39. Vallet, J. L., Bazer, F. W., Fliss, M. F., Thatcher, W. W. 1988. Effect of ovine conceptus secretory proteins and purified ovine trophoblast protein-1 on interoestrous interval and plasma concentrations of prostaglandins F-2 alpha and E and of 13,14-dihydro-15-keto prostaglandin F-2 alpha in cyclic ewes. *J Reprod Fertil* 84:493-504.
 40. Ott, T. L., Van Heeke, G., Hostetler, C. E., Schaule, T. K., Olmsted, J. J. 1993. Intrauterine injection of recombinant ovine interferon-tau extends the interoestrous interval in sheep. *Theriogenology* 40:757-69.
 41. Imakawa, K., Anthony, R. V., Kazemi, M., Marotti, K. R., Polites, H. G., Roberts, R. M. 1987. Interferon-like sequence of ovine trophoblast protein secreted by embryonic trophoblast. *Nature* 330:377-9.
 42. Pontzer, C. H., Torres, B. A., Vallet, J. L., Bazer, F. W., Johnson, H. M. 1988. Antiviral activity of the pregnancy recognition hormone ovine trophoblast protein-1. *Biochem Biophys Res Commun* 152:801-7.
 43. Pontzer, C. H., Bazer, F. W. and Johnson, H. M. 1991. Antiproliferative activity of a pregnancy recognition hormone, ovine trophoblast protein-1. *Cancer Res* 51:5304-7.
 44. Roberts, R. M., Imakawa, K., Niwano, Y., Kazemi, M. and Malathy, P. V. 1989. Interferon production by the preimplantation sheep embryo. *J Interferon Res* 9:175-87.
 45. Roberts, R. M., Cross, J. C. and Leaman, D. W. 1992. Interferons as hormones of pregnancy. *Endocr Rev* 13:432-52.
 46. Roberts, R. M., Leaman, D. W. and Cross, J. C. 1992. Role of interferons in maternal recognition of pregnancy in ruminants. *Proc Soc Exp Biol Med* 200:7-18.
 47. Bazer, F. W., Spencer, T. E. and Ott, T. L. 1996. Placental interferons. *Am J Reprod Immunol* 35:297-308.
 48. Hansen, T. R., Imakawa, K., Polites, H. G., Marotti, K. R. and Anthony, R. V., Roberts, R. M. 1988. Interferon RNA of embryonic origin is expressed transiently during early pregnancy in the ewe. *J Biol Chem* 263:12801-4.
 49. Fillion, C., Chaouat, G., Reinaud, P., Charpigny, J. C. and Martal, J. 1991. Immunoregulatory effects of ovine trophoblastin protein (oTP): all five isoforms suppress PHA-induced lymphocyte proliferation. *J Reprod Immunol* 19: 237-49.
 50. Tennakoon, D. K., Smith, R., Stewart, M. D., Spencer, T. E., Nayak, M., Welsh, C. J. 2001. Ovine IFN tau modulates the

- expression of MHC antigens on murine cerebrovascular endothelial cells and inhibits replication of Theiler's virus. *J Interferon Cytokine Res* 21:785-92.
51. Ealy, A. D., Alexenko, A. P., Keisler, D. H., Roberts, R. M. 1998. Loss of the signature six carboxyl amino acid tail from ovine interferon-tau does not affect biological activity. *Biol Reprod* 58:1463-8.
 52. Nephew, K. P., Whaley, A. E., Christenson, R. K., Imakawa, K. 1993. Differential expression of distinct mRNAs for ovine trophoblast protein-1 and related sheep type I interferons. *Biol Reprod* 48:768-78.
 53. Baumbach, G. A., Duby, R. T., Godkin, J. D. 1990. N-glycosylated and unglycosylated forms of caprine trophoblast protein-1 are secreted by preimplantation goat conceptuses. *Biochem Biophys Res Commun* 172:16-21.
 54. Guillomot, M., Michel, C., Gaye, P., Charlier, N., Trojan, J., Martal, J. 1990. Cellular localization of an embryonic interferon, ovine trophoblastin and its mRNA in sheep embryos during early pregnancy. *Biol Cell* 68:205-11.
 55. Spencer, T. E., Ott, T. L., Bazer, F. W. 1996. tau-Interferon: pregnancy recognition signal in ruminants. *Proc Soc Exp Biol Med* 213:215-29.
 56. Novick, D., Cohen, B., Rubinstein, M. 1994. The human interferon alpha/beta receptor: characterization and molecular cloning. *Cell* 77:391-400.
 57. Domanski, P., Yan, H., Witte, M. M., Krolewski, J., Colamonici, O. R. 1995. Homodimerization and intermolecular tyrosine phosphorylation of the Tyk-2 tyrosine kinase. *FEBS Lett* 374:317-22.
 58. Rosenfeld, C. S., Han, C. S., Alexenko, A. P., Spencer, T. E., Roberts, R. M. 2002. Expression of interferon receptor subunits, IFNAR1 and IFNAR2, in the ovine uterus. *Biol Reprod* 67:847-53.
 59. Darnell, J. E. Jr. 1997. STATs and gene regulation. *Science* 277:1630-5.
 60. Darnell, J. E. Jr., Kerr, I. M., Stark, G. R. 1994. Jak STAT pathways and transcriptional activation in response to IFNs and other extracellular signaling proteins. *Science* 264:1415-21.
 61. Stark, G. R., Kerr, I. M., Williams, B. R., Silverman, R. H., Schreiber, R. D. 1998. How cells respond to interferons. *Annu Rev Biochem* 67:227-64.
 62. Gauzzi, M. C., Velazquez, L., McKendry, R., Mogensen, K. E., Fellous, M., Pellegrini, S. 1996. Interferon-alpha-dependent activation of Tyk2 requires phosphorylation of positive regulatory tyrosines by another kinase. *J Biol Chem* 271:20494-500.
 63. Muller, M., Briscoe, J., Laxton, C., Guschin, D., Ziemiecki, A. 1993. The protein tyrosine kinase JAK1 complements defects in interferon-alpha/beta and -gamma signal transduction. *Nature* 366:129-35.
 64. Colamonici, O. R., Yan, H., Domanski, P., Handa, R., Smalley, D. 1994. Direct binding to and tyrosine phosphorylation of the alpha subunit of the type I interferon receptor by p135tyk2 tyrosine kinase. *Mol Cell Biol* 14:8133-42.
 65. Colamonici, O. R., Uyttendaele, H., Domanski, P., Yan, H., Krolewski, J. J. 1994. p135tyk2, an interferon-alpha-activated tyrosine kinase, is physically associated with an interferon-alpha receptor. *J Biol Chem* 269:3518-22.
 66. Yan, H., Krishnan, K., Greenlund, A. C., Gupta, S., Lim, J. T. 1996. Phosphorylated interferon-alpha receptor 1 subunit (IFNAR1) acts as a docking site for the latent form of the 113 kDa STAT2 protein. *Embo J* 15:1064-74.
 67. Silvennoinen, O., Ihle, J. N., Schlessinger, J., Levy, D. E. 1993. Interferon-induced nuclear signalling by Jak protein tyrosine kinases. *Nature* 366:583-5.
 68. Shuai, K., Horvath, C. M., Huang, L. H., Qureshi, S. A., Cowburn, D., Darnell, J. E. Jr. 1994. Interferon activation of the transcription factor Stat91 involves dimerization through SH2-phosphotyrosyl peptide interactions. *Cell* 76:821-8.
 69. Haque, S. J. and Williams, B. R. 1994. Identification and characterization of an interferon (IFN)-stimulated response element-IFN-stimulated gene factor 3-independent signaling pathway for IFN-alpha. *J Biol Chem* 269:19523-9.
 70. Bhattacharya, S., Eckner, R., Grossman, S., Oldread, E., Arany, Z. 1996. Cooperation of Stat2 and p300/CBP in signalling induced by interferon-alpha. *Nature* 383:344-7.
 71. Pine, R., Canova, A., Schindler, C. 1994. Tyrosine phosphorylated p91 binds to a single element in the ISGF2/IRF-1 promoter to mediate induction by IFN alpha and IFN gamma, and is likely to autoregulate the p91 gene. *Embo J* 13:158-67.
 72. Choi, Y., Johnson, G. A., Burghardt, R. C., Berghman, L. R., Joyce, M. M. 2001. Interferon regulatory factor-two restricts expression of interferon-stimulated genes to the endometrial stroma and glandular epithelium of the ovine uterus. *Biol Reprod* 65:1038-49.
 73. Johnson, G. A., Spencer, T. E., Hansen, T. R., Austin, K. J., Burghardt, R. C., Bazer, F.W. 1999. Expression of the interferon tau inducible ubiquitin cross-reactive protein in the ovine uterus. *Biol Reprod* 61:312-8.

74. Choi, Y., Johnson, G. A., Spencer, T. E., Bazer, F. W. 2003. Pregnancy and interferon tau regulate major histocompatibility complex class I and beta2-microglobulin expression in the ovine uterus. *Biol Reprod* 68:1703-10.
75. Johnson, G. A., Spencer, T. E., Burghardt, R. C., Joyce, M. M., Bazer, F. W. 2000. Interferon-tau and progesterone regulate ubiquitin cross-reactive protein expression in the ovine uterus. *Biol Reprod* 62:622-7.
76. Johnson, G. A., Stewart, M. D., Gray, C. A., Choi, Y., Burghardt, R. C. 2001. Effects of the estrous cycle, pregnancy, and interferon-tau on 2',5'-oligoadenylate synthetase expression in the ovine uterus. *Biol Reprod* 64:1392-9.
77. Johnson, G. A., Burghardt, R. C., Newton, G. R., Bazer, F. W., Spencer, T. E. 1999. Development and characterization of immortalized ovine endometrial cell lines. *Biol Reprod* 61:1324-30.
78. Stewart, D. M., Johnson, G. A., Vyhldal, C. A., Burghardt, R. C., Safe, S. H. 2001. Interferon-tau activates multiple signal transducer and activator of transcription proteins and has complex effects on interferon-responsive gene transcription in ovine endometrial epithelial cells. *Endocrinology* 142:98-107.
79. Spencer, T. E., Ott, T. L., Bazer, F. W. 1998. Expression of interferon regulatory factors one and two in the ovine endometrium: effects of pregnancy and ovine interferon tau. *Biol Reprod* 58:1154-62.
80. Stewart, D. M., Johnson, G. A., Vyhldal, C. A., Burghardt, R. C., Safe, S. H. 2001. Interferon-tau activates multiple signal transducer and activator of transcription proteins and has complex effects on interferon-responsive gene transcription in ovine endometrial epithelial cells. *Endocrinology* 142:98-107.
81. Ott, T. L., Yin, J., Wiley, A. A., Kim, H. T., Gerami-Naini, B. 1998. Effects of the estrous cycle and early pregnancy on uterine expression of Mx protein in sheep (*Ovis aries*). *Biol Reprod* 59:784-94.
82. Mirando, M. A., Short, E. C. Jr., Geisert, R. D., Vallet, J. L., Bazer, F. W. 1991. Stimulation of 2',5'-oligoadenylate synthetase activity in sheep endometrium during pregnancy, by intrauterine infusion of ovine trophoblast protein-1, and by intramuscular administration of recombinant bovine interferon-alpha II. *J Reprod Fertil* 93:599-607.
83. Johnson, G. A., Stewart, M. D., Gray, C. A., Choi, Y., Burghardt, R. C. 2001. Effects of the estrous cycle, pregnancy, and interferon-tau on 2',5'-oligoadenylate synthetase expression in the ovine uterus. *Biol Reprod* 64:1392-9.
84. Vallet, J. L., Barker, P. J., Lamming, G. E., Skinner, N., Huskisson, N. S. 1991. A low molecular weight endometrial secretory protein which is increased by ovine trophoblast protein-1 is a beta 2-microglobulin-like protein. *J Endocrinol* 130:R1-4.
85. Han, C. S., Mathialagan, N., Klemann, S. W., Roberts, R. M. 1997. Molecular cloning of ovine and bovine type I interferon receptor subunits from uteri, and endometrial expression of messenger ribonucleic acid for ovine receptors during the estrous cycle and pregnancy. *Endocrinology* 138:4757-67.
86. Stewart, M. D., Johnson, G. A., Bazer, F. W., Spencer, T. E. 2001. Interferon-tau (IFNtau) regulation of IFN-stimulated gene expression in cell lines lacking specific IFN-signaling components. *Endocrinology* 142:1786-94.
87. Decker, T., Lew, D. J., Mirkovitch, J., Darnell, J. E. Jr. 1991. Cytoplasmic activation of GAF, an IFN-gamma-regulated DNA-binding factor. *Embo J* 10:927-32.
88. Schindler, C., Shuai, K., Prezioso, V. R., Darnell, J. E. Jr. 1992. Interferon-dependent tyrosine phosphorylation of a latent cytoplasmic transcription factor. *Science* 257:809-13.
89. Levy, D. E., Kessler, D. S., Pine, R., Reich, N., Darnell, J. E. Jr. 1988. Interferon-induced nuclear factors that bind a shared promoter element correlate with positive and negative transcriptional control. *Genes Dev* 2:383-93.
90. Reich, N., Evans, B., Levy, D., Fahey, D., Knight, E. Jr., Darnell, J. E. Jr. 1987. Interferon-induced transcription of a gene encoding a 15-kDa protein depends on an upstream enhancer element. *Proc Natl Acad Sci USA* 84:6394-8.
91. Kim, S., Choi, Y., Bazer, F. W., Spencer, T. E. 2003. Identification of genes in the ovine endometrium regulated by interferon tau independent of signal transducer and activator of transcription 1. *Endocrinology* 144:5203-14.
92. Gray, C. A., Adelson, D. L., Bazer, F. W., Burghardt, R. C., Meeusen, E. N., Spencer, T. E. 2004. Discovery and characterization of an epithelial-specific galectin in the endometrium that forms crystals in the trophectoderm. *Proc Natl Acad Sci USA* 101:7982-7.
93. Farrell, P. J., Broeze, R. J., Lengyel, P. 1979. Accumulation of an mRNA and protein in interferon-treated Ehrlich ascites tumour cells. *Nature* 279:523-5.
94. Korant, B. D., Blomstrom, D. C., Jonak, G. J., Knight, E. Jr. 1984. Interferon-induced proteins. Purification and characterization of a 15,000-dalton protein from human and bovine cells induced by interferon. *J Biol Chem* 259:14835-9.
95. Haas, A. L., Ahrens, P., Bright, P. M., Ankel, H. 1987.

- Interferon induces a 15-kilodalton protein exhibiting marked homology to ubiquitin. *J Biol Chem* 262:11315-23.
96. Johnson, G. A., Austin, K. J., Collins, A. M., Murdoch, W. J., Hansen, T. R. 1999. Endometrial ISG17 mRNA and a related mRNA are induced by interferon-tau and localized to glandular epithelial and stromal cells from pregnant cows. *Endocrine* 10:243-52.
 97. Austin, K. J., Ward, S. K., Teixeira, M. G., Dean, V. C., Moore, D. W., Hansen, T. R. 1996. Ubiquitin cross-reactive protein is released by the bovine uterus in response to interferon during early pregnancy. *Biol Reprod* 54:600-6.
 98. Johnson, G. A., Burghardt, R. C., Newton, G. R., Bazer, F. W., Spencer, T. E. 1999. Development and characterization of immortalized ovine endometrial cell lines. *Biol Reprod* 61:1324-30.
 99. Samuel, C. E. 1991. Antiviral actions of interferon. Interferon-regulated cellular proteins and their surprisingly selective antiviral activities. *Virology* 183:1-11.
 100. Lengyel, P. 1993. Tumor-suppressor genes: news about the interferon connection. *Proc Natl Acad Sci USA* 90:5893-5.
 101. Salzberg, S., Hyman, T., Turm, H., Kinar, Y., Schwartz, Y. 1997. Ectopic expression of 2-5A synthetase in myeloid cells induces growth arrest and facilitates the appearance of a myeloid differentiation marker. *Cancer Res* 57:2732-40.
 102. Chin, K. C. and Cresswell, P. 2001. Viperin (cig5), an IFN-inducible antiviral protein directly induced by human cytomegalovirus. *Proc Natl Acad Sci U S A* 98:15125-30.
 103. Sun, B. J. and Nie, P. 2004. Molecular cloning of the viperin gene and its promoter region from the mandarin fish *Siniperca chuatsi*. *Vet Immunol Immunopathol* 101:161-70.
 104. Olofsson, P. S., Jatta, K., Wagsater, D., Gredmark, S., Hedin, U. 2005. The antiviral cytomegalovirus inducible gene 5/viperin is expressed in atherosclerosis and regulated by proinflammatory agents. *Arterioscler Thromb Vasc Biol* 25:e113-6.
 105. Helbig, K. J., Lau, D. T., Semendric, L., Harley, H. A., Beard, M. R. 2005. Analysis of ISG expression in chronic hepatitis C identifies viperin as a potential antiviral effector. *Hepatology* 42:702-10.
 106. Kang, D. C., Gopalkrishnan, R. V., Wu, Q., Jankowsky, E., Pyle, A. M., Fisher, P. B. 2002. mda-5: An interferon-inducible putative RNA helicase with double-stranded RNA-dependent ATPase activity and melanoma growth-suppressive properties. *Proc Natl Acad Sci USA* 99:637-42.
 107. Kang, D. C., Gopalkrishnan, R. V., Lin, L., Randolph, A., Valerie, K. 2004. Expression analysis and genomic characterization of human melanoma differentiation associated gene-5, mda-5: a novel type I interferon-responsive apoptosis-inducing gene. *Oncogene* 23:1789-800.
 108. Yoneyama, M., Kikuchi, M., Matsumoto, K., Imaizumi, T., Miyagishi, M. 2005. Shared and unique functions of the DExD/H box helicases RIG-I, MDA5, and LGP2 in antiviral innate immunity. *J Immunol* 175:2851-8.
 109. Andrejeva, J., Childs, K. S., Young, D. F., Carlos, T. S., Stock, N. 2004. The V proteins of paramyxoviruses bind the IFN-inducible RNA helicase, mda-5, and inhibit its activation of the IFN-beta promoter. *Proc Natl Acad Sci U S A* 101:17264-9.
 110. Kawai, T., Takahashi, K., Sato, S., Coban, C., Kumar, H. 2005. IPS-1, an adaptor triggering RIG-I- and Mda5-mediated type I interferon induction. *Nat Immunol* 6:981-8.
 111. Mohamed, O. A., Jonnaert, M., Labelle-Dumais, C., Kuroda, K., Clarke, H. J., Dufort, D. 2005. Uterine Wnt/beta-catenin signaling is required for implantation. *Proc Natl Acad Sci USA* 102:8579-84.
 112. Spencer, T. E., Johnson, G. A., Bazer, F. W., Burghardt, R. C., Palmarini, M. 2007. Pregnancy recognition and conceptus implantation in domestic ruminants: roles of progesterone, interferons and endogenous retroviruses. *Reproduction, Fertility and Development* 19:65-78.
 113. Cooper, D. N. and Barondes, S. H. 1999. God must love galectins; he made so many of them. *Glycobiology* 9:979-84.
 114. Barondes, S. H., Castronovo, V., Cooper, D. N., Cummings, R. D., Drickamer, K. 1994. Galectins: a family of animal beta-galactoside-binding lectins. *Cell* 76:597-8.
 115. Kazemi, M., Amann, J. F., Keisler, D. H., Ing, N. H., Roberts, R. M. 1990. A progesterone-modulated, low-molecular-weight protein from the uterus of the sheep is associated with crystalline inclusion bodies in uterine epithelium and embryonic trophectoderm. *Biol Reprod* 43:80-96.
 116. Gray, C. A., Dunlap, K. A., Burghardt, R. C., Spencer, T. E. 2005. Galectin-15 in ovine uteroplacental tissues. *Reproduction* 130:231-40.
 117. Kirschke, H., Barrett, A. J. and Rawlings, N. D. 1998. *Lysosomal Cysteine Proteases*. Oxford: Oxford University Press
 118. Afonso, S., Romagnano, L., Babiarz, B. 1997. The expression and function of cystatin C and cathepsin B and cathepsin L during mouse embryo implantation and placentation. *Development* 124:3415-25.
 119. Elangovan, S. and Moulton, B. C. 1980. Blastocyst

- implantation in the rat and the immunohistochemical distribution and rate of synthesis of uterine lysosomal cathepsin D. *Biol Reprod* 23:663-8.
120. Li, W. G., Jaffe, R. C. and Verhage, H. G. 1992. Immunocytochemical localization and messenger ribonucleic acid levels of a progesterone-dependent endometrial secretory protein (cathepsin L) in the pregnant cat uterus. *Biol Reprod* 47:21-8.
121. Li, W. G., Jaffe, R. C., Fazleabas, A. T., Verhage, H. G. 1991. Progesterone-dependent cathepsin L proteolytic activity in cat uterine flushings. *Biol Reprod* 44:625-31.
122. Verhage, H. G., Boomsma, R. A., Mavrogianis, P. A., Li, W., Fazleabas, A. T. and Jaffe, R. C. 1989. Immunological characterization and immunocytochemical localization of a progesterone-dependent cat endometrial secretory protein. *Biol Reprod* 41:347-54.
123. Geisert, R. D., Blair, R. M., Pratt, T., Zavy, M. T. 1997. Characterization and proteolytic activity of a cathepsin L-like polypeptide in endometrium and uterine flushings of cycling, pregnant and steroid-treated ovariectomized gilts. *Reprod Fertil Dev* 9:395-402.
124. Roberts, R. M., Bazer, F. W., Baldwin, N., Pollard, W. E. 1976. Progesterone induction of lysozyme and peptidase activities in the porcine uterus. *Arch Biochem Biophys* 177: 499-507.
125. Jokimaa, V., Oksjoki, S., Kujari, H., Vuorio, E. and Anttila, L. 2001. Expression patterns of cathepsins B, H, K, L and S in the human endometrium. *Mol Hum Reprod* 7:73-8.
126. Abrahamson, M., Barrett, A. J., Salvesen, G. and Grubb, A. 1986. Isolation of six cysteine proteinase inhibitors from human urine. Their physicochemical and enzyme kinetic properties and concentrations in biological fluids. *J Biol Chem* 261:11282-9.
127. Abrahamson, M., Olafsson, I., Palsdottir, A., Ulvsback, M., Lundwall, A. 1990. Structure and expression of the human cystatin C gene. *Biochem J* 268:287-94.
128. Hall, A., Hakansson, K., Mason, R. W., Grubb, A. and Abrahamson, M. 1995. Structural basis for the biological specificity of cystatin C. Identification of leucine 9 in the N-terminal binding region as a selectivity-conferring residue in the inhibition of mammalian cysteine peptidases. *J Biol Chem* 270:5115-21.
129. Grubb, A. and Lofberg, H. 1982. Human gamma-trace, a basic microprotein: amino acid sequence and presence in the adenohypophysis. *Proc Natl Acad Sci U S A* 79:3024-7.
130. Uddin, S., Majchrzak, B., Woodson, J., Arunkumar, P., Alsayed, Y. 1999. Activation of the p38 mitogen-activated protein kinase by type I interferons. *J Biol Chem* 274: 30127-31.
131. Katsoulidis, E., Li, Y., Mears, H., Plataniias, L. C. 2005. The p38 mitogen-activated protein kinase pathway in interferon signal transduction. *J Interferon Cytokine Res* 25:749-56.
132. Plataniias, L. C. 2005. Mechanisms of type-I- and type-II-interferon-mediated signalling. *Nat Rev Immunol* 5:375-86.
133. Uddin, S., Lekmine, F., Sharma, N., Majchrzak, B., Mayer, I. 2000. The Rac1/p38 mitogen-activated protein kinase pathway is required for interferon alpha-dependent transcriptional activation but not serine phosphorylation of Stat proteins. *J Biol Chem* 275:27634-40.
134. Doualla-Bell, F. and Koromilas, A. E. 2001. Induction of PG G/H synthase-2 in bovine myometrial cells by interferon-tau requires the activation of the p38 MAPK pathway. *Endocrinology* 142:5107-15.
135. Kaur, S., Uddin, S. and Plataniias, L. C. 2005. The PI3' kinase pathway in interferon signaling. *J Interferon Cytokine Res* 25:780-7.

(Received September 2, 2009; Revised September 23, 2009;

Accepted October 6, 2009)