

# Age-dependent extreme event exposure

Wim Thiery<sup>1,2</sup>, Stefan Lange<sup>3</sup>, Joeri Rogelj<sup>4,5</sup>, Carl-Friedrich Schleussner<sup>6,7</sup>,  
Lukas Gudmundsson<sup>2</sup>, Sonia I. Seneviratne<sup>2</sup>, Marina Andrijevic<sup>6,7</sup>, Katja  
Frieler<sup>3</sup>, Kerry Emanuel<sup>8</sup>, Tobias Geiger<sup>3,9</sup>, David N. Bresch<sup>10,11</sup>, Fang  
Zhao<sup>12,3</sup>, Sven N. Willner<sup>3</sup>, Matthias Büchner<sup>3</sup>, Jan Volkholz<sup>3</sup>, Nico Bauer<sup>3</sup>,  
Jinfeng Chang<sup>13,14</sup>, Philippe Ciais<sup>14</sup>, Marie Dury<sup>15</sup>, Louis François<sup>15</sup>, Manolis  
Grillakis<sup>16</sup>, Simon N. Gosling<sup>17</sup>, Naota Hanasaki<sup>18</sup>, Thomas Hickler<sup>19,20</sup>,  
Veronika Huber<sup>21</sup>, Akihiko Ito<sup>18</sup>, Jonas Jägermeyr<sup>22,23,3</sup>, Nikolay Khabarov<sup>5</sup>,  
Aristeidis Koutroulis<sup>24</sup>, Wenfeng Liu<sup>14,25</sup>, Wolfgang Lutz<sup>26,5</sup>, Matthias  
Mengel<sup>3</sup>, Christoph Müller<sup>3</sup>, Sebastian Ostberg<sup>3</sup>, Christopher P. O. Reyer<sup>3</sup>,  
Tobias Stacke<sup>27</sup>, Yoshihide Wada<sup>5</sup>

<sup>1</sup>Vrije Universiteit Brussel, Department of Hydrology and Hydraulic Engineering, Brussels, Belgium

<sup>2</sup>ETH Zurich, Institute for Atmospheric and Climate Science, Zurich, Switzerland

<sup>3</sup>Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, Potsdam,  
Germany

<sup>4</sup>Imperial College London, Grantham Institute for Climate Change and the Environment, London, UK

<sup>5</sup>International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

<sup>6</sup>Climate Analytics, Berlin, Germany

<sup>7</sup>Humboldt University, Integrative Research Institute on Transformations of Human-Environment  
Systems, Berlin, Germany

<sup>8</sup>Massachusetts Institute of Technology, Lorenz Center, Cambridge, MA, USA

<sup>9</sup>Deutscher Wetterdienst (DWD), Climate and Environment Consultancy, Stahnsdorf, Germany

<sup>10</sup>ETH Zurich, Institute for Environmental Decisions, Zurich, Switzerland

<sup>11</sup>Federal Office of Meteorology and Climatology MeteoSwiss, Zurich, Switzerland

<sup>12</sup>East China Normal University, School of Geographic Sciences, Shanghai, China

<sup>13</sup>Zhejiang University, College of Environmental and Resource Sciences, Hangzhou, China

<sup>14</sup>Institut Pierre Simon Laplace, Laboratoire des Sciences du Climat et de l'Environnement, CEA  
CNRS UVSQ, Gif sur Yvette, France

<sup>15</sup>University of Liège, Unit for Modelling of Climate and Biogeochemical Cycles, Liège, Belgium

<sup>16</sup>Foundation for Research and Technology Hellas, Institute for Mediterranean Studies, Rethimno,  
Greece

<sup>17</sup>University of Nottingham, School of Geography, Nottingham, UK

<sup>18</sup>National Institute for Environmental Studies, Tsukuba, Japan

<sup>19</sup>Goethe University Frankfurt, Institute of Physical Geography, Frankfurt, Germany

<sup>20</sup>Senckenberg Biodiversity and Climate Research Centre (SBIK-F), Frankfurt, Germany

<sup>21</sup>Universidad Pablo de Olavide, Department of Physical, Chemical and Natural Systems, Sevilla, Spain

<sup>22</sup>NASA Goddard Institute for Space Studies, New York, NY, USA

<sup>23</sup>Columbia University, Center for Climate Systems Research, New York, NY, USA

<sup>24</sup>Technical University of Crete, School of Environmental Engineering, Chania, Greece

40 <sup>25</sup>China Agricultural University, College of Water Resources and Civil Engineering, Beijing, China

<sup>26</sup>University of Vienna, Wittgenstein Centre for Demography and Global Human Capital, Vienna, Austria

<sup>27</sup>Helmholtz-Zentrum Geesthacht, Institute of Coastal Research, Geesthacht, Germany

45 **With the emergence of a global climate youth movement, questions of inter-**  
**generational justice regarding climate change mitigation have come to the**  
**fore. However, a scientific perspective on intergenerational climate impacts**  
**is still lacking. Here we show that newborns in 2020 are projected to expe-**  
**rience 2–7 times more extreme events globally under current climate pledges**  
**than someone born in 1960, using a novel framework that quantifies impacts**  
50 **as they are experienced along a person’s lifetime. Limiting warming to 1.5 °C**  
**consistently reduces that burden while still leaving younger generations with**  
**unavoidable impacts that are unmatched by those experienced by older gener-**  
**ations. Our results provide a scientific basis to understand the position from**  
**which younger generations challenge the present shortfall of adequate climate**  
55 **action.**

Young people around the world have been leading climate demonstrations since late 2018. This surge in climate protests has received explicit support from many climate scientists (1) and has been accompanied by the emergence of climate change litigation. Meanwhile, government actions are falling short of achieving the emission reductions required to halt global warming  
60 at the safe levels agreed upon under the UN Paris Agreement (2). This failure to adequately act implies that global warming could easily exceed 3 °C by the end of the century (2), and projec-

tions of future climate change impacts under such scenarios have far-reaching implications (3). This situation raises important questions about solidarity and fairness across generations (1, 4).

Under continued global warming, extreme events such as heatwaves will continue to rise in  
65 frequency, intensity, duration and spatial extent over the next decades (5–8). Current younger generations that will experience those future decades are therefore expected to face more impacts compared to their (grand)parents. However, the ruling paradigm to study climate change impacts assesses change in discrete time windows or at discrete levels of warming (3). Such an approach, which we term the Eulerian perspective (in demography called period perspective),  
70 inhibits quantification of exactly how much more impacts from climate change a particular birth cohort will experience compared to another generation.

Meteorological extremes, hazards, or climate change impacts are so far mostly studied as they evolve over time under varying emission scenarios and socio-economic pathways (6, 8, 9). For instance, applying a heatwave indicator (10) (table S1) to four bias-adjusted global climate  
75 models indicates that the land area annually affected by such heatwaves will increase from ~15 % around 2020 to ~22 % by 2100 under a 1.5 °C-compatible scenario and to ~46 % under a scenario in line with current emission reduction pledges (fig. 1a). Recent studies extended this approach by studying aspects of climate change as a function of global mean temperature (GMT) anomalies, highlighting the scenario-independence of several extreme event indicators  
80 (5, 7, 9) but remaining, in essence, a comparison of two time windows.

Instead, we here take a Lagrangian (or cohort) perspective to climate change impacts that measures extreme event exposure over the course of a person’s lifetime. Our intention is thereby to quantify, in the most robust way possible, the changes in lifetime exposure to climate extremes across generations. To this end, we perform a birth cohort analysis by combining an un-  
85 precedented collection of multi-model impact projections (7) with country-scale life expectancy information (11), gridded population data (12), and future global temperature trajectories (13)

from the Intergovernmental Panel on Climate Change (IPCC) Special Report on Global Warming of 1.5 °C (see Methods). By integrating the exposure of an average person in a country or region to extreme events across their lifetime, we encapsulate spatio-temporal changes in climate hazards, population density, cohort size, and life expectancy (fig. 1).

**Extreme event exposure** Our results allow for comparing lifetime exposure to climate extremes across birth cohorts globally. For example, a person born in 1960 will on average experience around  $4 \pm 2$  ( $1 \sigma$ ) heatwaves across their lifetime according to our extreme heatwave definition (fig. 1b). The lifetime heatwave exposure of this cohort is largely insensitive to the three future temperature scenarios considered here. A child born in 2020 will, in contrast, experience  $30 \pm 9$  heatwaves under a scenario following current climate pledges, which could be reduced to  $22 \pm 7$  heatwaves if warming is limited to 2 °C or  $18 \pm 8$  heatwaves if it is limited to 1.5 °C. In any case, that is 7, 6, or 4 times more compared to a person born in 1960. Repeating this analysis for all cohorts born between 1960 and 2020 highlights clear differences in lifetime exposure to heatwaves between older and younger cohorts globally (fig. 1c). The effect of alternative future temperature trajectories on the lifetime exposure multiplication factor becomes discernible only for cohorts younger than 40 years in 2020, with the largest differences for the youngest cohorts.

The previous example only uses one impact indicator and a subset of all possible future temperature pathways. We now expand this approach and consider six extreme event categories: wildfires, crop failures, droughts, river floods, heatwaves, and tropical cyclones (table S1), which we analyse under a wide range of temperature pathways that result in future warming ranging from constant present-day levels up to 3.5 °C by 2100 (see Methods and fig. S1). To this end, we generated a total of 273 global-scale projections with 15 impact models forced by four bias-adjusted global climate models (table S2). Inspired by the IPCC's Reasons for Con-

cern Framework (3), we visualise the exposure multiplication factors relative to a hypothetical reference person living under pre-industrial climate conditions as a function of the 2100 GMT anomaly and cohort (fig. 2). Life expectancy varies with the cohort, whereas the hypothetical reference person is given the same life expectancy as the oldest cohort in our figures. Therefore, 115 in contrast to the previous comparison of lifetime exposure across generations given historical and climate conditions (Fig. 1), we from now on assess how projected lifetime exposure of birth cohorts is affected by climate change since pre-industrial and by increased life expectancy since 1960.

Our results highlight that lifetime exposure to each of the considered extreme events consistently increases for higher warming levels and younger cohorts. Changes in extreme event 120 frequencies have had relatively little effect on lifetime exposure for cohorts above age 55 in 2020, but this rapidly changes for younger cohorts as they start experiencing extreme events in the coming years and decades (fig. 2). For a 3 °C global warming pathway, a 6-year old in 2020 will experience twice as many wildfires and tropical cyclones, 3 times more river floods, 125 4 times more crop failures, 5 times more droughts, and 36 times more heatwaves relative to the reference person. Such conditions clearly pose a severe threat to the safety of young cohorts. While qualitatively consistent, quantitative exposure changes differ among categories: for wildfires and tropical cyclones, increases in exposure remain limited relative to the other categories, whereas heatwave exposure increases much more strongly, up to a factor 44 for newborns under 130 3.5 °C of global warming. Aggregating the exposure multiplication factors across the six categories shows that people younger than 10 in 2020 will experience about a fourfold increase in extreme events under 1.5 °C of global warming, an increase that older cohorts will never experience, even if a scenario towards 3.5 °C warming is followed (fig. S2a). Under a 3 °C global warming pathway, children under 8 will face an almost fivefold increase in extreme event exposure. These exposure multiplication factors scale robustly with the warming pathway and cohort 135

across a range of aggregation methods, despite some variation in the factor values (fig. S2).

We then calculate the probability of each person's lifetime exposure occurring under pre-industrial climate conditions. Lives with an accumulated exposure that would occur with less than 0.01 % probability under pre-industrial climate (that is, with less than a 1-in-10 000 chance) are thereby classified as unprecedented. We find that cohorts above age 55 in 2020 will on average live an unprecedented life only for heatwaves and crop failures, while cohorts aged 0-40 in 2020 will additionally face unprecedented exposure to droughts and flooding above 1.5 °C warming (fig. 2a-f). Aggregated across all the impact categories, lifetime exposure to extremes is unprecedented at all warming levels and cohorts (fig. S2a).

**Regional patterns** Behind this global average picture, there are important spatial variations. Repeating the analysis for a selection of world regions (fig. S3) reveals marked differences between regions (figs. S4–S5), while a country-level assessment highlights even stronger spatial disparities (Supplementary Note 1; figs. S6–7). We find a particularly strong increase in lifetime exposure across the Middle East and North Africa, with on average at least 7 times higher exposure for all cohorts younger than 25 years in 2020 under current pledges (fig. S8a). In sub-Saharan Africa, newborns will on average experience 5.9 times more extreme events compared to a reference person living under pre-industrial climate, while newborns in other regions will on average experience 3.7–5.3 times more extremes. This burden on newborns in terms of additional exposure to extreme events is substantially reduced when limiting global warming to 1.5 °C: the strongest reductions in exposure are found in the Middle East and North Africa (-39%), Europe and Central Asia (-28%), and North America (-26%), while benefits in Sub-Saharan Africa, East Asia and the Pacific roughly correspond to the global average (-24%).

Grouping countries by income category instead of by region highlights that young generations in low-income countries will face by far the strongest increases in lifetime exposure

160 with a more than fivefold increase for newborns under current pledges (fig. S8b). High-income  
countries, on the other hand, face the smallest increases for younger cohorts and the smallest  
variation across generations. However, while 60-year old people in high-income countries rep-  
resent 22% of their cohort globally, this fraction has reduced to 10% for newborns (fig. S9-10).  
The corresponding relative cohort sizes of low-income countries, on the contrary, increased  
165 from 5% to 18%. Thus, children born in the present and future are much more likely to be born  
in regions facing the highest increase in lifetime extreme event exposure. For example, 53 mil-  
lion children born in Europe and Central Asia between 2016 and 2020 will experience 3.8–4.0  
times more extreme events under current pledges, but 172 million children of the same age in  
sub-Saharan Africa face a factor 5.5–5.9 increase in lifetime extreme event exposure, including  
170 a factor 50–54 increase in lifetime heatwave exposure (fig. S9-10). This combined rapid growth  
in cohort size and extreme event exposure (fig. S9-12) highlights a disproportionate climate  
change burden for young generations in the Global South.

Improvements in life expectancy (fig. 1a; fig. S13) represent a confounding factor in the  
signal of increasing exposure to extreme events over a person’s lifetime. However, we find  
175 that globally, climate change explains 98% of a newborn’s exposure change under the current  
pledges scenario (fig. S14, see Methods). In high-income countries, the enhanced exposure  
of a newborn is almost entirely attributable to climate change (99%), whereas in low, lower-  
middle and upper-middle income countries, climate change contributes 98% of the total expo-  
sure change (figs. S14–S15).

180 **Discussion** Analysis of climate risks to humans has traditionally considered impacts as they  
evolve over time. Our Lagrangian approach provides a more intuitive account of climate im-  
pacts by tracking exposure to extreme events across a person’s lifetime and comparing changes  
across generations. While we comprehensively account for hazards and exposure using an

unprecedented climate impact modeling effort (7, 12) and within-country population density  
185 variability, there are multiple reasons to think that our approach, if anything, underestimates  
intergenerational differences in exposure (Supplementary Note 2).

Further work could aim at extending this novel approach to include further demographic  
dimensions and vulnerability (14). Vulnerability to extreme events depends on a range of socio-  
economic factors but may also evolve over the course of a lifetime. A young person may for  
190 instance experience little health impacts from a heatwave compared to older people (15), but  
schooling infrastructure destroyed by a tropical cyclone may have a disproportionate detrimen-  
tal effect on children's education which could persist throughout their entire lifetime. And while  
communities may possibly become less vulnerable to extreme events over time, limits to adap-  
tive capacity remain even under optimistic pathways beyond mid-century (14) (Supplementary  
195 Note 3).

Climate change impacts may also engender migration and ultimately even affect life ex-  
pectancy via increased mortality, two aspects which are not considered in this study. Even  
though climate change increases mortality (15), it is currently not included in life expectancy  
estimates and population projections like the ones we use here. Likewise, migration triggered  
200 by environmental degradation may change both exposure and vulnerability to extreme events.  
Further analysis should therefore aim at systematically integrating population dynamics and  
climate risk assessments to better understand the long-term impacts of extreme climate events  
and to improve socioeconomic scenario development.

Our results overall highlight the strong benefits of aligning policies with the Paris Agree-  
205 ment for safeguarding the future of current young generations. While all generations younger  
than 60 in 2020 will live unprecedented lives in terms of extreme event exposure, lifetime expo-  
sure to climate change impacts drastically increases for younger generations as global warming  
progresses. Heatwaves dominate the escalating exposure to hazards, but a consistent rise is



also observed for droughts, wildfires, tropical cyclones, crop failures as well as flooding. The  
210 strongest increases occur in low income countries where strongly rising extreme events (8) af-  
fect a rapidly growing young population. Overall, limiting global warming to 1.5 °C instead  
of following the current pledges scenario nearly halves (-40%) the additional exposure of new-  
borns to extreme heatwaves and substantially reduces the burden for wildfires (-11%), crop  
failures (-27%), droughts (-28%), tropical cyclones (-29%), and river floods (-34%). These  
215 findings have direct implications for climate litigation and call for ambitious mitigation efforts  
to improve intergenerational and international justice.

## References

1. G. Hagedorn, *et al.*, *Science* **364**, 139.2 (2019).
2. J. Rogelj, *et al.*, *Nature* **534**, 631 (2016).
- 220 3. IPCC, *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, C. Field, *et al.*, eds. (2014), pp. 1–32.
4. J. Hansen, *et al.*, *Earth System Dynamics* **8**, 577 (2017).
5. R. A. Betts, *et al.*, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **376**, 20160452 (2018).
- 225 6. J. Sillmann, V. V. Kharin, F. W. Zwiers, X. Zhang, D. Bronaugh, *Journal of Geophysical Research: Atmospheres* **118**, 2473 (2013).
7. S. Lange, *et al.*, *Earth's Future* **8**, 1 (2020).
8. L. J. Harrington, *et al.*, *Environmental Research Letters* **11**, 055007 (2016).
- 230 9. N. W. Arnell, *et al.*, *Environmental Research Letters* **14** (2019).
10. S. Russo, J. Sillmann, E. M. Fischer, *Environmental Research Letters* **10**, 124003 (2015).
11. United Nations Department of Economic and Social Affairs Population Division, *World Population Prospects 2019* (2019).
12. K. Frieler, *et al.*, *Geoscientific Model Development* **10**, 4321 (2017).
- 235 13. J. Rogelj, *et al.*, *Mitigation Pathways Compatible with 1.5C in the Context of Sustainable Development. In: Global Warming of 1.5C. An IPCC Special Report on the impacts*

*of global warming of 1.5C above pre-industrial levels and related global greenhouse gas emission pathw*, V. Masson-Delmotte, *et al.*, eds. (2018), pp. 93–174.

14. M. Andrijevic, J. Crespo Cuaresma, R. Muttarak, C. F. Schleussner, *Nature Sustainability* **3**, 35 (2020).

240

15. N. Watts, *et al.*, *The Lancet* **394**, 1836 (2019).


**Acknowledgements** We greatly thank Iliusi Vega, Lila Warszawski, Ted Veldkamp, Christian Folberth, Alexandra Henrot, Dieter Gerten, Sibyll Schaphoff, Hannes Müller Schmied, Kazuya Nishina, Jörg Steinkamp, Matthew Forrest, Hong Yang and Chao Yue for running impact simulations for ISIMIP2b, for coordinating an ISIMIP sector, and/or for developing and interpreting the data repository. W.T. thanks Jan Steyaert and Veerle Leroy for sparking the idea that led to this study. Urs Beyerle and the ETH Zurich cluster team are thanked for support with the CLM4.5 simulations. We also thank the National Center for Atmospheric Research (NCAR) for maintaining CLM and making the source code publicly available. T.G. and S.W. acknowledge support by the German Federal Ministry of Education and Research (BMBF) under the research projects SLICE (FKZ: 01LA1829A) and CLIC (FKZ: 01LA1817C), respectively. C.-F.S. acknowledges support by the German Federal Ministry of Education and Research (FKZ: 01LS1905A). N.H. was supported by Environment Research and Technology Development Fund (JPMEERF20182R02). S.I.S. acknowledges partial support from the European Research Council through the ERC Proof-of-concept grant MESMER-X under H2020-EU.1.1 (Grant number: 964013). V.H. received support from the Spanish Ministry of Economy, Industry and Competitiveness (Grant ID: PCIN-2017-046). C.P.O.R. acknowledges funding from the EU Horizon 2020 research and innovation program under grant agreement number 821010 (CASCADES).

**Author Contributions** W.T. coordinated and led the study. W.T., J.R., C.-F.S., L.G. and S.I.S. designed the analysis. S.L. processed the raw ISIMIP2b simulation output to generate the exposed land fraction data with help from T.G., D.N.B., J.V., and M.B.. K.E. generated the tropical cyclone tracks. F.Z. and S.W. ran the CaMa-Flood simulations and flood processing chain for generating the flood maps. W.T. performed all further analyses and wrote the manuscript with help from J.R., C.-F.S., S.I.S., S.N.G., and S.L.. K.F., M.A., M.M., N.B. and W.L. contributed

further ideas to the data analysis. W.T., J.C., M.D., L.F., N.H., T.H., A.I., N.K., W.L., C.M., S.O., T.S., M.G., A.K., and Y.W. performed ISIMIP2b simulations, a simulation round coordinated by K.F. and M.M. with W.T., P.C., S.N.G., V.H., J.J., T.H., C.M., and C.P.O.R. as sector coordinators. All authors read and approved the manuscript.

270 **Data Availability** All materials that have contributed to the reported results are available upon request, including the postprocessed ISIMIP2b data. Correspondence and requests for materials should be addressed to W.T. (wim.thiery@vub.be). The ISIMIP2b impact simulations and gridded population data are available at <https://esg.pik-potsdam.de/search/isimip/>, the life expectancy data at <https://population.un.org/wpp/Download/Standard/Mortality/>, the cohort size data at <http://dataexplorer.wittgensteincentre.org/wcde-v2/>, and the IPCC SR1.5 Scenario data at <https://data.ene.iiasa.ac.at/iamc-1.5c-explorer>.

**Code Availability** All codes used for the analyses are available through the github repository of the Department of Hydrology and Hydraulic Engineering at VUB ([https://github.com/VUB-HYDR/2020\\_Thiery\\_etal](https://github.com/VUB-HYDR/2020_Thiery_etal)).



**Figure 1: From an Eulerian (period) to a Lagrangian (cohort) perspective on climate change impacts.** (a) Historical and future evolution of global land area fraction annually exposed to heatwaves under a 1.5 °C, 2 °C, and current pledges scenario. Each line represents the multi-model mean of a heatwave metric calculated from the four bias-adjusted global climate models available in ISIMIP2b (see Methods and table S1). All lines were smoothed using a 10-point moving average and the uncertainty band spans 1 standard deviation across the model ensemble. Horizontal grey arrows indicate the global-average lifespan of a person born in 1960 and 2020, respectively. (b) Lifetime heatwave exposure for the 1960 and 2020 birth cohort, respectively, under the three scenarios. The rotated factors above each bar indicate the exposure multiplication factor relative to the 1960 birth cohort under the respective scenario. (c) Exposure multiplication factors for lifetime heatwave exposure across birth cohorts relative to the exposure of the 1960 birth cohort under the respective scenario. Uncertainty bands represent model uncertainty as the inter-quartile range for the 2020 birth cohort exposure relative to the multi-model mean exposure of the 1960 birth cohort.




Figure 2: **lifetime exposure on the rise.** Exposure multiplication factors across birth cohorts (x-axis) under a range of global warming trajectories (fig. S1) reaching 0.87 °C to 3.5 °C global mean temperature (GMT) anomalies in 2100 relative to the pre-industrial (PI) reference period (1850–1900; y-axis) for (a) wildfires, (b) crop failures, (c) droughts, (d) river floods, (e) heatwaves, and (f) tropical cyclones. All factors are computed relative to the mean exposure of a hypothetical reference person living under pre-industrial climate conditions with year-1960 life expectancy, and therefore incorporate the effect of historical as well as projected climate change. The grey contour delineates lifetime extreme event exposure with 0.01 % probability of occurrence under pre-industrial climate conditions; absence of the contour indicates that this probability is lower for all cases covered. Note the different colour bar range for heatwaves.

## **Supplementary materials**

Materials and Methods

Supplementary Notes

Figs. S1 to S14

<sup>285</sup> Tables S1 to S2

References (*16-96*)



# Supplementary Materials for Age-dependent extreme event exposure

Thiery et al.

## Materials and Methods

5 The aim of this study is to quantify, in the most robust way possible, the global- and regional-scale increase in exposure to climate extremes for younger generations. To this end, we integrate the exposure of an average member of a birth cohort to extreme events across their lifetime and subsequently compare these results across cohorts and regions. This is achieved by combining five sources of data, which are explained hereafter.

10 **Employed data** First, we developed the largest multi-model climate impact projections database available to date as part of the Inter-Sectoral Impact Model Intercomparison Project phase 2b (ISIMIP2b) (16). Following the ISIMIP2b protocol, we performed simulations with 15 impact models that represent variables relevant for the six extreme event categories described below (CARAIB (17), LPJ-GUESS (18), LPJmL (19, 20), ORCHIDEE (21), VISIT (22, 23),  
15 GEPIC (24), PEPIC (25, 26), CLM4.5 (27, 28), H08 (29), JULES-W1 (30), MPI-HM (31, 32), PCR-GLOBWB (33, 34), WaterGAP2 (35, 36), HWMIId-humidex (37), and KE-TG (38)). These process-based models represent the state of the art of global-scale hydrological, vegetation, agricultural, land surface, heat stress, and tropical cyclone modeling (39–45). Each model provides relevant biophysical impact variables such as runoff, crop yields, or soil moisture at a  
20 spatial resolution of  $0.5^\circ \times 0.5^\circ$  and at daily to annual time scales. The impact models are

each driven by up to four downscaled and bias-adjusted (46) global climate models (GCMs; GFDL-ESM2M (47), HadGEM2-ES (48), IPSL-CM5A-LR (49), and MIROC5 (50)) participating in the fifth phase (51) of the Coupled Model Intercomparison Project (CMIP5) under pre-industrial, historical, and RCP 2.6, 6.0 and 8.5 climate forcings. GCMs are designed to capture the spatially-explicit climate response to rising greenhouse gas concentrations and other anthropogenic forcings; as such, they are the cornerstone of scientific knowledge on future climate change. Here, the four GCMs were selected based on data availability and representativeness of the entire CMIP5 ensemble (see Supplementary Note 4 for a detailed description of the selection, downscaling, and bias adjustment procedure). Besides transient historical and future climate information, our simulations represent other human influences via input data mimicking historical socioeconomic development until 2005 and assuming fixed year-2005 conditions thereafter. Overall, our impact data set consists of 273 global-scale impact projections spanning the period 1861–2099 (table S2) and 101 pre-industrial control simulations covering on average 542 years each. A more detailed description of the data is provided in (37).

Second, we employ data on life expectancy at the age of 5 available from the United Nations World Population Prospects (52), indicating the number of years a 5-year old would be expected to live if mortality patterns prevailing at the time of observation (year to which this period indicator pertains) were to remain constant throughout their lifetimes. The data is available at the country, regional, and global scale (fig. S3) in 5-year blocks for 1950-1955 to 2015-2020. From this data, we calculate cohort life expectancy at birth which adjusts for child mortality 0-4 because child mortality distorts the pattern that shall be studied. First, we translate the data to annual values using linear interpolation, assume the life expectancy value is representative for the middle year of the 5-year block, and linearly extrapolate life expectancy from 2017 to 2025 in every country. To capture the entire length of the life span starting with birth, we subsequently add 5 years and assign the value to the birth year of the respective cohort, thereby assuming that

this value represents life expectancy at birth excluding child mortality. Finally, we translate this indicator from period to cohort life expectancy by adding 6 years to the value of the period life expectancy estimate at birth (53). This is based on a rather conservative assumption of future increases in life expectancy given the current uncertainty about future mortality trends. We note that that this life expectancy data ignores impacts from climate change. Our approach thus follows UN fertility and mortality projections, but omits climate change feedbacks on population dynamics.

Third, future GMT trajectories are derived from scenarios compiled in support of the IPCC Special Report on Global Warming of 1.5 °C (SR1.5) (54, 55) and were subsequently made available through the Integrated Assessment Modeling Consortium and the International Institute for Applied Systems Analysis (56, 57). We select three marker scenarios: two scenarios limiting global warming to 1.5 °C and 2.0 °C above pre-industrial levels (58), respectively, and a third scenario consistent with current (2020) Nationally Determined Contributions (NDCs) – also referred to as the current pledges scenario (fig. S1). The 1.5 °C and NDC scenarios (originally labelled ‘MESSAGEix-GLOBIOM 1.0\_LowEnergyDemand’ and ‘MESSAGE-GLOBIOM 1.0\_ADVANCE\_INDC’, respectively) were developed with the Integrated Assessment Model MESSAGE-GLOBIOM version 1.0 (59). The 2.0 °C scenario was derived using IMAGE version 3.0.1 (60). The 2 °C compatible scenario is assessed by the IPCC SR1.5 to keep warming below 2 °C with at least 66 % probability, whereas the 1.5 °C scenario limits warming to 1.5 °C with 50 % probability but potentially exceeds this level temporarily by less than 0.1 °C (54). The GMT anomalies in 2091–2100 compared to the pre-industrial reference period (1850–1900) are 1.4 °C, 1.7 °C and 2.4 °C for the 1.5 °C, the 2.0 °C and the current pledges scenario, respectively (fig. S1).

Fourth, gridded population reconstructions and projections are obtained from the ISIMIP2b input data repository. Historical reconstructions are based on version 3.2 of the History Database

of the Global Environment (HYDE3.2) (61) while future projections are derived from a gravity-based downscaling model (62) under the Middle-of-the-Road Shared Socioeconomic Pathway 2 (SSP2) (63). Social and economic trends of the SSP2 scenario do not markedly shift from the historical trends. Most countries complete the demographic transition, and the population  
75 growth levels off in the second half of the century. Under SSP2, all countries in the world are projected to develop with medium fertility, mortality and migration trends (64). The gridded future population projections account for population growth, urbanization level, and spatial urbanization pattern by incorporating variations of these patterns across regions and SSPs (62). For population change, countries are categorized according to fertility and income into three  
80 groups (high fertility, low fertility with high incomes, and low fertility), whereas for urbanization, countries are grouped based on income alone (Low, Medium and High income). While population density evolves over time according to these drivers, climate-induced changes in migration, urbanisation, fertility, and mortality are not considered in this data set, and therefore not in our approach. We analysed the uncertainty associated with the gridded population data by  
85 testing the sensitivity of our results to using gridded population projections under an alternative SSP. Using SSP3 – a pathway considered inconsistent with RCP2.6 – instead of SSP2 showed little sensitivity of the results to the SSP choice, reflecting the fact that our analysis builds on within-country relative population density variability rather than on absolute population totals.

Finally, we use country-scale cohort size data provided by the Wittgenstein Centre and avail-  
90 able through its Human Capital Data Explorer (65). We consider cohort sizes of the year 2020 and linearly interpolate the 5-year block data to annual time scale, assuming the cohort size value to be valid for the center year of the block.

**Extreme event definition.** A detailed description of the processing of the ISIMIP2b simulations is provided in Ref. (37) and summarised hereafter. We consider 6 extreme event categories:

95 wildfires, crop failure, droughts, river floods, heatwaves, and tropical cyclones. We select these six extreme event categories because we know from existing studies (16, 66) that these hazards (i) will increase in frequency, intensity, and/or duration with projected climate change, (ii) can lead to strong impacts when they occur, and (iii) can be tackled comprehensively in a modelling framework such as ISIMIP. For each category and simulated calendar year, we compute the land  
100 fraction per grid cell exposed at least once to an extreme event as defined in table S1. As such, the land fraction annually exposed to extreme events becomes a comparable quantity across event categories.

Since the 8 considered global hydrological models do not provide flood extent and apply different routing schemes (67), we employ the global-scale river routing model CaMa-Flood (68)  
105 to compute the land area exposed to river flooding using daily gridded runoff from the global hydrological models as input (table S2). For tropical cyclones, we use the average exposure over a 100-member ensemble of tropical cyclone tracks downscaled from GCM output (38). While the strongest increases in tropical cyclone impacts are expected from increasing cyclone intensities (38, 69), our projections also show a rise in tropical cyclone frequency under con-  
110 tinued warming, consistent with (37, 38). For wildfires, we quantify the pure climate change effect on burned area using a suite of global vegetation models (see Supplementary Note 5 for more details). A grid cell is considered to be exposed to a heatwave in a given year if the Heat Wave Magnitude Index daily (HWMId) (70, 71) of that year exceeds the 99<sup>th</sup> percentile of the HWMId distribution under pre-industrial climate conditions of that grid cell. For droughts,  
115 heatwaves, and crop failure, we define the extreme event occurrence based on the exceedance of a pre-industrial percentile threshold (table S1). While the exact percentile value is an arbitrary choice, the approach allows for a robust estimation of the threshold values thanks to the long time span of the pre-industrial control simulations. Moreover, a sensitivity analysis with multiple heatwave definitions showed only little sensitivity of the relative exposure changes at the

120 global scale, suggesting a limited influence of the choice of the pre-industrial percentile value on the analysis of the historical and future impact model simulations. Finally, we also analyse lifetime cold spell exposure, which we define as the counterpart of heatwaves (see Supplementary Note 6).

**Exposure calculation.** In this study, we integrate the exposure of an average person in any 125 country or region to climate impacts across their lifetime. This cohort analysis considers land areas only and is performed across 178 countries (fig. S6), 11 regions (fig. S3) and the globe.

We first compute country-scale spatial averages of annual land area exposure weighted by population totals of the corresponding year over all available historical, RCP 2.6, 6.0 and 8.5 simulations. This way, our exposure assessment accounts for temporal changes in population 130 density under a Middle-of-the-Road scenario. The resulting time series is then mapped onto the SR1.5 scenarios (1.5 °C, 2 °C and NDCs, respectively) by selecting from the concatenated historical-RCP series the year with the GMT anomaly closest to the annual anomaly in each SR1.5 scenario, thereby effectively using the ISIMIP2b impact simulations as damage functions (time-shift approach (72, 73); fig. 1a). While the analysis can also be performed directly 135 on the RCPs, we decide to apply the time-shift approach because (i) of the increasing policy relevance of low-end warming scenarios like the 1.5° and 2°-compatible scenarios; (ii) the ISIMIP2b framework only samples a small set of greenhouse gas concentration pathways (i.e. three RCPs), whereas we here analyse a range of potential warming scenarios (see below), (iii) to better align biophysical impact projections arising from climate models with different 140 transient climate response to cumulative carbon emissions (TCRE). Both in ISIMIP2b and the SR1.5 scenarios, GMT anomalies are computed using the 1850-1900 historical period (51-year average) as reference (37, 74). Simulations whereby the absolute GMT difference with the SR1.5 scenario in any year exceeds 0.2 °C are excluded to avoid that low-end RCP projections

(e.g. RCP 2.6) inform high-end warming scenarios (e.g. current pledges). Our assumption  
145 that the simulated impacts are scenario-independent is generally valid for the considered extreme event categories (37) and allows us to maximise the impact information considered in each SR1.5 scenario. That said, some aspects of these extreme events show a lagged response to global mean temperature increase, making our assessment conservative (see Supplementary Note 2).

150 Next, we accumulate for each simulation, country and birth year within the period 1960-2020 the extreme event exposure across an average life span in that country. As life expectancy extends up to the year 2113 in some countries and birth cohorts, we assume that beyond 2099, annually exposed land fractions, GMT anomalies and gridded population densities are constant at the 2090-2099 average. In contrast, for some spatial units and early birth years, the life  
155 expectancy at birth may not extend until 2020; for those individuals still alive in 2020 the lifetime exposure accounts for the average, not actual life span. To obtain lifetime exposure values at the regional and global scale, we compute for each birth cohort the weighted spatial average of the country-scale exposure using the size of that particular cohort in each country as weighing factor. Analogous to the well-established distinction between the Eulerian and  
160 Lagrangian perspective in atmospheric science and between the period and cohort approach in demography, we suggest that the resulting lifetime extreme event exposure values represent the Lagrangian/cohort view on climate change impacts.

Computing the multi-model arithmetic mean per extreme impact category then enables the comparison of different birth cohorts (fig. 1b), whereby the results combine the effect of changes  
165 in extreme event occurrence as a consequence of climate change and the change in life expectancy in that spatial entity.

**Exposure multiplication factor.** To analyse the lifetime exposure data, we use the exposure multiplication factor ( $EMF$ ), which is defined as

$$EMF = \frac{E_{\text{new}}}{E_{\text{ref}}} \quad (1)$$

170 where  $E_{\text{ref}}$  is the lifetime exposure of a person born in the reference year 1960 – that is, all people being 60 years old on 31 December 2020 – and  $E_{\text{new}}$  is the lifetime exposure of a person born in a later year. This metric allows us to compare birth cohorts across a range of birth years (fig. 1c). For instance, an  $EMF$  of 2 for a newborn and  $E_{\text{ref}}$  of 3.5 heatwaves implies that a person born in a given country in 1960 will on average face 3.5 heatwaves across their  
175 lifetime, whereas a person born in 2020 on will on average face 7 heatwaves. To avoid  $EMF$  values being infinite, we assign the value of 100 in the exceptional cases when extreme events emerge in a country or region. The  $EMF$  metric relates to the probability ratio metric used in previous studies (75–77), where the probability ratio is generally used as a ratio of frequencies of occurrence with probabilities limited to  $[0, 1]$  by definition. However, the  $EMF$  metric is  
180 a ratio of event counts (not of event probabilities), and explicitly includes exposure next to hazards, thereby moving towards more comprehensive risk definitions (76).

We consider three approaches to aggregate the information across extreme event categories. The first method computes the geometric mean across the  $EMF$  per event category. In this approach, percent changes in each of the categories equally contribute to the total change, but the  
185 approach yields conservative estimates in spatial units not affected by one or several categories under past and future conditions (e.g., tropical cyclones in high-latitude countries). Uncertainties in this approach are computed as the geometric mean of the per-category  $EMF$  computed based on the ensemble’s 25<sup>th</sup> and 75<sup>th</sup> percentile lifetime exposure relative to the multi-model mean exposure under pre-industrial climate conditions. The second approach is to calculate the  
190 geometric mean of the lifetime exposure across the six considered event categories and sub-



sequently compute the *EMF*. The third approach is to compute the harmonic mean across the *EMF* per event category. The harmonic mean is suited for computing the mean across ratios (such as the *EMF*) and is the most conservative of the Pythagorean means. Note that the arithmetic mean or the sum are not considered here because the results would be dominated  
195 by the strong increase in heatwave occurrence. The sensitivity to the aggregation procedure is visualised in fig. S2; while the first and third aggregation method yield consistent results, the second approach yields substantially higher exposure estimates. Taking a conservative approach, all further cross-category results are based on the first aggregation method.

**Exposure scaling with GMT.** To derive the EMFs shown in fig. 2, we first construct 28  
200 stylised GMT trajectories. The trajectories are obtained through piecewise linear interpolation between five scenarios: a present-day constant temperature (taken here as the 2009 GMT anomaly of 0.87 °C), a linear temperature increase from 0.87 °C in 2009 to 3.5 °C by 2100, and the three SR1.5 scenarios (1.5 °C, 2 °C, NDC). The resulting scenarios thereby cover, in 2100, the 0.87 – 3.5 °C GMT anomaly range with a 0.1 °C increment (fig. S1).

205 For each pathway, we subsequently compute the lifetime exposure per spatial unit and event category following the methodology explained above. As reference for the *EMF* calculation, we consider the average exposure of a person with year-1960 life expectancy under pre-industrial climate conditions. To this end, we first compute, for each of the 101 pre-industrial control simulations, the lifetime exposure for 100 bootstrapped time series. We then pool the  
210 resulting exposure values for all available simulations within that extreme event category, and calculate the arithmetic mean from the resulting distribution. The pre-industrial control exposure thereby samples from one distribution comprising uncertainty from both internal variability and structural climate and impact model deficiencies.

Once the *EMF* is calculated per extreme event category, the multi-event *EMF* is obtained

215 by computing the geometric mean across the  $EMF$  values per category (see above). The re-  
sulting  $EMF$  values are subsequently smoothed using a three-element moving average along  
the vertical and visualised in fig. 2. Note that in fig. 2, uncertainty increases along the y-axis  
as fewer impact projections are available to sample from towards higher warming levels. This  
sampling artefact explains the apparent reduction in droughts and river floods EMF for some  
220 cohorts above 3°C warming relative to pre-industrial (fig. 2c-d).

Next to the  $EMF$ , we also analyse the probability of experiencing, under pre-industrial  
climate conditions, the lifetime exposure values obtained under the stylised pathways. To this  
end, we calculate the empirical inverse percentiles from the pre-industrial control distribution of  
the lifetime exposure under each of the stylised GMT trajectories. To obtain the pre-industrial  
225 exposure distribution aggregated across the six extreme event categories, we first select 1 000  
random combinations of one simulation per extreme event category and subsequently com-  
pute in each combination the geometric mean  $EMF$  across the categories. Since each random  
combination consists of 100 lifetime exposure values obtained via bootstrapping (see above),  
this yields a distribution of 100 000 lifetime exposure values. Like with the  $EMF$  fields, the  
230 resulting probability fields are smoothed using a three-element moving average along the ver-  
tical, except for tropical cyclones, where a fourth-order polynomial is fitted to the threshold  
probability contour to account for the higher uncertainties in these projections obtained from a  
single impact model. The results are visualised as grey contours in fig. 2 and denote the 99.99<sup>th</sup>  
percentile, that is, above and right of these contours, one has less than 0.01 % probability of  
235 living such life under pre-industrial climate conditions. We refer to the latter case as living an  
unprecedented life.

With the exception of wildfires and tropical cyclones, our extreme events are defined based  
on extreme percentiles estimated from the pre-industrial control simulations (table S1) (37).  
Due to a statistical artefact (78, 79), the expected relative frequency of exceedance of those

240 percentiles (or of falling below for the 2.5<sup>th</sup> crop yield and soil moisture percentiles) may potentially be larger in data that was not used to estimate the percentiles (such as data from the historical simulations and the future scenario simulations) than in the pre-industrial control data. However, we believe that this issue can be disregarded in our analysis, because (i) we use a very large sample of pre-industrial control simulations (542 years on average) to estimate the  
245 percentile values in each simulation, and (ii) our analysis consists of relative changes between cohort lifespans that all fall entirely outside of the base period.

While fig. 2 is inspired by the burning ember diagrams shown in various IPCC reports (80–82), we acknowledge that our results cannot be directly translated into this risk framework, primarily because we only consider 2 dimensions of risk in our assessment, that is, hazard and  
250 exposure. Further work could aim at including vulnerability into the assessment, for instance by incorporating vulnerability projections and associated adaptation potentials (83) in the analysis.

**Life expectancy versus climate change.** To isolate the contribution of life expectancy change to the total change in lifetime exposure, we repeat the lifetime exposure calculation but apply it to the pre-industrial control simulations (see details below; figs. S14–S15). Assuming this term  
255 corresponds to the pure life expectancy effect in the absence of climate change, the residual represents the contribution from climate change.

## Supplementary Notes

### Supplementary Note 1 – Country-level analysis

Breaking the analysis down to country scale highlights strong spatial disparities (figs. S6–7).

260 Lifetime heatwave exposure for the 2020 birth cohort relative to the 1960 birth cohort increases in every country in the world and under all future scenarios considered (fig. S6a-b). However, in several countries the 2020 birth cohort will face more than 10 times as many extreme heatwaves compared to the 1960 birth cohort under current pledges. This is for instance the case in countries in Central Africa, the Middle East and West and Southeast Asia. Under a 1.5 °C  
265 scenario, these multiplication factors are often substantially reduced. Similar patterns emerge for the other extreme event categories (fig. S7), though the multiplication factors are subject to higher uncertainty and in some cases indicate reduced exposure, for instance in exposure to crop failures in several Eurasian countries under 1.5 °C warming. Aggregated across all categories, most countries ultimately show a decrease in the exposure multiplication factor going from the  
270 current pledges scenario to 1.5 °C of global warming (fig. S6c-d), highlighting a clear incentive for younger generations of limiting global warming to 1.5 °C instead of the 2.6–3.1 °C expected from current pledges (84, 85).

Our drought definition is based on the frequency of occurrence of extremely low monthly soil moisture values during at least seven consecutive months (table S1) as computed by 8 global  
275 vegetation models and global hydrological models from the ISIMIP biomes and water sector, respectively (CLM4.5, H08, LPJmL, JULES-W1, MPI-HM, ORCHIDEE, PCR-GLOBWB, and WaterGAP2; table S2). Thus, we mechanistically account for changes in both precipitation and evapotranspiration in our drought projections. If precipitation increases in a region according to a GCM projection, the impact simulation driven by this GCM might project less  
280 droughts. This is, for instance, the case in Russia under 1.5°C warming and current pledges,

and in parts of Scandinavia under 1.5°C warming (fig. S7e-f). But if the evapotranspiration increase is larger than the precipitation increase (especially during the dry season), drought frequency may increase despite projected increases in precipitation, like is the case in East Africa (fig. S7e-f) (86, 87). This also explains why one region may face an increase in exposure to both  
285 droughts and river floods (e.g. Southern Africa and large parts of Asia; fig. S7e-h).

## **Supplementary Note 2 – Why our estimates may be considered conservative**

Our approach yields conservative lifetime exposure estimates for at least six reasons. First, it treats consecutive extreme events (88) affecting a specific location within a calendar year as one,  
290 leading to an underestimation of the number of events in present-day as well as their increase in frequency. Second, it only considers changes in the frequency of extreme events, neglecting possible increases in event intensity and duration (71). For tropical cyclones, for instance, projected increases in storm intensity can be considered equally important (38), whereas heatwave duration and intensity are increasing next to heatwave frequency (89). Third, we do not take  
295 into account the effects of compounding extremes, even though, for instance, severe droughts, heatwaves and crop failures tend to co-occur (88, 90–93). Fourth, we only consider exposure to local hazards, yet extreme events such as crop failures may lead to regional or even global food price instability when occurring in isolation or concurrently (81, 92, 94). Fifth, we employ stringent definitions of extreme events, with for instance heatwaves occurring only about four times  
300 in a lifetime on average for the 1960 birth cohort (fig. 1b). Several extreme event categories therefore occur only over part of the globe (37), leading to an underrepresented risk when aggregating across extreme event categories. Finally, some aspects of the extreme event categories we consider demonstrate a lagged response to global warming. This notably applies for tropical cyclones, which cause substantial impacts via the storm surge they generate. These storm surges

305 are amplified by background sea level rise (95) which lags the global mean temperature increase by decades to centuries (96, 97). This lagged response further augments inter-generational inequity, and this to the extent that even the already committed sea level rise will enhance lifetime exposure of generations well beyond the ones we consider in this study (96, 97). On shorter  
310 time scales, this reasoning also applies to wildfires, as fuel aridity may build up over several years in response to a long-term warming trend (98). Overall, these six reasons highlight that our current results may underestimate changes in actual extreme event exposure and thereby underscore the benefits of climate action for current and future young generations.

### **Supplementary Note 3 – Vulnerability, impacts and adaptation**

Our study is deliberately limited to *exposure* to climate *hazards*, given that both adaptation over  
315 time and age-dependent vulnerability are extremely difficult to quantify. In that sense, we compute changes in the hazards that people face throughout their lifetime (e.g. a heatwave or a river flood), but make no call about the *risk* or *impact* which they generate (e.g. mortality, infrastructure damage). By altering their vulnerabilities, communities can also adapt to the changes in exposure to hazards. For the extreme event categories considered here, these adaptation options  
320 can take many forms: e.g. changing crop types, agricultural management (irrigation, fertiliser), flood protection, reservoir deployment and management, fire management, and improved warning systems. Depending on the adaptation options that communities will (be able to) choose, the level of risk arising from the hazard exposure will vary.

While adaptive capacities may increase over time, extreme events result in detrimental im-  
325 pacts already today, even in developed countries with very high adaptive capacity (see, for instance, the 2019-2020 wildfires in California and Australia). Moreover, for many developing countries, increased adaptive capacity is needed to address today's climate risks – it does not imply that these countries can cope with unprecedented future hazards even under very

optimistic scenarios of socio-economic development. Recent research showed that adaptive ca-  
330 pacities are far from uniformly rising Andrijevic2020a and that it will take until well into the  
21st century for many developing countries to reach current OECD levels. Overall, there are  
huge differences in adaptive capacities between countries with no signs that this gap will drasti-  
cally reduce in the next decades, and even in countries with high adaptive capacities, it is unsure  
whether entire populations will be able to adapt to severe climate change impacts. Finally, we  
335 note the existence of quite hard (physiological) limits to adaptation for several of the extreme  
events considered in this study. For instance, a wet-bulb temperature of 35°C defines the limit  
of human survivability (99–101). In other cases, migration may be the final adaptation option,  
which may in turn change both exposure and vulnerability to (a set of) hazards. For instance, a  
poor Ugandan farmer migrating to the Capital to escape from increasing crop failure may end  
340 up settling in one of Kampala’s informal settlements which are very prone to flooding, heat  
stress, and vector-borne diseases (102, 103).

Overall, the aspect of changing vulnerability and adaptive capacity requires careful consid-  
eration and the absence of a framework to quantitatively integrate future vulnerabilities into  
climate risk scenarios is why our analysis focuses explicitly on exposure to climate hazards  
345 instead of climate risk.

A cohort-based approach raises the question about its relationship to discounting over time  
to adequately account and weigh intergenerational interests. However, the concept of discount-  
ing does not apply to our study given its focus on extreme event exposure instead of climate  
change impacts. Discounting is commonly used as a methodological approach within cost-  
350 benefit analysis (CBA). Yet, our cohort-based extreme event exposure differs from the CBA  
frameworks, because it only maps out the consequences and uncertainties of different future  
climate change pathways in terms of extreme event exposure. Incorporating an exposure per-  
spective, such as ours, within a CBA would require to value the exposures and aggregate them

into a cost or welfare metric, since discounting can only be applied to value and welfare metrics.

355 Such aggregation – including the discounting – implicitly assumes that exposures between different groups are comparable and, therefore, trade-offs can be resolved. In principle, the results of the extreme event exposures can serve as input to such analysis, but it requires to value the exposures and aggregate them.

Next to informing a CBA, the extreme event exposure results could also inform a rights-  
360 based approach. Rights-based approaches argue for “a general right against risking” (104). Rights-based approaches are related to the precautionary principle and relate future risks implied by current action to the infringement on future opportunities and the exercising of basic rights.

In recent years, climate change and the risks for future generations has been increasingly  
365 brought to courts. The cohort-based extreme event exposure approach can help to inform this debate. However, we refrain from suggesting criteria or requirements that should be applied to derive the level of climate change. A comprehensive consideration of competing rights would need to take a broader set of rights into account and could not only rely on extreme event exposure. The priorities of competing rights and the weighing of competing rights requires a  
370 broader decision analysis framework that is beyond the scope of the present analysis. Finally, the methodological difference between CBA- and rights-based approaches does not necessarily imply a stronger or weaker level of future climate change and therefore higher or lower near term emissions. This is partly due to the sensitivity of CBA results to the choice of the discount rate (105) and the role competing rights can play.

#### 375 **Supplementary Note 4 – GCM selection and bias-adjustment procedure**

Global Climate Models (GCMs) allow to simulate climate and weather extremes under historical and future atmospheric conditions. The ISIMIP2b climate forcing builds on a selection of



GCM output from the fifth phase of the Coupled Model Intercomparison Project (CMIP5) (106). The ISIMIP2b forcing data covers four of the CMIP5 GCMs (IPSL-CM5A-LR, HadGEM2-ES, 380 MIROC5, GFDL-ESM2M). Uncertainty of future greenhouse gas emissions is spanned through scenarios, which are the Representative Concentration Pathways (RCPs) (107) used in CMIP5 and consequently in the IPCC's Fifth Assessment Report. In support for the IPCC Special Report on global warming of 1.5 °C, ISIMIP2b covered initially RCP2.6 and RCP6.0, a low emission and an intermediate stabilization scenario, with the high emission RCP8.5 scenario 385 added at a later stage, leading to a lower impact simulation availability (see table S2 for the scenarios covered by each impact model). The four GCMs were selected by availability of variables necessary for impact modeling and their position in the distribution of equilibrium climate sensitivity (ECS) in the CMIP5 ensemble. With an ECS of 4.1 °C for IPSL-CM5A-LR, 4.6 °C for HadGEM2-ES, 2.7 °C for MIROC5 and 2.4 °C for GFDL-ESM2M, the GCM 390 selection includes two models at the lower and two at the upper end of the CMIP5 ensemble range (2.1 °C to 4.7 °C). The climate model data is regridded from its original resolution to the ISIMIP impact model grid at a spatial resolution of 0.5° × 0.5°. The climate model data is bias-adjusted (16, 46) to better represent the statistical distribution of observational weather data while preserving simulated trends. In addition, we use sub-daily output of the GCMs listed 395 above that is not bias adjusted to force the high-resolution tropical-cyclone model.

## **Supplementary Note 5 – Wildfire simulations**

The global burned area has seen a decrease in recent years, with the decrease explained mostly by a decrease in the number of fires associated with agricultural expansion (108, 109). In this study, the wildfire simulations are designed to quantify the pure climate change effect on wild- 400 fire activity. To this end, we compute the change in burned area purely from a change in climate implemented via the GCM forcing, while keeping cropland area, population levels and other

socioeconomic factors constant at 2005 levels. The vegetation models used here are suited for this type of analyses since their fire modules generally do not include human ignition (except for ORCHIDEE in which ignition depends on population density, with population density kept constant after 2005 in our simulation design). In addition, a historical land use-induced negative trend in burned area does not preclude that exposure to wildfire activity may rise in the next decades. Regional increases in wildfire activity have already been attributed to anthropogenic climate change (98) and fire weather is projected to substantially intensify in the coming decades (110). Moreover, exposure to fire may increase due to population expansion in fire-prone regions, an effect which may even outweigh changes in burned area (111). We account for this effect thanks to the use of annual gridded population density projections when spatially averaging the hazard maps. Combining our simulated climate-induced burned area changes with projected population density data suggest a clear increase wildfire exposure despite substantial uncertainties, corroborating a recent IPCC assessment of rising wildfire damage risk under continued global warming (81, 82).


## Supplementary Note 6 – Cold spells

Next to the six extreme event categories considered in this study, the influence on climate change on cold spell exposure could also be considered. To this end, we consider a grid cell to be exposed to a cold spell in a given year if the Cold Wave Magnitude Index daily (CWMId) of that year exceeds the 99<sup>th</sup> percentile of the CWMId distribution under pre-industrial climate conditions of that grid cell. We define the CWMId as the maximum magnitude of all cold periods occurring in a year, where a cold period is a period of at least six consecutive days with daily maximum temperature falling below a threshold value  $T_{p10}$  which is defined as the 10<sup>th</sup> percentile of daily maximum temperatures under pre-industrial climate conditions, centered on a 31-day window. The magnitude of each cold period in a year is the sum of the daily


magnitudes on the consecutive days composing the cold period, with daily magnitude calculated according to  $M_d(T_d) = 0$  if  $T_d \geq T_{pi75}$  else  $(T_{pi75} - T_d)/(T_{pi75} - T_{pi25})$ , where  $T_d$  is the daily maximum temperature on day  $d$  of the cold period and  $T_{pi25}$  and  $T_{pi75}$  are the 25<sup>th</sup> and 75<sup>th</sup> percentile, respectively, of the annual minimum of the daily maximum temperature under pre-  
430 industrial climate conditions. To estimate  $T_{pi10}$ ,  $T_{pi25}$ , and  $T_{pi75}$ , we use more than 400 years of daily maximum temperature data at  $0.5^\circ \times 0.5^\circ$  spatial resolution representing pre-industrial climate conditions as available from the ISIMIP2b climate input data set. Based on these more than 400 years of temperature data we subsequently derive the 99<sup>th</sup> percentile of the CWMId distribution under pre-industrial climate conditions. As such, we characterise cold spells as the  
435 mirror of heatwave events (except for the period which we require to be six instead of three days).

The results of the analysis indicate an overall reduction of exposure to cold spells under increasing global mean temperature levels and for younger cohorts (fig. S16a). Except for a few countries in West Asia, lifetime cold spell exposure reduces consistently across most countries,  
440 with the strongest reductions in Africa, the Middle East, Europe, Canada, and parts of South America (fig. S16b).


## Supplementary Figures




**Supplementary Figure 1 | Global mean temperature pathways.** Historical and future evolution of global mean temperature (GMT) anomalies relative to the pre-industrial (PI) reference scenario, taken here as the 1850–1900 average. Shown are three marker scenarios taken from the IPCC Special Report on Global Warming of 1.5°C (54), and 25 additional stylised pathways used for constructing fig. 2 (see Methods).




**Supplementary Figure 2 | Cross-category lifetime exposure and influence of aggregation method on exposure sensitivity. (a)** Same as fig. 2, but with exposure aggregated by computing the geometric mean across the exposure multiplication factor ( $EMF$ ) per category, **(b)** Same as fig. 2, but with exposure aggregated by computing the geometric mean across the lifetime exposure ( $EXP$ ), **(c)** Same as fig. 2, but with exposure aggregated by computing the harmonic mean across the  $EMF$  per category. We note that the second aggregation method (geometric mean on exposure) yields higher  $EMF$  values because the absence of events in one extreme event category results in zero cross-category exposure in a given country. As this occurs more frequently under pre-industrial control conditions, this leads to unrealistically low global-average pre-industrial control exposure values and hence artificially high  $EMF$  values.



**Supplementary Figure 3 | World Regions.** Groups of countries based on (a) geographical location and (b) income category for which life expectancy data is available (52). The income groups are defined based on present-day conditions. The region definitions are taken from the World Bank (112) and abbreviated as follows (see e.g. fig. S10-11): East Asia & Pacific (EASP), Europe & Central Asia (EUCA), Latin America & Caribbean (LAMC), Middle East & North Africa (MENA), North America (NAM), South Asia (SAS), Sub-Saharan Africa (SSA).




**Supplementary Figure 4 | Regional lifetime exposure.** Same as fig. S2a, but for world regions shown in fig. S3a.




**Supplementary Figure 5 | Income-based lifetime exposure.** Same as fig. S2a, but for income group regions shown in fig. S3b.







**Supplementary Figure 6 | Local value of global mitigation.** (a,b) Heatwave and (c,d) all-category exposure multiplication factors at the country scale for the 2020 birth cohort relative to the 1960 birth cohort under (a,c) the current pledges scenario and (b,d) the 1.5 °C scenario. Country-scale exposure multiplication factors aggregate within-country variability in population density and land fraction affected by extreme events.




**Supplementary Figure 7 | Country-scale exposure multiplication factors.** Same as fig. S6, but for the extreme event categories (a,b) wildfires, (c,d) crop failures, (e,f) droughts, (g,h) river floods, and (i,j) tropical cyclones. Country-scale exposure multiplication factors aggregate within-country variability in population density and land fraction affected by extreme events. Note that the large tropical cyclone multiplication factors for some world regions with no or very low numbers of observed tropical cyclone landfalls (e.g. West and Southwest Africa, Western South America, Western Europe) are based on a small number of simulated tropical cyclones and should therefore be treated with caution.




**Supplementary Figure 8 | Uneven distribution of lifetime exposure.** All-category exposure multiplication factors across birth cohorts under the current pledges scenario for a range of **(a)** geographic regions and **(b)** income groups (see fig. S3 for the region definitions (112)). The factors are computed relative to the mean exposure of a reference person with year-1960 life expectancy under pre-industrial climate conditions. The kink in the curve for the Middle East and North Africa for cohorts below 20 years in 2020 can be explained by the sudden drop in life expectancy for these cohorts in Iraq and Syria.




**Supplementary Figure 9 | Exposure increases across world regions.** Regional EMFs relative to the hypothetical pre-industrial reference cohort under 1.5 °C, 2.0 °C, and current pledges pathways (colored radial distance) per world region for all extreme event categories (rows) and cohorts 60, 40, 20, and 0 years old in 2020 (columns). The upper row and angle show the relative cohort size per region. Note the different radial scale for wildfires and heatwaves.




**Supplementary Figure 10 | Exposure increase across income country groups.** Same as fig. S9 but for income country groups defined in fig. S3b. Note the different radial scale for wildfires and heatwaves.



**Supplementary Figure 11 | Exposure increase<sup>20</sup> relative to the 1960 birth cohort across income country groups.** Same as fig. S9 but using the 1960 birth cohort as reference for the EMF calculation. Note the different radial scale for heatwaves.




**Supplementary Figure 12 | Exposure increase relative to the 1960 birth cohort across income country groups.** Same as fig. S10 but using the 1960 birth cohort as reference for the EMF calculation. Note the different radial scale for heatwaves.




**Supplementary Figure 13 | Life expectancy on the rise.** Assumed increase in global cohort life expectancy at birth (adjusted for child mortality 0-4 -- see methods section).






**Supplementary Figure 14 | Drivers of increasing exposure across world regions.** All-category exposure multiplication factors across birth cohorts separated by driver under the current pledges scenario for a range of geographic regions (see fig. S3a for the region definitions (112)). As the figure is based on exposure instead of EMF, we use the arithmetic mean instead of the geometric mean to aggregate the information across the categories.



**Supplementary Figure 15 | Drivers of increasing exposure across income country groups.** Same as fig. S14, but for income country groups defined in fig. S3b. As the figure is based on exposure instead of EMF, we use the arithmetic mean instead of the geometric mean to aggregate the information across the categories.



**Supplementary Figure 16 | Cold spell exposure change.** (a) Exposure multiplication factors across birth cohorts (x-axis) under a range of global warming trajectories (fig. S1) reaching 0.87 °C to 3.5 °C global mean temperature (GMT) anomalies in 2100 relative to the pre-industrial (PI) reference period (1850–1900; y-axis) for coldwaves. All factors are computed relative to the mean exposure of a hypothetical reference person living under pre-industrial climate conditions with year-1960 life expectancy. (b) Same as fig. S6a, but for CWMId.

**Supplementary Tables**

**Supplementary Table 1** | Definition of extreme events. For each category, the number of impact models (M.) and future projections (P.) is reported. Further details are provided in the Methods section and in (37)

<b>Event</b>	<b>M.</b>	<b>P.</b>	<b>Definition of land area exposed</b>
<b>Wildfire</b>	5	53	Annual aggregate of monthly burned land area simulated by global vegetation models
<b>Crop failure</b>	3	24	Fraction of grid cell where one of the considered crops (maize, wheat, soy or rice) is grown and the corresponding crop yield falls short of the 2.5 <sup>th</sup> percentile of the pre-industrial reference distribution; crop-specific land area fractions are added up.
<b>Drought</b>	8	86	Entire grid cell if monthly soil moisture falls short of the 2.5 <sup>th</sup> percentile of the preindustrial reference distribution for at least 7 consecutive months.
<b>River Flood</b>	8	86	Flooding is assumed to occur whenever daily discharge ( $0.5^\circ \times 0.5^\circ$ resolution) exceeds the preindustrial 100-year return level (i.e. the 99 <sup>th</sup> percentile); to derive the associated land area affected per grid cell, simulated runoff is translated into inundation areas ( $2.5' \times 2.5'$ resolution) by CaMa-Flood (68).
<b>Heatwave</b>	1	12	Entire grid cell if the Heat Wave Magnitude Index daily (HWMId) (70, 71) of that year exceeds the 99 <sup>th</sup> percentile of the HWMId distribution under pre-industrial climate conditions of that grid cell. The HWMId is defined as the maximum magnitude of all hot periods occurring in a year, where a hot period is a period of at least 3 consecutive days with daily maximum temperature exceeding a threshold value $T_{pi90}$ which is defined as the 90 <sup>th</sup> percentile of daily maximum temperatures under pre-industrial climate conditions, centered on a 31-day window. The magnitude of each hot period in a year is the sum of the daily magnitudes on the consecutive days composing the hot period, with daily magnitude calculated according to $M_d(T_d) = 0$ if $T_d \leq T_{pi25}$ else $(T_d - T_{pi25}) / (T_{pi75} - T_{pi25})$ , where $T_d$ is the daily maximum temperature on day $d$ of the hot period and $T_{pi25}$ and $T_{pi75}$ are the 25 <sup>th</sup> and 75 <sup>th</sup> percentile, respectively, of the annual maximum of the daily maximum temperature under pre-industrial climate conditions. To estimate $T_{pi90}$ , $T_{pi25}$ , and $T_{pi75}$ , we use more than 400 years of daily maximum temperature data at $0.5^\circ \times 0.5^\circ$ spatial resolution representing pre-industrial climate conditions as available from the ISIMIP2b climate input data set. Based on these >400 years of temperature data we subsequently derive the 99 <sup>th</sup> percentile of the HWMId distribution under pre-industrial climate conditions.
<b>Tropical cyclone</b>	1	12	Fraction of grid cell exposed to 1-minute sustained hurricane-force winds ( $\geq 64$ kt) at least once a year ( $0.1^\circ \times 0.1^\circ$ resolution); information required about wind fields is derived from center location and minimum pressure/maximum wind speed (113, 114).

**Supplementary Table 2** | ISIMIP2b model simulations used for the analysis. Each model simulation consists of a concatenated historical and future (RCP2.6, 6.0 or 8.5) simulation done with one impact model (IM) and one global climate model (GCM). The last column indicates the number of simulation years available from the pre-industrial control simulation (PIcontrol). For tropical cyclones, each simulation represents the average of a 100-member ensemble of tropical cyclone tracks downscaled from GCM output (38).

<b>Extreme</b>	<b>IM</b>	<b>GCM</b>	<b>RCP</b>	<b>PIcontrol (years)</b>
Wildfires	CARAIB	GFDL-ESM2M	2.6, 6.0	439
Wildfires	CARAIB	HadGEM2-ES	2.6, 6.0	639
Wildfires	CARAIB	IPSL-CM5A-LR	2.6, 6.0	639
Wildfires	CARAIB	MIROC5	2.6, 6.0	639
Wildfires	LPJ-GUESS	GFDL-ESM2M	2.6, 6.0, 8.5	439
Wildfires	LPJ-GUESS	HadGEM2-ES	2.6, 6.0, 8.5	639
Wildfires	LPJ-GUESS	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Wildfires	LPJ-GUESS	MIROC5	2.6, 6.0, 8.5	639
Wildfires	LPJmL	GFDL-ESM2M	2.6, 6.0, 8.5	439
Wildfires	LPJmL	HadGEM2-ES	2.6, 6.0, 8.5	639
Wildfires	LPJmL	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Wildfires	LPJmL	MIROC5	2.6, 6.0, 8.5	639
Wildfires	ORCHIDEE	GFDL-ESM2M	2.6, 6.0, 8.5	239
Wildfires	ORCHIDEE	HadGEM2-ES	2.6, 6.0, 8.5	239
Wildfires	ORCHIDEE	IPSL-CM5A-LR	2.6, 6.0, 8.5	439
Wildfires	ORCHIDEE	MIROC5	2.6, 6.0, 8.5	239
Wildfires	VISIT	GFDL-ESM2M	2.6, 6.0, 8.5	439
Wildfires	VISIT	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Wildfires	VISIT	MIROC5	2.6, 6.0, 8.5	639
Crop failures	GEPIC	GFDL-ESM2M	2.6, 6.0	439
Crop failures	GEPIC	HadGEM2-ES	2.6, 6.0	639
Crop failures	GEPIC	IPSL-CM5A-LR	2.6, 6.0	639
Crop failures	GEPIC	MIROC5	2.6, 6.0	639
Crop failures	LPJmL	GFDL-ESM2M	2.6, 6.0	439
Crop failures	LPJmL	HadGEM2-ES	2.6, 6.0	639
Crop failures	LPJmL	IPSL-CM5A-LR	2.6, 6.0	639
Crop failures	LPJmL	MIROC5	2.6, 6.0	639
Crop failures	PEPIC	GFDL-ESM2M	2.6, 6.0	439
Crop failures	PEPIC	HadGEM2-ES	2.6, 6.0	639
Crop failures	PEPIC	IPSL-CM5A-LR	2.6, 6.0	639
Crop failures	PEPIC	MIROC5	2.6, 6.0	639
Droughts	CLM4.5	GFDL-ESM2M	2.6, 6.0, 8.5	239

**Table 2 continued from previous page**

<b>Extreme</b>	<b>IM</b>	<b>GCM</b>	<b>RCP</b>	<b>PIcontrol (years)</b>
Droughts	CLM4.5	HadGEM2-ES	2.6, 6.0, 8.5	239
Droughts	CLM4.5	IPSL-CM5A-LR	2.6, 6.0, 8.5	239
Droughts	CLM4.5	MIROC5	2.6, 6.0, 8.5	239
Droughts	H08	GFDL-ESM2M	2.6, 6.0, 8.5	439
Droughts	H08	HadGEM2-ES	2.6, 6.0, 8.5	639
Droughts	H08	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Droughts	H08	MIROC5	2.6, 6.0, 8.5	639
Droughts	LPJmL	GFDL-ESM2M	2.6, 6.0, 8.5	439
Droughts	LPJmL	HadGEM2-ES	2.6, 6.0, 8.5	639
Droughts	LPJmL	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Droughts	LPJmL	MIROC5	2.6, 6.0, 8.5	639
Droughts	JULES-W1	GFDL-ESM2M	2.6, 6.0, 8.5	439
Droughts	JULES-W1	HadGEM2-ES	2.6, 6.0, 8.5	639
Droughts	JULES-W1	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Droughts	JULES-W1	MIROC5	2.6, 6.0, 8.5	639
Droughts	MPI-HM	GFDL-ESM2M	2.6, 6.0	439
Droughts	MPI-HM	IPSL-CM5A-LR	2.6, 6.0	639
Droughts	MPI-HM	MIROC5	2.6, 6.0	639
Droughts	ORCHIDEE	GFDL-ESM2M	2.6, 6.0, 8.5	439
Droughts	ORCHIDEE	HadGEM2-ES	2.6, 6.0, 8.5	439
Droughts	ORCHIDEE	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Droughts	ORCHIDEE	MIROC5	2.6, 6.0, 8.5	439
Droughts	PCR-GLOBWB	GFDL-ESM2M	2.6, 6.0	439
Droughts	PCR-GLOBWB	HadGEM2-ES	2.6, 6.0	639
Droughts	PCR-GLOBWB	IPSL-CM5A-LR	2.6, 6.0	639
Droughts	PCR-GLOBWB	MIROC5	2.6, 6.0	639
Droughts	WaterGAP2	GFDL-ESM2M	2.6, 6.0, 8.5	439
Droughts	WaterGAP2	HadGEM2-ES	2.6, 6.0, 8.5	639
Droughts	WaterGAP2	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Droughts	WaterGAP2	MIROC5	2.6, 6.0, 8.5	639
River floods	CLM4.5	GFDL-ESM2M	2.6, 6.0, 8.5	439
River floods	CLM4.5	HadGEM2-ES	2.6, 6.0, 8.5	439
River floods	CLM4.5	IPSL-CM5A-LR	2.6, 6.0, 8.5	439
River floods	CLM4.5	MIROC5	2.6, 6.0, 8.5	439
River floods	H08	GFDL-ESM2M	2.6, 6.0, 8.5	439
River floods	H08	HadGEM2-ES	2.6, 6.0, 8.5	639
River floods	H08	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
River floods	H08	MIROC5	2.6, 6.0, 8.5	639

**Table 2 continued from previous page**

<b>Extreme</b>	<b>IM</b>	<b>GCM</b>	<b>RCP</b>	<b>PIcontrol (years)</b>
River floods	LPJmL	GFDL-ESM2M	2.6, 6.0, 8.5	439
River floods	LPJmL	HadGEM2-ES	2.6, 6.0, 8.5	639
River floods	LPJmL	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
River floods	LPJmL	MIROC5	2.6, 6.0, 8.5	639
River floods	JULES-W1	GFDL-ESM2M	2.6, 6.0, 8.5	439
River floods	JULES-W1	HadGEM2-ES	2.6, 6.0, 8.5	439
River floods	JULES-W1	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
River floods	JULES-W1	MIROC5	2.6, 6.0, 8.5	439
River floods	MPI-HM	GFDL-ESM2M	2.6, 6.0	439
River floods	MPI-HM	IPSL-CM5A-LR	2.6, 6.0	639
River floods	MPI-HM	MIROC5	2.6, 6.0	639
River floods	ORCHIDEE	GFDL-ESM2M	2.6, 6.0, 8.5	439
River floods	ORCHIDEE	HadGEM2-ES	2.6, 6.0, 8.5	439
River floods	ORCHIDEE	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
River floods	ORCHIDEE	MIROC5	2.6, 6.0, 8.5	439
River floods	PCR-GLOBWB	GFDL-ESM2M	2.6, 6.0	439
River floods	PCR-GLOBWB	HadGEM2-ES	2.6, 6.0	639
River floods	PCR-GLOBWB	IPSL-CM5A-LR	2.6, 6.0	639
River floods	PCR-GLOBWB	MIROC5	2.6, 6.0	639
River floods	WaterGAP2	GFDL-ESM2M	2.6, 6.0, 8.5	439
River floods	WaterGAP2	HadGEM2-ES	2.6, 6.0, 8.5	639
River floods	WaterGAP2	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
River floods	WaterGAP2	MIROC5	2.6, 6.0, 8.5	639
Heatwaves	HWMId-humidex	GFDL-ESM2M	2.6, 6.0, 8.5	439
Heatwaves	HWMId-humidex	HadGEM2-ES	2.6, 6.0, 8.5	639
Heatwaves	HWMId-humidex	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Heatwaves	HWMId-humidex	MIROC5	2.6, 6.0, 8.5	639
Tropical cyclones	KE-TG-meanfield	GFDL-ESM2M	2.6, 6.0, 8.5	439
Tropical cyclones	KE-TG-meanfield	HadGEM2-ES	2.6, 6.0, 8.5	439
Tropical cyclones	KE-TG-meanfield	IPSL-CM5A-LR	2.6, 6.0, 8.5	639
Tropical cyclones	KE-TG-meanfield	MIROC5	2.6, 6.0, 8.5	639



## Supplementary References

- 445 16. K. Frieler, *et al.*, *Geoscientific Model Development* **10**, 4321 (2017).
17. M. Dury, *et al.*, *iForest - Biogeosciences and Forestry* **4**, 82 (2011).
18. B. Smith, *et al.*, *Biogeosciences* **11**, 2027 (2014).
19. S. Schaphoff, *et al.*, *Geoscientific Model Development* **11**, 1343 (2018).
20. S. Schaphoff, *et al.*, *Geoscientific Model Development* **11**, 1377 (2018).
- 450 21. M. Guimberteau, *et al.*, *Geoscientific Model Development* **11**, 121 (2018).
22. A. Ito, T. Oikawa, *Ecological Modelling* **151**, 143 (2002).
23. A. Ito, M. Inatomi, *Journal of Hydrometeorology* **13**, 681 (2012).
24. C. Folberth, T. Gaiser, K. C. Abbaspour, R. Schulin, H. Yang, *Agriculture, Ecosystems Environment* **151**, 21 (2012).
- 455 25. W. Liu, *et al.*, *Agricultural and Forest Meteorology* **221**, 164 (2016).
26. W. Liu, *et al.*, *Science of The Total Environment* **572**, 526 (2016).
27. D. M. Lawrence, *et al.*, *Journal of Advances in Modeling Earth Systems* **3**, M03001 (2011).
28. W. Thiery, *et al.*, *Journal of Geophysical Research: Atmospheres* **122**, 1403 (2017).
29. N. Hanasaki, S. Yoshikawa, Y. Pokhrel, S. Kanae, *Hydrology and Earth System Sciences*  
460 **22**, 789 (2018).
30. M. J. Best, *et al.*, *Geoscientific Model Development* **4**, 677 (2011).
31. S. Hagemann, L. D. Gates, *Climate Dynamics* **21**, 349 (2003).

32. T. Stacke, S. Hagemann, *Hydrology and Earth System Sciences* **16**, 2915 (2012).
33. Y. Wada, D. Wisser, M. F. P. Bierkens, *Earth System Dynamics* **5**, 15 (2014).
- 465 34. Y. Wada, I. E. M. de Graaf, L. P. H. van Beek, *Journal of Advances in Modeling Earth Systems* **8**, 735 (2016).
35. H. Müller Schmied, *et al.*, *Hydrology and Earth System Sciences* **18**, 3511 (2014).
36. H. Müller Schmied, *et al.*, *Hydrology and Earth System Sciences* **20**, 2877 (2016).
37. S. Lange, *et al.*, *Earth's Future* **8**, 1 (2020).
- 470 38. K. A. Emanuel, *Proceedings of the National Academy of Sciences* **110**, 12219 (2013).
39. Y. Pokhrel, *et al.*, *Nature Climate Change* **11**, 226 (2021).
40. L. Gudmundsson, *et al.*, *Science* **371**, 1159 (2021).
41. R. Reinecke, *et al.*, *Hydrology and Earth System Sciences* **25**, 787 (2021).
42. C.-E. Telteu, *et al.*, *Geoscientific Model Development* **14**, 3843 (2021).
- 475 43. I. Vanderkelen, *et al.*, *Geophysical Research Letters* **47** (2020).
44. A. Ito, *et al.*, *Environmental Research Letters* **15**, 044006 (2020).
45. R. Wartenburger, *et al.*, *Environmental Research Letters* **13**, 075001 (2018).
46. S. Lange, *Earth System Dynamics* **9**, 627 (2018).
47. J. P. Dunne, *et al.*, *Journal of Climate* **25**, 6646 (2012).
- 480 48. C. D. Jones, *et al.*, *Geoscientific Model Development* **4**, 543 (2011).

49. J.-L. Dufresne, *et al.*, *Climate Dynamics* **40**, 2123 (2013).
50. M. Watanabe, *et al.*, *Journal of Climate* **23**, 6312 (2010).
51. K. E. Taylor, R. J. Stouffer, G. A. Meehl, *Bulletin of the American Meteorological Society* **93**, 485 (2012).
- 485 52. United Nations Department of Economic and Social Affairs Population Division, *World Population Prospects 2019* (2019).
53. J. R. Goldstein, K. W. Wachter, *Population Studies* **60**, 257 (2006).
54. J. Rogelj, *et al.*, *Mitigation Pathways Compatible with 1.5C in the Context of Sustainable Development*. In: *Global Warming of 1.5C. An IPCC Special Report on the impacts of global warming of 1.5C above pre-industrial levels and related global greenhouse gas emission pathw*, V. Masson-Delmotte, *et al.*, eds. (2018), pp. 93–174.
- 490
55. J. Rogelj, *et al.*, *Nature Climate Change* **8**, 325 (2018).
56. D. Huppmann, J. Rogelj, E. Kriegler, V. Krey, K. Riahi, *Nature Climate Change* **8**, 1027 (2018).
- 495 57. D. Huppmann, *et al.*, IAMC 1.5C Scenario Explorer and Data hosted by Integrated Assessment Modeling Consortium International Institute for Applied Systems Analysis (2019).
58. A. Grubler, *et al.*, *Nature Energy* **3**, 515 (2018).
59. V. Krey, *et al.*, *MESSAGE-GLOBIOM 1.0 Documentation* (International Institute for Applied Systems Analysis (IIASA), 2016).

- 500 60. E. Stehfest, *et al.*, *Integrated Assessment of Global Environmental Change with IMAGE 3.0: Model description and policy applications* (PBL Netherlands Environmental Assessment Agency, The Hague, 2014).
61. K. K. Goldewijk, A. Beusen, J. Doelman, E. Stehfest, *Earth System Science Data* **9**, 927 (2017).
- 505 62. B. Jones, B. C. O'Neill, *Environmental Research Letters* **11**, 084003 (2016).
63. O. Fricko, *et al.*, *Global Environmental Change* **42**, 251 (2017).
64. S. KC, M. Potancokova, R. Bauer, A. Goujon, E. Striessnig, Summary of Data, Assumptions and Methods for New Wittgenstein Centre for Demography and Global Human Capital ( WIC ) Population Projections by Age, Sex and Level of Education for 195 Countries to 2100, *Tech. rep.*, International Institute for Applied Systems Analysis (2013).
- 510 65. W. Lutz, A. Goujon, S. KC, M. Stonawski, N. Stilianakis, *Demographic and human capital scenarios for the 21st century: 2018 assessment for 201 countries* (Publications Office of the European Union, 2018).
66. S. I. Seneviratne, *et al.*, *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation*, C. Field, *et al.*, eds. (Cambridge University Press, Cambridge, UK, 2012), pp. 109–230.
- 515 67. F. Zhao, *et al.*, *Environmental Research Letters* **12**, 075003 (2017).
68. D. Yamazaki, S. Kanae, H. Kim, T. Oki, *Water Resources Research* **47**, 9726 (2011).
69. IPCC, *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* pp. 1447–1466 (2013).
- 520

70. S. Russo, J. Sillmann, E. M. Fischer, *Environmental Research Letters* **10**, 124003 (2015).
71. S. Russo, J. Sillmann, A. Sterl, *Scientific Reports* **7**, 7477 (2017).
72. N. Herger, B. M. Sanderson, R. Knutti, *Geophysical Research Letters* **42**, 3486 (2015).
- 525 73. R. James, R. Washington, C.-F. Schleussner, J. Rogelj, D. Conway, *Wiley Interdisciplinary Reviews: Climate Change* **8**, e457 (2017).
74. M. Allen, *et al.*, *Global Warming of 1.5C. An IPCC Special Report on the impacts of global warming of 1.5C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of*  
530 *climate change*, V. Masson-Delmotte, *et al.*, eds. (2018), pp. 49–91.
75. P. a. Stott, D. a. Stone, M. R. Allen, *Nature* **432**, 610 (2004).
76. E. M. Fischer, R. Knutti, *Nature Climate Change* **5**, 560 (2015).
77. W. Thiery, *et al.*, *Nature Communications* **11**, 290 (2020).
78. X. Zhang, G. Hegerl, F. W. Zwiers, J. Kenyon, *Journal of Climate* **18**, 1641 (2005).
- 535 79. S. Sippel, *et al.*, *Geophysical Research Letters* **42**, 9990 (2015).
80. IPCC, *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, C. Field, *et al.*, eds. (2014), pp. 1–32.
81. M. Hurlbert, *et al.*, *Climate Change and Land: an IPCC special report on climate change, desertification, land degradation, sustainable land management, food security, and green-*  
540 *house gas fluxes in terrestrial ecosystems*, P. Shukla, *et al.*, eds. (IPCC, 2019), pp. 673–800.

82. Z. Zommers, *et al.*, *Nature Reviews Earth Environment* (2020).
83. M. Andrijevic, J. Crespo Cuaresma, T. Lissner, A. Thomas, C. F. Schleussner, *Nature Communications* **11**, 1 (2020).
- 545 84. J. Rogelj, *et al.*, *Nature* **534**, 631 (2016).
85. UNEP, Emissions Gap Report 2019, *Tech. rep.*, United Nations Environment Programme (2019).
86. W. Thiery, *et al.*, *Nature Communications* **7**, 12786 (2016).
87. N. Souverijns, W. Thiery, M. Demuzere, N. P. M. V. Lipzig, *Environmental Research Letters* **11**, 114011 (2016).
- 550
88. J. Zscheischler, *et al.*, *Nature Reviews Earth and Environment* **1**, 333 (2020).
89. S. E. Perkins-Kirkpatrick, S. C. Lewis, *Nature Communications* **11**, 3357 (2020).
90. F. Piontek, *et al.*, *Proceedings of the National Academy of Sciences* **111**, 3233 (2014).
91. J. Zscheischler, S. I. Seneviratne, *Science Advances* **3**, e1700263 (2017).
- 555 92. K. Kornhuber, *et al.*, *Nature Climate Change* **10**, 48 (2020).
93. C.-F. Schleussner, *et al.*, *Environmental Research Letters* **13**, 064007 (2018).
94. F. Gaupp, J. Hall, S. Hochrainer-Stigler, S. Dadson, *Nature Climate Change* **10**, 54 (2020).
95. A. J. Reed, *et al.*, *Proceedings of the National Academy of Sciences* **112**, 12610 (2015).
96. M. Mengel, A. Nauels, J. Rogelj, C.-F. Schleussner, *Nature Communications* **9**, 601 (2018).
- 560 97. A. Nauels, *et al.*, *Proceedings of the National Academy of Sciences* **116**, 23487 (2019).

98. J. T. Abatzoglou, A. P. Williams, *Proceedings of the National Academy of Sciences* (2016).
99. J. S. Pal, E. A. B. Eltahir, *Nature Climate Change* **6**, 197 (2016).
100. E.-s. Im, J. S. Pal, E. A. B. Eltahir **2010**, 1 (2017).
101. F. Saeed, C. F. Schleussner, M. Ashfaq, *Geophysical Research Letters* **48** (2021).
- 565 102. O. Brousse, *et al.*, *International Journal of Climatology* **40**, 4586 (2020).
103. O. Brousse, *et al.*, *Environmental Research Letters* **15**, 124051 (2020).
104. L. H. Meyer, F. Schuppert, H. Stelzer, A. Placani, *Living with Uncertainty* (Cambridge University Press, Cambridge, 2009), pp. 72–117.
105. W. Nordhaus, *Science* **317**, 1682 (2007).
- 570 106. K. E. Taylor, R. J. Stouffer, G. A. Meehl, *Bulletin of the American Meteorological Society* **93**, 485 (2012).
107. M. Meinshausen, *et al.*, *Climatic Change* **109**, 213 (2011).
108. N. Andela, *et al.*, *Science* **356**, 1356 (2017).
109. V. K. Arora, J. R. Melton, *Nature Communications* **9**, 1326 (2018).
- 575 110. J. T. Abatzoglou, A. P. Williams, R. Barbero, *Geophysical Research Letters* **46**, 326 (2019).
111. W. Knorr, A. Arneeth, L. Jiang, *Nature Climate Change* **6**, 781 (2016).
112. The World Bank, Life expectancy at birth (2020).
113. K. Emanuel, *Nature* **436**, 686 (2005).

580 114. T. Geiger, K. Frieler, D. N. Bresch, *Earth System Science Data* **10**, 185 (2018).