

REVIEW ARTICLE

MECHANISMS OF DISEASE

Interleukin-17 and Type 17 Helper T Cells

Pierre Miossec, M.D., Ph.D., Thomas Korn, M.D., and Vijay K. Kuchroo, Ph.D.

From the Department of Immunology and Rheumatology, Hôpital Edouard Herriot, University of Lyon, Lyon, France (P.M.); the Department of Neurology, Technical University of Munich, Munich, Germany (T.K.); and the Center for Neurologic Diseases, Brigham and Women's Hospital, Harvard Medical School, Boston (V.K.K.). Address reprint requests to Dr. Miossec at the Clinical Immunology Unit, Department of Immunology and Rheumatology, Hôpital Edouard Herriot, 5, Place d'Arsonval, 69003 Lyon, France, or at miossec@univ-lyon1.fr.

Drs. Miossec and Korn contributed equally to this article.

N Engl J Med 2009;361:888-98.

Copyright © 2009 Massachusetts Medical Society.

IN 1986, MOSMANN AND COFFMAN INTRODUCED THE CONCEPT OF DISTINCT types of helper T cells, which was based on the types of cytokines that T cells produce when they are stimulated to differentiate. They named these lymphocytes type 1 helper T cells (Th1 cells) and type 2 helper T cells (Th2 cells).¹ Th1 cells produce large quantities of interferon- γ , induce delayed hypersensitivity reactions, activate macrophages, and are essential for the defense against intracellular pathogens (Fig. 1). Th2 cells produce mainly interleukin-4 and are important in inducing IgE production, recruiting eosinophils to sites of inflammation, and helping to clear parasitic infections (Fig. 1). These distinctions allowed the assignment of a specific functional phenotype to helper T cells based on the effector cytokines that they produce. The production of effector cytokines underlies the term “effector helper T cells.”

ORIGINS AND FUNCTIONS OF Th17 CELLS

HELPER T-CELL SUBGROUPS

More recently, T cells were shown to produce cytokines that could not be classified according to the Th1–Th2 scheme. Interleukin-17 was among these cytokines,² and the T cells that preferentially produce interleukin-17, but not interferon- γ or interleukin-4, were named Th17 cells.^{3,4} Since these T cells constitute a distinct lineage, we now have three types of effector helper T cells: Th1, Th2, and Th17⁵ (Fig. 1).

The mechanism of induction and the effector functions of this new class of effector helper T cells are just beginning to be understood. Their function in clearing specific types of infectious organisms, their role in inducing inflammation, and the molecular events that cause them to differentiate are the focus of important studies in immunology. Like Th1 and Th2 cells, Th17 cells produce a group of distinctive cytokines — interleukin-17 (also called interleukin-17A), interleukin-17F, interleukin-22, and interleukin-21 — all of which participate in orchestrating a specific kind of inflammatory response (Table 1).

DIFFERENTIATION OF Th17 CELLS

We will first describe Th17 cells in mice, since the first discoveries involving these cells were made in mice. Although there are major analogies to the development of Th17 cells in humans, some differences have been observed. Cytokines produced by cells of the innate immune system govern the differentiation of helper T cells. The cells of the innate immune system are the first line of defense against pathogens; their pattern-recognition receptors, which are not specific for any particular epitope, allow them to respond to a wide variety of microbial invaders by producing cytokines that activate T cells of the adaptive immune system. Interferon- γ and interleukin-12 drive naive T cells into the Th1 pathway, whereas interleukin-4 initiates the differentiation of naive T cells into Th2 cells⁶ (Fig. 2). At the molecular







Th Group	Cell Products	Cell Target	Infectious Agents
Th1	 Interleukin-12R Interferon- γ Interleukin-2	 Macrophages Dendritic cells	Intracellular bacteria Fungi Viruses
Th17	 Interleukin-23R Interleukin-17A Interleukin-17F Interleukin-21 Interleukin-22	 Neutrophils	Extracellular bacteria Fungi
Th2	 Interleukin-4R Interleukin-4 Interleukin-13 Interleukin-5	 Eosinophils Basophils	Parasites

Figure 1. Helper T-Cell (Th) Subgroups and Effector Functions.
The cytokine profile (including key cytokine receptors as denoted by R), the effector cell type that is activated, and the corresponding types of infections are shown for each Th subgroup.

level, the differentiation of Th1 and Th2 cells requires lineage-specific transcription factors: T-bet for Th1 cells⁷ and GATA3 and c-Maf for Th2 cells.^{8,9} These factors activate the hallmark Th1 and Th2 cytokine genes, *IFN- γ* and *IL-4*, respectively (Fig. 2).

Interleukin-12 is also important for the differentiation of Th1 cells. It has two subunits, p35 and p40. Another protein, p19, which has no activity of its own, combines with the p40 subunit of interleukin-12 to form a unique heterodimeric cytokine called interleukin-23.¹⁰ Thus, interleukin-12 and interleukin-23 have in common the p40 subunit, but they also have unique subunits, p35 (interleukin-12) and p19 (interleukin-23). In studies of genetically deficient mice that specifically lacked interleukin-23 or interleukin-12, the loss of interleukin-23 made the animals highly resistant to the development of autoimmunity and inflammation, whereas the loss of interleukin-12 did not.^{11,12} These results suggest that it is not interleukin-12 and Th1 cells that are required for the induction of autoimmune-mediated inflammation but rather interleukin-23. They also suggest that T cells that differentiate under the influence of interleukin-23 are key players in

the induction of autoimmunity. This concept is supported by experiments in which the induction of inflammation and autoimmunity in mice was made possible by injecting interleukin-17-producing T cells that had been induced to differentiate and proliferate in vitro by interleukin-23.¹³ Thus, it appears that the interleukin-23–Th17 axis is a predominant pathway to the induction of autoimmune disease.

Interleukin-23 can expand a population of Th17 cells in vitro even when they are rendered genetically deficient in the master transcription factors of Th1 and Th2 cells^{3,4}; this observation confirms that Th17 cells are a lineage that is distinct from Th1 and Th2 cells. The finding that interleukin-23 is a differentiating cytokine for Th17 cells poses a major conceptual problem, however, because naive T cells do not express receptors for interleukin-23; thus, highly purified naive T cells cannot differentiate into Th17 cells in the presence of interleukin-23.¹⁴

The problem took a new turn when three independent groups simultaneously discovered that a combination of transforming growth factor β (TGF- β) plus interleukin-6 induced the differentiation of naive T cells into Th17 cells.¹⁴⁻¹⁶ This

Table 1. Sources and Functions of Key Cytokines.*

Cytokine	Main Cell Source	Function
Interferon- γ	Th1 cells, natural killer cells, natural killer T cells	Cell-mediated immunity; control of intracellular pathogens; inhibition of Th17 pathway
Interleukin-1	Monocytes, other cells	Proinflammatory cytokine; induction of Th17 cells
Interleukin-4	Th2 cells, natural killer T cells	Antibody-mediated immunity; control of parasitic infections; antiinflammatory effect by inhibition of interleukin-1, TNF, and interleukin-6 production by monocytes; inhibition of Th17 pathway
Interleukin-6	Monocytes, other cells	Induction of acute-phase proteins; effects on B cells; induction of Th17 cells
Interleukin-8	Monocytes, other cells	Major chemokine for neutrophils
Interleukin-12	Monocytes, dendritic cells	Induction of Th1 pathway; acts in synergy with interleukin-18
Interleukin-17	Th17 cells, natural killer cells, natural killer T cells	Proinflammatory cytokine; control of extracellular pathogens; induction of matrix destruction; synergy with TNF and interleukin-1
Interleukin-18	Monocytes, dendritic cells	Induction of Th1 pathway; acts in synergy with interleukin-12
Interleukin-21	Th17 cells	Amplification of Th17 pathway in autocrine fashion
Interleukin-22	Th17 cells	Induction of epithelial-cell proliferation and of antimicrobial proteins in keratinocytes
Interleukin-23	Monocytes, dendritic cells	Th17 expansion and stabilization
Interleukin-25	Th2 cells	Interleukin-17 family member; induction of Th2-associated cytokines; inhibition of interleukin-1 and interleukin-23
TGF- β	Many cells	Induction of Foxp3 ⁺ regulatory T cells in the absence of interleukin-6; together with interleukin-6, interleukin-21, and interleukin-1 β , induction of Th17 cells
TNF	Monocytes, dendritic cells	Proinflammatory cytokine; acts synergistically with interleukin-17

* TGF- β denotes transforming growth factor β , and TNF tumor necrosis factor.

finding was surprising, since TGF- β had been classified as an immunosuppressive cytokine, not as an inducer of T-cell differentiation. Furthermore, naive T cells that are exposed to TGF- β alone express forkhead box P3 (Foxp3), the master transcription factor that induces regulatory T cells — T cells that suppress inflammation and inhibit autoimmunity.¹⁷ A relevant finding is that interleukin-6 is a potent inhibitor of TGF- β -driven induction of Foxp3⁺ regulatory T cells.¹⁶ Interleukin-6 not only suppresses the generation of these cells, but together with TGF- β , it also forces naive T cells to express interleukin-17 and to become Th17 cells. Thus, Th17 cells and Foxp3⁺ regulatory T cells are reciprocally related: TGF- β induces naive T cells to develop into suppressor regulatory T cells, whereas interleukin-6 switches the transcriptional program initiated by TGF- β in a way that induces the development of Th17 cells.

Th17 cells express a unique transcription fac-

tor, ROR- γ t,¹⁸ which induces transcription of the *IL-17* gene in naive helper T cells and is required for the development of interleukin-17 producing cells in the presence of interleukin-6 and TGF- β .¹⁹ ROR- γ t must act in cooperation with other transcription factors, including ROR- α , signal transducer and activator of transcription 3 (STAT3), IRF-4, and runt-related transcription factor 1 (Runx1), for full commitment of precursors to the interleukin-17 lineage.²⁰⁻²³ Activation of ROR- γ t also causes expression of the receptor for interleukin-23, indicating that interleukin-23 acts on T cells that are already committed to the Th17 lineage. Exposure of developing Th17 cells to interleukin-23 not only enhances the expression of interleukin-17 but also induces interleukin-22 and suppresses interleukin-10 and interferon- γ , which are not normally associated with the Th17 phenotype.²⁴ Thus, interleukin-23 is essential for stabilizing the Th17 phenotype.

The interferon- γ and interleukin-4 produced




Figure 2. Differentiation of Mouse Th17 Cells.

Naive mouse T cells can differentiate into one of three effector helper T-cell (Th) subgroups. Each pathway is under the control of a different set of cytokines. The Th17 pathway is under the control of transforming growth factor β (TGF- β) plus interleukin-6 and interleukin-1 or TGF- β plus interleukin-21 followed by interleukin-23. This pathway is inhibited by interferon- γ and interleukin-4. The transcription factor (T-bet, ROR- γ t, or GATA3) characteristic of each pathway is shown. R denotes receptor.


by Th1 and Th2 cells, respectively, amplify the differentiation of these cells in an autocrine loop. Interleukin-17, by contrast, is neither a growth factor nor a differentiation factor for Th17 cells; thus, it cannot amplify Th17 responses. However, a member of the interleukin-2 cytokine family — interleukin-21, which is produced in large amounts by mature Th17 cells — can, together with TGF- β , amplify Th17-cell differentiation²⁵⁻²⁷ (Fig. 2); in the absence of interleukin-21, the expansion of Th17 cells is defective. In short, there is also an autocrine loop for Th17 cells, but in this loop, TGF- β and interleukin-21 are major factors.

HUMAN Th17 CELLS

Initially, TGF- β plus interleukin-6 were not considered to be differentiation factors for human Th17 cells. On the contrary, it was thought that the generation of human Th17 cells from naive precursors was inhibited by TGF- β and promoted by interleukin-6 plus interleukin-1 β .^{28,29} The studies underlying these ideas, however, did not use genuinely naive T cells as a starting popula-

tion and did not control for endogenous sources of TGF- β such as serum and platelets. When naive T cells from cord blood were cultured in serum-free medium, the generation of Th17 cells as a result of the interaction between TGF- β and an “inflammatory” cytokine was confirmed in human T cells.^{30,31} It appears that TGF- β plus interleukin-21,³⁰ TGF- β plus a combination of interleukin-6 and interleukin-23, or interleukin-6 plus interleukin-21³¹ can induce the expression of ROR-c, the human counterpart of murine ROR- γ t (Fig. 3).

As with Th1 and Th2 cells, no single surface marker is specific for Th17 cells. However, coexpression of the chemokine receptors CCR4 and CCR6³² or expression of CCR2 in the absence of CCR5³³ appears to define human Th17 cells. (Chemokines induce chemotactic responses in neighboring cells that display receptors for chemokines, of which there are four main types: CXC, CC, CX3C, and XC.) Some memory helper T cells produce both interferon- γ and interleukin-17,³⁴ and these cells express CXCR3 in addi-



tion to CCR4 and CCR6. They also appear to express both the T-bet and ROR-c transcription factors. In reactive lymph nodes and inflamed tissues, large cells with a resemblance to plasma cells produce interleukin-17 (Fig. 3), suggesting that Th17 cells acquire an activated phenotype at the tissue site.³⁵

INTERLEUKIN-17 AND Th17 CELLS IN DISEASE

RESPONSES TO INFECTIOUS AGENTS

Th17 cells can rapidly initiate an inflammatory response that is dominated by neutrophils (Fig. 1); indeed, acute inflammation in which neutrophils are prominent is typical of Th17-driven inflammation. Immunity mediated by Th17 cells is particularly important at epithelial and mucosal surfaces, as indicated by the pattern of expression of their chemokine receptors and effector cytokines.^{36,37}

A number of pathogens induce mainly Th17 responses (Fig. 1). They include gram-positive

Propionibacterium acnes, gram-negative *Citrobacter rodentium*, *Klebsiella pneumoniae*, bacteroides species and borrelia species, *Mycobacterium tuberculosis*, and fungi such as *Candida albicans*.^{14,38-42} To rid the body of fungi and certain extracellular bacteria requires inflammation of the type engendered by Th17 cells. The role of interleukin-17 and Th17 cells in clearing infections has been shown in the hyper-IgE syndrome, in which a mutation in *STAT3*, one of a family of transcription activators, nullifies the ability to mount Th17 responses. Patients with this disorder have recurrent *C. albicans* and *Staphylococcus aureus* infections in the skin and lungs.^{43,44}

THE INTERLEUKIN-23–Th17 PATHWAY IN CHRONIC INFLAMMATION AND AUTOIMMUNITY

Unregulated Th17 responses or overwhelming interleukin-17 production from T cells and other sources is associated with chronic inflammation and severe immunopathologic conditions. Interleukin-17 was first shown to induce interleukin-6 in fibroblasts² and in cultured synoviocytes in






Target-Cell Type	Products Released	Biologic Effect	Condition
 Macrophage, dendritic cell	Interleukin-1 TNF Interleukin-6 CRP	Inflammation	Infections Psoriasis Graft rejection
 Endothelial cell	Interleukin-6 Coagulation MMP	Vessel activation	Reperfusion injury Thrombosis Atherosclerosis
 Fibroblast	Interleukin-6 Chemokines Growth factors MMP	Matrix destruction	Multiple sclerosis Crohn's disease
 Osteoblast	RANKL MMP Osteoclastogenesis	Bone erosion	Prosthesis loosening Periodontal disease Rheumatoid arthritis
 Chondrocyte	MMP	Cartilage damage	

Figure 4. Effects of Interleukin-17 on Cell Functions and Its Role in the Pathophysiology of Diseases.
 For each key effect of interleukin-17, the target-cell type involved and the products released in response to interleukin-17 are shown. Each biologic effect is linked to examples of conditions in which an association with the presence of interleukin-17 has been observed. CRP denotes C-reactive protein, MMP matrix metalloproteinase, RANKL receptor activator of nuclear factor- κ B ligand, and TNF tumor necrosis factor.

patients with rheumatoid arthritis⁴⁵ (Fig. 4). Most parenchymal cells express interleukin-17 receptors⁴⁶ (Table 2), and signaling through these receptors induces target cells to produce pro-inflammatory factors such as interleukin-6, interleukin-1, tumor necrosis factor (TNF), CXCL8 (interleukin-8), and matrix metalloproteinases.^{2,3,56} Through the production of matrix proteinases, interleukin-17 can also destroy extracellular matrix and cause bone resorption. In bone, interleukin-17 stimulates osteoblasts to express the receptor activator of nuclear factor- κ B (RANK) ligand (RANKL).⁵⁷ Such osteoblasts can activate osteoclasts, which express the membrane protein RANK, a receptor for RANKL, at their surface. Th17 cells also express RANKL, but they may not activate osteoclasts by a RANKL-RANK

interaction. Rather, secretion of interleukin-17 by Th17 cells and induction of RANKL on cells such as osteoblasts that support the activation of osteoclasts appear to be required for bone loss. Through the RANKL-RANK system, interleukin-17 may have a role in rheumatoid arthritis, periodontal disease, and loosening of joint prostheses.⁵⁸ In rheumatoid arthritis, the production of TNF, interleukin-1, and interleukin-17 by synovial cells is predictive of joint destruction.⁵⁹

Interleukin-6 is both a target of interleukin-17 and a differentiation factor for Th17 cells. By inducing the production of interleukin-6 or interleukin-1 β , interleukin-17 activates a positive feedback loop that commits naive T cells to the Th17 lineage. Moreover, by inducing chemokine production, Th17 cells attract numerous effector

Table 2. Receptors of Th17-Associated Cytokines.*

Receptor and Its Structure, Distribution, and Function	Reference
Interleukin-17A receptor	
Interleukin-17RA, the cognate receptor for interleukin-17A, is highly expressed on hematopoietic cells and expressed at lower levels on osteoblasts, fibroblasts, and endothelial and epithelial cells	Yao et al. ²
Human interleukin-17RC binds human interleukin-17A with high affinity, but mouse interleukin-17RC does not bind mouse interleukin-17A	Kuestner et al. ⁴⁷
Human interleukin-17RA and interleukin-17RC can form a heterodimer that binds human interleukin-17A	Toy et al. ⁴⁸
Interleukin-17RA appears to be part of the functional interleukin-25 receptor (a heterodimer consisting of interleukin-17RA and interleukin-17RB)	Rickel et al. ⁴⁹
Interleukin-17F receptor	
Interleukin-17RC, the cognate receptor for interleukin-17F, is expressed at low levels on hematopoietic cells and at high levels on nonhematopoietic cells	Toy et al. ⁴⁸
Human interleukin-17RA–interleukin-17RC heterodimers can bind human interleukin-17F	Toy et al. ⁴⁸
Interleukin-21 receptor	
This receptor is a heterodimer consisting of the common cytokine-receptor γ chain (γ_c) and interleukin-21R	Leonard and Spolski ⁵⁰
γ_c is expressed on lymphoid cells	Takeshita et al. ⁵¹
Interleukin-21R is restricted to hematopoietic cells (but not only lymphoid cells) with highest levels of expression on B cells, but also on T cells, natural killer cells, and some populations of myeloid cells	Parrish-Novak et al. ⁵²
Interleukin-22 receptor	
This receptor is a heterodimer of interleukin-22R1 and interleukin-10R2	Kotenko et al. ⁵³
Interleukin-10R2 is ubiquitously expressed on hematopoietic and nonhematopoietic cells	Moore et al. ⁵⁴
Interleukin-22R1 is expressed on a variety of epithelial and parenchymal tissues (skin, liver, kidney, pancreas, intestine, lung)	Wolk et al. ⁵⁵

* Some Th17 cytokines, such as interleukin-21, might exclusively target lymphoid cells, whereas the Th17-associated effector cytokines interleukin-17, interleukin-17F, and interleukin-22 have widespread effects on many tissues.

T cells into inflamed tissue; acting in synergy with TNF and interleukin-1 β , interleukin-17 is a potent inducer of the chemokine CCL20, which is strongly chemotactic for lymphocytes, including Th17 cells. These Th17 cells are drawn to sites of interleukin-17–driven inflammation through CCR6, the receptor for CCL20, which is detected on Th17 cells.³² In addition, chemokines such as interferon-inducible protein-10 (IP-10), a member of the CXC chemokine family, attract other immune cells such as Th1 cells and monocytes into inflamed tissues.⁴¹

There is evidence that, apart from rheumatoid arthritis,⁵⁹ Th17 cells are involved in psoriasis,⁶⁰ multiple sclerosis,⁶¹ and inflammatory bowel disease.⁶² They may also participate in the development of corticosteroid-resistant asthma.^{63,64} Genetic studies have linked certain sequence variants of the interleukin-23–receptor gene to suscepti-

bility to Crohn's disease, psoriasis, and psoriatic arthritis.^{62,65,66}

The predominant T-cell population that can be isolated from the skin lesions of patients with psoriasis has a Th17 phenotype,⁶⁷ which accords with the attraction of inflammatory cells to epithelial tissues by CCL20–CCR6 signaling. In preclinical models of inflammatory hyperkeratosis, the pathogenic role of interleukin-22 production by Th17 cells, driven by interleukin-23, has been shown conclusively.⁶⁸

In multiple sclerosis, the role of Th17 cells, if any, has been difficult to explore. *IL-17* and *IL-6* are among the most highly expressed genes in brain lesions in patients with the disease,⁶⁹ and elevated levels of interleukin-17 have been detected in serum and cerebrospinal fluid from patients with multiple sclerosis.⁶¹ In patients with multiple sclerosis in whom lesions are restricted

to the optic nerves and spinal cord, the levels of interleukin-17 and interleukin-8 (CXCL8) in serum and cerebrospinal fluid are higher than in conventional multiple sclerosis,⁷⁰ and the level of interleukin-17 in the cerebrospinal fluid correlates with the extent of spinal lesions as measured by means of magnetic resonance imaging.⁷⁰ An *in vitro* study suggested that Th17 cells have the capacity to breach the blood-brain barrier and infiltrate the parenchyma of the central nervous system.⁷¹ The expression of the p19 chain of interleukin-23 is increased in monocyte-derived dendritic cells in blood samples from patients with multiple sclerosis, and this increase correlates with an augmented capacity of the dendritic cells to induce the production of interleukin-17 by T cells.⁷²

THERAPEUTIC POTENTIAL

Targeting the interleukin-6 receptor with a monoclonal antibody (e.g., tocilizumab, a humanized monoclonal antibody against the receptor) and preempting the interleukin-1 receptor with an interleukin-1-receptor antagonist (e.g., anakinra, a recombinant human interleukin-1-receptor antagonist) are two effective approaches to the treatment of rheumatoid arthritis and other autoimmune inflammatory diseases.^{73,74} Given that interleukin-6 regulates the balance between Th17 and regulatory T cells,^{75,76} it is possible that blocking interleukin-6–induced intracellular signals by a monoclonal antibody against the interleukin-6 receptor ameliorates the function of regulatory T cells, thereby bringing the immune system into physiologic balance.

Since interleukin-23 enhances interleukin-17 production and induces production of other effector cytokines in Th17 cells, inhibition of interleukin-23 is another way to control Th17 cells. Treatment with a monoclonal antibody against p40, a polypeptide common to interleukin-12 and interleukin-23 (e.g., treatment with ustekinumab), has been shown to have efficacy in psoriasis and Crohn's disease.^{77,78} In Crohn's disease, the antibody caused a local reduction of the levels of interleukin-12 and interleukin-23. However, since it neutralizes both interleukin-12 and interleukin-23, the effects cannot be attributed specifically to the interleukin-23–Th17 axis. Nevertheless, studies of a psoriasis-like skin disease and preclinical models of inflammatory bowel disease

in mice suggest that interleukin-23–driven inflammation dominates these models of psoriasis and Crohn's disease rather than the interleukin-12–interferon- γ axis.^{67,68,79,80}

The most direct way to control the biologic effects of Th17 cells would be to target the effector cytokines that they produce. Monoclonal antibodies against interleukin-17 or the interleukin-17 receptor and a soluble interleukin-17 receptor (Table 2) have been developed for clinical application. Phase 2 trials of a monoclonal antibody against interleukin-17 (AIN457) for psoriasis, rheumatoid arthritis, Crohn's disease, and psoriatic arthritis are under way. So far, inhibitors of interleukin-21 and interleukin-22 have been tested only in preclinical models of autoimmune diseases,^{81,82} and we will have to await further results to determine whether these inhibitors are clinically useful.

Some cytokines have anti-interleukin-17 properties and control the development of Th17 cells. Interleukin-4, for example, inhibits the production and functions of interleukin-17,³ and interleukin-25, which is produced by Th2 cells, also inhibits the production of interleukin-17 by downregulating interleukin-23, interleukin-1, and interleukin-6.⁸³ Treatment with interleukin-25 can suppress autoimmune inflammation of the brain in mice.⁸³ In the inflamed central nervous system, resident microglial cells are the major source of interleukin-25.

Another cytokine, interleukin-27, a member of the interleukin-12–interleukin-23 family and a heterodimer of p28 and EB13 (a glycoprotein related to p40), specifically inhibits the development of Th17 cells.^{84,85} In addition, interleukin-27 participates in the induction and differentiation of Tr1 cells, which resemble regulatory T cells and produce interleukin-10 and interferon- γ . The interaction with Tr1 cells may allow interleukin-27 to suppress inflammation indirectly.^{86–88}

Future treatments could target the effector functions of Th17 cells. Interleukin-17 induces the production of interleukin-1 and TNF- α . Interleukin-1 and TNF- α may not directly block the generation of Foxp3+ regulatory T cells; however, they inhibit the functions of regulatory T cells. This phenomenon may explain the recurrence of immunoinflammatory disease when treatment with TNF inhibitors is discontinued.⁸⁹ A combination of interleukin-17 and TNF inhibitors, administered either simultaneously or sequentially,

might better control inflammation and might even restore the function of regulatory T cells^{90,91} (see figure in the Supplementary Appendix, available with the full text of this article at NEJM.org).

CONCLUSIONS

In 1995, the newly discovered interleukin-17 was thought to be of minimal importance because it lacked immediate effects on T cells and B cells.⁵⁷ Interest in this molecule began to emerge when its role in inducing inflammation was established. Now, there is agreement that interleukin-17 has an important role in providing protection against infection and in inducing and maintaining chronic inflammatory diseases.

Molecules involved in the induction of Th17 cells and their effector functions (i.e., interleukins 6, 17, 21, 22, and 23) have been identified, and this knowledge will allow the rational development of strategies for modulating Th17 cells. The reciprocal relationship between regulatory T cells and Th17 cells suggests possibilities for shifting the balance between them in a manner that restores the function of regulatory T cells in autoimmune diseases.

Supported by grants from the Deutsche Forschungsgemeinschaft (KO2964/2-1, to Dr. Korn) and from the National Institutes of Health (NIH), the National Multiple Sclerosis Society, and the Juvenile Diabetes Research Foundation Center at Harvard Medical School (all to Dr. Kuchroo), and by a Javits Neuroscience Investigator Award from the NIH (to Dr. Kuchroo).

No potential conflict of interest relevant to this article was reported.

REFERENCES

- Mosmann TR, Coffman RL. TH1 and TH2 cells: different patterns of lymphokine secretion lead to different functional properties. *Annu Rev Immunol* 1989;7:145-73.
- Yao Z, Fanslow WC, Seldin MF, et al. Herpesvirus Saimiri encodes a new cytokine, IL-17, which binds to a novel cytokine receptor. *Immunity* 1995;3:811-21.
- Park H, Li Z, Yang XO, et al. A distinct lineage of CD4 T cells regulates tissue inflammation by producing interleukin 17. *Nat Immunol* 2005;6:1133-41.
- Harrington LE, Hatton RD, Mangan PR, et al. Interleukin 17-producing CD4+ effector T cells develop via a lineage distinct from the T helper type 1 and 2 lineages. *Nat Immunol* 2005;6:1123-32.
- Bettelli E, Korn T, Kuchroo VK. Th17: the third member of the effector T cell trilogy. *Curr Opin Immunol* 2007;19:652-7.
- Glimcher LH, Murphy KM. Lineage commitment in the immune system: the T helper lymphocyte grows up. *Genes Dev* 2000;14:1693-711.
- Szabo SJ, Kim ST, Costa GL, Zhang X, Fathman CG, Glimcher LH. A novel transcription factor, T-bet, directs Th1 lineage commitment. *Cell* 2000;100:655-69.
- Ho IC, Hodge MR, Rooney JW, Glimcher LH. The proto-oncogene c-maf is responsible for tissue-specific expression of interleukin-4. *Cell* 1996;85:973-83.
- Zheng W, Flavell RA. The transcription factor GATA-3 is necessary and sufficient for Th2 cytokine gene expression in CD4 T cells. *Cell* 1997;89:587-96.
- Oppmann B, Lesley R, Blom B, et al. Novel p19 protein engages IL-12p40 to form a cytokine, IL-23, with biological activities similar as well as distinct from IL-12. *Immunity* 2000;13:715-25.
- Cua DJ, Sherlock J, Chen Y, et al. Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune inflammation of the brain. *Nature* 2003;421:744-8.
- Murphy CA, Langrish CL, Chen Y, et al. Divergent pro- and antiinflammatory roles for IL-23 and IL-12 in joint autoimmune inflammation. *J Exp Med* 2003;198:1951-7.
- Langrish CL, Chen Y, Blumenschein WM, et al. IL-23 drives a pathogenic T cell population that induces autoimmune inflammation. *J Exp Med* 2005;201:233-40.
- Mangan PR, Harrington LE, O'Quinn DB, et al. Transforming growth factor-beta induces development of the T(H)17 lineage. *Nature* 2006;441:231-4.
- Veldhoen M, Hocking RJ, Atkins CJ, Locksley RM, Stockinger B. TGFbeta in the context of an inflammatory cytokine milieu supports de novo differentiation of IL-17-producing T cells. *Immunity* 2006;24:179-89.
- Bettelli E, Carrier Y, Gao W, et al. Reciprocal developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells. *Nature* 2006;441:235-8.
- Li MO, Wan YY, Sanjabi S, Robertson AK, Flavell RA. Transforming growth factor-beta regulation of immune responses. *Annu Rev Immunol* 2006;24:99-146.
- Ivanov II, McKenzie BS, Zhou L, et al. The orphan nuclear receptor RORgamma directs the differentiation program of proinflammatory IL-17(+) T helper cells. *Cell* 2006;126:1121-33.
- Zhou L, Lopes JE, Chong MM, et al. TGF-beta-induced Foxp3 inhibits T(H)17 cell differentiation by antagonizing RORgamma function. *Nature* 2008;453:236-40.
- Yang XO, Pappu BP, Nurieva R, et al. T helper 17 lineage differentiation is programmed by orphan nuclear receptors RORalpha and RORgamma. *Immunity* 2008;28:29-39.
- Yang XO, Panopoulos AD, Nurieva R, et al. STAT3 regulates cytokine-mediated generation of inflammatory helper T cells. *J Biol Chem* 2007;282:9358-63.
- Brüstle A, Heink S, Huber M, et al. The development of inflammatory T(H)-17 cells requires interferon-regulatory factor 4. *Nat Immunol* 2007;8:958-66.
- Zhang F, Meng G, Strober W. Interactions among the transcription factors Runx1, RORgamma and Foxp3 regulate the differentiation of interleukin 17-producing T cells. *Nat Immunol* 2008;9:1297-306. [Erratum, *Nat Immunol* 2009;10:223.]
- McGeachy MJ, Bak-Jensen KS, Chen Y, et al. TGF-beta and IL-6 drive the production of IL-17 and IL-10 by T cells and restrain T(H)-17 cell-mediated pathology. *Nat Immunol* 2007;8:1390-7.
- Korn T, Bettelli E, Gao W, et al. IL-21 initiates an alternative pathway to induce proinflammatory T(H)17 cells. *Nature* 2007;448:484-7.
- Nurieva R, Yang XO, Martinez G, et al. Essential autocrine regulation by IL-21 in the generation of inflammatory T cells. *Nature* 2007;448:480-3.
- Zhou L, Ivanov II, Spolski R, et al. IL-6 programs T(H)-17 cell differentiation by promoting sequential engagement of the IL-21 and IL-23 pathways. *Nat Immunol* 2007;8:967-74.
- Acosta-Rodriguez EV, Napolitani G, Lanzavecchia A, Sallusto F. Interleukins 1beta and 6 but not transforming growth factor-beta are essential for the differentiation of interleukin 17-producing human T helper cells. *Nat Immunol* 2007;8:942-9.
- Wilson NJ, Boniface K, Chan JR, et al.

- Development, cytokine profile and function of human interleukin 17-producing helper T cells. *Nat Immunol* 2007;8:950-7.
30. Yang L, Anderson DE, Baecher-Allan C, et al. IL-21 and TGF-beta are required for differentiation of human T(H)17 cells. *Nature* 2008;454:350-2.
31. Manel N, Unutmaz D, Littman DR. The differentiation of human T(H)-17 cells requires transforming growth factor-beta and induction of the nuclear receptor RORgamma. *Nat Immunol* 2008;9:641-9.
32. Acosta-Rodriguez EV, Rivino L, Geginat J, et al. Surface phenotype and antigenic specificity of human interleukin 17-producing T helper memory cells. *Nat Immunol* 2007;8:639-46.
33. Sato W, Aranami T, Yamamura T. Cutting edge: human Th17 cells are identified as bearing CCR2+CCR5- phenotype. *J Immunol* 2007;178:7525-9.
34. Annunziato F, Cosmi L, Liotta F, Maggi E, Romagnani S. The phenotype of human Th17 cells and their precursors, the cytokines that mediate their differentiation and the role of Th17 cells in inflammation. *Int Immunol* 2008;20:1361-8.
35. Page G, Sattler A, Kersten S, Thiel A, Radbruch A, Miossec P. Plasma cell-like morphology of Th1-cytokine-producing cells associated with the loss of CD3 expression. *Am J Pathol* 2004;164:409-17.
36. Aujla SJ, Chan YR, Zheng M, et al. IL-22 mediates mucosal host defense against Gram-negative bacterial pneumonia. *Nat Med* 2008;14:275-81.
37. Ouyang W, Kolls JK, Zheng Y. The biological functions of T helper 17 cell effector cytokines in inflammation. *Immunity* 2008;28:454-67.
38. Ye P, Rodriguez FH, Kanaly S, et al. Requirement of interleukin 17 receptor signaling for lung CXC chemokine and granulocyte colony-stimulating factor expression, neutrophil recruitment, and host defense. *J Exp Med* 2001;194:519-27.
39. Chung DR, Kasper DL, Panzo RJ, et al. CD4+ T cells mediate abscess formation in intra-abdominal sepsis by an IL-17-dependent mechanism. *J Immunol* 2003;170:1958-63.
40. Infante-Duarte C, Horton HF, Byrne MC, Kamradt T. Microbial lipopeptides induce the production of IL-17 in Th cells. *J Immunol* 2000;165:6107-15.
41. Khader SA, Bell GK, Pearl JE, et al. IL-23 and IL-17 in the establishment of protective pulmonary CD4+ T cell responses after vaccination and during *Mycobacterium tuberculosis* challenge. *Nat Immunol* 2007;8:369-77.
42. Huang W, Na L, Fidel PL, Schwarzenberger P. Requirement of interleukin-17A for systemic anti-*Candida albicans* host defense in mice. *J Infect Dis* 2004;190:624-31.
43. Milner JD, Brenchley JM, Laurence A, et al. Impaired T(H)17 cell differentiation in subjects with autosomal dominant hyper-IgE syndrome. *Nature* 2008;452:773-6.
44. Ma CS, Chew GY, Simpson N, et al. Deficiency of Th17 cells in hyper IgE syndrome due to mutations in STAT3. *J Exp Med* 2008;205:1551-7.
45. Chabaud M, Fossiez F, Taupin JL, Miossec P. Enhancing effect of IL-17 on IL-1-induced IL-6 and leukemia inhibitory factor production by rheumatoid arthritis synoviocytes and its regulation by Th2 cytokines. *J Immunol* 1998;161:409-14.
46. Kolls JK, Lindén A. Interleukin-17 family members and inflammation. *Immunity* 2004;21:467-76.
47. Kuestner RE, Taft DW, Haran A, et al. Identification of the IL-17 receptor related molecule IL-17RC as the receptor for IL-17F. *J Immunol* 2007;179:5462-73.
48. Toy D, Kugler D, Wolfson M, et al. Cutting edge: interleukin 17 signals through a heteromeric receptor complex. *J Immunol* 2006;177:36-9.
49. Rickel EA, Siegel LA, Yoon BR, et al. Identification of functional roles for both IL-17RB and IL-17RA in mediating IL-25-induced activities. *J Immunol* 2008;181:4299-310.
50. Leonard WJ, Spolski R. Interleukin-21: a modulator of lymphoid proliferation, apoptosis and differentiation. *Nat Rev Immunol* 2005;5:688-98.
51. Takeshita T, Asao H, Ohtani K, et al. Cloning of the gamma chain of the human IL-2 receptor. *Science* 1992;257:379-82.
52. Parrish-Novak J, Dillon SR, Nelson A, et al. Interleukin 21 and its receptor are involved in NK cell expansion and regulation of lymphocyte function. *Nature* 2000;408:57-63.
53. Kotenko SV, Izotova LS, Mirochnitchenko OV, et al. Identification of the functional interleukin-22 (IL-22) receptor complex: the IL-10R2 chain (IL-10Rbeta) is a common chain of both the IL-10 and IL-22 (IL-10-related T cell-derived inducible factor, IL-TIF) receptor complexes. *J Biol Chem* 2001;276:2725-32.
54. Moore KW, de Waal Malefyt R, Coffman RL, O'Garra A. Interleukin-10 and the interleukin-10 receptor. *Annu Rev Immunol* 2001;19:683-765.
55. Wolk K, Kunz S, Witte E, Friedrich M, Asadullah K, Sabat R. IL-22 increases the innate immunity of tissues. *Immunity* 2004;21:241-54.
56. Fossiez F, Djossou O, Chomarat P, et al. T cell interleukin-17 induces stromal cells to produce proinflammatory and hematopoietic cytokines. *J Exp Med* 1996;183:2593-603.
57. Page G, Miossec P. RANK and RANKL expression as markers of dendritic cell-T cell interactions in paired samples of rheumatoid synovium and lymph nodes. *Arthritis Rheum* 2005;52:2307-12.
58. Sato K, Suematsu A, Okamoto K, et al. Th17 functions as an osteoclastogenic helper T cell subset that links T cell activation and bone destruction. *J Exp Med* 2006;203:2673-82.
59. Kirkham BW, Lassere MN, Edmonds JP, et al. Synovial membrane cytokine expression is predictive of joint damage progression in rheumatoid arthritis: a two-year prospective study (the DAMAGE study cohort). *Arthritis Rheum* 2006;54:1122-31.
60. Krueger GG, Langley RG, Leonardi C, et al. A human interleukin-12/23 monoclonal antibody for the treatment of psoriasis. *N Engl J Med* 2007;356:580-92.
61. Matusiewicz D, Kivisäkk P, He B, et al. Interleukin-17 mRNA expression in blood and CSF mononuclear cells is augmented in multiple sclerosis. *Mult Scler* 1999;5:101-4.
62. Duerr RH, Taylor KD, Brant SR, et al. A genome-wide association study identifies IL23R as an inflammatory bowel disease gene. *Science* 2006;314:1461-3.
63. Molet S, Hamid Q, Davoine F, et al. IL-17 is increased in asthmatic airways and induces human bronchial fibroblasts to produce cytokines. *J Allergy Clin Immunol* 2001;108:430-8.
64. Barczyk A, Pierzchala W, Sozanska E. Interleukin-17 in sputum correlates with airway hyperresponsiveness to methacholine. *Respir Med* 2003;97:726-33.
65. Cargill M, Schrodi SJ, Chang M, et al. A large-scale genetic association study confirms IL12B and leads to the identification of IL23R as psoriasis-risk genes. *Am J Hum Genet* 2007;80:273-90.
66. Liu Y, Helms C, Liao W, et al. A genome-wide association study of psoriasis and psoriatic arthritis identifies new disease loci. *PLoS Genet* 2008;4(3):e1000041.
67. Pène J, Chevalier S, Preisser L, et al. Chronically inflamed human tissues are infiltrated by highly differentiated Th17 lymphocytes. *J Immunol* 2008;180:7423-30.
68. Zheng Y, Danilenko DM, Valdez P, et al. Interleukin-22, a T(H)17 cytokine, mediates IL-23-induced dermal inflammation and acanthosis. *Nature* 2007;445:648-51.
69. Lock C, Hermans G, Pedotti R, et al. Gene-microarray analysis of multiple sclerosis lesions yields new targets validated in autoimmune encephalomyelitis. *Nat Med* 2002;8:500-8.
70. Ishizu T, Osogawa M, Mei FJ, et al. Intrathecal activation of the IL-17/IL-8 axis in opticospinal multiple sclerosis. *Brain* 2005;128:988-1002.
71. Kebir H, Kreyenborg K, Ifergan I, et al. Human T(H)17 lymphocytes promote blood-brain barrier disruption and central nervous system inflammation. *Nat Med* 2007;13:1173-5.
72. Vaknin-Dembinsky A, Balashov K,

- Weiner HL. IL-23 is increased in dendritic cells in multiple sclerosis and down-regulation of IL-23 by antisense oligos increases dendritic cell IL-10 production. *J Immunol* 2006;176:7768-74. [Erratum, *J Immunol* 2006;177:2025.]
73. Yokota S, Miyamae T, Imagawa T, et al. Therapeutic efficacy of humanized recombinant anti-interleukin-6 receptor antibody in children with systemic-onset juvenile idiopathic arthritis. *Arthritis Rheum* 2005;52:818-25.
74. Dinarello CA. Blocking IL-1 in systemic inflammation. *J Exp Med* 2005;201:1355-9.
75. Serada S, Fujimoto M, Mihara M, et al. IL-6 blockade inhibits the induction of myelin antigen-specific Th17 cells and Th1 cells in experimental autoimmune encephalomyelitis. *Proc Natl Acad Sci U S A* 2008;105:9041-6.
76. Korn T, Mitsdoerffer M, Croxford AL, et al. IL-6 controls Th17 immunity in vivo by inhibiting the conversion of conventional T cells into Foxp3+ regulatory T cells. *Proc Natl Acad Sci U S A* 2008;105:18460-5.
77. Leonardi CL, Kimball AB, Papp KA, et al. Efficacy and safety of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with psoriasis: 76-week results from a randomised, double-blind, placebo-controlled trial (PHOENIX 1). *Lancet* 2008;371:1665-74.
78. Sandborn WJ, Feagan BG, Fedorak RN, et al. A randomized trial of ustekinumab, a human interleukin-12/23 monoclonal antibody, in patients with moderate-to-severe Crohn's disease. *Gastroenterology* 2008;135:1130-41.
79. Kullberg MC, Jankovic D, Feng CG, et al. IL-23 plays a key role in *Helicobacter hepaticus*-induced T cell-dependent colitis. *J Exp Med* 2006;203:2485-94.
80. Uhlig HH, McKenzie BS, Hue S, et al. Differential activity of IL-12 and IL-23 in mucosal and systemic innate immune pathology. *Immunity* 2006;25:309-18.
81. Young DA, Hegen M, Ma HL, et al. Blockade of the interleukin-21/interleukin-21 receptor pathway ameliorates disease in animal models of rheumatoid arthritis. *Arthritis Rheum* 2007;56:1152-63.
82. Ma HL, Liang S, Li J, et al. IL-22 is required for Th17 cell-mediated pathology in a mouse model of psoriasis-like skin inflammation. *J Clin Invest* 2008;118:597-607.
83. Kleinschek MA, Owyang AM, Joyce-Shaikh B, et al. IL-25 regulates Th17 function in autoimmune inflammation. *J Exp Med* 2007;204:161-70.
84. Stumhofer JS, Laurence A, Wilson EH, et al. Interleukin 27 negatively regulates the development of interleukin 17-producing T helper cells during chronic inflammation of the central nervous system. *Nat Immunol* 2006;7:937-45.
85. Batten M, Li J, Yi S, et al. Interleukin 27 limits autoimmune encephalomyelitis by suppressing the development of interleukin 17-producing T cells. *Nat Immunol* 2006;7:929-36.
86. Awasthi A, Carrier Y, Peron JP, et al. A dominant function for interleukin 27 in generating interleukin 10-producing anti-inflammatory T cells. *Nat Immunol* 2007;8:1380-9.
87. Fitzgerald DC, Zhang GX, El-Behi M, et al. Suppression of autoimmune inflammation of the central nervous system by interleukin 10 secreted by interleukin 27-stimulated T cells. *Nat Immunol* 2007;8:1372-9. [Erratum, *Nat Immunol* 2008;9:105.]
88. Stumhofer JS, Silver JS, Laurence A, et al. Interleukins 27 and 6 induce STAT3-mediated T cell production of interleukin 10. *Nat Immunol* 2007;8:1363-71.
89. Feldmann M, Brennan FM, Foxwell BM, Taylor PC, Williams RO, Maini RN. Anti-TNF therapy: where have we got to in 2005? *J Autoimmun* 2005;25:Suppl:26-8.
90. Zaba LC, Cardinale I, Gilleaudeau P, et al. Amelioration of epidermal hyperplasia by TNF inhibition is associated with reduced Th17 responses. *J Exp Med* 2007;204:3183-94.
91. Nadkarni S, Mauri C, Ehrenstein MR. Anti-TNF-alpha therapy induces a distinct regulatory T cell population in patients with rheumatoid arthritis via TGF-beta. *J Exp Med* 2007;204:33-9. [Erratum, *J Exp Med* 2007;204:205.]

Copyright © 2009 Massachusetts Medical Society.

FULL TEXT OF ALL JOURNAL ARTICLES ON THE WORLD WIDE WEB

Access to the complete text of the *Journal* on the Internet is free to all subscribers. To use this Web site, subscribers should go to the *Journal's* home page (NEJM.org) and register by entering their names and subscriber numbers as they appear on their mailing labels. After this one-time registration, subscribers can use their passwords to log on for electronic access to the entire *Journal* from any computer that is connected to the Internet. Features include a library of all issues since January 1993 and abstracts since January 1975, a full-text search capacity, and a personal archive for saving articles and search results of interest. All articles can be printed in a format that is virtually identical to that of the typeset pages. Beginning 6 months after publication, the full text of all Original Articles and Special Articles is available free to nonsubscribers.