

9-2013

## INTERMITTENT HYPOXIA-INDUCED RENAL ANTIOXIDANTS AND OXIDA- TIVE DAMAGE IN MALE MICE: HORMETIC DOSE RESPONSE

Weixia Sun

*The First Hospital of Jilin University, China and University of Louisville, USA*

Xia Yin

*The First Hospital of Jilin University, China and University of Louisville, USA*

Yuehui Wang

*The Second Hospital of Jilin University, China*

Yi Tan

*The First Hospital of Jilin University, China and University of Louisville, USA*

Lu Cai

*The First Hospital of Jilin University, China and University of Louisville, USA*

*See next page for additional authors*

Follow this and additional works at: [https://scholarworks.umass.edu/dose\\_response](https://scholarworks.umass.edu/dose_response)

---

### Recommended Citation

Sun, Weixia; Yin, Xia; Wang, Yuehui; Tan, Yi; Cai, Lu; Wang, Bo; Cai, Jun; and Fu, Yaowen (2013) "INTERMITTENT HYPOXIA-INDUCED RENAL ANTIOXIDANTS AND OXIDA- TIVE DAMAGE IN MALE MICE: HORMETIC DOSE RESPONSE," *Dose-Response: An International Journal*: Vol. 11 : Iss. 3 , Article 9.

Available at: [https://scholarworks.umass.edu/dose\\_response/vol11/iss3/9](https://scholarworks.umass.edu/dose_response/vol11/iss3/9)

This Article is brought to you for free and open access by ScholarWorks@UMass Amherst. It has been accepted for inclusion in Dose-Response: An International Journal by an authorized editor of ScholarWorks@UMass Amherst. For more information, please contact [scholarworks@library.umass.edu](mailto:scholarworks@library.umass.edu).

---

# INTERMITTENT HYPOXIA-INDUCED RENAL ANTIOXIDANTS AND OXIDATIVE DAMAGE IN MALE MICE: HORMETIC DOSE RESPONSE

## **Authors**

Weixia Sun, Xia Yin, Yuehui Wang, Yi Tan, Lu Cai, Bo Wang, Jun Cai, and Yaowen Fu

## INTERMITTENT HYPOXIA-INDUCED RENAL ANTIOXIDANTS AND OXIDATIVE DAMAGE IN MALE MICE: HORMETIC DOSE RESPONSE

**Weixia Sun\***, **Xia Yin\*** □ The First Hospital of Jilin University, China and KCHRI at the Department of Pediatrics, University of Louisville, USA

**Yuehui Wang\*** □ The Second Hospital of Jilin University, China

**Yi Tan, Lu Cai** □ KCHRI at the Department of Pediatrics, University of Louisville, USA and The Chinese-American Research Institute for Diabetic Complications, China

**Bo Wang** □ The Department of Pathology, Inner Mongolia Forestry General Hospital, Yakeshi, China

**Jun Cai** □ KCHRI at the Department of Pediatrics, University of Louisville, USA

**Yaowen Fu** □ The First Hospital of Jilin University, China

□ Obstructive sleep apnea causes cardiovascular disease via chronic intermittent hypoxia (IH), which may be related to oxidative stress. Nuclear factor-erythroid 2-related factor 2 (Nrf2) is an important cellular defense mechanism against oxidative stress by regulating its down-stream multiple antioxidants. The present study was to define whether IH can induce renal pathogenic damage and if so, whether Nrf2 and its down-stream antioxidants are involved in IH-induced pathogenic changes. Mice were culled for exposure to intermittent air as control or IH that consisted of 20.9% O<sub>2</sub>/ 8% O<sub>2</sub>F<sub>1</sub>O<sub>2</sub> alternation cycles (30 episodes per h) with 20 seconds at the nadir F<sub>1</sub>O<sub>2</sub> for 12 h a day during daylight. Short-term IH exposure (3 – 7 days) induced significant increases in renal inflammatory response and antioxidant levels along with a reduction of the spontaneous content of malondialdehyde while long-term IH exposure (8 weeks) induced a significant decrease of antioxidant levels and significant increases of renal inflammation, oxidative damage, cell death, and fibrosis. This study suggests that IH induces a hormetic response, i.e.: short-term IH exposure is able to induce a protective response to protect the kidney from oxidative damage while long-term IH exposure is able to induce a damage effect on the kidney.

*Key words: Intermittent hypoxia, kidney hypoxic damage, Nrf2, metallothionein*

\* These authors contribute equally to this work.

Address correspondence to Dr. Lu Cai, MD, PhD, Department of Pediatrics, University of Louisville, 570 South Preston Street, Baxter I, Suite 321B or 304F, Louisville, KY 40202; Phone: (502) 852-2214; Fax: (502) 852-5634; E-mail: [10cai001@louisville.edu](mailto:10cai001@louisville.edu)

Dr. Yaowen Fu, MD, PhD, Center of Organ Transplantation, the First Hospital of Jilin University, 71 Xinmin Street, Changchun 130021, Phone: + 86-13756661667; E-mail: [fuyaowen@medmail.com.cn](mailto:fuyaowen@medmail.com.cn)

W. Sun and others

## INTRODUCTION

As a highly prevalent chronic disease, obstructive sleep apnoea (OSA) is characterized by recurrent episodes of partial or complete upper airway collapse and obstruction during sleep. OSA affects about 9% of adult women and 24% of adult men (Young *et al.* 1993). The intermittent hypoxia (IH) has been considered as a main cause of OSA pathogenesis since IH is able to induce oxidative stress, inflammation, atherosclerosis, endothelial dysfunction and hypertension (Dematteis *et al.* 2009).

In a cross-sectional study of 35 patients with chronic kidney diseases, the majority of patients (54%) had OSA (Markou *et al.* 2006). Despite a high overall oxygen supply, the tissue oxygen tension in the kidney is comparatively low that renders the kidney prone to hypoxic injury (Eckardt *et al.* 2003). Compared to subjects without nocturnal hypoxia, subjects with nocturnal hypoxia demonstrated an almost three-fold increase in the risk of accelerated loss of kidney function (Ahmed *et al.* 2011). However, there was also report that declining kidney function increased the prevalence of sleep apnea too (Nicholl *et al.* 2012). Therefore, it was unclear whether OSA-related IH induces kidney damage or kidney dysfunction causes sleep apnea. Can IH directly affect the kidney function in the subject who does not have any abnormality? To answer this question, we have to use animal studies. However, there was no a detail time-course study on the kidney response to IH in terms of structural and functional changes as well as cellular pathogenic changes with animal models.

Inflammation-related oxidative stress that has been assumed as a major cause for the cardiovascular damage and dysfunction in patients with OSA (Khayat *et al.* 2009) may also contribute to chronic kidney disease progression. Oxidative stress describes an imbalance between the production of reactive oxygen or nitrogen species (ROS or RNS) and the antioxidant capacity.

Cellular ROS and RNS levels and their effects are regulated by a variety of specific antioxidant systems. Nuclear factor-erythroid 2-related factor 2 (Nrf2) is one of the important cellular defense mechanisms against oxidative stress (Tkachev *et al.* 2011). It regulates many phase II detoxifying enzymes and proteins that detoxify xenobiotics and neutralize ROS and/or RNS to maintain cellular redox homeostasis. Hemeoxygenase-1 (HO-1), NAD(P)H quinone oxidoreductase-1 (NQO-1), superoxide dismutase (SOD), and metallothionein (MT) are among the well-studied Nrf2 target genes (Tkachev *et al.* 2011). It has been reported that hypoxia can induce Nrf2 activation (Papaiahgari *et al.* 2006).

The present study was to investigate the dynamic responses of the kidney to short- and long-term exposure to IH for 3 days to 8 weeks, which was followed by renal functional, structural and biochemical examina-

*Renal pathogenic effect of intermittent hypoxia*

tions. To study whether oxidative stress is involved in the pathophysiological changes, Nrf2 expression and its down-stream antioxidants were examined dynamically.

**MATERIALS AND METHODS****Animals**

FVB mice were used for this study and housed in the University of Louisville Research Resources Center at 22 °C with a 12-h light/dark cycle with free access to standard rodent chow and tap water. All animal procedures were approved by the Institutional Animal Care and Use Committee, which is certified by the American Association for Accreditation of Laboratory Animal Care.

**IH and sham exposure**

The murine model of IH exposures during sleep has been established and extensively utilized in the study of sleep apnea-associated morbidities (Cai *et al.* 2012). In this study, 8 – 10 weeks old male mice were culled for exposure by two groups – IH group and intermittent air control group. Selection of male mice only for the present study was because several measurements induced by IH were significantly protected in female mice (Li *et al.* 2012). The outcome from the present study will not reflect the response of female mice to the same challenge. The animals were randomly allocated into four control groups and four hypoxia groups: 3 days, 1 week, 2 weeks, and 8 weeks. Animal numbers were 3 in control groups (except for 2 weeks control, n = 5) and 5 in hypoxia groups (except for 2 weeks hypoxia group, n = 8). The IH paradigm consisted of 20.9% O<sub>2</sub>/8% O<sub>2</sub> F<sub>1</sub>O<sub>2</sub> alternation cycles (30 episodes per h) with 20 seconds at the nadir F<sub>1</sub>O<sub>2</sub> for 12 h a day during daylight. Pulse oxyhemoglobin saturation (SpO<sub>2</sub>) changed in a recurrent manner with the nadir hemoglobin oxygen saturations mainly ranging between 60% and 70% to mimic hypoxia/reoxygenation events occurring in moderate to severe OSA patients. The IH exposure lasted different times from 3 days to 8 weeks. All animals were assigned to identical custom-designed chambers (Oxycycler model A84XOV, BioSpherix, Lacona, NY, USA) to be operated under 12-h light-dark cycle. After IH exposures, mice were transferred to room air until sacrifice for tissue collection. At the end of study, animal renal function was measured by individually measuring 24-h urinary protein and then sacrificed to collect the kidney individually from each animal for perform various biochemical measurements.

W. Sun and others

### **Analysis of the kidney function**

Mice were placed in metabolic cages for 24-h urine collection. Urine protein was measured by Bio-Rad Protein Assay Kit (Bio-Rad Laboratories, Hercules, CA, USA).

### **Western blots**

MT expression was detected by a modified Western blotting protocol (Wang *et al.* 2006). Briefly, kidney proteins were treated with DTT at a final concentration of 20 mM at 56 °C for 30 min, followed by addition of iodoacetamide (Sigma Chemical Co. St. Louis, MO, USA) at 50 mM at room temperature for 1 h in the dark. In addition, proteins were electrophoretically separated and transferred to nitrocellulose membrane with the transfer buffer including 2 mM CaCl<sub>2</sub>. The monoclonal antibody against human's MT (Dako North America, Carpinteria, CA, USA) was used at 1:1000 dilutions in 3% BSA at 4 °C overnight. Since in the transfer buffer contains CaCl<sub>2</sub>, these blots for MT could not be stripped and reprobed for β-actin analysis. Therefore, another parallel gel was used for β-actin analysis with regular process as did for other proteins (see below).

The regular Western blotting was performed as described in our previous studies (Cai *et al.* 2005; Wang *et al.* 2006). Briefly, kidney tissue were homogenized and fractionated on 10% sodium dodecyl sulfate polyacrylamide gel electrophoresis gels, and proteins were transferred to a nitrocellulose membrane. The membrane was incubated overnight at 4 °C with primary antibodies, which included HO-1 (1:500), connective tissue growth factor (CTGF, 1:500), intracellular adhesion molecule-1 (ICAM-1, 1:250), vascular cell adhesion molecule-1 (VCAM-1, 1:250), hypoxia-inducible factor 1-α (HIF-1α, 1:500), extracellular signal-regulated kinases (ERK1/2, 1:1000), Nrf2 (1:500), NQO-1 (1:500), SOD-1 (1:1000, all from Santa Cruz, CA, USA), plasminogen activator inhibitor 1 (PAI-1, 1:2000, BD Biosciences, Franklin Lakes, NJ, USA), and β-actin (1:5000, Cell Signaling). The rest of Western blotting and quantitative densitometry procedure was described in previous studies (Cai *et al.* 2005; Wang *et al.* 2006).

### **Histopathology examination**

Kidneys were collected and immersion-fixed in 10% neutral formalin, embedded in paraffin and sectioned into 5-μm-thick sections onto glass slides. After deparaffinization, tissue sections were rehydrated and stained with 0.1% Sirius-red F3BA and 0.25% Fast green FCF. Sirius-red stained sections were assessed for the proportion of fibrosis (collagen) in the kidney tissues. Apoptotic cell death in the kidney was measured by terminal deoxynucleotidyl transferase-mediated dUTP nick end labelling

*Renal pathogenic effect of intermittent hypoxia*

(TUNEL) staining with the ApopTag Peroxidase In Situ Apoptosis Detection Kit (Chemicon, Temecula, CA, USA), as described before (Cai *et al.* 2005).

**Quantitative analysis of lipid peroxides**

The lipid peroxide concentration was determined by a previously described method (Cai *et al.* 2005) which measures the amount of thiobarbituric acid reactivity by the amount of malondialdehyde (MDA) formed during acid hydrolysis of the lipid peroxide compound. After weigh out approximately 20 mg of tissue into a 1.5 ml centrifuge tube, add 200  $\mu$ l of RIPA Buffer and sonicate for 15 seconds at 40 V over ice. Centrifuge the tube at 1600 $\times$ g for 10 minutes at 4°C. Use the supernatant for analysis. The reaction mixture contained 50  $\mu$ l sample, 20  $\mu$ l 8.1% sodium dodecyl sulfate, 150  $\mu$ l 20% acetic acid solution (buffered to pH3.5), and 210  $\mu$ l 0.571% thiobarbituric acid. The mixture was then incubated at 90 °C for 1 h, cooled in an ice bath, mixed with 100  $\mu$ l distilled water and then shaken vigorously. After centrifugation at 4000 rpm for 15 min., absorbance of the solvent layer was measured at 540 nm. Tetraethoxypropane was used as an external standard, and the lipid peroxide level expressed in terms of nanomoles MDA per gram wet weight.

**Statistical analysis**


Data were collected from multiple mice and presented as mean  $\pm$  SD. Two-way ANOVA was used for analysis for the differences exist and if so, a post hoc Tukey's test was used for analysis for the difference between groups, with Origin 7.5 laboratory data analysis and graphing software. Statistical significance was considered as  $P < 0.05$ .

**RESULTS****Effects of IH on the kidney weight, renal function, and morphological change**

The ratio of kidney weight to body weight significantly decreased at the 3<sup>rd</sup> day after IH and did not change at the 1<sup>st</sup>, 2<sup>nd</sup> and 8<sup>th</sup> weeks after IH (Fig. 1A). Renal function analysis was examined by measuring 24-h urinary albumin level for mice exposed to IH for 8 weeks, which was not significantly changed (Fig. 1B). Morphological examination showed a significant increase of blood cells in the kidney of mice treated with IH for 8 weeks (Fig. 1C). In addition, there was also a significant increase of renal VCAM-1 expression, examined by Western blotting (Fig. 1D).

Next a detail time-course effect of IH on renal inflammatory responses was investigated by examining inflammatory cytokines such as ICAM-1 since it is one of the major markers of inflammation in tissue and PAI-1

W. Sun and others




**FIGURE 1.** Effects of IH on kidney weight, 24-h urine protein, morphological changes and renal VCAM-1 expression. Eight-10 weeks old male FVB mice were exposed to intermittent air (Control, C) or IH for 3 days, 1 week, 2 weeks and 8 weeks. Body weight and kidney weight was measured at the end of experiments at indicated exposure times. The ratio of kidney weight to body weight (A) was calculated and presented as mg kidney weight to g body weight. Urine was collected in metabolic cages during a 24-h time interval at the end of experiments (8 weeks). Microalbuminuria (B) was detected. Kidney morphology was examined with hematoxylin & eosin staining (C). Original magnification  $\times 400$ . Bars: 50  $\mu\text{m}$ . Renal tissues were subject to Western blotting assay for VCAM-1 expression (D).  $\beta$ -actin was used for loading control. Data are presented as mean  $\pm$  SD ( $n = 3 - 8$ ). \*,  $p < 0.05$  vs control group.

since it acts as both pro-fibrotic mediator and inflammatory mediator (Cesari *et al.* 2010; Samarakoon *et al.* 2010). We found that IH induced significant increase of the renal expression of ICAM-1 (Fig. 2A) and PAI-1 (Fig.2B) after exposure to IH for 3 and 7 days (1 week), no change for 2 weeks, but significant increase again for 8 weeks. Therefore it appears that there were two phases of inflammatory responses in the kidney in response to acute exposure and chronic exposure to IH.

#### Long-term exposure to IH induced renal fibrotic effect

The increased expression of PAI-1 as both inflammatory and pro-fibrotic cytokine in the kidney of mice exposed to IH for 8 weeks suggests the possible existence of renal fibrosis. Sirius-red staining for collagen, followed by semi-quantitative analysis, showed no significantly fibrotic




*Renal pathogenic effect of intermittent hypoxia*

**FIGURE 2.** Effects of IH on renal inflammatory responses. Eight-10 weeks old male FVB mice were exposed to intermittent air (C) or IH for 3 days, 1 week, 2 weeks and 8 weeks. Renal tissue was collected for examining ICAM-1 (A) and PAI-1 (B) expression by Western blotting assay.  $\beta$ -actin was used for loading control. Data are presented as mean  $\pm$  SD ( $n = 3 - 8$ ). \*,  $p < 0.05$  vs corresponding controls.

response in the kidney of mice exposed to IH for 3 days until 2 weeks, but a significant positive staining in interstitial area of the kidney of mice exposed to IH for 8 weeks (Fig. 3A,B). The fibrotic effect of 8-week IH exposure on the kidney was confirmed by the increased expression of CTGF, an important pro-fibrotic mediator, by Western blotting assay (Fig.3C).

W. Sun and others




**FIGURE 3.** Effects of IH on renal fibrosis. Kidney tissue was stained with Sirius-red for collagen (A). Original magnification  $\times 200$ . Bars: 50  $\mu\text{m}$ . Semi-quantitative analysis was done by computer imaging system (B). Renal expression of CTGF was examined by Western blotting assay (C). Data are presented as mean  $\pm$  SD ( $n = 3 - 8$ ). \*,  $p < 0.05$  vs corresponding controls.

### Effects of IH on renal antioxidant contents, oxidative damage and cell death

The renal Nrf2 expression and its downstream target genes HO-1, NQO1, MT and SOD1 expression were examined in the following studies with Western blotting assay. Renal expressions of Nrf2 (Fig. 4A) and HO-1 (Fig. 4B) significantly increased at early as the 3<sup>rd</sup> day and then gradually returned to normal level at the 8<sup>th</sup> week after exposure to IH. Renal expression of MT was increased at the 3<sup>rd</sup> day and significantly decreased at the 8<sup>th</sup> week (Fig. 4C). Neither renal NQO-1 (Fig. 4D) nor SOD1 (Fig. 4E) expression was changed during the period of IH exposure from 3 days to 8 weeks.

To functionally evaluate the effect of increased antioxidants in the kidney of IH-treated mice, MDA as an index of lipid peroxidation in response to oxidative stress was measured and showed that spontaneous


*Renal pathogenic effect of intermittent hypoxia*

**FIGURE 4.** Effects of IH on renal antioxidants and renal lipid peroxidation. Renal tissue from FVB mouse exposed to intermittent air (C) or IH were subject to Western blotting assay for the expression of Nrf2 (A), HO-1 (B), MT (C), NQO-1 (D), and SOD1 (E). Concentration of renal lipid peroxides (F) was determined by TBA assay, which measures the amount of thiobarbituric acid reactivity with malondialdehyde (MDA) formed during the acid hydrolysis of lipid peroxide compound. Lipid peroxide level was expressed in terms of nanomoles MDA per gram wet weight. Data are presented as mean  $\pm$  SD (n = 3 – 8). \*, p < 0.05 vs corresponding controls.

contents of renal MDA was significantly decreased in the kidney of mice exposed to IH for 3 days to 2 weeks, but there was a significant increase in the kidney of mice exposed to IH for 8 weeks (Fig. 4F).

Next apoptotic cell death in the kidney of IH-treated mice was also examined by TUNEL staining (Fig. 5A,B) and caspase-3 cleavage with Western blotting assay (Fig. 5C). Both measurements showed that exposure to IH for 3 days to 2 weeks did not induce, but exposure to IH for 8 weeks induced a significant increase of apoptotic cell death.

W. Sun and others



**FIGURE 5.** Effects of IH on renal apoptotic cell death. Renal apoptotic cell death was examined by TUNEL staining (A), followed by semi-quantitative analysis with expression of an average number of TUNEL positive cells per five high-power fields (B). Arrows indicate TUNEL positive cells, which are expressed as average number of TUNEL positive stained cells per five high power fields. \*  $P < 0.05$  vs. control group. Original magnification  $\times 200$ . Bars:  $50 \mu\text{m}$ . Renal tissue was also subject to Western blotting assay for cleaved-caspase-3 expression (C). Data are presented as mean  $\pm$  SD ( $n = 3 - 8$ ). \*,  $p < 0.05$  vs corresponding controls.

### Effects of IH on renal ERK1/2 phosphorylation and HIF-1 $\alpha$


To explore signal mechanism, renal expression of phosphorylated ERK1/2 was examined with Western blotting assay, which revealed that ERK1/2 phosphorylation was significantly increased in the kidney of mice exposed to IH for 3 days or 1 week as early phase and also for 8 weeks as the late phase (Fig. 6A), which is comparable inflammatory response (Fig. 2).

HIF-1 $\alpha$  is a key mediator of cellular responses to hypoxia. Therefore, we examined renal expression of HIF-1 $\alpha$  protein with Western blotting assay (Fig. 6B) and found that exposure of mice to IH for 3 days to 2 weeks did not significantly affect, but to IH for 8 weeks significantly increased renal HIF-1 $\alpha$  expression.

### DISCUSSION

In the present study, we have provided the first evidence that short-term IH exposure can induce a renal inflammatory response without significant pathophysiological abnormality while chronic IH exposure induced a significant renal inflammation along with significant increases

## Renal pathogenic effect of intermittent hypoxia



**FIGURE 6.** Effects of IH on renal ERK1/2 and HIF-1 $\alpha$  phosphorylation. Renal tissue from FVB mouse exposed to intermittent air (C) or IH were subject to Western blotting assay for total or phosphorylated ERK1/2 (A) and HIF-1 $\alpha$  (B) expression. Data are presented as mean  $\pm$  SD (n = 3 – 8). \*, p < 0.05 vs corresponding controls.

of renal cell death and fibrosis. The early inflammatory response was accompanied with significant up-regulation of antioxidants, including Nrf2 and its down-stream antioxidants such as HO-1 and MT, while the late inflammatory response was accompanied with decreased antioxidants and increased oxidative damage. In terms of signaling pathways, the early protective inflammatory response and up-regulation of antioxidant may be related to the ERK1/2 phosphorylation while the late detrimental inflammation along with cell death and fibrosis may be related to the late up-regulated both ERK1/2 phosphorylation and HIF-1 $\alpha$  expression.

Induction of a hormetic response in the heart by short-term IH has been documented (Beguin *et al.* 2007), but there was no any informa-

W. Sun and others

tion for the renal response to IH. Hypoxia can induce inflammation (Hartmann *et al.* 2000; Eltzschig *et al.* 2011). In the present study we demonstrated that quick induction of renal inflammation after 3-day to 1-week IH exposure, shown by increased expression of ICAM and PAI-1, was associated with a quick up-regulation of renal antioxidants including Nrf2, HO-1 and MT. This may represent body's stress response to the sudden IH, and the acute stress responses may include acute inflammatory response and quick up-regulation of renal antioxidant contents. The up-regulated expression of antioxidants (Fig. 4A-C) was associated with a significant decrease of renal lipid peroxidation (Fig. 4F). However, when mice were long-termly exposed to IH, the kidney would become incapable to tolerate IH-induced changes, reflected by the second increase of renal inflammation along with lipid peroxidation, cell death and mild renal fibrosis. Although there was no similar study to the present study, a previous study has shown the up-regulation of Nrf2 expression and activation of its down-stream genes in the kidney of mice subjected to renal ischemia and reperfusion (Leonard *et al.* 2006). Short IH exposure also up-regulated pro-inflammatory and antioxidant genes such as Nrf2 and HO-1 in human aortic endothelial cells, to provide a protective effect against oxidative stress (Polotsky *et al.* 2010). MT as a potent antioxidant plays an important role in preventing oxidative damage in the kidney under different conditions (Dorian *et al.* 1995; Sharma *et al.* 2002). However, when mice were exposed to a long-term IH (8 weeks), renal antioxidants such as Nrf2 and HO-1 were not up-regulated anymore and MT protein expression was even decreased compared to control, which may be the major mechanism for the induction of renal oxidative damage and cell death that both can lead to the mild fibrotic response in the kidney of exposed to 8-week IH.

Signaling mechanisms for the early adaptive response may be related to the activation of ERK1/2. Emerging evidence has indicated that ERK1/2, one of the best-characterized members of the mitogen-activated protein kinase (MAPK) family, mediates a range of activity from metabolism, motility, and inflammation to cell death and survival. The phosphorylated ERK1/2 level is usually increased in response to various stresses, but whether an increase in ERK1/2 phosphorylation is protective or detrimental is highly variable. The intensity and duration of stress along with the individual response determine the final outcome. Several studies indicated that activation of ERK/MAPK signaling pathways with protective genes during short-term hypoxia might represent a rapid activation of anti-apoptotic pathways (Milton *et al.* 2008) while the delayed and sustained activation of ERK/MAPK might contribute to detrimental effects (Shinozaki *et al.* 2006; Lu *et al.* 2011). There was a report that exposure to IH for 24 h induced a significant increase of Erk1/2 phosphorylation

*Renal pathogenic effect of intermittent hypoxia*

(Beguin *et al.* 2007). In agreement with these early studies, we found here that the early activation of ERK1/2 MAPK is associated with a prevention of oxidative damage and up-regulation of several antioxidants, including Nrf2, HO-1 and MT. Indeed, induction of HO-1 and even Nrf2 is also dependent on ERK1/2 activation under many conditions (Manandhar *et al.* 2007; Yang *et al.* 2011).

However, long-term and sustained activation of ERK /MAPK signaling pathway has been reported to contribute to oxidative damage and apoptotic cell death (Shinozaki *et al.* 2006; Lu *et al.* 2011). Furthermore, HIF-1 $\alpha$  was found to be activated by an ERK-dependent pathway in response to hypoxia under certain conditions (Minet *et al.* 2000); therefore, we assumed that the later increase of ERK1/2 MAPK phosphorylation activates HIF-1 $\alpha$  that leads to the pathogenic effects we observed in the present study in the kidney of mice exposed to IH.

HIF-1 is the major regulator of oxygen homeostasis within the cell, affecting and regulating dozens of genes as cellular oxygen concentrations change. HIF-1 activity is induced when mice or cultured cells are subjected to IH, an effect that is related to oxidative stress (Semenza *et al.* 2007). For instance, carotid bodies from mice that are heterozygous for a null allele at the locus encoding HIF-1 appeared histologically normal but did not respond to continuous hypoxia or CIH. In contrast to wild-type littermates, when heterozygous-null mice are subjected to CIH, they do not develop hypertension or increased levels of HIF-1 and ROS. This report suggests the existence of a feed-forward mechanism in which CIH-induced ROS activate HIF-1, which then promotes persistent oxidative stress and damage (Semenza and Prabhakar 2007). In addition, a recent study has established a pathogenic mechanism linking HIF-1, ROS generation, and cardiovascular pathology in response to IH, i.e.: HIF-1 mediates increased expression of NADPH oxidase-2 in response to IH to generate ROS and RNS (Yuan *et al.* 2011). Overexpression of HIF-1 in alveolar epithelial cells resulted in increased apoptosis (Krick *et al.* 2005). Several studies also demonstrated the induction of PAI-1 and CTGF by HIF-1 $\alpha$  in response to hypoxia and other stresses (Higgins *et al.* 2007; Kimura *et al.* 2008).

There may be a limitation that the matured younger (8 – 10 weeks old) animals were used in the present study whereas it is well known that conditions like sleep apnea increase or worsen with age (Hwang *et al.* 1994; Martin *et al.* 2002). For instance, the influence of a short-time isobaric hypoxia as well as reoxygenation on markers of oxidative stress (MDA, total SOD, GSH) and on the mRNA expression of the antioxidative enzymes (Cu/Zn- and Mn-SOD, catalase, GSH reductase and GSH peroxidase) were differentially affected by age of animals between liver and kidney (Martin *et al.* 2002). Therefore, animals with different ages will be used to investigate the effect of age on the hormetic dose response

W. Sun and others

in terms of renal antioxidant expression and oxidative damage in response to different period of IH, as performed in the present study.

In summary, here we provided the first evidence that short-term IH exposure induced renal acute inflammatory response with significant up-regulation of antioxidants including Nrf2, HO-1 and MT with a significant inhibition of oxidative damage while long-term IH exposure induced significant renal inflammatory and down-regulation of antioxidants such as MT along with significant increases of renal oxidative damage, cell death and fibrosis. This study suggests that IH is able to induce a renal hormetic response, i.e.: short-term IH induces a protective response against the renal oxidative damage, but long-term IH exposure induces a damage effect on the kidney.

### ACKNOWLEDGEMENT

The work was supported in part by grants from American Diabetes Association (1-11-BS-17 to Dr. L. Cai), Sleep Research Society Foundation/J. Christian Gillin M.D. Research Grant (001GN09 to Dr. J. Cai), and from NSF of China (No.81070189 to Dr. Y. Wang), and also by Start-Up fund for Chinese-American Research Institute for Diabetic Complications from Wenzhou Medical College (to Drs. Y. Tan & L. Cai).

### REFERENCES

- Ahmed, S.B., Ronksley, P.E., Hemmelgarn, B.R., Tsai, W.H., Manns, B.J., Tonelli, M., Klarenbach, S.W., Chin, R., Clement, F.M. and Hanly, P.J. 2011. Nocturnal hypoxia and loss of kidney function. *PLoS One* 6(4): e19029.
- Beguín, P.C., Belaidi, E., Godin-Ribuot, D., Levy, P. and Ribuot, C. 2007. Intermittent hypoxia-induced delayed cardioprotection is mediated by PKC and triggered by p38 MAP kinase and Erk1/2. *J Mol Cell Cardiol* 42(2): 343-351.
- Cai, J., Tuong, C.M., Zhang, Y., Shields, C.B., Guo, G., Fu, H. and Gozal, D. 2012. Mouse intermittent hypoxia mimicking apnoea of prematurity: effects on myelinogenesis and axonal maturation. *J Pathol* 226(3): 495-508.
- Cai, L., Wang, J., Li, Y., Sun, X., Wang, L., Zhou, Z. and Kang, Y.J. 2005. Inhibition of superoxide generation and associated nitrosative damage is involved in metallothionein prevention of diabetic cardiomyopathy. *Diabetes* 54(6): 1829-1837.
- Cesari, M., Pahor, M. and Incalzi, R.A. 2010. Plasminogen activator inhibitor-1 (PAI-1): a key factor linking fibrinolysis and age-related subclinical and clinical conditions. *Cardiovasc Ther* 28(5): e72-91.
- Dematteis, M., Godin-Ribuot, D., Arnaud, C., Ribuot, C., Stanke-Labesque, F., Pepin, J.L. and Levy, P. 2009. Cardiovascular consequences of sleep-disordered breathing: contribution of animal models to understanding the human disease. *ILAR J* 50(3): 262-281.
- Dorian, C. and Klaassen, C.D. 1995. Protection by zinc-metallothionein (ZnMT) against cadmium-metallothionein-induced nephrotoxicity. *Fundam Appl Toxicol* 26(1): 99-106.
- Eckardt, K.U., Rosenberger, C., Jurgensen, J.S. and Wiesener, M.S. 2003. Role of hypoxia in the pathogenesis of renal disease. *Blood Purif* 21(3): 253-257.
- Eltzschig, H.K. and Carmeliet, P. 2011. Hypoxia and inflammation. *N Engl J Med* 364(7): 656-665.
- Hartmann, G., Tschop, M., Fischer, R., Bidlingmaier, C., Riepl, R., Tschop, K., Hautmann, H., Endres, S. and Toepfer, M. 2000. High altitude increases circulating interleukin-6, interleukin-1 receptor antagonist and C-reactive protein. *Cytokine* 12(3): 246-252.



*Renal pathogenic effect of intermittent hypoxia*

- Higgins, D.F., Kimura, K., Bernhardt, W.M., Shrimanker, N., Akai, Y., Hohenstein, B., Saito, Y., Johnson, R.S., Kretzler, M., Cohen, C.D., Eckardt, K.U., Iwano, M. and Haase, V.H. 2007. Hypoxia promotes fibrogenesis in vivo via HIF-1 stimulation of epithelial-to-mesenchymal transition. *J Clin Invest* 117(12): 3810-3820.
- Hwang, S.M., Wilson, P.D., Laskin, J.D. and Denhardt, D.T. 1994. Age and development-related changes in osteopontin and nitric oxide synthase mRNA levels in human kidney proximal tubule epithelial cells: contrasting responses to hypoxia and reoxygenation. *J Cell Physiol* 160(1): 61-68.
- Khayat, R., Patt, B. and Hayes, D., Jr. 2009. Obstructive sleep apnea: the new cardiovascular disease. Part I: Obstructive sleep apnea and the pathogenesis of vascular disease. *Heart Fail Rev* 14(3): 143-153.
- Kimura, K., Iwano, M., Higgins, D.F., Yamaguchi, Y., Nakatani, K., Harada, K., Kubo, A., Akai, Y., Rankin, E.B., Neilson, E.G., Haase, V.H. and Saito, Y. 2008. Stable expression of HIF-1 $\alpha$  in tubular epithelial cells promotes interstitial fibrosis. *Am J Physiol Renal Physiol* 295(4): F1023-1029.
- Krick, S., Eul, B.G., Hanze, J., Savai, R., Grimminger, F., Seeger, W. and Rose, F. 2005. Role of hypoxia-inducible factor-1 $\alpha$  in hypoxia-induced apoptosis of primary alveolar epithelial type II cells. *Am J Respir Cell Mol Biol* 32(5): 395-403.
- Leonard, M.O., Kieran, N.E., Howell, K., Burne, M.J., Varadarajan, R., Dhakshinamoorthy, S., Porter, A.G., O'Farrelly, C., Rabb, H. and Taylor, C.T. 2006. Reoxygenation-specific activation of the antioxidant transcription factor Nrf2 mediates cytoprotective gene expression in ischemia-reperfusion injury. *FASEB J* 20(14): 2624-2626.
- Li, Q.Y., Li, M., Feng, Y., Guo, Q., Gu, S.Y., Liu, J.L., Zhang, R.F. and Wan, H.Y. 2012. Chronic Intermittent Hypoxia Induces Thioredoxin System Changes in a Gender-Specific Fashion in Mice. *AM J MED SCI* 343(6): 458-461.
- Lu, T.H., Hsieh, S.Y., Yen, C.C., Wu, H.C., Chen, K.L., Hung, D.Z., Chen, C.H., Wu, C.C., Su, Y.C., Chen, Y.W., Liu, S.H. and Huang, C.F. 2011. Involvement of oxidative stress-mediated ERK1/2 and p38 activation regulated mitochondria-dependent apoptotic signals in methylmercury-induced neuronal cell injury. *Toxicol Lett* 204(1): 71-80.
- Manandhar, S., Cho, J.M., Kim, J.A., Kensler, T.W. and Kwak, M.K. 2007. Induction of Nrf2-regulated genes by 3H-1, 2-dithiole-3-thione through the ERK signaling pathway in murine keratinocytes. *Eur J Pharmacol* 577(1-3): 17-27.
- Markou, N., Kanakaki, M., Myrianthefs, P., Hadjiyanakos, D., Vlassopoulos, D., Damianos, A., Siamopoulos, K., Vasiliou, M. and Konstantopoulos, S. 2006. Sleep-disordered breathing in non-dialyzed patients with chronic renal failure. *Lung* 184(1): 43-49.
- Martin, R., Fitzl, G., Mozet, C., Martin, H., Welt, K. and Wieland, E. 2002. Effect of age and hypoxia/reoxygenation on mRNA expression of antioxidative enzymes in rat liver and kidneys. *Exp Gerontol* 37(12): 1481-1487.
- Milton, S.L., Dirk, L.J., Kara, L.F. and Prentice, H.M. 2008. Adenosine modulates ERK1/2, PI3K/Akt, and p38MAPK activation in the brain of the anoxia-tolerant turtle *Trachemys scripta*. *J Cereb Blood Flow Metab* 28(8): 1469-1477.
- Minet, E., Arnould, T., Michel, G., Roland, I., Mottet, D., Raes, M., Remacle, J. and Michiels, C. 2000. ERK activation upon hypoxia: involvement in HIF-1 activation. *FEBS Lett* 468(1): 53-58.
- Nicholl, D.D., Ahmed, S.B., Loewen, A.H., Hemmelgarn, B.R., Sola, D.Y., Beecroft, J.M., Turin, T.C. and Hanly, P.J. 2012. Declining Kidney Function Increases the Prevalence of Sleep Apnea and Nocturnal Hypoxia. *Chest* 141(6):1422-1430.
- Papiahgari, S., Zhang, Q., Kleeberger, S.R., Cho, H.Y. and Reddy, S.P. 2006. Hyperoxia stimulates an Nrf2-ARE transcriptional response via ROS-EGFR-PI3K-Akt/ERK MAP kinase signaling in pulmonary epithelial cells. *Antioxid Redox Signal* 8(1-2): 43-52.
- Polotsky, V.Y., Savransky, V., Bevans-Fonti, S., Reinke, C., Li, J., Grigoryev, D.N. and Shimoda, L.A. Intermittent and Sustained Hypoxia Induce a Similar Gene Expression Profile in the Human Aortic Endothelial Cells. *Physiol Genomics* 2010 Mar 2. [Epub ahead of print] PMID: 20197421.
- Samarakoon, R., Goppelt-Struebe, M. and Higgins, P.J. 2010. Linking cell structure to gene regulation: signaling events and expression controls on the model genes PAI-1 and CTGF. *Cell Signal* 22(10): 1413-1419.
- Semenza, G.L. and Prabhakar, N.R. 2007. HIF-1-dependent respiratory, cardiovascular, and redox responses to chronic intermittent hypoxia. *Antioxid Redox Signal* 9(9): 1391-1396.

*W. Sun and others*

- Sharma, R., Sharma, M., Datta, P.K. and Savin, V.J. 2002. Induction of metallothionein-I protects glomeruli from superoxide-mediated increase in albumin permeability. *Exp Biol Med* (Maywood) 227(1): 26-31.
- Shinozaki, Y., Koizumi, S., Ohno, Y., Nagao, T. and Inoue, K. 2006. Extracellular ATP counteracts the ERK1/2-mediated death-promoting signaling cascades in astrocytes. *Glia* 54(6): 606-618.
- Tkachev, V.O., Menshchikova, E.B. and Zenkov, N.K. 2011. Mechanism of the Nrf2/Keap1/ARE signaling system. *Biochemistry (Mosc)* 76(4): 407-422.
- Wang, J., Song, Y., Elsherif, L., Song, Z., Zhou, G., Prabhu, S.D., Saari, J.T. and Cai, L. 2006. Cardiac metallothionein induction plays the major role in the prevention of diabetic cardiomyopathy by zinc supplementation. *Circulation* 113(4): 544-554.
- Yang, Y.C., Lii, C.K., Lin, A.H., Yeh, Y.W., Yao, H.T., Li, C.C., Liu, K.L. and Chen, H.W. 2011. Induction of glutathione synthesis and heme oxygenase 1 by the flavonoids butein and phloretin is mediated through the ERK/Nrf2 pathway and protects against oxidative stress. *Free Radic Biol Med* 51(11): 2073-2081.
- Young, T., Palta, M., Dempsey, J., Skatrud, J., Weber, S. and Badr, S. 1993. The occurrence of sleep-disordered breathing among middle-aged adults. *N Engl J Med* 328(17): 1230-1235.
- Yuan, G., Khan, S.A., Luo, W., Nanduri, J., Semenza, G.L. and Prabhakar, N.R. 2011. Hypoxia-inducible factor 1 mediates increased expression of NADPH oxidase-2 in response to intermittent hypoxia. *J Cell Physiol* 226(11): 2925-2933.