

## VU Research Portal

### **Interpersonal psychotherapy of depression: A meta-analysis.**

Cuijpers, P.; Geraedts, A.S.; van Oppen, P.C.; Andersson, G.; Markowitz, J.C.; van Straten, A.

#### ***published in***

American Journal of Psychiatry  
2011

#### ***DOI (link to publisher)***

[10.1176/appi.ajp.2010.10101411](https://doi.org/10.1176/appi.ajp.2010.10101411)

#### ***document version***

Publisher's PDF, also known as Version of record

#### **[Link to publication in VU Research Portal](#)**

#### ***citation for published version (APA)***

Cuijpers, P., Geraedts, A. S., van Oppen, P. C., Andersson, G., Markowitz, J. C., & van Straten, A. (2011). Interpersonal psychotherapy of depression: A meta-analysis. *American Journal of Psychiatry*, 168, 581-592. <https://doi.org/10.1176/appi.ajp.2010.10101411>

#### **General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal ?

#### **Take down policy**

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

#### **E-mail address:**

[vuresearchportal.ub@vu.nl](mailto:vuresearchportal.ub@vu.nl)

## Interpersonal Psychotherapy for Depression: A Meta-Analysis

Pim Cuijpers, Ph.D.

Anna S. Geraedts, M.A.

Patricia van Oppen, Ph.D.

Gerhard Andersson, Ph.D.

John C. Markowitz, M.D.

Annemieke van Straten, Ph.D.

**Objective:** Interpersonal psychotherapy (IPT), a structured and time-limited therapy, has been studied in many controlled trials. Numerous practice guidelines have recommended IPT as a treatment of choice for unipolar depressive disorders. The authors conducted a meta-analysis to integrate research on the effects of IPT.

**Method:** The authors searched bibliographical databases for randomized controlled trials comparing IPT with no treatment, usual care, other psychological treatments, and pharmacotherapy as well as studies comparing combination treatment using pharmacotherapy and IPT. Maintenance studies were also included.

**Results:** Thirty-eight studies including 4,356 patients met all inclusion criteria. The overall effect size (Cohen's *d*) of the 16 studies that compared IPT and a control group was 0.63 (95% confidence interval [CI]=0.36 to 0.90), corresponding to a number needed to treat of 2.91. Ten stud-

ies comparing IPT and other psychological treatments showed a nonsignificant differential effect size of 0.04 (95% CI=-0.14 to 0.21; number needed to treat=45.45) favoring IPT. Pharmacotherapy (after removal of one outlier) was more effective than IPT (*d*=-0.19, 95% CI=-0.38 to -0.01; number needed to treat=9.43), and combination treatment was not more effective than IPT alone, although the paucity of studies precluded drawing definite conclusions. Combination maintenance treatment with pharmacotherapy and IPT was more effective in preventing relapse than pharmacotherapy alone (odds ratio=0.37; 95% CI=0.19 to 0.73; number needed to treat=7.63).

**Conclusions:** There is no doubt that IPT efficaciously treats depression, both as an independent treatment and in combination with pharmacotherapy. IPT deserves its place in treatment guidelines as one of the most empirically validated treatments for depression.

(*Am J Psychiatry Cuijpers et al.; AiA:1-12*)

Unipolar depressive disorders have a high prevalence (1, 2) and incidence (3), and they meaningfully impair quality of life for patients and their relatives (4, 5). Moreover, depressive disorders are linked with increased mortality rates (6), high levels of health service use, and huge economic costs (7-9). Major depression ranks fourth in disease burden worldwide, and it is expected to rank first in high-income countries by 2030 (10).

Practice guidelines recommend both pharmacological and psychological interventions for depressive disorders (11-14). Interpersonal psychotherapy (IPT) is recommended in these guidelines as one of the two psychological treatments of choice, the other being cognitive-behavioral therapy (CBT). IPT is a structured, time-limited psychological intervention based on interpersonal theory (15-18) and specifically developed for the treatment of major depression (19, 20).

Although numerous randomized controlled trials have examined the effects of IPT, only one meta-analysis has been conducted to evaluate IPT for depression (21). That analysis included a total of 13 studies and found signifi-

cant and large effects for IPT compared with placebo or no treatment, and superior effects of IPT compared with CBT. A substantial number of studies of IPT have been published since then. Furthermore, the earlier meta-analysis did not examine heterogeneity, possible effect moderators that may explain heterogeneity, publication bias, or the quality of included studies. We therefore decided to conduct a new meta-analysis to examine whether IPT is an efficacious treatment and deserves the prominent place it currently holds in treatment guidelines.

### Method

#### Search Strategy

We searched the literature using several methods. First, we used our existing database on psychological treatment of depression in adults, which is continuously updated and currently contains 1,122 full-text papers. This database, described in detail elsewhere (22), has been used in a series of earlier meta-analyses ([www.evidencebasedpsychotherapies.org](http://www.evidencebasedpsychotherapies.org)). We also conducted a comprehensive literature search (from 1966 to January 2010) in PubMed, PsycINFO, EMBASE, the Cochrane Central Register of

Controlled Trials, and Dissertation Abstracts International. We identified abstracts by combining terms indicative of psychological treatment and depression. We also collected the primary studies from 42 meta-analyses of psychological treatment for depression. Second, we carried out an additional literature search (from 1966 to January 2010) in PubMed, PsycINFO, and EMBASE to retrieve studies of IPT in adolescents. Third, we collected the primary studies from the previous meta-analysis of IPT (21) and checked the reference lists of included studies.

### Inclusion and Exclusion Criteria

We included studies of randomized controlled trials in which IPT for adults or adolescents with a unipolar depressive disorder or an elevated level of depressive symptoms was compared with waiting list, usual care, placebo, psychological treatment, pharmacological treatment, or combination treatment with IPT and pharmacotherapy. We also included maintenance studies in which patients were successfully treated during the acute treatment phase and then randomized to receive IPT or another treatment condition in the continuation phase.

We excluded studies that examined interpersonal counseling for subthreshold depression (23–26) because interpersonal counseling differs considerably from full IPT and is not intended for individuals with full-syndrome depressive disorders. No language restrictions were applied. Eligibility judgment was performed independently by two reviewers (A.S.G. and P.C.). In case of inconsistency, a third reviewer (A.v.S.) was consulted.

### Quality Assessment

We assessed the validity of the studies according to the basic criteria suggested by the *Cochrane Handbook for Systematic Reviews of Interventions* (27): adequate sequence generation (the randomization scheme was generated correctly); allocation to conditions by an independent (third) party; blinding of assessors of outcomes; completeness of follow-up data; and no selective outcome reporting. (We omitted the criterion of adequacy of random allocation concealment to respondents because concealment of random allocation is impossible in psychological treatment.)

### Analyses

We calculated effect sizes (Cohen's *d*) for all individual studies on the basis of postscore analysis using the formula  $d = (M_c - M_e) / SD_{ec}$ , where  $M_c$  is the mean of the outcome measures in the control group,  $M_e$  is the mean of the outcome measures in the experimental groups, and  $SD_{ec}$  is the pooled standard deviation of the posttest scores of the two groups (28, 29). An effect size of 0.8 can be considered large, 0.5 moderate, and 0.2 small (30). For studies that reported more than one outcome, a single pooled effect size was calculated for each study.

The individual effect sizes were pooled in the Comprehensive Meta-Analysis (CMA) software program ([www.meta-analysis.com](http://www.meta-analysis.com)). Pooled effect sizes were calculated separately for posttest comparisons of IPT with usual care, no treatment, or placebo; with other psychotherapy; and with pharmacotherapy; in addition, combination treatment with IPT and pharmacotherapy was compared with pharmacotherapy alone.

Because we expected considerable heterogeneity, we calculated pooled effect sizes with the random-effects model. However, we first tested the heterogeneity under the fixed model using the  $I^2$  statistic (31).  $I^2$  describes the variance between studies as a proportion of the total variance. A value of 25% indicates low heterogeneity, 50% moderate heterogeneity, and 75% high heterogeneity. We also report the *p* value of the *Q* statistic. A significant *Q* value rejects the null hypothesis of homogeneity.

The standardized mean difference is not easy to interpret from a clinical viewpoint. Therefore, we transformed the standardized mean differences into the numbers needed to treat, using the

formulas Kraemer and Kupfer provide (32). The number needed to treat indicates the number of patients who must receive treatment to generate an additional positive outcome in the experimental group relative to the comparison group (33).

For the IPT maintenance studies, we calculated the odds ratio of recurrence of depression in maintenance IPT compared with a control condition, as well as the number needed to treat (in this case as the reverse of the risk difference).

We also performed subgroup analyses to test for significant differences between effect sizes in different categories of studies. In these analyses, we used the mixed-effects model, which pooled studies within subgroups with the random-effects model but tested for significant differences between subgroups with the fixed-effects model. If a subgroup contained fewer than three studies, we did not conduct the subgroup analysis. Because there are indications that psychological treatments of dysthymia are less effective (33), we decided to repeat all analyses after removing studies specifically targeting patients with dysthymia.

Publication bias was examined by inspecting the funnel plot. A funnel plot is a plot of a measure of study size (the standard error) on the vertical axis as a function of effect size on the horizontal axis. Large studies appear at the top of the graph and tend to cluster near the mean effect size. Smaller studies appear toward the bottom of the graph. As there is greater sampling variation in effect size estimates in the smaller studies, they will be dispersed across a range of values (34). Visual inspection of a funnel plot can give an indication of publication bias. The studies can be expected to spread symmetrically about the pooled effect size when publication bias is absent. When bias exists, the bottom of the plot will show a higher concentration of studies on one side of the mean than the other. This is because smaller studies are more likely to be published if they have larger than average effects, which makes them more likely to meet the criterion for statistical significance.

We also examined possible publication bias using Duval and Tweedie's trim-and-fill procedure (34). If a meta-analysis has included all relevant studies, the funnel plot should be symmetric and dispersed equally on either side of the mean effect. If there is publication bias, the funnel plot will be asymmetric, with more studies to the right side of the mean effect size (studies with large effect sizes) than to the left of the mean (studies with small or nonsignificant effect sizes, which can be expected to be harder to publish). Duval and Tweedie developed a method for imputing missing studies based on the assumption that studies should be equally distributed on both sides of the mean effect size. This procedure yields an estimate of the effect size after accounting for publication bias (adjusted effect size).

### Power Calculation

We expected that several comparisons (e.g., IPT compared with pharmacotherapy, or pharmacotherapy compared with combination treatment) would involve only a limited number of studies. We therefore conducted a power calculation to assess whether the included studies had sufficient statistical power to detect small effect sizes. Although there are no clear definitions for what constitutes a small effect size, we defined a small effect as  $d=0.2$ , according to the indications of Cohen (30), but we also examined how many studies would be needed to find an effect size of 0.3.

We conducted the power calculation according to the procedures described by Borenstein and colleagues (35). A power calculation indicated that we would need to include at least 32 studies with a mean sample size of 50 (25 participants per condition) to be able to detect an effect size of 0.2 (conservatively assuming a high level of between-study variance [ $\tau^2$ ], a statistical power of 0.80, and an alpha of 0.05). Alternatively, we would need 20 studies with 80 participants apiece to detect an effect size of 0.2, or 16 studies with 100 participants. To detect an effect size of 0.3, we

would need 14 studies with 50 participants, nine studies with 80 participants, or seven studies with 100 participants.

## Results

### Selection and Inclusion of Studies

Having examined a total of 10,487 abstracts, we retrieved 1,209 full-text papers for further study. Of these, we excluded 1,171 papers that did not meet inclusion criteria (Figure 1). A total of 38 studies met all inclusion criteria and were included in this meta-analysis (36–73).

### Characteristics of Included Studies

The 38 studies included 4,356 patients (1,338 in the IPT conditions, 812 in control conditions, 713 in pharmacotherapy conditions, 468 in other psychotherapy conditions, 510 in combination treatment with IPT and pharmacotherapy, and 515 in maintenance studies). Selected characteristics of the included studies are presented in Table 1.


Thirty-three of the 38 studies examined the effects of IPT as an acute treatment, and the remaining five examined IPT as a maintenance treatment after successful recovery from a depressive disorder. Sixteen studies compared IPT with a control condition (waiting list, usual care, placebo, other), 10 compared IPT with another psychotherapy, 10 contrasted IPT with pharmacotherapy, and 10 compared a combination treatment (IPT plus pharmacotherapy) with pharmacotherapy alone. Of the 16 studies comparing IPT with a control condition, eight used usual care as the control condition, three used a waiting list control group, two used a pill placebo, and three used another control group (monthly 30-minute nontherapeutic sessions; a parenting education control program; and nonscheduled treatment).

In 29 studies patients met criteria for a depressive disorder according to a diagnostic interview (four studies specifically targeted dysthymic patients); in four studies, patients had scored above a cutoff on a depression questionnaire. Seventeen studies treated adults in general, six treated adolescents, four treated older adults, four treated patients with somatic disorders, two treated women with postpartum depression, and the remaining five treated other, more specific target groups. Fourteen studies used the original IPT manual, and 19 reported having adapted the manual to the needs of the study's target population. Adaptations were minor and included adapting the number of sessions, addressing specific needs of the target groups, and changing the individual format to a group format. The 38 studies were conducted in 13 countries, with most in the United States (N=21, including two in Puerto Rico) and Europe (N=7).

### Quality Assessment

The quality of the studies varied. Nineteen studies reported an adequate sequence generation, while the other

**FIGURE 1. Flow Chart of the Selection of Studies of Interpersonal Psychotherapy (IPT) for the Treatment of Unipolar Depression**



19 did not report a sequence generation method. Twelve studies reported allocation to conditions by an independent (third) party. Twenty-five studies reported using blinded outcome assessors, 10 did not report blinding of assessors, and three used self-report outcome measures. In 28 studies intent-to-treat analyses (completeness of follow-up data) were conducted. Nine studies (24%) met all quality criteria.

### IPT Compared With Standard or No Treatment

We were able to compare the effects of IPT with a waiting list, usual care, or placebo control condition in 16 studies (Table 2). The mean effect size (Cohen's d) was 0.63 (95% confidence interval [CI]=0.36 to 0.90), which corresponds to a number needed to treat of 2.91. Heterogeneity was high ( $I^2=82.96\%$ ). After removal of a possible outlier, the mean effect size decreased to 0.52 (95% CI=0.36 to 0.68; number needed to treat=3.50), with low to moderate heterogeneity ( $I^2=42.84\%$ ). Meta-analyses based on the two most commonly used instruments (the 17-item Hamilton Depression Rating Scale [HAM-D] and the Beck Depression Inventory [BDI]) yielded comparable outcomes (Table 2). Figure 2 presents the effect sizes and confidence intervals.

The 16 studies had on average 92 participants (46 in the IPT and 46 in the control conditions). This generated suffi-

**TABLE 1. Selected Characteristics of 38 Randomized Controlled Trials of Interpersonal Psychotherapy in Patients With Depressive Disorders**

Study Authors (Reference Number)	IPT Type <sup>a</sup>	Diagnosis (Basis of Diagnosis) <sup>b</sup>	Target Group
Bellino et al. (36)	O	Major depression (clinical judgment)	Adults with borderline personality disorder
Blom et al. (37)	O	Major depression (SCID) and HAM-D ≥14	Adult outpatients
Bodenmann et al. (38)	O	Major depression or dysthymia (SCID)	Adults over age 60 (and their partners)
Bolton et al. (39)	A	Major depression or subsyndromal depression (clinical interview)	Adults
Bolton et al. (40)	A	Major depression or subsyndromal depression (clinical interview)	Adolescent survivors of war and displacement
Browne et al. (41)	O	Dysthymia (SCID) with or without major depression	Adults
De Mello et al. (42)	A	Dysthymia (clinical interview), major depression permitted	Adults
Elkin et al. (43)	O	Major depression (RDC) and HAM-D ≥14	Adults
Finkenzeller et al. (44)	A	HAM-D ≥14	Stroke patients
Frank et al. (45)	M	Recovered from major depression (RDC) after acute treatment (PHA+weekly IPT) <sup>c</sup>	Adults
Lespérance et al. (46)	O	Major depression (SCID) and HAM-D-24 ≥20	Patients with coronary artery disease
Klerman et al. (47)	M	Clinical improvement after acute treatment (PHA <sup>c</sup> ) for neurotic depression (DSM-II)	Adults
Levkovitz et al. (48)	M	Recovered from major depression (DSM-IV) after acute treatment (PHA <sup>c</sup> )	Adults
Luty et al. (49)	O	Major depression (SCID; DSM-III-R)	Adults
Markowitz et al. (50)	A	HAM-D-24 ≥15 and clinical judgment of depressive symptoms	HIV patients
Markowitz et al. (51)	A	Dysthymia (SCID) and HAM-D-24 >13	Adults
Markowitz et al. (52)	A	Dysthymia (SCID) and HAM-D-24 >13	Adults with secondary alcohol use disorder
Marshall et al. (53)	O	Major depression (SCID) and HAM-D ≥10	Adults
Martin et al. (54)	O	Major depression (clinical judgment) and HAM-D ≥18	Adults
Mufson et al. (55)	A	HAM-D-24 ≥15	Adolescents
Mufson et al. (56)	A	Mood disorder (K-SADS)	Adolescents
Mulcahy et al. (57)	A	Major depression (DSM-IV, MCMI-III)	Women with postpartum depression
Neimeyer and Feixas (58)	O	Major depression (RDC) and BDI >15	Adults
O'Hara et al. (59)	A	Major depression (SCID) and HAM-D ≥12	Women with postpartum depression
Ransom et al. (60)	A	Major depression or dysthymia (PRIME-MD)	Rural HIV patients
Reynolds et al. (61)	A	Major depression (SADS, SCID)	Bereaved adults (age 50 or older)
Reynolds et al. (62)	M	Recovered from major depression after 16-week acute treatment (PHA + weekly IPT) <sup>c</sup>	Older adults
Reynolds et al. (63)	M	Recovered from major depression after 22-week acute treatment (PHA + weekly IPT) <sup>c</sup>	Older adults
Rosselló and Bernal (64)	A	Major depression and/or dysthymia (DISC)	Adolescents
Rosselló et al. (65)	A	Major depression (DISC) or CDI ≥13	Adolescents
Schramm et al. (66)	A	Major depression (SCID) and HAM-D ≥16	Adult inpatients
Schulberg et al. (67)	O	Major depression (DIS) and HAM-D ≥13	Adults
Sloane et al. (68)	O	Major depression (RDC) and HAM-D ≥17	Older adults
Spinelli and Endicott (69)	A	Major depression (SCID) and HAM-D-24 ≥12	Pregnant women
Swartz et al. (70)	A	Major depression (SCID) and HAM-D ≥15	Mothers whose children receive psychiatric treatment
Tang et al. (71)	A	BDI >19	Adolescents
Van Schaik et al. (72)	O	Major depression (PRIME-MD)	Older adults (age 55 or older)
Weissman et al. (73)	O	Major depression (SADS) and RTADS ≥7	Adults

<sup>a</sup> A=acute treatment using an adapted manual; M=maintenance treatment; O=acute treatment using original manual.

<sup>b</sup> BDI=Beck Depression Inventory; CDI=Children's Depression Inventory; CDRS=Cornell Dysthymia Rating Scale; CES-D=Center for Epidemiologic Studies Depression Scale; DIS=Diagnostic Interview Schedule; DISC=Diagnostic Interview Schedule for Children; EPDS=Edinburgh Postnatal Depression Scale; GDS=Geriatric Depression Scale; HADS=Hospital Anxiety and Depression Scale; HAM-D=17-item Hamilton Depression Rating Scale; HAM-D-24=24-item Hamilton Depression Rating Scale; K-SADS=Schedule for Affective Disorders and Schizophrenia for School-Age Children; MADRS=Montgomery-Åsberg Depression Rating Scale; MCMI-III=Millon Clinical Multiaxial Inventory-III; PRIME-MD=Primary Care Evaluation of Mental Disorders; RDC=Research Diagnostic Criteria; RTADS=Raskin Three Area Depression Scale; SADS=Schedule for Affective Disorders and Schizophrenia; SCID=Structured Clinical Interview for DSM-IV Axis I Disorders.

Study Interventions <sup>c</sup> (N)	Format <sup>d</sup>	Sessions (N)	ITT <sup>e</sup>	Outcome Measures <sup>b</sup>	Country <sup>f</sup>
IPT+PHA (N=20); PHA (N=19)	IND	24	N	HAM-D	IT
IPT (N=50); IPT+PHA (N=49); PHA (N=47); IPT+PLA (N=47)	IND	12	Y	HAM-D, MADRS	NL
IPT (N=20); COCT (N=20); CBT (N=20)	IND	20	Y	HAM-D, BDI	SW
IPT (N=107); UC (N=117)	GRP	16	Y	Specially developed measures for local use	UG
IPT (N=105); WL (N=104); creative play (N=105)	GRP	16	Y	Specially developed measures for local use	UG
IPT (N=178); IPT+PHA (N=212); PHA (N=196)	IND	10	N	MADRS	CA
IPT+PHA (N=16); PHA (N=19)	IND	16	N	HAM-D, MADRS	BR
IPT (N=61); CBT (N=59); PHA (N=57); PLA (N=62)	IND	16	Y	HAM-D, BDI	US
IPT (N=27); IPT+PHA (N=23); PHA (N=24)	GRP	8	Y	HAM-D, HADS	GE
M-IPT+M-PHA (N=25); M-PHA (N=28); M-IPT (N=26); M-IPT+PLA (N=26); PLA (N=23)	IND	Monthly	Y	Recurrence rate after 3 years	US
IPT+PHA (N=67); PHA (N=75); IPT+PLA (N=75); PLA (N=67)	IND	12	Y	HAM-D	CA
M-IPT+M-PHA (N=25); M-IPT+PLA (N=25); M-IPT (N=25); M-PHA (N=25); PLA (N=25); clinical management only (N=25)	IND	Weekly	Y	Recurrence rate after 8 months	US
M-IPT+M-PHA (N=7); M-PHA (N=7)	GRP	18	Y	HAM-D score at 6-month follow-up	IS
IPT (N=91); CBT (N=86)	IND	16	Y	MADRS	NZ
IPT (N=24); SUP (N=24); CBT (N=27); SUP+PHA (N=26)	IND	11	Y	BDI, HAM-D	US
IPT (N=23); IPT+PHA (N=21); SUP (N=26); PHA (N=24)	IND	16	Y	BDI, CDRS, HAM-D	US
IPT (N=14); SUP (N=12)	IND	16	Y	BDI, CDRS, HAM-D	US
IPT (N=35); CBT (N=37); PHA (N=30)	IND	16	N	HAM-D	CA
IPT (N=13); PHA (N=15)	IND	16	Y	BDI, HAM-D	UK
IPT (N=24); monitoring (N=24)	IND	12	Y	BDI, HAM-D	US
IPT (N=34); UC (N=29)	IND	12	Y	BDI, HAM-D	US
IPT (N=29); UC (N=28)	GRP	10	N	EPDS, BDI, HAM-D	AU
IPT (N=39); CBT+homework (N=32); CBT, no homework (N=31)	GRP	10	N	HAM-D	US
IPT (N=60); WL (N=60)	IND	12	Y	BDI, HAM-D	US
IPT (N=41); UC (N=38)	TEL	10	Y	BDI	US
IPT+PHA (N=16); PHA (N=25)	IND	16	Y	HAM-D	US
M-PHA+M-IPT (N=25); M-PHA (N=28); M-IPT+PLA (N=25); PLA (N=29)	IND	Monthly	Y	Recurrence rate after 3 years	US
M-PHA+M-IPT (N=28); M-PHA (N=35); M-IPT+PLA (N=35); PLA (N=18)	IND	Monthly	Y	Recurrence rate after 2 years	US
IPT (N=23); CBT (N=25); WL (N=23)	IND	12	Y	CDI	PR
IPT (N=60); CBT (N=52)	IND+ GRP	12	Y	CDI	PR
IPT+PHA (N=63); PHA (N=61)	IND+ GRP	23	Y	BDI, HAM-D	GE
IPT (N=93); PHA (N=91); UC (N=92)	IND	16	Y	HAM-D	US
IPT (N=19); PHA (N=10); PLA (N=14)	IND	6	N	BDI, HAM-D	US
IPT (N=21); parenting education (N=17)	IND	10	Y	BDI, EPDS, HAM-D	US
IPT (N=26); UC (N=21)	IND	9	N	BDI, HAM-D	US
IPT (N=35); UC (N=38)	IND	12	Y	BDI	TW
IPT (N=69); UC (N=74)	IND	8	Y	GDS, MADRS, PRIME-MD	NL
IPT (N=17); IPT+PHA (N=23); on-demand telephone support (N=21); PHA (N=20)	IND	16	N	RTADS	US

<sup>c</sup> CBT=cognitive-behavioral therapy; COCT=coping-oriented couples therapy; IPT=interpersonal psychotherapy; PHA=pharmacotherapy; PLA=placebo; SUP=supportive psychotherapy; UC=usual care; WL=waiting list.

<sup>d</sup> GRP=Group treatment; IND=individual treatment; TEL=telephone treatment.

<sup>e</sup> ITT=intent to treat; N=no; Y=yes.

<sup>f</sup> AU=Australia; BR=Brazil; CA=Canada; GE=Germany; IS=Israel; IT=Italy; NL=Netherlands; NZ=New Zealand; PR=Puerto Rico; SW=Sweden; TW=Taiwan; UG=Uganda; UK=United Kingdom; US=United States.

**TABLE 2. Effect Sizes in Meta-Analysis of Studies Comparing Interpersonal Psychotherapy With Control Conditions, Pharmacotherapy, and Other Psychotherapies<sup>a</sup>**

Comparison	Number of Comparisons	d	95% CI	Heterogeneity <sup>b</sup>	p <sup>c</sup>	Number Needed to Treat
<b>IPT versus control condition</b>						
All studies	16	0.63***	0.36 to 0.90	82.96***		2.91
Outlier removed <sup>d</sup>	15	0.52***	0.36 to 0.68	42.84*		3.50
HAM-D only	9	0.60***	0.39 to 0.81	40.47		3.05
BDI only	7	0.60***	0.32 to 0.89	59.62*		3.05
Subgroup analyses <sup>e</sup>						
Adapted manual						
No	5	0.29	0.12 to 0.47	0.00	0.004	6.17
Yes	10	0.67	0.48 to 0.85	27.52		2.75
Target group						
Adults in general	3	0.37**	0.15 to 0.58	0.00	0.214	4.85
Adolescents	5	0.63***	0.43 to 0.82	0.00		2.91
Specific target group	7	0.53**	0.19 to 0.87	68.11**		3.42
Basis of diagnosis						
Diagnostic interview	12	0.57***	0.40 to 0.73	29.74	0.424	3.18
Self-report	3	0.38	-0.04 to 0.80	67.73*		4.72
Control group						
Usual care	6	0.50***	0.28 to 0.72	32.20	0.182	3.62
Waiting list	3	0.81***	0.42 to 1.20	59.63		2.30
Other	5	0.38**	0.14 to 0.62	0.00		4.72
Intent-to-treat analysis						
Yes	10	0.57***	0.38 to 0.76	51.56*	0.718	3.18
No	4	0.50**	0.16 to 0.84	12.97		3.62
<b>IPT versus other psychotherapy<sup>f</sup></b>						
All studies	13	0.04	-0.14 to 0.21	39.81		45.45
One effect size per study (highest)	10	0.01	-0.20 to 0.21	44.59		166.67
One effect size per study (lowest)	10	0.00	-0.20 to 0.19	41.11		-
HAM-D only	10	0.12	-0.07 to 0.31	19.04		14.71
BDI only	7	0.28	-0.02 to 0.57	45.46		6.41
Studies on dysthymia excluded	11	0.01	-0.18 to 0.20	42.09		166.67
Subgroup analyses <sup>e</sup>						
Adapted manual						
Yes	7	-0.09	-0.25 to 0.07	0.00	0.098	20.00
No	6	0.25	-0.12 to 0.63	62.27*		7.14
Comparison group						
CBT	9	-0.04	-0.23 to 0.16	39.72	0.113	45.45
Other psychotherapy	4	0.27	-0.06 to 0.59	4.68		6.58
Target group						
Adults	8	-0.08	-0.24 to 0.08	0.00	0.116	21.74
More specific group	5	0.31	-0.15 to 0.77	69.51*		5.75
Intent-to-treat analysis						
Yes	9	0.16	-0.06 to 0.38	35.74	0.106	11.11
No	3	-0.13	-0.40 to 0.15	0.00		13.51
<b>IPT versus pharmacotherapy<sup>f</sup></b>						
All studies	10	-0.12	-0.36 to 0.12	61.98**		14.71
Outlier removed <sup>g</sup>	9	-0.19*	-0.38 to -0.01	30.95		9.43
HAM-D only	8	-0.16	-0.43 to 0.11	57.39*		11.11
BDI only	3	-0.31	-0.72 to 0.09	40.39		5.75
Studies on dysthymia excluded	7	-0.12	-0.33 to 0.09	23.78		14.71
Subgroup analyses <sup>e</sup>						
Medication <sup>h</sup>						
SSRI or SNRI	3	-0.39**	-0.68 to -0.11	14.18	0.037	4.59
Tricyclic antidepressant	4	-0.02	-0.22 to 0.19	0.00		83.33
Intent-to-treat analysis						
Yes	5	-0.15	-0.42 to 0.13	45.10	0.450	11.90
No	4	-0.28*	-0.50 to -0.06	0.00		6.41

**TABLE 2. Effect Sizes in Meta-Analysis of Studies Comparing Interpersonal Psychotherapy With Control Conditions, Pharmacotherapy, and Other Psychotherapies<sup>a</sup> (continued)**

Comparison	Number of Comparisons	d	95% CI	Heterogeneity <sup>b</sup>	p <sup>c</sup>	Number Needed to Treat
<b>Pharmacotherapy versus combination treatment<sup>f</sup></b>						
All studies	10	0.16	-0.03 to 0.36	39.26		11.11
HAM-D only	8	0.21	-0.11 to 0.52	64.64**		8.47
BDI only	2	0.10	-0.20 to 0.41	0.00		17.86
Studies on dysthymia excluded	7	0.22	-0.03 to 0.47	44.17		8.06
<b>Subgroup analyses</b>						
<b>Adapted manual</b>						
Yes	5	0.23	-0.04 to 0.51	50.15	0.494	7.69
No	5	0.08	-0.25 to 0.41	40.50		21.74
<b>Medication</b>						
SSRI	5	0.07	-0.27 to 0.41	64.96*	0.562	25.00
Tricyclic antidepressant	2	0.33	-0.11 to 0.76	0.00		5.43
Other/protocol	3	0.27	-0.00 to 0.54	0.00		6.58
<b>Target group</b>						
Adults	5	0.07	-0.12 to 0.27	0.00	0.569	25.00
More specific group	5	0.19	-0.16 to 0.54	61.35*		9.43
<b>Intent-to-treat analysis</b>						
Yes	6	0.12	-0.10 to 0.34	23.03	0.391	14.71
No	4	0.35	-0.12 to 0.82	63.84*		5.10

<sup>a</sup> These data are from analysis using the random effects model. BDI=Beck Depression Inventory; HAM-D=Hamilton Depression Rating Scale; IPT=interpersonal psychotherapy; SNRI=serotonin/norepinephrine reuptake inhibitor; SSRI=selective serotonin reuptake inhibitor.

<sup>b</sup> Variance between studies as a proportion of the total variance; heterogeneity tested using the I<sup>2</sup> statistic. Low heterogeneity=25%; moderate heterogeneity=50%; high heterogeneity=75%. The p values indicated in this column refer to whether the Q statistic is significant (the I<sup>2</sup> statistic does not include a test of significance).

<sup>c</sup> The p values in this column indicate whether the difference between the effect sizes in the subgroups is significant.

<sup>d</sup> Bolton et al. (39).

<sup>e</sup> Subgroup analyses are conducted without outliers.

<sup>f</sup> Positive effect sizes indicate superior effects of IPT.

<sup>g</sup> Finkenzeller et al. (44).

<sup>h</sup> Two studies in which other pharmacotherapies were used (nefazodone, protocolized treatment) were excluded from these analyses.

<sup>i</sup> Positive effect sizes indicate superior effects of combination treatment.

\*p<0.05. \*\*p<0.01. \*\*\*p<0.001.

cient statistical power to detect an effect size of 0.21 (number needed to treat=8.47).

Inspection of the funnel plot and Duval and Tweedie's trim and fill procedure did not indicate possible publication bias: the effect size adjusted for publication bias exactly equaled the unadjusted effect size. None of the studies in this subsample specifically targeted dysthymic disorder.

We examined some basic moderators in subgroup analyses. We found no indication that type of target group (adults, adolescents, more specific target group), method of diagnosing depressive disorder (diagnostic interview or other), or use of intent-to-treat analyses (yes or no) were significantly associated with effect size, although the number of studies was small in several subgroups. We did find that the studies using the original manual produced significantly lower effect sizes (d=0.29; number needed to treat=6.17) than did studies that used an adapted manual (d=0.67; number needed to treat=2.75) (p<0.01). Heterogeneity was low and not significant in both subgroups.

Studies that used a waiting list control group yielded larger effect sizes than studies that employed usual care or other control conditions. That this difference was not sta-

tistically significant (p>0.05) may reflect the small number of studies using a waiting list control group (N=3).

### ***IPT Compared With Other Psychotherapies***

Ten studies (13 comparisons) compared posttest effects of IPT to another psychotherapy (Table 2; see also Figure S1 in the data supplement that accompanies the online edition of this article). On average, the 13 comparisons included 74 patients (37 per condition), which sufficed to detect an effect size of 0.25 (number needed to treat=7.14).

The overall effect size for the 13 comparisons was 0.04 (95% CI=-0.14 to 0.21; number needed to treat=45.45) favoring IPT, which was not statistically significant (p=0.40). Heterogeneity was low to moderate (I<sup>2</sup>=39.81%).

In these analyses we included three studies that compared two psychological treatments with the same control group. This means that multiple comparisons from these three studies were included in the same analysis. The multiple comparisons, however, are not independent of one another, which may have resulted in artificially reduced heterogeneity and affected the pooled effect size. We examined such possible effects by conducting an analysis in which we included only one effect size per study. First, we



included only the comparison with the largest effect size from the studies with multiple comparisons. Then, in another analysis, we included only the smallest effect size. As illustrated in Table 2, the resulting effect sizes were almost identical to those of the overall analyses. Heterogeneity did not increase considerably and remained moderate to high in these analyses.

The mean effect size based on the HAM-D resulted in comparable outcomes. The effect size based on the BDI was somewhat larger ( $d=0.28$ ; number needed to treat=6.41) but did not reach statistical significance. Removal of the studies specifically treating dysthymic disorder had little effect on the overall effect size. There were no outlier studies.

Again, we examined possible moderators in subgroup analyses (Table 2). We did not find that the effect size differed significantly between studies that used an adapted manual and those that used the original manual; between studies in which IPT was compared with CBT and those in which IPT was compared with other psychotherapies; between studies treating adults in general and those targeting more specific groups (e.g., adolescents, people with somatic illnesses); between studies using different treatment formats; and between studies using intent-to-treat analyses and those using per-protocol analyses.

There were some indications for publication bias. After adjustment for publication bias, the effect size decreased to  $-0.11$  (95% CI= $-0.31$  to  $0.09$ ; number of imputed studies=4; number needed to treat=16.13).

### ***IPT Compared With Pharmacotherapy***

Ten studies compared IPT with pharmacotherapy. These studies included on average 82 participants and had sufficient power to detect an effect size of 0.28 (number needed to treat=6.41).

A nonsignificant differential overall effect size of  $-0.12$  (95% CI= $-0.36$  to  $0.12$ ; number needed to treat=14.71) favored pharmacotherapy (see Figure S2 in the online data supplement). Heterogeneity was moderate to high ( $I^2=61.98\%$ ). After removal of one possible outlier (Finkenzeller et al. [44]), the overall effect size became significant ( $d=-0.19$ ; 95% CI= $-0.38$  to  $-0.01$ ; number needed to treat=9.43,  $p<0.05$ ;  $I^2=30.95\%$ ), indicating a significant superior effect of pharmacotherapy. Removal of the studies aimed at dysthymia resulted in comparable outcomes.

Subgroup analyses indicated that selective serotonin reuptake inhibitors (SSRIs) were significantly more effective than IPT ( $N=3$ ;  $d=-0.39$ ,  $p<0.01$ ; number needed to treat=4.59), whereas tricyclic antidepressants were not ( $N=4$ ;  $d=-0.02$ ,  $p>0.1$ ; number needed to treat=83.33), and the studies comparing SSRIs with IPT differed significantly from those examining tricyclics ( $p<0.05$ ). Two of the three studies comparing IPT with SSRIs, however, involved patients with dysthymia. Furthermore, the number of studies in each of these subgroups was very small, so these results should be considered with caution.

These analyses gave no indication of publication bias; the unadjusted and adjusted effect sizes were identical, with zero imputed studies.

### ***IPT Compared With Combination Treatment***

In 10 studies the combination of IPT and pharmacotherapy was compared with pharmacotherapy alone. These 10 comparisons had on average 80 participants, which yielded enough statistical power to detect an effect size of 0.28 (number needed to treat=6.41). The mean effect size indicating the difference between these two types of treatment was 0.16 (95% CI= $-0.03$  to  $0.36$ ; number needed to treat=11.11) in favor of combination treatment (see Figure S3 in the online data supplement). This difference was not statistically significant, perhaps reflecting the small number of studies and consequent low statistical power. Heterogeneity was low to moderate ( $I^2=39.26\%$ ). Removing the studies on dysthymic disorder did not yield a significant difference either. The subgroup analyses (original manual versus adapted manual; adults versus more specific group; intent-to-treat analyses versus per-protocol analyses) identified no significant differences between subgroups. There were some indications for publication bias: the effect size adjusted for publication bias was somewhat smaller than the unadjusted effect size ( $d=0.07$ ; 95% CI= $-0.13$  to  $0.27$ ; number of imputed studies=3; number needed to treat=25.00).

### ***IPT as Maintenance Treatment***

We were able to compare maintenance pharmacotherapy in patients who had recovered from a depressive disorder with combination treatment with IPT and pharmacotherapy maintenance treatment in five studies. Four of these presented recurrence rates. The fifth study reported only means and standard deviations for patients; in this study, the odds ratio was calculated using the procedures integrated in the CMA software. This resulted in an odds ratio of 122.77, which was considered incredible (the other odds ratios ranged from 1 to 3.75). We therefore did not use the study in these analyses. The remaining four studies generated an odds ratio of 0.37 (95% CI=0.19 to 0.73,  $p<0.01$ ;  $I^2=0\%$ ; number needed to treat=7.63), which indicates that maintenance IPT combined with pharmacotherapy significantly reduced the recurrence rate compared with pharmacotherapy alone after successful treatment of acute depression (see Figure S4 in the online data supplement). Because of the small number of studies, we did not conduct additional analyses.

We were also able to compare the combination of maintenance IPT and pill placebo with pill placebo alone in four studies. The resulting odds ratio was 0.47 (95% CI=0.25 to 0.87;  $I^2=0\%$ ; number needed to treat=5.95), indicating that maintenance IPT was more protective against relapse than pill placebo alone. As none of the other possible comparisons (maintenance IPT versus control groups; maintenance IPT versus pharmacotherapy) had more than two comparisons, we decided not to perform a meta-analysis.

## Discussion

We identified 38 randomized trials (with a total of 4,356 patients) examining the effects of IPT. Compared with control groups, we found a moderate to large effect of IPT in the acute treatment of depression. We also found some indications that IPT had less efficacy than SSRI pharmacotherapy. However, the overall difference was small, not all analyses were significant, and the number of studies in this subsample was small.

We found indications that combination treatment with IPT and pharmacotherapy was somewhat more efficacious than pharmacotherapy alone, although this difference reached significance only when the HAM-D was used as an outcome measure. However, the effect size was also small, and again this subsample of studies was relatively small, limiting statistical power. In a larger meta-analysis of studies comparing combination treatments with psychotherapy and pharmacotherapy and pharmacotherapy alone, we found that combination treatments were significantly better than pharmacotherapy alone (74). Furthermore, combination treatment may have greater efficacy for patients with more severe or chronic major depression (75).

We did not find that IPT had greater efficacy than other psychotherapies, including CBT, although the number of studies was too small to draw definite conclusions. IPT is the only type of psychotherapy for depression aside from CBT that has been compared with control groups, other psychotherapies, antidepressant medication, and combination treatments. Other types of psychotherapy are far less well examined. At present, therefore, IPT and CBT may be considered the best options for psychological treatments for depression. There is no indication that IPT is superior to CBT, and the two seem equally effective overall. Whether to prescribe CBT or IPT should depend on patient preference and moderating factors based on differential therapeutics (76).

Although the number of studies examining the effects of maintenance IPT was small, these studies had relatively high methodological quality. Analysis of these studies indicated that maintenance IPT combined with pharmacotherapy reduced the relapse rate considerably compared with pharmacotherapy alone. We also found that placebo plus IPT was more effective than placebo alone in reducing relapse rates.

Our finding that pharmacotherapy has greater efficacy than IPT is consistent with results from earlier meta-analytic research (77); this should not come as a surprise, since there was considerable overlap of included studies. The superior effect of combination treatment over pharmacotherapy alone also accords with earlier meta-analytic research (74). This suggests that IPT has an additional effect on depression beyond the effects of pharmacotherapy, although the effect size was small.

That IPT is efficacious compared with control conditions and probably augments pharmacotherapy is impor-

tant. Pharmacotherapy may have limited benefit in situations such as complicated grief, where IPT can be crucial. Medication and psychotherapies presumably work by different mechanisms, and they generally relieve symptoms in different temporal patterns. Effective psychotherapies such as IPT are therefore among the most important instruments available to clinicians.

We found that studies using the original IPT manual produced significantly lower effect sizes than studies that used an adapted manual. This may reflect the fact that the original manual has been examined by several groups other than the inventors of IPT, whereas the adapted versions of the manual have been examined mainly by the researchers who developed them. Yet the original manual would inevitably have received more use than later adaptations, and differences in outcomes might also derive from different treatment populations, therapist skills, and the adaptations themselves. Nonetheless, the larger effect sizes of the adapted versions should be considered with caution.

This study has several limitations. First, the quality of the included studies was not optimal. Only nine of 38 (24%) studies met all quality criteria. Although this proportion is relatively high compared with other studies of psychotherapy for depression (we previously found [77] that only 11 of 115 [10%] controlled trials of psychotherapy for adult depression met all quality criteria), it is still too low. We recommend that future research use and report adequate randomization methods, correct blinding of outcome assessors, and intent-to-treat analyses.

Second, the number of studies in several subanalyses was relatively small and may have lacked statistical power to detect smaller effect sizes. A third limitation is that we found indications for publication bias in some analyses, although the mean effect sizes did not decrease considerably after adjustment for publication bias, and none of the resulting effect sizes differed significantly from the unadjusted effect sizes. This contradicts a recent meta-analysis of publication bias in psychotherapy for adult depression (78), which found no indications for publication bias of IPT studies. The present analysis included more studies, and its results can be assumed to be more up-to-date.

Despite these limitations, we found clear indications for the efficacy of IPT for unipolar depression. IPT is one of the best empirically validated psychological treatments for depression currently available, and its inclusion in treatment guidelines is justified.

---

Received Oct. 1, 2010; revision received Oct. 24, 2010; accepted Dec. 13, 2010 (doi: 10.1176/appi.ajp.2010.10101411). From the Department of Clinical Psychology and the Department of Psychiatry, Vrije Universiteit (VU University) Amsterdam; EMGO Institute for Health and Care Research, VU University Amsterdam and VU University Medical Center Amsterdam; Department of Behavioral Sciences and Learning, Linköping University, Linköping, Sweden; Department of Clinical Neuroscience, Psychiatry Section, Karolinska Institutet, Stockholm; New York State Psychiatric Institute; and Department of Psychiatry, Columbia University College of Physicians and Surgeons, New York. Address correspondence and reprint requests to Dr. Cuijpers, Department of

Clinical Psychology, VU University Amsterdam, Van der Boechorststraat 1, 1081 BT Amsterdam; p.cuijpers@psy.vu.nl (e-mail).

Dr. Markowitz receives funding for an IPT-related study from the National Institute of Mental Health and minor royalties from books on psychotherapy, including IPT, and receives a stipend from Elsevier Press as associate editor of a journal. The other authors report no financial relationships with commercial interests.

References

1. Kessler RC, McGonagle KA, Zhao S, Nelson CB, Hughes M, Eshleman S, Wittchen HU, Kendler KS: Lifetime and 12-month prevalence of DSM-III-R psychiatric disorders in the United States: results from the National Comorbidity Survey. *Arch Gen Psychiatry* 1994; 51:8–19
2. Alonso J, Angermeyer MC, Bernert S, Bruffaerts R, Brugha TS, Bryson H, de Girolamo G, Graaf R, Demyttenaere K, Gasquet I, Haro JM, Katz SJ, Kessler RC, Kovess V, Lépine JP, Ormel J, Polidori G, Russo LJ, Vilagut G, Almansa J, Arbabzadeh-Bouchez S, Autonell J, Bernal M, Buist-Bouwman MA, Codony M, Domingo-Salvany A, Ferrer M, Joo SS, Martínez-Alonso M, Matschinger H, Mazzi F, Morgan Z, Morosini P, Palacín C, Romera B, Taub N, Vollebergh WA; ESEMeD/MHEDEA 2000 Investigators, European Study of the Epidemiology of Mental Disorders (ESEMeD) Project: Prevalence of mental disorders in Europe: results from the European Study of the Epidemiology of Mental Disorders (ESEMeD) project. *Acta Psychiatr Scand Suppl* 2004; 109:S21–S27
3. Waraich P, Goldner EM, Somers JM, Hsu L: Prevalence and incidence studies of mood disorders: a systematic review of the literature. *Can J Psychiatry* 2004; 49:124–138
4. Ustun TB, Ayuso-Mateos JL, Chatterji S, Mathers C, Murray CJL: Global burden of depressive disorders in the year 2000. *Br J Psychiatry* 2004; 184:386–392
5. Saarni SI, Suvisaari J, Sintonen H, Pirkola S, Koskinen S, Aromaa A, Lönnqvist J: Impact of psychiatric disorders on health-related quality of life: general population survey. *Br J Psychiatry* 2007; 190:326–332
6. Cuijpers P, Smit F: Excess mortality in depression: a meta-analysis of community studies. *J Affect Disord* 2002; 72:227–236
7. Berto P, D’Ilario D, Ruffo P, Di Virgilio RF: Depression: cost-of-illness studies in the international literature: a review. *J Ment Health Pol Econ* 2000; 3:3–10
8. Greenberg PE, Birnbaum HG: The economic burden of depression in the US: societal and patient perspectives. *Exp Opin Pharmacother* 2005; 6:369–376
9. Smit F, Cuijpers P, Oostenbrink J, Batelaan N, de Graaf R, Beekman A: Excess costs of common mental disorders: population-based cohort study. *J Ment Health Pol Econ* 2006; 9:193–200
10. Mathers CD, Loncar D: Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* 2006; 3:e442
11. American Psychiatric Association: Practice guideline for the treatment of patients with major depressive disorder, 2nd ed. *Am J Psychiatry* 2000; 157(Apr suppl)
12. Asberg M, Bengtsson F, Hagberg B, Henriksson F, Jonsson B, Karlsson I, Kimbre S, von Knorring AL, Krakau I, Mathe A, Martensson B, Ornander H, Thelander S: Treatment of Depression: A Systematic Review. Stockholm, Swedish Council on Technology Assessment in Health Care, 2004
13. National Institute for Clinical Excellence (NICE): Depression: Management of Depression in Primary and Secondary Care (National Clinical Practice Guideline 23). London, NICE, 2004, amended 2007
14. Ellis P; Royal Australian and New Zealand College of Psychiatrists Clinical Practice Guidelines Team for Depression: Australian and New Zealand clinical practice guidelines for the treatment of depression. *Aust NZ J Psychiatry* 2004; 38:389–407
15. Bowlby J: Attachment and Loss, vol 2, Separation: Anxiety and Anger. New York, Basic Books, 1973
16. Klerman GL, Weissman MM, Rounsaville BJ, Chevron ES: Interpersonal Psychotherapy of Depression. New York, Basic Books, 1984
17. Meyer A: Psychobiology: A Science of Man. Springfield, Ill, Charles C Thomas, 1957
18. Sullivan HS: The Interpersonal Theory of Psychiatry. New York, WW Norton, 1953
19. Weissman MM, Markowitz JW, Klerman GL: Comprehensive Guide to Interpersonal Psychotherapy. New York, Basic Books, 2000
20. Markowitz JC, Svartberg M, Swartz HA: Is IPT time-limited psychodynamic psychotherapy? *J Psychother Pract Res* 1998; 7:185–195
21. De Mello MF, de Jesus Mari J, Bacaltchuk J, Verdelli H, Neugebauer R: A systematic review of research findings on the efficacy of interpersonal therapy for depressive disorders. *Eur Arch Psychiatry Clin Neurosci* 2005; 255:75–82
22. Cuijpers P, van Straten A, Warmerdam L, Andersson G: Psychological treatment of depression: a meta-analytic database of randomized studies. *BMC Psychiatry* 2008; 8:36
23. Forsyth KM: The Design and Implementation of a Depression Prevention Program (doctoral dissertation). Kingston, RI, University of Rhode Island, Department of Psychology, 2000
24. Mossey JM, Knott KA, Higgins M, Talerico K: Effectiveness of a psychosocial intervention, interpersonal counselling, for subsyndromal depression in medically ill elderly. *J Gerontol A Biol Sci Med Sci* 1996; 51:M172–M178
25. Neugebauer R, Kline J, Markowitz JC, Bleiberg KL, Baxi L, Rosing MA, Levin B, Keith J: Pilot randomized controlled trial of interpersonal counseling for subsyndromal depression following miscarriage. *J Clin Psychiatry* 2006; 67:1299–1304
26. Young JF, Mufson L, Davies M: Efficacy of interpersonal psychotherapy-adolescent skills training: an indicated preventive intervention for depression. *J Child Psychol Psychiatry* 2006; 47:1254–1262
27. Higgins JPT, Green S (eds): Cochrane Handbook for Systematic Reviews of Interventions, 5.0.2. London, UK, Cochrane Collaboration, 2009 ([www.cochrane-handbook.org](http://www.cochrane-handbook.org))
28. Cooper H, Hedges LV: The Handbook of Research Synthesis. New York, Russell Sage Foundation Press, 1994
29. Hedges LV, Olkin I: Statistical Methods for Meta-Analysis. San Diego, Academic Press, 1985
30. Cohen J: Statistical Power Analysis for the Behavioral Sciences, 2nd ed. Hillsdale, NJ, Erlbaum, 1988
31. Higgins JPT, Thompson SG, Deeks JJ, Altman DG: Measuring inconsistency in meta-analyses. *BMJ* 2003; 327:557–560
32. Kraemer HC, Kupfer DJ: Size of treatment effects and their importance to clinical research and practice. *Biol Psychiatry* 2006; 59:990–996
33. Cuijpers P, van Straten A, Schuurmans J, van Oppen P, Hollon SD, Andersson G: Psychotherapy for chronic major depression and dysthymia: a meta-analysis. *Clin Psychol Rev* 2010; 30:51–62
34. Duval DL, Tweedie R: Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 2000; 56:455–463
35. Borenstein M, Hedges LV, Higgins JPT, Rothstein HR: Introduction to Meta-Analysis. Chichester, UK, John Wiley & Sons, 2009
36. Bellino S, Zizza M, Rinaldi C, Bogetto F: Combined treatment of major depression in patients with borderline personality disorder: a comparison with pharmacotherapy. *Can J Psychiatry* 2006; 51:453–460
37. Blom MJB, Jonker K, Dusseldorp E, Spinhoven P, Hoencamp E, Haffmans J, van Dyck R: Combination treatment for acute depression is superior only when psychotherapy is added to medication. *Psychother Psychosom* 2007; 76:289–297

38. Bodenmann G, Plancherel B, Beach SRH, Widmer K, Gabriel B, Meuwly N, Charvoz L, Hautzinger M, Schramm E: Effects of coping-oriented couples therapy on depression: a randomized clinical trial. *J Consult Clin Psychol* 2008; 76:944–954
39. Bolton P, Bass J, Neugebauer R, Verdelli H, Clougherty KF, Wickramaratne P, Speelman L, Ndogoni L, Weissman M: Group interpersonal psychotherapy for depression in rural Uganda. *JAMA* 2003; 289:3117–3124
40. Bolton P, Bass J, Betancourt T, Speelman L, Onyango G, Clougherty KF, Neugebauer R, Murray L, Verdelli H: Interventions for depression symptoms among adolescent survivors of war and displacement in northern Uganda: a randomized controlled trial. *JAMA* 2007; 298:519–527
41. Browne G, Steiner M, Roberts J, Gafni A, Byrne C, Dunn E, Bell B, Mills M, Chalklin L, Wallik D, Kraemer J: Sertraline and/or interpersonal psychotherapy for patients with dysthymic disorder in primary care: 6-month comparison with longitudinal 2-year follow-up of effectiveness and costs. *J Affect Dis* 2002; 68:317–330
42. De Mello MF, Myczcowski LM, Menezes PR: A randomized controlled trial comparing moclobemide and moclobemide plus interpersonal psychotherapy in the treatment of dysthymic disorder. *J Psychother Pract Res* 2001; 10:117–123
43. Elkin I, Shea MT, Watkins JT, Imber SD, Sotsky SM, Collins JF, Glass DR, Pilkonis PA, Leber WR, Docherty JP, Fiester SJ, Parloff MB: National Institute of Mental Health Treatment of Depression Collaborative Research Program: general effectiveness of treatments. *Arch Gen Psychiatry* 1989; 46:971–982
44. Finkenzeller W, Zobel I, Rietz S, Schramm E, Berger M: Interpersonelle Psychotherapie und Pharmakotherapie bei Post-Stroke-Depression: Machbarkeit und Effektivität (Interpersonal psychotherapy and pharmacotherapy for post-stroke depression: feasibility and effectiveness). *Nervenarzt* 2009; 80:805–812
45. Frank E, Kupfer DJ, Perel JM, Cornes C, Jarrett DB, Mallinger AG, Thase ME, McEachran AB, Grochocinski VJ: Three-year outcomes for maintenance therapies in recurrent depression. *Arch Gen Psychiatry* 1990; 47:1093–1099
46. Lespérance F, Frasur-Smith N, Koszycki D, Laliberté MA, van Zyl LT, Baker B, Swenson JR, Ghatavi K, Abramson BL, Dorian P, Guertin MC: Effects of citalopram and interpersonal psychotherapy on depression in patients with coronary artery disease: the Canadian Cardiac Randomized Evaluation of Antidepressant and Psychotherapy Efficacy (CREATE) trial. *JAMA* 2007; 297:367–379
47. Klerman GL, DiMascio A, Weissman M, Prusoff B, Paykel ES: Treatment of depression by drugs and psychotherapy. *Am J Psychiatry* 1974; 131:186–191
48. Levkovitz Y, Shahar G, Native G, Hirsfeld E, Treves I, Krieger I, Fennig S: Group interpersonal psychotherapy for patients with major depression disorder: pilot study. *J Affect Dis* 2000; 60:191–195
49. Luty SE, Carter JD, McKenzie JM, Rae AM, Frampton CM, Mulder RT, Joyce PR: Randomised controlled trial of interpersonal psychotherapy and cognitive-behavioural therapy for depression. *Br J Psychiatry* 2007; 190:496–502
50. Markowitz JC, Kocsis JH, Fishman B, Spielman LA, Jacobsberg LB, Frances AJ, Klerman GL, Perry SW: Treatment of depressive symptoms in human immunodeficiency virus-positive patients. *Arch Gen Psychiatry* 1998; 55:452–457
51. Markowitz JC, Kocsis JH, Bleiberg KL, Christos PJ, Sacks M: A comparative trial of psychotherapy and pharmacotherapy for “pure” dysthymic patients. *J Affect Dis* 2005; 89:167–175
52. Markowitz JC, Kocsis JH, Christos P, Bleiberg K, Karlin C: Pilot study of interpersonal psychotherapy for dysthymic patients with secondary alcohol abuse or dependence. *J Nerv Ment Dis* 2008; 196:468–474
53. Marshall MB, Zuroff DC, McBride C, Bagby RM: Self-criticism predicts differential response to treatment for major depression. *J Clin Psychology* 2008; 64:231–244
54. Martin SD, Martin E, Rai SS, Richardson MA, Royall R, Eng C: Brain blood flow changes in depressed patients treated with interpersonal psychotherapy or venlafaxine hydrochloride: preliminary findings. *Arch Gen Psychiatry* 2001; 58:641–648
55. Mufson L, Weissman MM, Moreau D, Garfinkel R: Efficacy of interpersonal psychotherapy for depressed adolescents. *Arch Gen Psychiatry* 1999; 56:573–579
56. Mufson L, Pollack DK, Wickramaratne P, Nomura Y, Olfson M, Weissman MM: A randomized effectiveness trial of interpersonal psychotherapy for depressed adolescents. *Arch Gen Psychiatry* 2004; 61:577–584
57. Mulcahy R, Reay RE, Wilkinson RB, Owen C: A randomised control trial for the effectiveness of group interpersonal psychotherapy for postnatal depression. *Arch Womens Ment Health* 2010; 13:125–139
58. Neimeyer RA, Feixas G: The role of homework and skill acquisition in the outcome of group cognitive therapy for depression. *Behav Ther* 1990; 21:281–292
59. O'Hara MW, Stuart S, Gorman LL, Wenzel A: Efficacy of interpersonal psychotherapy for postpartum depression. *Arch Gen Psychiatry* 2000; 57:1039–1045
60. Ransom D, Heckman TG, Anderson T, Garske J, Holroyd K, Basta T: Telephone-delivered interpersonal psychotherapy for HIV-infected rural persons with depression: a pilot trial. *Psychiatr Serv* 2008; 59:871–877
61. Reynolds CF III, Miller MD, Pasternak RE, Frank E, Perel JM, Cornes C, Houck PR, Mazumdar S, Dew MA, Kupfer DJ: Treatment of bereavement-related major depressive episodes in later life: a controlled study of acute and continuation treatment with nortriptyline and interpersonal psychotherapy. *Am J Psychiatry* 1999; 156:202–208
62. Reynolds CF III, Frank E, Perel JM, Imber SD, Cornes C, Miller MD, Mazumdar S, Houck PR, Dew MA, Stack JA, Pollock BG, Kupfer DJ: Nortriptyline and interpersonal psychotherapy as maintenance therapies for recurrent major depression: a randomized controlled trial in patients older than 59 years. *JAMA* 1999; 281:39–45
63. Reynolds CF III, Dew MA, Pollock BG, Mulsant BH, Frank E, Miller MD, Houck PR, Mazumdar S, Butters MA, Stack JA, Schlernitzauer MA, Whyte EM, Gildengers A, Karp J, Lenze E, Szanto K, Bensasi S, Kupfer DJ: Maintenance treatment of major depression in old age. *N Engl J Med* 2006; 354:1130–1138
64. Rosselló J, Bernal G: The efficacy of cognitive-behavioral and interpersonal treatments for depression in Puerto Rican adolescents. *J Consult Clin Psychol* 1999; 67:734–745
65. Rosselló J, Bernal G, Rivera-Medina C: Individual and group CBT and IPT for Puerto Rican adolescents with depressive symptoms. *Cult Div Ethn Minor Psychol* 2008; 14:234–245
66. Schramm E, van Calker D, Dykieriek P, Lieb K, Kech S, Zobel I, Leonhart R, Berger M: An intensive treatment program of interpersonal psychotherapy plus pharmacotherapy for depressed inpatients: acute and long-term results. *Am J Psychiatry* 2007; 164:768–777
67. Schulberg HC, Block MR, Madonia MJ, Scott CP, Rodriguez E, Imber SD, Perel J, Lave J, Houck PR, Coulehan JL: Treating major depression in primary care practice: eight-month clinical outcomes. *Arch Gen Psychiatry* 1996; 53:913–919
68. Sloane RB, Staples FR, Schneider LS: Interpersonal therapy versus nortriptyline for depression in the elderly, in *Clinical and Pharmacological Studies in Psychiatric Disorders*. Edited by Burrows G, Norman TR, Dermerstein L. London, John Libbey, 1985, pp 344–346

69. Spinelli MG, Endicott J: Controlled clinical trial of interpersonal psychotherapy versus parenting education program for depressed pregnant women. *Am J Psychiatry* 2003; 160:555–562
70. Swartz HA, Frank E, Zuckoff A, Cyranowski JM, Houck PR, Cheng Y, Fleming MA, Grote NK, Brent DA, Shear MK: Brief interpersonal psychotherapy for depressed mothers whose children are receiving psychiatric treatment. *Am J Psychiatry* 2008; 165:1155–1162
71. Tang TC, Jou SH, Ko CH, Huang SY, Yen CF: Randomized study of school-based intensive interpersonal psychotherapy for depressed adolescents with suicidal risk and parasuicide behavior. *Psychiatry Clin Neurosci* 2009; 63:463–470
72. Van Schaik A, van Marwijk H, Adèr H, van Dyck R, de Haan M, Penninx B, van der Kooij K, van Hout H, Beekman A: Interpersonal psychotherapy for elderly patients in primary care. *Am J Geriatr Psychiatry* 2006; 14:777–786
73. Weissman MM, Prusoff BA, DiMascio A, Neu C, Goklaney M, Klerman GL: The efficacy of drugs and psychotherapy in the treatment of acute depressive episodes. *Am J Psychiatry* 1979; 136:555–558
74. Cuijpers P, Dekker J, Hollon SD, Andersson G: Adding psychotherapy to pharmacotherapy in the treatment of depressive disorders in adults: a meta-analysis. *J Clin Psychiatry* 2009; 70:1219–1229
75. Rush AJ, Thase ME: Psychotherapies for depressive disorders: a review, in *Depressive Disorders: WPA Series Evidence and Experience in Psychiatry*. Edited by Maj M, Sartorius N. Chichester, UK, John Wiley & Sons, 1999, pp 161–206
76. Sotsky SM, Glass DR, Shea MT, Pilkonis PA, Collins JF, Elkin I, Watkins JT, Imber SD, Leber WR, Moyer J, Oliveri ME: Patient predictors of response to psychotherapy and pharmacotherapy: findings in the NIMH Treatment of Depression Collaborative Research Program. *Am J Psychiatry* 1991; 148:997–1008
77. Cuijpers P, van Straten A, van Oppen P, Andersson G: Are psychological and pharmacologic interventions equally effective in the treatment of adult depressive disorders? a meta-analysis of comparative studies. *J Clin Psychiatry* 2008; 69:1675–1685
78. Cuijpers P, Smit F, Bohlmeijer E, Hollon SD, Andersson G: Efficacy of cognitive-behavioural therapy and other psychological treatments for adult depression: meta-analytic study of publication bias. *Br J Psychiatry* 2010; 196:173–178