



**Cochrane**  
**Library**

Cochrane Database of Systematic Reviews

## Interventions for waterpipe smoking cessation (Review)

Maziak W, Jawad M, Jawad S, Ward KD, Eissenberg T, Asfar T

Maziak W, Jawad M, Jawad S, Ward KD, Eissenberg T, Asfar T.

Interventions for waterpipe smoking cessation.

*Cochrane Database of Systematic Reviews* 2015, Issue 7. Art. No.: CD005549.

DOI: 10.1002/14651858.CD005549.pub3.

[www.cochranelibrary.com](http://www.cochranelibrary.com)

## TABLE OF CONTENTS

HEADER . . . . .	1
ABSTRACT . . . . .	1
PLAIN LANGUAGE SUMMARY . . . . .	2
SUMMARY OF FINDINGS FOR THE MAIN COMPARISON . . . . .	4
BACKGROUND . . . . .	4
OBJECTIVES . . . . .	5
METHODS . . . . .	5
RESULTS . . . . .	6
Figure 1. . . . .	7
Figure 2. . . . .	9
Figure 3. . . . .	10
DISCUSSION . . . . .	10
AUTHORS' CONCLUSIONS . . . . .	11
ACKNOWLEDGEMENTS . . . . .	12
REFERENCES . . . . .	12
CHARACTERISTICS OF STUDIES . . . . .	14
DATA AND ANALYSES . . . . .	20
Analysis 1.1. Comparison 1 Intervention versus control, Outcome 1 Prolonged Cessation. . . . .	20
APPENDICES . . . . .	20
WHAT'S NEW . . . . .	23
HISTORY . . . . .	23
CONTRIBUTIONS OF AUTHORS . . . . .	23
DECLARATIONS OF INTEREST . . . . .	24
SOURCES OF SUPPORT . . . . .	24
DIFFERENCES BETWEEN PROTOCOL AND REVIEW . . . . .	24
INDEX TERMS . . . . .	24

[Intervention Review]

# Interventions for waterpipe smoking cessation

Wasim Maziak<sup>1,2</sup>, Mohammed Jawad<sup>3,4</sup>, Sena Jawad<sup>5</sup>, Kenneth D Ward<sup>6,7</sup>, Thomas Eissenberg<sup>7,8</sup>, Taghrid Asfar<sup>7,9</sup>

<sup>1</sup>Department of Epidemiology, Robert Stempel College of Public Health and Social Work, Florida International University, Miami, Florida, USA. <sup>2</sup>Department of Epidemiology, Robert Stempel College of Public Health and Social Work, Florida International University, Miami, Florida, USA. <sup>3</sup>Department of Primary Care and Public Health, Imperial College London, London, UK. <sup>4</sup>Academic Unit of Primary Care and Population Sciences, University of Southampton, Southampton, UK. <sup>5</sup>Nuffield Department of Primary Care Health Sciences, University of Oxford, Oxford, UK. <sup>6</sup>Center for Community Health, School of Public Health, Memphis, Tennessee, USA. <sup>7</sup>Syrian Center for Tobacco Studies, Aleppo, Syrian Arab Republic. <sup>8</sup>Department of Psychology and Center for the Study of Tobacco Products, Virginia Commonwealth University, Richmond, Virginia, USA. <sup>9</sup>Department of Public Health Sciences, University of Miami Miller School of Medicine, Miami, Florida, USA

Contact address: Wasim Maziak, Department of Epidemiology, Robert Stempel College of Public Health and Social Work, Florida International University, Miami, Florida, USA. [wmaziak@fiu.edu](mailto:wmaziak@fiu.edu).

**Editorial group:** Cochrane Tobacco Addiction Group.

**Publication status and date:** New search for studies and content updated (conclusions changed), published in Issue 7, 2015.

**Review content assessed as up-to-date:** 19 June 2015.

**Citation:** Maziak W, Jawad M, Jawad S, Ward KD, Eissenberg T, Asfar T. Interventions for waterpipe smoking cessation. *Cochrane Database of Systematic Reviews* 2015, Issue 7. Art. No.: CD005549. DOI: 10.1002/14651858.CD005549.pub3.

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

## ABSTRACT

### Background

Waterpipe tobacco smoking is a traditional method of tobacco use, especially in the Eastern Mediterranean Region (EMR), but its use is now spreading worldwide. Recent epidemiological data, for example, show that waterpipe smoking has become the most prevalent tobacco use method among adolescents in the EMR, and the second most prevalent in the US. Waterpipes are used socially, often being shared between friends or family at home, or in dedicated bars and cafes that provide waterpipes to patrons. Because the smoke passes through a reservoir of water, waterpipe tobacco smoking is perceived as being less harmful than other methods of tobacco use. At least in some cultures, women and girls are more likely to use a waterpipe than to use other forms of tobacco, and it is popular among younger smokers. Accumulating evidence suggests that some waterpipe smokers become addicted, have difficulty quitting, and experience similar health risks as cigarette smokers.

### Objectives

To evaluate the effectiveness of tobacco cessation interventions for waterpipe users.

### Search methods

We searched the Cochrane Tobacco Addiction Review Group specialized register in June 2015. We also searched MEDLINE, EMBASE, PsycINFO and CINAHL, using variant terms and spellings ('waterpipe' or 'narghile' or 'arghile' or 'shisha' or 'goza' or 'narkeela' or 'hookah' or 'hubble bubble'). We searched for trials, published or unpublished, in any language, and especially in regions where waterpipe use is widespread.

### Selection criteria

We sought randomized, quasi-randomized or cluster-randomized controlled trials of smoking cessation interventions for waterpipe smokers of any age or gender. The primary outcome of interest was abstinence from tobacco use, measured at six months post-cessation or longer, regardless of whether abstinence was biochemically verified. We included interventions that were pharmacological

---

**Interventions for waterpipe smoking cessation (Review)**

Copyright © 2015 The Cochrane Collaboration. Published by John Wiley & Sons, Ltd.

1

(for example, nicotine replacement therapy (NRT) or bupropion) or behavioural, or both, and could be directed at individual waterpipe users or at groups of users. We only included tobacco cessation interventions, and did not consider trials of prevention of uptake.

### **Data collection and analysis**

Two review authors assessed abstracts of the studies retrieved by the search strategy, for possible inclusion in the review. We retrieved full-text articles for all abstracts that any of the authors believed might be suitable. Two review authors then extracted data and assessed trial quality independently in accordance with standard Cochrane Collaboration methodologies. We aimed to pool groups of studies that we considered to be sufficiently similar, provided there was no evidence of substantial statistical heterogeneity, and aimed to estimate a pooled risk ratio (RR) using the Mantel-Haenszel fixed-effect method. Where meta-analysis was not possible, we presented summary and descriptive statistics.

### **Main results**

Our search retrieved 1311 unique citations, of which 1289 were excluded after title and abstract screening. Of the remaining 22, we excluded 19 because they were empirical studies that were not randomized, quasi-randomized or cluster-randomized controlled trials ( $n = 12$ ), because they were review articles ( $n = 3$ ), because they described protocols only ( $n = 2$ ), they were conducted among cigarette smokers only ( $n = 1$ ), or they had only a three-month follow-up ( $n = 1$ ).

We identified three controlled trials which tested cessation interventions for waterpipe smokers. Studies were carried out in Egypt (Mohlman 2013), Pakistan (Dogar 2014), and the US (Lipkus 2011). One was a randomized controlled trial and two were cluster-randomized trials. Two studies tested individual-level interventions, and one tested a community-level intervention. Two studies included only behavioural interventions, and one study (Dogar 2014) included two intervention groups: one behavioural, and the other behavioural with bupropion. The Lipkus and Mohlman studies delivered waterpipe-specific interventions, and the Dogar study delivered a non-specific tobacco intervention. Due to study variation we did not pool results, and intervention effects are reported descriptively. Compared to control groups, waterpipe smoking cessation rates were higher in the intervention groups in all three studies, with a significant difference in two studies. For the Dogar study, the RRs for waterpipe smoking abstinence at 25 weeks among waterpipe-only smokers were 2.2 (95% confidence interval (CI) 1.3 to 3.8; 180 participants) in the behavioural group, and 2.5 (95% CI 1.3 to 4.7; 84 participants) in the behavioural plus bupropion group. In our analysis we have combined both groups, to give a RR of 2.28 (95% CI 1.36 to 3.83; 200 participants). The Mohlman study delivered a RR in male waterpipe-smokers at one year in favour of the intervention of 3.25 (95% CI 1.19 to 8.89).

### **Authors' conclusions**

Although the literature on waterpipe cessation interventions remains sparse, the reviewed studies provide a basis for developing interventions in this area. The lack of statistically significant effects in one of the three studies is not unexpected, given the small and pilot nature of the studies. The studies highlight important design and content issues that need to be considered for future cessation trials in waterpipe smokers. These include building on the vast experience developed in the study of smoking cessation interventions in cigarette smokers, whilst including components and assessment tools that address the specific aspects of waterpipe smoking, such as its social dimension, unique experiences, and cues.

## **PLAIN LANGUAGE SUMMARY**

### **Can users of waterpipes be helped to quit through smoking cessation interventions?**

#### **Background**

Waterpipe smoking is a traditional method of tobacco use, especially in the Eastern Mediterranean Region, but its use is now spreading worldwide. It is smoked socially and often shared between friends or family at home, or in bars and cafes that provide waterpipes to patrons. In the absence of relevant data, many waterpipe tobacco smokers believe this form of tobacco use is less lethal and addictive than other methods of tobacco smoking, because the smoke passes through water on its way to the user. At least in some cultures, women and girls are more likely to use a waterpipe than other forms of tobacco, and it is popular among younger smokers. Current evidence suggests that waterpipe smoking may be as addictive as other forms of tobacco use, that some users have difficulty quitting on their own and that they may experience similar risks to health as cigarette smokers.

#### **Study characteristics**

We searched for controlled trials in the Cochrane Tobacco Addiction Review Group specialized register, in June 2015. We also searched a number of electronic databases, including MEDLINE, EMBASE, PsycINFO and CINAHL, using a variety of names and spellings for waterpipe use ('waterpipe' or 'narghile' or 'arghile' or 'shisha' or 'goza' or 'narkeela' or 'hookah' or 'hubble bubble'). We searched for published and unpublished trials in any language, and especially in areas where waterpipe use is widespread. We identified three studies that tested behavioural methods to help waterpipe smokers to quit. Two were waterpipe-specific interventions and one was a non-specific tobacco intervention. One small, pilot study was set in the USA, and delivered a Powerpoint presentation online to 91 college students who were using waterpipe. One study was a secondary analysis of data from 264 waterpipe smokers who were part of a trial that enrolled people suspected of having tuberculosis from 33 healthcare clinics in Pakistan. Clinics were randomly assigned to deliver a behavioural intervention versus control (usual care), or a behavioural intervention plus medication (bupropion) versus control (usual care). The third study, set in Egypt, targeted both cigarette and waterpipe smokers, and was a community-based programme.

### **Key results**

In all three trials, the percentage of participants who stopped smoking waterpipe was higher in the intervention groups than in the control groups, although this was a statistically significant finding in only two of the trials. People who received either behavioural treatment or behavioural treatment plus bupropion were more likely to quit waterpipe smoking at six months follow-up than those who received usual care. Men smoking waterpipe in the Egyptian study were more likely to have quit at one year follow-up in the intervention villages than in the control villages. These studies provide support to suggest that cessation interventions may help waterpipe smokers to quit. However, further larger studies are needed to build on this.

### **Quality of the evidence**

The trials were all rated at very low quality of evidence, as they were relatively small studies, with at least one high risk of bias.

## SUMMARY OF FINDINGS FOR THE MAIN COMPARISON *[Explanation]*

Waterpipe intervention compared with a control for waterpipe cessation				
Outcomes	Impact	Number of Participants (Studies)	Certainty of the evidence (GRADE)*	Comments
<b>Prolonged Cessation</b>	(Dogar 2014) RR 2.48 (95% CI: 1.36 to 3.83) for 25 weeks cessation (Lipkus 2011) RR 1.46 (95% CI: 0.81 to 2.62) for 6 months (Mohlman 2013) RR 3.25 (95% CI: 1.19 to 2.12) for 12 months cessation	200 (1) 91 (1) 540 (1)	⊕○○○ <b>very low</b>	The studies were not pooled as the interventions were not sufficiently similar in design and participant demographics. There is no pooled effect estimate
<b>CI:</b> Confidence interval; <b>RR:</b> Risk Ratio				
GRADE Working Group grades of evidence <b>High quality:</b> Further research is very unlikely to change our confidence in the estimate of effect. <b>Moderate quality:</b> Further research is likely to have an important impact on our confidence in the estimate of effect and may change the estimate. <b>Low quality:</b> Further research is very likely to have an important impact on our confidence in the estimate of effect and is likely to change the estimate. <b>Very low quality:</b> We are very uncertain about the estimate.				
*The certainty of the evidence was very low as all the studies had at least one high risk of bias in accordance with the GRADE framework (see <a href="#">Figure 2</a> ).				

## BACKGROUND

Estimates suggest that by 2030 there will be more than 10 million tobacco-related deaths a year worldwide, with 70% of them occurring in developing countries (Peto 2001). Patterns of tobacco usage and uptake are of increasing concern, as the tobacco industry concentrates its marketing in developing countries, paying particular attention to women and girls and to a wide range of tobacco products (GYTS 2003).

One traditional method of smoking tobacco, especially in the Eastern Mediterranean Region (EMR), is the waterpipe, in which smoke passes through a reservoir of water before inhalation by the smoker. The waterpipe, known as *narjeela* in formal Arabic, goes

by various local names such as *shisha*, *narghile*, *arghile*, and *hookah* (Maziak 2004). Although waterpipe use was uncommon in most of the world before the 1990s, it has enjoyed a recent resurgence, and is now spreading into areas where there was no previous tradition of use (Ward 2015). In most countries of the EMR, waterpipe smoking has become the most common tobacco use method among youth, and the trend is spreading to other world regions such as the US, where waterpipe smoking became the second most popular tobacco use method among college, high-, and middle-school students (Arrazola 2015; Maziak 2015; Primack 2009). While solitary waterpipe use is quite common, waterpipe use is predominantly a social phenomenon, occurring among friends or family, and often in dedicated cafés and bars (Akl 2015; Martinasek

2011).

It is hard to establish all of the potential factors responsible for the global spread of an addictive behaviour such as waterpipe smoking. An addictive behaviour will tend to spread gradually unless it is countered by effective policies and regulations. It is believed that the resurgence in the popularity of waterpipe was sparked by the introduction of flavoured, sweetened tobacco called Maassel in the Middle East during the early 1990s, whilst the global economy, advancements in communication and social media, emigration and tourism have helped to spread the practice globally (Maziak 2015). The lack of effective policies to deal with this relatively new trend is certainly contributing to the vacuum within which this tobacco use method is allowed to thrive (Jawad 2015).

Many waterpipe smokers believe that waterpipe smoking is a safer alternative to cigarettes; an apparent misperception given the available evidence (Akl 2015; Asfar 2008; El-Zaatari 2015; Martinasek 2011). This evidence demonstrates the wide-ranging potential harm of waterpipe smoking, as well as its addictive nature (Aboaziza 2015; El-Zaatari 2015; Shihadeh 2015). Many waterpipe users become dependent, evidenced by urges to smoke waterpipe and other withdrawal symptoms when they abstain, relief of these symptoms when they smoke waterpipe, and difficulty in quitting (Aboaziza 2015). Several studies have reported that between 25% and 64% of waterpipe users want to quit (Akl 2013; Anjum 2008; Ward 2005) and that at least 25% make a quit attempt in any given year (Anjum 2008; Ward 2005; Ward 2006). Quit rates, however, are very low (Ward 2006).

The most identified behavioural association with waterpipe smoking is cigarette smoking. Many studies from around the world have documented the salience of cigarette smoking among waterpipe smokers, and cigarette smoking has been shown to be a major predictor of waterpipe smoking among youth. In the US Monitoring the Future survey, waterpipe use among high-school seniors was associated with current and former cigarette smoking (Maziak 2015). Dual smoking, however, tends to decrease with age, as older smokers are usually more loyal to a single tobacco use method. For example, in a study alluded to earlier, comparing novice with established waterpipe smokers in Syria, the prevalence of dual smoking was 47.7% for novice smokers compared to 26.5% for established ones (Maziak 2015).

Given the global increase in waterpipe smoking, and evidence that many users become dependent and have difficulty quitting on their own, waterpipe-specific tobacco cessation interventions are required. This review aims to summarize the evidence available regarding smoking cessation interventions for waterpipe smokers.

## OBJECTIVES

To evaluate the effectiveness of tobacco cessation interventions for waterpipe users.

## METHODS

### Criteria for considering studies for this review

#### Types of studies

Randomized controlled trials, quasi-randomized controlled trials and cluster-randomized controlled trials.

#### Types of participants

Current (past month) users of waterpipes for tobacco smoking, of any age and either gender.

#### Types of interventions

We included interventions directed at waterpipe users that were pharmacological (for example, nicotine replacement therapy (NRT) or bupropion) or behavioural, or both. These could be directed at individual users or groups of users. We only included cessation interventions, and did not consider trials of prevention of smoking uptake.

#### Types of outcome measures

The primary outcome was abstinence from any tobacco waterpipe use for six months or more from the beginning of intervention. We report abstinence at longest follow-up, and prefer the strictest definition of abstinence (continuous or prolonged over point prevalence, as defined by Hughes 2003). We prefer biochemically-validated abstinence over self-reported abstinence.

### Search methods for identification of studies

We searched the Cochrane Tobacco Addiction Review Group specialized register for trials, using the terms 'waterpipe' or 'narghile' or 'arghile' or 'shisha' or 'goza' or 'narkeela' or 'hookah' or 'hubble bubble', plus variant spellings of these terms, and 'smoking' in the title or abstract, or as keywords. This register was developed from electronic searching of MEDLINE, EMBASE and PsycINFO, together with handsearching of specialist journals, conference proceedings and reference lists of previous trials and overviews. We also searched MEDLINE (1946 to present), EMBASE (1980 to present), CINAHL (1981 to present) and PsycINFO (1806 to present), using the above free-text terms combined with MeSH or free-text smoking-related terms (smok\* or tobacco or cigar\* or nicotine). We searched for trials, published or unpublished, in any language, and especially in regions where waterpipe use is widespread. We also used our existing bibliography, compiled from earlier exhaustive reviews of the literature on waterpipe smoking

(e.g. Aboaziza 2015; El-Zaatari 2015; Jawad 2015; Maziak 2004; Maziak 2015; Shihadeh 2015). The most recent search was completed on 19th June 2015.

### Data collection and analysis

Two review authors (MJ and SJ) assessed the abstracts of studies retrieved by the search strategy, for possible inclusion in the review. We retrieved full-text articles for all abstracts which either review author believed might be suitable.

### Assessment of full articles

Two review authors (MJ and SJ) assessed each full-text article independently, using the agreed inclusion criteria. Where there was ambiguity in trial reporting or a lack of data, we contacted investigators for clarification where possible. If we could not retrieve missing data we considered exclusion on that basis.

We rated the overall methodological quality of studies as being at low, moderate, or high risk of bias for each of the following criteria to assess risk of bias:

1. Random sequence generation
2. Concealment of allocation
3. Blinding of participants and personnel
4. Blinding of outcome assessment
5. Incomplete outcome data (attrition rates and losses to follow-up)
6. Biochemical verification of smoking status

We maintained a full list of excluded studies.

### Data collection

We extracted and reported the following information, where available, concerning each study:

- Country and study setting
- Dates study was conducted
- Theoretical framework (including a brief description of the intervention)
- Focus of the intervention (e.g. any tobacco use, waterpipe smoking)

- Type of intervention, its duration, intensity, delivery format
- Length of follow-up
- Number of participants or number of clusters and participants
  - Age range, socio-economic status, gender and ethnicity (if relevant) of participants
  - Definition of smoking status used (e.g. level of waterpipe use, concurrent use of other tobacco)
  - Definition of abstinence
  - Biochemical validation (if present)
  - Differential effects post-intervention relating to age, gender, ethnicity and intensity of intervention
  - Adverse effects of the intervention
  - Sources of funding

We aimed to pool groups of studies that we considered to be sufficiently similar in their interventions, comparison groups, setting and participants, provided that there was no evidence of substantial statistical heterogeneity as assessed by the  $I^2$  statistic (Higgins 2003). We aimed to estimate a pooled risk ratio (RR) using the Mantel-Haenszel fixed-effect method, based on the quit rates at longest follow-up for trials with at least six months follow-up from the start of the intervention. Where meta-analysis was not possible, we present a descriptive summary and descriptive statistics. We include a glossary of tobacco-specific terms (Appendix 1) as an additional table in this review.


## RESULTS

### Description of studies

Our search retrieved 1311 unique citations, of which 1289 were excluded after title and abstract screening. Of the remaining 22, we excluded 19 for the following reasons: they were non-randomized studies ( $n = 12$ ), review articles ( $n = 3$ ), a protocol only ( $n = 2$ ), conducted among cigarette smokers only ( $n = 1$ ), or had only a three-month follow-up ( $n = 1$ ). The flow of studies is illustrated in Figure 1.



**Figure 1. Study flow diagram.**



We identified three eligible studies which tested cessation interventions for waterpipe smokers. One was a randomized controlled trial (Lipkus 2011) and two were cluster-randomized trials (Dogar 2014; Mohlman 2013). Two were individual-level interventions (Dogar 2014; Lipkus 2011) and one was a community-level intervention (Mohlman 2013). Interventions were carried out in Egypt (Mohlman 2013), Pakistan (Dogar 2014), and the US (Lipkus 2011). All interventions were behavioural interventions, although one study included two intervention groups: one behavioural, and the other behavioural with bupropion (Dogar 2014). One study was based on a theoretical framework (Dogar 2014). Further characteristics of included studies can be found in [Characteristics of included studies](#).

The first study took place in Pakistan between 2010 and 2011 and was a three-arm, cluster-randomized non-inferiority trial among a mix of cigarette-only smokers ( $n = 1181$ ), waterpipe-only smokers ( $n = 200$ ) and mixed smokers ( $n = 460$ ) (Dogar 2014). The three arms were standard care (control group), a brief behavioural intervention (BSS) and a brief behavioural intervention plus bupropion for seven weeks (BSS+). The behavioural intervention was adapted from evidence-based treatments used for cigarette smokers and involved two structured sessions (the first 30 minutes long, the second 10 minutes long) one week apart. It was delivered by tuberculosis DOTS (directly observed treatment, short course) paramedics. The bupropion regimen was 75 mg/day for the first week and 150 mg/day for the next six weeks. The control group received a leaflet with standard health messages about the harms of tobacco. The clusters were primary and secondary healthcare centres registered as diagnostic centres by a tuberculosis programme. The study authors adjusted for the effect of clustering by conducting a multi-level analysis. The study was funded by the International Development Research Centre, Canada.

The second study took place in the US between 2009 and 2010, and was a randomized controlled web-based intervention among waterpipe smoking college/university students (Lipkus 2011). Ninety-one students were randomized to non-health-related information about waterpipe (control group) or to both non-health-related and health-related information about waterpipe (intervention group). The study was funded by grants from the US National Cancer Institute and National Institute on Drug Abuse.

The final study took place in Egypt between 2004 and 2005, and was a cluster-randomized controlled community-level intervention (Mohlman 2013). The clusters were villages in the Qalyubia governorate. Villages, with a total of 7657 participants, were randomised to receive a behavioural intervention through a variety of activities engaging school students, places of worship, and adult women, and delivered by teachers, religious leaders and female social-change agents respectively. Primary school students partook in activities to prevent the initiation of tobacco use through its deglamorisation and teaching of health effects. Preparatory and

secondary school students were taught social skills to handle peer pressure to smoke. Religious communities were informed of the health effects of tobacco use/secondhand smoke, and the sinful nature of smoking. Adult women at home were taught about the health effects of tobacco use/secondhand smoke and how to protect themselves and their children from it in a culture-specific way. Control villages received no intervention but had access to Egypt's National Tobacco Control Program during the study. The study authors adjusted for the effect of clustering in the analysis. The study was funded by The Fogarty International Center of the US National Institutes of Health.

Studies were not comparable in terms of participants' smoking status, intervention type, and outcome measures. For example, participants in Dogar 2014 smoked a local form of unflavoured waterpipe tobacco a median of 10 times per day, participants in Mohlman 2013 smoked two to three times per day, and participants in Lipkus 2011 smoked flavoured Maassel monthly. With regards to the delivery method of interventions, one study provided a web-based intervention (Lipkus 2011), one provided group intervention (Mohlman 2013), and one provided interventions aimed at individuals (Dogar 2014). With regards to outcomes, only one study (Dogar 2014) biochemically validated abstinence by expired carbon monoxide ( $\text{CO} \leq 9$  ppm), while the remaining two trials relied on only self report of smoking status (Lipkus 2011; Mohlman 2013). Follow-up length ranged from six months to one year post-intervention.

### Risk of bias in included studies

We considered two studies (Dogar 2014; Lipkus 2011) to be at low risk of bias for random sequence generation as they adequately described a simple randomization process. Mohlman 2013 used a randomized design but the details were not reported. We deemed only one study to be at low risk of bias for adequate allocation concealment (Dogar 2014) as it was concealed by a researcher blinded to centre identity. The remaining two studies did not provide information on allocation concealment (Lipkus 2011; Mohlman 2013). We rated none of the studies at low risk of bias for blinding of participants and personnel; Dogar 2014 was an open-label trial, while in Lipkus 2011 and Mohlman 2013 blinding was not mentioned. Although none of the studies reported the presence of blinding, we considered one study (Dogar 2014) to be at low risk of bias for blinding of outcome assessment, as it was biochemically verified with a carbon monoxide measurement of 9 ppm or less. We rated all three at low risk of bias for completeness of data, as missing data ranged from 7% (Dogar 2014) to 23% (Lipkus 2011). Mohlman 2013 excluded women from the dataset because smoking prevalence was very low in this group. No studies were at low risk of bias for selective reporting. In Dogar 2014 outcomes at five weeks were not reported, whereas in Lipkus 2011 and

Mohlman 2013 cessation data were not reported using appropriate effect estimates. A summary of the 'Risk of bias' assessment for the included studies can be found in [Figure 2](#).

**Figure 2. Risk of bias summary: review authors' judgements about each risk of bias item for each included study.**

	Random sequence generation (selection bias)	Allocation concealment (selection bias)	Blinding of participants and personnel (performance bias)	Blinding of outcome assessment (detection bias)	Incomplete outcome data (attrition bias)	Selective reporting (reporting bias)	Other bias
Dogar 2014	+	+	-	+	+	-	+
Lipkus 2011	+	?	?	-	+	-	-
Mohlman 2013	?	?	?	-	+	-	-


### Effects of interventions

See: [Summary of findings for the main comparison](#)

Due to the variation in studies outlined above, we did not pool the results of studies using meta-analysis or conduct statistical tests for heterogeneity. We report the intervention effects descriptively

and present abstinence data for individual trials in [Figure 3](#).

**Figure 3. Forest plot of comparison: I Raw Data, outcome: I.I Prolonged Cessation.**



**Footnotes**

(1) Combined behavioural and (behavioural + bupropion) groups

For [Dogar 2014](#) we considered the data provided by waterpipe smokers only (n = 200). Compared to the control group, the risk ratio (RR) of smoking abstinence at 25 weeks was 2.2 (95% CI 1.3 to 3.8) for the BSS group and 2.5 (95% CI 1.3 to 4.7) for the BSS+ group. In [Figure 3](#) we have combined the BSS and BSS+ groups to create a single intervention group.

For [Lipkus 2011](#), when comparing the intervention to the control group, the RR for waterpipe cessation at six months was 1.46 (95% CI 0.81 to 2.62; n = 70).

For the third study ([Mohlman 2013](#)), the RR of waterpipe cessation at one year was 3.25 (95% CI 1.19 to 8.89; n = 540) for the intervention group compared to the control group.

## DISCUSSION

Waterpipe use is a growing phenomenon associated with substantial toxicant exposure, numerous health risks, and development of dependence in a sizeable proportion of users ([Aboaziza 2015](#); [El-Zaatari 2015](#); [Maziak 2015](#); [Shihadeh 2015](#)). Despite these adverse consequences, development and evaluation of cessation interventions for the waterpipe are scarce. In our review, we found only three studies which met our inclusion criteria, covering 831 participants, that have examined interventions to help waterpipe users quit smoking. All three studies tested behavioural interventions, and one study also included a combined behavioural/pharmacological (bupropion) intervention group ([Dogar 2014](#)). Due to lack of comparability across the three studies in terms of participants' smoking status, intervention type, and outcome assessment, we did not conduct statistical tests for heterogeneity and meta-analysis, but present intervention effects descriptively. Two trials were conducted among adults in the Middle East ([Dogar 2014](#); [Mohlman 2013](#)), and one study was conducted among young adults in the US ([Lipkus 2011](#)). This should be considered for

the generalizability of findings from these studies. Compared to control groups, smoking cessation rates were higher in the intervention groups in all three studies; however, the difference was not statistically significant in one study ([Lipkus 2011](#)). These findings suggest that waterpipe smokers may be more likely to stop smoking successfully when using a community or a tailored smoking cessation intervention than usual care; however they should be treated with caution due to the paucity and limitations of the available data.

The lack of a statistically significant effect in the American trial is not unexpected, given the small, pilot nature of the study. A Cochrane review of individual behavioural interventions for cigarette smoking cessation demonstrated an RR of 1.39 (95% CI 1.2 to 1.57) ([Lancaster 2005](#)). At least two of the studies ([Dogar 2014](#); [Lipkus 2011](#)) were unlikely to have the power to detect a comparable RR. Suboptimal length of follow-up (less than one year) was another limitation for two of the three included studies, as well as a reliance on self-reported data in all but one study ([Dogar 2014](#)). Abstinence verification methods should also be suitable for waterpipe smoking. Expired breath CO, which is good for the detection of smoking in the past 24 hours only, may not accurately verify abstinence in intermittent waterpipe smokers, but can be used as a 'bogus pipeline' ([Asfar 2014](#); [Murray 1987](#); [Patrick 1994](#)) to encourage truthful reporting of abstinence violations among intermittent users. The absence of standard definitions for waterpipe smoking status for inclusion in the trials was also an issue that affected comparability of the reviewed studies. For example, two of the reviewed studies ([Lipkus 2011](#); [Mohlman 2013](#)) recruited current waterpipe smokers who smoked waterpipe in the past month, and one study ([Dogar 2014](#)) recruited regular waterpipe smokers who "smoke  $\geq 1$  waterpipe per day." Cigarette cessation trials typically enrol daily smokers and outcome evaluation is often focused on whether participants have returned to daily

smoking. We noted the same inconsistency for the definition and verification of abstinence in the three studies. In Dogar 2014, the primary outcome was continuous abstinence at six-month follow-up, while in Lipkus 2011 abstinence was defined as reporting no longer using waterpipe at the six-month follow-up, and finally, in Mohlman 2013 abstinence was defined as not smoking waterpipe in the last month before the 12-month follow-up. Not only are these outcomes hard to compare, they are not consistent with the relevant scientific recommendations for cessation trials and with common patterns of waterpipe smoking. Because many waterpipe smokers are cigarette smokers as well, cessation outcomes should be standardised to allow comparison with the cigarette literature as well as to accommodate waterpipe smokers' usage patterns. For example, the definition of *prolonged abstinence* defined as; no smoking, not even a puff, after a grace period of two weeks after quit date, and relapse as smoking at least once a week on two consecutive weeks (SRNT 2002), are standard cigarette-based definitions that would seem to be suitable for waterpipe.

Out of the three included studies, only one evaluated a combined intervention (behavioural plus bupropion), and did not show any apparent additional benefit of adding bupropion to behavioural support in achieving cessation (Dogar 2014), although these two conditions were not directly tested against one another. The efficacy of other pharmacological cessation modalities such as nicotine replacement therapy (NRT) or varenicline, which have been shown to be useful in dependent cigarette smokers, have not yet been tested in waterpipe smokers. Given that some waterpipe smokers exhibit signs and symptoms of dependence (Aboaziza 2015), pharmacotherapy may be useful during cessation. This can be particularly relevant to highly quit-motivated dual waterpipe/cigarette users (Ward 2014). However, individuals who are less dependent, have smoked for shorter periods of times, and who cite social stigma (e.g. family disapproval) as a reason to stop smoking make up the majority of waterpipe smokers interested in quitting (Borgan 2013; Ward 2005). These individuals may be less likely to benefit from pharmacological treatments. Such considerations require having some standard, waterpipe-specific measure of dependence that allows for the variability in both individual smoking habits and nicotine content of different waterpipe tobacco brands to be captured in a standardized manner. A recently developed scale (Lebanon Waterpipe Dependence Scale-11) to characterize waterpipe dependence has shown that self-reported dependence level correlates with measurements of nicotine metabolites in flavoured waterpipe tobacco users (Salameh 2008). This measure was developed based on cigarette smoking instruments and without input from waterpipe smokers, but could be the first step in accurately measuring dependence among waterpipe smokers.

Offering behavioural support adapted from a validated cigarette-smoking cessation programme could be a useful starting point for waterpipe smokers who are interested in quitting. Of the three studies reviewed, two offered behavioural interventions utilizing

similar strategies to those shown to be effective for cigarette smokers (Lipkus 2011; Mohlman 2013). Process evaluation data from one of our excluded studies (Asfar 2014) indicates that the methods they used, which were adapted from traditional smoking cessation methods, were acceptable to waterpipe smokers. However, such approaches will miss dealing with the strong social dimension of waterpipe use, as it shapes use patterns, cues for smoking, and the attitudes and preferences of waterpipe smokers (Aboaziza 2015; Jawad 2013; Maziak 2015).

Risks associated with the social use of waterpipe, such as the potential to contract infectious diseases through sharing the same waterpipe and using it repeatedly without proper sanitation in café settings, can also provide powerful drives for cessation. Future cessation efforts should consider introducing and examining new methods of cessation intervention, such as group smoking cessation. Results from the process evaluation in Asfar 2014 indicate that one-third of waterpipe smokers were interested in participating in a group counselling intervention. As the social context of waterpipe smoking frequently involves family members (Akl 2015; Maziak 2015), family-based cessation interventions could also be a promising avenue to pursue (Asfar 2014).

Despite the fact that the waterpipe epidemic is most pronounced among youth and young adults (such as college students), only one of the three reviewed studies was conducted among college students. The study provided preliminary evidence that minimally intensive interventions delivered online to educate college waterpipe smokers of the harm, addiction, and toxicant exposure associated with waterpipe smoking can increase understanding of the harms of waterpipe use, perceptions of risk, desire to quit, and eventually cessation (Lipkus 2011). Utilizing youth-oriented technology such as smart phones, text messaging, social networks, or multimedia may provide promising cessation approaches for this at-risk population.

Although the small number and methodological limitations of waterpipe cessation trials to date do not allow firm recommendations to be made on the comparative efficacy of various cessation methods, they do provide a new evidence base on which to build further. They highlight important design and content issues that need to be considered for future cessation trials in waterpipe smokers. These include building on the vast experience of cigarette smoking cessation interventions whilst introducing intervention components and assessment tools that address the specific aspects of waterpipe smoking. It also highlights some of the challenges of future waterpipe cessation trials that relate to their specific set-up, usage patterns and context, and adaptability of cigarette-based definitions and measures to the waterpipe.

## **AUTHORS' CONCLUSIONS**

### **Implications for practice**

- Waterpipe smoking is spreading globally, and carries considerable health risks. Due to its addictive nature, waterpipe users who want to quit find it difficult to do so. This review suggests that waterpipe smokers may be more likely to stop smoking successfully when using a smoking cessation intervention than usual care, but this needs to be treated with caution due to the paucity and limitations of the data. Relying on behavioural cessation approaches from the cigarette literature seems to be a good starting point, but these should be adapted to the specific nature of waterpipe smoking as a predominantly social and intermittent behaviour with prolonged sessions. Adding a pharmacological agent (bupropion) did not seem to have an additional benefit to behavioural support in achieving cessation, again based on the limited data available.

### Implications for research

- Standard definitions and assessments of waterpipe use, dependence and cessation need to be adopted. For example, including smokers with “regular” waterpipe use, defined as smoking three or more waterpipes a week, in smoking cessation trials, will allow smoking cessation efforts to focus on those most in need (most dependent), and at the same time allow the use of standard cigarette-based definitions and verification of abstinence (e.g. prolonged abstinence; saliva cotinine). Since cotinine offers a window of four to five days for detection of nicotine exposure, less frequent waterpipe smoking/abstinence can be hard to verify biochemically. Most of these definitions are already developed and need to be adopted by waterpipe cessation research (e.g. [Jarvis 1988](#); [SRNT 2002](#)).

- As waterpipe cessation trials are still in their infancy, it will be helpful to develop and adopt consistent standards for reporting outcomes to facilitate comparing study results (e.g.

standard definition of regular smoker, definition of abstinence including duration, self report, and cut-off point of various biochemical verification procedures).

- Since many waterpipe smokers are cigarette smokers as well, such measures need to be consistent with the cigarette literature and at the same time accommodate waterpipe smokers’ intermittent usage patterns.

- Waterpipe dependence measures need to be developed and adopted that can capture the common as well as unique (e.g. social dimension) aspects of tobacco dependence in waterpipe smokers.

- Efforts to develop and test behavioural strategies that fit the unique features of waterpipe smoking (e.g. the social cues, and intermittent usage patterns) and address waterpipe-specific facilitators and barriers to quitting are much needed ([Maziak 2015](#)).

- Given how little is currently known about who will use cessation treatment, which treatments they will use, and what specific methods work, it is imperative that large-scale, randomized controlled trials be conducted to rigorously test behavioural, pharmacological and combined cessation approaches.

### ACKNOWLEDGEMENTS

None

### REFERENCES

#### References to studies included in this review

##### Dogar 2014 *{published data only}*

\* Dogar O, Jawad M, Shah SK, Newell JN, Kanaan M, Khan MA, et al. Effect of cessation interventions on hookah smoking: post-hoc analysis of a cluster-randomized controlled trial. *Nicotine & Tobacco Research* 2014;**16**(6): 682–8.

Siddiqi K, Khan A, Ahmad M, Dogar O, Kanaan M, Newell JN, et al. Action to stop smoking in suspected tuberculosis (ASSIST) in Pakistan: a cluster randomized, controlled trial. *Annals of Internal Medicine* 2013;**158**(9):667–75.

##### Lipkus 2011 *{published and unpublished data}*

Lipkus IM, Eissenberg T, Schwartz-Bloom RD, Prokhorov AV, Levy J. Affecting perceptions of harm and addiction

among college waterpipe tobacco smokers. *Nicotine & Tobacco Research* 2011;**13**(7):599–610.

##### Mohlman 2013 *{published data only}*

Mohlman MK, Boulos DN, El Setouhy M, Radwan G, Makambi K, Jillson I, et al. A randomized, controlled community-wide intervention to reduce environmental tobacco smoke exposure. *Nicotine & Tobacco Research* 2013;**15**(8):1372–81.

#### References to studies excluded from this review

##### Asfar 2014 *{published data only}*

Asfar T, Al Ali R, Rastam S, Maziak W, Ward KD. Behavioral cessation treatment of waterpipe smoking: the first pilot randomized controlled trial. *Addictive Behaviors* 2014;**39**(6):1066–74.

## Additional references

### Aboaziza 2015

Aboaziza E, Eissenberg T. Waterpipe tobacco smoking: what is the evidence that it supports nicotine/tobacco dependence?. *Tobacco Control* 2015;**24**(Suppl):i44–53.

### Akl 2013

Akl EA, Jawad M, Lam WY, Co CN, Obeid R, Irani J. Motives, beliefs and attitudes towards waterpipe tobacco smoking: a systematic review. *Harm Reduction Journal* 2013;**10**(1):12.

### Akl 2015

Akl EA, Ward KD, Breddini D, Khaliel R, Alexander AC, Loutfi T, Alaoui H, Afifi RA. The allure of the waterpipe: a narrative review of factors affecting the epidemic rise in waterpipe smoking among young persons globally. *Tobacco Control* 2015;**24**(Suppl):i13–21.

### Anjum 2008

Anjum Q, Ahmed F, Ashfaq T. Knowledge, attitude and perception of water pipe smoking (Shisha) among adolescents aged 14–19 years. *The Journal of the Pakistan Medical Association* 2008;**58**(6):312.

### Arrazola 2015

Arrazola RA, Singh T, Corey CG, Husten CG, Neff LJ, Apelberg BJ, et al. Tobacco use among middle and high school students - United States, 2011–2014. *Morbidity and Mortality Weekly Report* 2015;**64**(14):381–5.

### Asfar 2008

Asfar T, Weg MV, Maziak W, Hammal F, Eissenberg T, Ward KD. Outcomes and adherence in Syria's first smoking cessation trial. *American Journal of Health Behavior* 2008;**32**(2):146–56.

### Borgan 2013

Borgan SM, Marhoon ZA, Whitford DL. Beliefs and perceptions toward quitting waterpipe smoking among café waterpipe tobacco smokers in Bahrain. *Nicotine & Tobacco Research* 2013;**15**(11):1816–21.

### El-Zaatari 2015

El-Zaatari ZM, Chami HA, Zaatari GS. Health effects associated with waterpipe smoking. *Tobacco Control* 2015;**24**(Suppl):i31–i43.

### GYTS 2003

Global Youth Tobacco Survey Collaborating Group. Differences in worldwide tobacco use by gender: findings from the global youth tobacco survey. *Journal of School Health* 2003;**73**(6):207–15.

### Higgins 2003

Higgins JB, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. *BMJ* 2003;**327**(7414):557.

### Hughes 2003

Hughes JR, Keely JP, Niaura RS, Ossip-Klein DJ, Richmond RL, Swan GE. Measures of abstinence in clinical trials: issues and recommendations. *Nicotine & Tobacco Research* 2003;**5**(1):13–25.

### Jarvis 1988

Jarvis MJ, Russell MA, Benowitz NL, Feyerabend, C. Elimination of cotinine from body fluids: implications for noninvasive measurement of tobacco smoke exposure. *American Journal of Public Health* 1988;**78**(6):696–8.

### Jawad 2013

Jawad M, Jawad S, Mehdi A, Sardar A, Jawad AM, Hamilton FL. A qualitative analysis among regular waterpipe tobacco smokers in London universities. *The International Journal of Tuberculosis and Lung Disease* 2013;**17**(10):1364–9.

### Jawad 2015

Jawad M, El Kadi L, Mugharbil S, Nakkash R. Waterpipe tobacco smoking legislation and policy enactment: a global analysis. *Tobacco Control* 2015;**24**(Suppl):i60–i65.

### Lancaster 2005

Lancaster T, Stead LF. Individual behavioural counselling for smoking cessation. *Cochrane Database of Systematic Reviews* 2005, Issue 2. [DOI: 10.1002/14651858.CD001292.pub2]

### Martinasek 2011

Martinasek MP, McDermott RJ, Martini L. Waterpipe (hookah) tobacco smoking among youth. *Current Problems in Pediatric and Adolescent Health Care* 2011;**41**(2):34–57.

### Maziak 2004

Maziak W, Ward KD, Afifi Soweid RA, Eissenberg T. Tobacco smoking using a waterpipe: a re-emerging strain in a global epidemic. *Tobacco Control* 2004;**13**(4):327–33.

### Maziak 2015

Maziak W, Taleb ZB, Bahelah R, Islam F, Jaber R, Auf R, et al. The global epidemiology of waterpipe smoking. *Tobacco Control* 2015;**24**(Suppl):i3–i12.

### Murray 1987

Murray DM, O'Connell CM, Schmid LA, Perry CL. The validity of smoking self-reports by adolescents: a reexamination of the bogus pipeline procedure. *Addictive Behaviors* 1987;**12**(1):7–15.

### Patrick 1994

Patrick DL, Cheadle A, Thompson DC, Diehr P, Koepsell T, Kinne S. The validity of self-reported smoking: a review and meta-analysis. *American Journal of Public Health* 1994;**84**(7):1086–93.

### Peto 2001

Peto R, Lopez AD. Future worldwide health effects of current smoking patterns. In: Koop CE, Pearson CE, Schwarz MR editor(s). *Critical Issues in Global Health*. San Francisco CA: Jossey-Bass, 2001.

### Primack 2009

Primack BA, Walsh M, Bryce C, Eissenberg T. Water-pipe tobacco smoking among middle and high school students in Arizona. *Pediatrics* 2009;**123**(2):e282–e288.

### Salameh 2008

Salameh P, Waked M, Aoun Z. Waterpipe smoking: construction and validation of the Lebanon Waterpipe Dependence Scale (LWDS-11). *Nicotine and Tobacco Research* 2008;**10**(1):149–58.

**Shihadeh 2015**

Shihadeh A, Schubert J, Klaiany J, El Sabban M, Luch A, Saliba NA. Toxicant content, physical properties and biological activity of waterpipe tobacco smoke and its tobacco-free alternatives. *Tobacco Control* 2015;**24**(Suppl):i22–i30.

**SRNT 2002**

SRNT Subcommittee on Biochemical Verification. Biochemical verification of tobacco use and cessation. *Nicotine & Tobacco Research* 2002;**4**(2):149–59.

**Ward 2005**

Ward KD, Hammal F, VanderWeg MW, Eissenberg T, Asfar T, Rastam S, et al. Are waterpipe users interested in quitting?. *Nicotine & Tobacco Research* 2005;**7**(1):149–56.

**Ward 2006**

Ward KD, Eissenberg T, Rastam S, Asfar T, Mzayek F, Fouad MF, et al. The tobacco epidemic in Syria. *Tobacco Control* 2006;**15**(Suppl):24–9.

**Ward 2014**

Ward KD, Ahn S, Mzayek F, Al Ali R, Rastam S, Asfar T, et al. The relationship between waterpipe smoking and body weight: population-based findings from Syria. *Nicotine & Tobacco Research* 2014;**17**(1):34–40.

**Ward 2015**

Ward KD. The waterpipe: an emerging global epidemic in need of action. *Tobacco Control* 2015;**24**(Suppl):i1–i2.

\* Indicates the major publication for the study



## CHARACTERISTICS OF STUDIES

### Characteristics of included studies [ordered by study ID]

#### Dogar 2014

Methods	<p>Year(s) of study: 2010 - 2011</p> <p>Study design: 3-arm cluster-randomized controlled non-inferiority trial</p> <p>Country: Pakistan</p> <p>Region: Jhang and Sarghoda districts</p> <p>Setting: 33 primary and secondary health centres</p> <p>Theoretical framework: based on the World Health Organization's '5As Approach'</p>
Participants	<p>Adults aged over 18 years with suspected tuberculosis (cough <math>\geq</math> 3 weeks, of unknown cause)</p> <p>Excluded: those requiring hospitalization or urgent medical attention</p> <p>Recruitment method: patients attending primary and secondary healthcare centres registered as diagnostic centres by a tuberculosis program in 2 Pakistani districts</p> <p>33 clusters, 200 adults</p> <p>Mean age 51.5 (SD 13.8), median household income USD 81.4 (IQR 69.8), 21% women</p> <p>Definition of smoking status: <math>\geq</math> 1 waterpipe/day</p>
Interventions	<p>Focus of intervention: any smoking use</p> <p>Type of intervention: behavioural and pharmacological</p> <p>Description of the intervention:</p> <p>Control group: given a leaflet with standard health messages about the harms of tobacco</p> <p>Intervention group 1: 2 brief behavioural support cessations (1st visit 30 mins, 2nd on quit day 10 mins)</p> <p>Intervention group 2: 2 brief behavioural support cessations (as above) plus bupropion for 7 weeks (75 mg/day for 1st week, 150 mg/day for next 6 weeks)</p> <p>Intervention delivered by: tuberculosis DOTS (directly observed treatment, short course) paramedics</p>
Outcomes	<p>Continuous waterpipe smoking abstinence</p> <p>Length of follow-up: six months</p> <p>Biochemical validation: CO verified (<math>&lt;</math> 10 ppm)</p>
Notes	<p>Differential effects post-intervention: none reported</p> <p>Adverse effects of intervention: none reported</p> <p>The study was funded by the International Development Research Centre, Canada</p>

#### *Risk of bias*

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Low risk	Not mentioned in sufficient detail in this paper; full trial methodology found in Siddiqi 2013:

**Dogar 2014** (Continued)

		Quote: “a researcher who was blinded to center identity used computer-generated random-number lists to generate allocation sequence” Comment: probably done
Allocation concealment (selection bias)	Low risk	Not mentioned in sufficient detail in this paper; full trial methodology found in Siddiqi 2013: Quote: “a researcher who was blinded to center identity used computer-generated random-number lists to generate allocation sequence” Comment: probably done
Blinding of participants and personnel (performance bias) All outcomes	High risk	Not mentioned in this paper; full trial methodology found in Siddiqi 2013: Quote: “the lack of blinding also meant that a degree of observer bias was possible” Comment: probably not done
Blinding of outcome assessment (detection bias) All outcomes	Low risk	Not mentioned in this paper; full trial methodology found in Siddiqi 2013: Quote: “the lack of blinding also meant that a degree of observer bias was possible” Comment: outcome measurement is biochemically verified, and is unlikely to be affected by blinding
Incomplete outcome data (attrition bias) All outcomes	Low risk	Primary outcome data missing for 7.0% of waterpipe-only smokers. Reasons for missing data unlikely to be related to true outcome. No exclusions reported
Selective reporting (reporting bias)	High risk	Outcomes not reported at 5 weeks
Other bias	Low risk	Biochemical verification of outcome

**Lipkus 2011**

Methods	Year(s) of study: 2009 - 2010 Study design: randomized controlled web-based behavioural intervention Country: USA Region: North Carolina Setting: 6 college and university campuses Theoretical framework: none reported
---------	---

Participants	Adults enrolled in a 4-year college or university course Recruitment method: newspaper advertisements, flyers posted around campuses, Craig's list, campus-wide Listserv 91 adults Mean age 20.4 (SD 2.0), 24.2% women, 76.7% white Definition of smoking status: past-month waterpipe smoking	
Interventions	Focus of intervention: waterpipe smoking Type of intervention: behavioural Description of the intervention: Control group: 8 MS PowerPoint slides on waterpipe mechanism of action, chemical composition, and epidemiology; average length of intervention 3.6 minutes Intervention group: 20 MS PowerPoint slides on waterpipe mechanisms of action, chemical composition, epidemiology, puff topography, toxicant exposure, and health outcomes; average length of intervention 7.5 minutes Intervention delivered by: online	
Outcomes	Ticking the survey item: "no longer smoking waterpipe" Length of follow-up: 6 months Biochemical validation: not present	
Notes	Differential effects post-intervention: none reported Adverse effects of intervention: none reported The study was funded by grants from the US National Cancer Institute and National Institute on Drug Abuse	
<b><i>Risk of bias</i></b>		
<b>Bias</b>	<b>Authors' judgement</b>	<b>Support for judgement</b>
Random sequence generation (selection bias)	Low risk	Quote: "those who logged on were randomized to either a control or an experimental group with equal probability by our program" Comment: probably done
Allocation concealment (selection bias)	Unclear risk	No information on allocation concealment Comment: insufficient confidence that allocation concealment was adequate
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No information on blinding Comment: insufficient confidence that blinding was adequate
Blinding of outcome assessment (detection bias) All outcomes	High risk	No information on blinding Comment: outcome measurement not biochemically verified.

**Lipkus 2011** (Continued)

Incomplete outcome data (attrition bias) All outcomes	Low risk	23.1% of participants did not conduct the 6-month follow-up. Unlikely to be related to outcome. No reasons given for loss to follow-up. 1 participant in the 6-month follow-up had missing data and was not analysed. No exclusions reported
Selective reporting (reporting bias)	High risk	Cessation data not appropriately presented as effect estimates
Other bias	High risk	No biochemical verification of outcome

**Mohlman 2013**

Methods	Year(s) of study: 2004 - 2005 Study design: cluster-randomized controlled behavioural intervention Country: Egypt Region: Qalyubia governorate Setting: Villages Theoretical framework: None reported
Participants	All household members aged over 12 years old, although results pertain only to adult men (n of women for self-reported smoking too small); Waterpipe smokers: Intervention villages 250, control villages: 290 Recruitment method: Systematic approach of households 6 clusters, 7657 residents Mean age 36.9, 41.8% illiterate, 87.3% employed, 55.3% women Definition of smoking status: Past-month waterpipe smoking
Interventions	Focus of intervention: cigarettes and/or waterpipe smoking Type of intervention: behavioural Description of the intervention: educational approach for primary/preparatory/secondary school students, mosques and churches, and key female social change agents ( <i>raedat refeyat</i> ) Intervention delivered by: teachers, religious leaders, female social change agents Control group: no intervention
Outcomes	Waterpipe smoking prevalence Length of follow-up: 12 months Biochemical validation: not present
Notes	Differential effects post-intervention: none reported Adverse effects of intervention: none reported The study was funded by the Fogarty International Center of the US National Institutes of Health
<b>Risk of bias</b>	

**Mohlman 2013** (Continued)

Bias	Authors' judgement	Support for judgement
Random sequence generation (selection bias)	Unclear risk	Quote: "the remaining six villages were randomly allocated to either the control group or the intervention group" Comment: insufficient confidence that the allocation sequence was genuinely randomised
Allocation concealment (selection bias)	Unclear risk	No information on allocation sequence Comment: insufficient confidence that allocation concealment was adequate
Blinding of participants and personnel (performance bias) All outcomes	Unclear risk	No information on blinding Comment: insufficient confidence that blinding was adequate
Blinding of outcome assessment (detection bias) All outcomes	High risk	No information on blinding Comment: outcome measurement is not biochemically verified
Incomplete outcome data (attrition bias) All outcomes	Low risk	77.5% retention rate. Women excluded due to very low self-reporting of tobacco use. All completed pre-intervention survey but not all completed post-intervention survey. No reasons given for loss to follow-up
Selective reporting (reporting bias)	High risk	Cessation data not appropriately presented as effect estimates
Other bias	High risk	No biochemical verification of outcome

**Characteristics of excluded studies** [ordered by study ID]

Study	Reason for exclusion
Asfar 2014	3-month follow up

## DATA AND ANALYSES

### Comparison 1. Intervention versus control


Outcome or subgroup title	No. of studies	No. of participants	Statistical method	Effect size
<a href="#">1 Prolonged Cessation</a>	3		Risk Ratio (M-H, Fixed, 95% CI)	Subtotals only

#### Analysis 1.1. Comparison 1 Intervention versus control, Outcome 1 Prolonged Cessation.

Review: Interventions for waterpipe smoking cessation

Comparison: 1 Intervention versus control

Outcome: 1 Prolonged Cessation



(1) Combined behavioural and (behavioural + bupropion) groups

## APPENDICES

### Appendix I. Glossary of tobacco-related terms

Term	Definition
Abstinence	A period of being quit, i.e. stopping the use of cigarettes or other tobacco products. May be defined in various ways; see also: point prevalence abstinence; prolonged abstinence; continuous/sustained abstinence
Biochemical verification	Also called 'biochemical validation' or 'biochemical confirmation': A procedure for checking a tobacco user's report that he or she has not smoked or used tobacco. It can be measured by testing levels of nicotine or cotinine or other chemicals in blood, urine, or saliva, or by measuring levels of carbon monoxide in exhaled breath or in blood
Bupropion	A pharmaceutical drug originally developed as an antidepressant, but now also licensed for smoking cessation; trade names Zyban, Wellbutrin (when prescribed as an antidepressant)
Carbon monoxide (CO)	A colourless, odourless highly poisonous gas found in tobacco smoke and in the lungs of people who have recently smoked, or (in smaller amounts) in people who have been exposed to tobacco smoke. May be used for biochemical verification of abstinence
Cessation	Also called 'quitting'. The goal of treatment to help people achieve abstinence from smoking or other tobacco use, also used to describe the process of changing the behaviour
Continuous abstinence	Also called 'sustained abstinence'. A measure of cessation often used in clinical trials involving avoidance of all tobacco use since the quit day until the time the assessment is made. The definition occasionally allows for lapses. This is the most rigorous measure of abstinence
'Cold Turkey'	Quitting abruptly, and/or quitting without behavioural or pharmaceutical support
Craving	A very intense urge or desire [to smoke]. See: Shiffman et al 'Recommendations for the assessment of tobacco craving and withdrawal in smoking cessation trials' Nicotine & Tobacco Research 2004; 6(4): 599-614
Dopamine	A neurotransmitter in the brain which regulates mood, attention, pleasure, reward, motivation and movement
Efficacy	Also called 'treatment effect' or 'effect size': The difference in outcome between the experimental and control groups
Harm reduction	Strategies to reduce harm caused by continued tobacco/nicotine use, such as reducing the number of cigarettes smoked, or switching to different brands or products, e.g. potentially reduced exposure products (PREPs), smokeless tobacco

(Continued)

Lapse/slip	Terms sometimes used for a return to tobacco use after a period of abstinence. A lapse or slip might be defined as a puff or two on a cigarette. This may proceed to relapse, or abstinence may be regained. Some definitions of continuous, sustained or prolonged abstinence require complete abstinence, but some allow for a limited number or duration of slips. People who lapse are very likely to relapse, but some treatments may have their effect by helping people recover from a lapse
nAChR	[neural nicotinic acetylcholine receptors]: Areas in the brain which are thought to respond to nicotine, forming the basis of nicotine addiction by stimulating the overflow of dopamine
Nicotine	An alkaloid derived from tobacco, responsible for the psychoactive and addictive effects of smoking
Nicotine Replacement Therapy (NRT)	A smoking cessation treatment in which nicotine from tobacco is replaced for a limited period by pharmaceutical nicotine. This reduces the craving and withdrawal experienced during the initial period of abstinence while users are learning to be tobacco-free. The nicotine dose can be taken through the skin, using patches, by inhaling a spray, or by mouth using gum or lozenges
Outcome	Often used to describe the result being measured in trials that is of relevance to the review. For example smoking cessation is the outcome used in reviews of ways to help smokers quit. The exact outcome in terms of the definition of abstinence and the length of time that has elapsed since the quit attempt was made may vary from trial to trial
Pharmacotherapy	A treatment using pharmaceutical drugs, e.g. NRT, bupropion.
Point prevalence abstinence (PPA)	A measure of cessation based on behaviour at a particular point in time, or during a relatively brief specified period, e.g. 24 hours, 7 days. It may include a mixture of recent and long-term quitters. cf. prolonged abstinence, continuous abstinence
Prolonged abstinence	A measure of cessation which typically allows a 'grace period' following the quit date (usually of about two weeks), to allow for slips/lapses during the first few days when the effect of treatment may still be emerging
Relapse	A return to regular smoking after a period of abstinence.
Secondhand smoke	Also called passive smoking or environmental tobacco smoke [ETS]. A mixture of smoke exhaled by smokers and smoke released from smouldering cigarettes, cigars, pipes, bidis, etc. The smoke mixture contains gases and particulates, including nicotine, carcinogens and toxins
Self-efficacy	The belief that one will be able to change one's behaviour, e.g. to quit smoking
SPC [Summary of Product Characteristics]	Advice from the manufacturers of a drug, agreed with the relevant licensing authority, to enable health professionals to prescribe and use the treatment safely and effectively



(Continued)

Tapering	A gradual decrease in dose at the end of treatment, as an alternative to abruptly stopping treatment
Titration	A technique of dosing at low levels at the beginning of treatment, and gradually increasing to full dose over a few days, to allow the body to get used to the drug. It is designed to limit side effects
Withdrawal	A variety of behavioural, affective, cognitive and physiological symptoms, usually transient, which occur after use of an addictive drug is reduced or stopped. See: Shiffman et al 'Recommendations for the assessment of tobacco craving and withdrawal in smoking cessation trials' . Nicotine & Tobacco Research 2004; 6(4): 599-614

## WHAT'S NEW

Last assessed as up-to-date: 19 June 2015.

Date	Event	Description
13 March 2015	New citation required and conclusions have changed	Three new citations identified. Conclusions updated.
14 December 2014	New search has been performed	Searches updated, three studies identified and included.

## HISTORY

Protocol first published: Issue 4, 2005

Review first published: Issue 4, 2007

Date	Event	Description
4 August 2008	Amended	Converted to new review format.
9 August 2007	New citation required and conclusions have changed	Substantive amendment

## CONTRIBUTIONS OF AUTHORS

All authors contributed to the conceptualisation and preparation of the review, and approved the final text. MJ and SJ assessed the studies retrieved by the search strategy, and updated the Methods and Results sections accordingly. TA updated the Discussion according to the new results. WM and KW updated all the manuscript in light of new literature on the subject, and the reviewed articles. TE reviewed the updated version, edited and finalized it.

## DECLARATIONS OF INTEREST

KW has no known conflicts of interest

MJ has no known conflicts of interest

SJ has no known conflicts of interest

TA has no known conflicts of interest

WM is funded by National Institute on Drug Abuse (NIDA) grant R01 DA035160. WM's research in tobacco control, including the waterpipe, has been funded by the US National Institutes of Health since 2002.

TE's research on waterpipe tobacco smoking and various other tobacco products, including electronic cigarettes, has been supported by grants from the U.S. National Institutes of Health and is currently supported by the U.S. National Institutes of Health's National Institute on Drug Abuse and the Center for Tobacco Products of the US Food and Drug Administration (P50DA036105). TE also occasionally receives honoraria for delivering invited lectures regarding tobacco product effects to a variety of university audiences.

The content of the work published herein is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the Food and Drug Administration.

## SOURCES OF SUPPORT

### Internal sources

- No sources of support supplied

### External sources

- US Public Health Service Grants TW05962, TW07233 USA; Initiative for Cardiovascular Health Research in the Developing Countries (IC-Health), India.

## DIFFERENCES BETWEEN PROTOCOL AND REVIEW

None.

## **INDEX TERMS**

### **Medical Subject Headings (MeSH)**

\*Water; Bupropion [therapeutic use]; Randomized Controlled Trials as Topic; Smoking [\*prevention & control]; Smoking Cessation [methods; statistics & numerical data]; Tobacco Use Cessation [\*methods; statistics & numerical data]

### **MeSH check words**

Female; Humans; Male