

Intestinal carriage of *Staphylococcus aureus*: how does its frequency compare with that of nasal carriage and what is its clinical impact?

D. S. Acton · M. J. Tempelmans Plat-Sinnige ·
W. van Wamel · N. de Groot · A. van Belkum

Received: 16 June 2008 / Accepted: 11 July 2008 / Published online: 8 August 2008
© The Author(s) 2008

Abstract The bacterial species *Staphylococcus aureus*, including its methicillin-resistant variant (MRSA), finds its primary ecological niche in the human nose, but is also able to colonize the intestines and the perineal region. Intestinal carriage has not been widely investigated despite its potential clinical impact. This review summarizes literature on the topic and sketches the current state of affairs from a microbiological and infectious diseases' perspective. Major findings are that the average reported detection rate of intestinal carriage in healthy individuals and patients is 20% for *S. aureus* and 9% for MRSA, which is approximately half of that for nasal carriage. Nasal carriage seems to predispose to intestinal carriage, but sole intestinal carriage occurs relatively frequently and is observed in 1 out of 3 intestinal carriers, which provides a rationale to include intestinal screening for surveillance or in outbreak settings. Colonization of the intestinal tract with *S. aureus* at a young age occurs at a high frequency and may affect the host's immune system. The frequency of intestinal carriage is generally underestimated and may significantly contribute to bacterial dissemination and subsequent risk of infections. Whether intestinal rather than nasal *S. aureus* carriage is a primary predictor for infections is still ill-defined.

Introduction

Nasal colonization by *Staphylococcus aureus* is a well-established risk factor for acute cutaneous infections, post-operative infections, as well as most other types of *S. aureus* infections [1–3]. Several recent studies have suggested that lasting colonization of *S. aureus* in the human intestinal tract also occurs and that this may have important clinical implications. Still, compared to nasal carriage, gastro-intestinal colonization by *S. aureus* has been sparsely studied.

In the 1950s and 1960s, intestinal *S. aureus* carriage was first defined and studied as a potential cause of antibiotic-associated diarrhea (AAD) [4]. However, after the identification of *Clostridium difficile* as the most common pathogen of hospital-acquired AAD in the 1970s [5], the role of gastro-intestinal colonization as a risk factor for (intestinal) *S. aureus* infection has been neglected for decades and the issue has only recently re-emerged. The pandemic rise in the incidence of methicillin-resistant *S. aureus* (MRSA) strains, as opposed to methicillin-sensitive *S. aureus* (MSSA), has contributed greatly to the renewed interest in intestinal *S. aureus* colonization. This covered investigations into risk factors for both AAD as well as health care-associated (HA) infections. Furthermore, community-acquired (CA) MRSA infection among individuals without “HA risk factors” was first recognized in the late 1990s. CA-MRSA is now emerging as an apparent epidemic. In recent studies, in addition to nasal carriage, rectal carriage of CA-MRSA has also been documented. In this review we will give an historical update and a systemic overview of the more recent studies related to intestinal and/or perineal carriage of *S. aureus* (MSSA, HA-MRSA, and CA-MRSA) and we will discuss its clinical implications.

D. S. Acton (✉) · M. J. Tempelmans Plat-Sinnige · N. de Groot
Mucovax B.V.,
Niels Bohrweg 11–13,
2333 CA Leiden, The Netherlands
e-mail: dacton@mucovax.nl

M. J. Tempelmans Plat-Sinnige · W. van Wamel · A. van Belkum
Department of Medical Microbiology
and Infectious Diseases, Erasmus MC,
Gravendijkwal 230,
3015 CE Rotterdam, The Netherlands

Culture methods and definition of perineal and intestinal carriage

Techniques to determine carriage or colonization can be based on classical culture assays using selective broths or agars for MSSA, as well as novel growth-based phenotypic and molecular methods for both MSSA and MRSA detection [6]. Methods used to define intestinal carriage include culture of stool, rectal swabs or anal swabs. Also, swabs from the perianal area (including the perineum and the groin or inguinal region) are generally accepted to define intestinal carriage. For these sites, however, it can be argued that these may also represent skin carriage. In a limited number of reports direct comparison of the frequency yields for these different sites of colonization within the same study have been presented. Rectal swabs were reported to have higher yields of *S. aureus* detection than stool cultures [7], but a more recent comparison [8] gave opposite results for MRSA. Other studies reported similar frequencies of carriage for both perianal and groin sites of carriage for nursing home residents and hospitalized patients respectively [9, 10]. Also, direct comparisons of the rectal and perineal sites gave similar frequencies of detection [11] and in another recent study screening of the perineal area, the rectum and the inguinal area gave similar frequencies of detection of MRSA [12], which legitimates the use of all of these sites to define intestinal carriage.

Intestinal carriage frequencies of *S. aureus* in adults

Most early studies on the intestinal carriage of *S. aureus* were performed on hospitalized patients and timely overviews of frequencies of intestinal carriage in adults upon admission to the hospital were reported [4, 13]. The frequencies found in these relatively ancient studies ranged from 8 to 31% (Table 1). Among adults, nasal *S. aureus* carriers were reported to yield *S. aureus* from their feces more often than non-nasal carriers did. Strain typing suggested involvement of the same strains for both colonization sites [14]. Vice versa, about 50–70% of perineal carriers were also nasal carriers of the same strain [15]. It was reported that of 50 healthy male students screened, 11 (22%) had perineal carriage of *S. aureus* and that 5 of these were non-nasal carriers (10%) [16]. This implied that non-nasal carriers might be susceptible to intestinal carriage, suggesting that mechanisms for nasal and intestinal carriage differ.

More recent data on the intestinal carriage of *S. aureus* in adults were generated as part of studies mostly aimed at identifying MRSA. In most of these studies the patients were screened for intestinal as well as nasal carriage, allowing a comparison of these sites of colonization

Table 1 Reported frequencies of intestinal carriage of *S. aureus* in adults upon admission to hospital in early studies

Reference	Percentage positives
[92]	18
[93]	17
[14]	23
[94]	31
[95]	8
[16]	22
[96]	21
[15]	12
[97]	13

(Table 2). For instance, in a study on 500 pregnant women attending an antenatal clinic a frequency of intestinal carriage of *S. aureus* of 12% (59 patients) and a nasal carriage frequency of 24% (120 patients) were reported [17]. Interestingly, 8% (41) of the patients had intestinal carriage in the absence of nasal carriage, covering 25% of all *S. aureus*-positive patients. Among patients in a skilled nursing facility, nasal and rectal swabs or stools were surveyed for MSSA and MRSA. Of these, 22% (76 patients) had intestinal carriage of *S. aureus* [18]. Ten percent of the patients ($n=34$) were intestinal carriers in the absence of nasal carriage.

The prevalence of *S. aureus* colonization of the perineum was examined in a longitudinal study on 84 community-dwelling adults with spinal cord dysfunction [19]. *S. aureus* was detected in 20 (24%) individuals. Nasal carriage was detected in 55% of all patients from a subset of 22 patients of whom 23% had intestinal carriage only. By follow-up of 30 patients with at least five cultures for 1 year, the perineal carriage pattern was assessed. They found that 10% had persistent carriage, 63% had intermittent carriage, and 27% were non-carriers, in agreement with nasal carriage patterns [1]. Paired perineal/nasal carriage was determined for 22 participants. Of the 16 perineal carriers in this group, 5 did not have nasal carriage. Among the 11 with both perineal and nasal carriage, all but 1 carried the same *spa* type at both sites. Many similar studies were performed, the results of which are summarized in Table 2.

Altogether the frequency of nasal carriage detection of *S. aureus* ranged from 24% to 61% and intestinal carriage frequencies in these studies ranged from 10% to 37%. These ranges probably reflect the different criteria used for the selection of patient groups and the differences in risk of *S. aureus* colonization or infection amongst these heterogeneous groups of patients from different geographical regions. Therefore, calculation of an overall frequency of *S. aureus* carriage from these studies by using actual total numbers of patients may be debated, but, nevertheless, results in approximately 20% of cases of intestinal carriage

Table 2 Frequencies of detection of *S. aureus* (including MRSA) with regard to intestinal and nasal carriage and intestinal carriage in the absence of nasal carriage in adults

Reference	Total number of patients screened	Intestinal carriage, percentage of total (<i>n</i> /total)	Nasal carriage, percentage of total (<i>n</i> /total ^a)	Intestinal carriage in the absence of nasal carriage, percentage of total (<i>n</i> /total ^a)	Intestinal carriage in the absence of nasal carriage, percentage of intestinal carriers (<i>n</i> /total ^a)	Method used	Patient group
[17]	500	12 (59)	24 (120)	8 (41)	70	Perineal swabs	Pregnant women
[77]	37	24 (9)	30 (15/50)	NS	NS	Rectal swab	Mothers 1 week after delivery
[55]	62	11 (7)	44 (27)	3 (2)	29	Perineal swabs	Healthy adults
[18]	354	22 (76)	61 (214)	10 (34)	45	Stools	Private SNF patients
[48]	204	29 (59)	47 (96)	3 (7)	12	Rectal swabs	ICU liver transplant patients
[24]	231	10 (24)	30 (70)	5 (11)	46	Perineal swabs	Intensive care patients
[90]	94	37 (35)	49 (47/96)	10 (9)	26	Rectal swabs	Liver transplant recipients
[44]	71	37 (26)	51 (36)	4 (3)	12	Stools	Selected inpatients
[10]	213	25 (54)	43 (91)	NS	NS	Perianal swabs	Nursing home residents
Totals	1,766	20 (349/1,766)	40 (716/1,781)	8 (112/1,538)	37 (112/302)		

NS, not specified; SNF, skilled nursing facilities; ICU, intensive care unit

^a Totals are indicated when frequencies are based on a different total number of patients screened

compared with nasal carriage of approximately 40%. When reported frequencies of the separate studies are plotted, it appears as if there is a linear correlation between the incidence of nasal and intestinal carriage (see Fig. 1). The slope of the linear regression is 0.55, which suggests that intestinal carriage is probably preceded by nasal carriage. However, sole intestinal carriage also seems to occur with frequencies ranging from 3–10%. When intestinal carriage in the absence of nasal carriage is calculated from these studies using actual numbers of patients a figure of 8% is obtained. Furthermore, on average, 1 out of 3 intestinal carriers (37%) does not seem to be colonized in the nares (Table 2).

Obviously, compared with nasal carriage the intestinal carriage of *S. aureus* is less frequent, but still may have important clinical consequences.

Intestinal MRSA in adult patients at high risk

During the last 2 decades, a number of studies have reported intestinal screening to determine carriage of MRSA beyond the nasal cavity, in patients at high risk of MRSA at admission or during a stay in a hospital (Table 3). Long-term care patients were analyzed and detection of MRSA was reported in stools from 29 out of 354 patients (8%) [18]. Nasal carriage frequency was 15% (52 patients). The detection of 2 MRSA patients on the basis of 10 stool cultures (20%), obtained during an outbreak in an institution for adults with developmental disabilities, was reported, whereas nasal screening detected MRSA in 16 out of 28 patients (57%) [20]. A frequency of intestinal carriage of 10% (8 out of 84) in community-dwelling

patients with spinal cord dysfunction was recorded [19]. Nasal screening of a subgroup of 22 of these patients revealed a nasal MRSA frequency of 55% (12 patients). Furthermore, 205 patients were analyzed who were known to be previously colonized and/or infected with MRSA. Intestinal carriage detection frequency was 33% and nasal carriage detection frequency was 52% [21]. Many other reports describing similar data are surveyed in Table 3. When totals are calculated from these reports, the average nasal carriage frequency in these patients at high risk was 12% and intestinal carriage frequency was 9%.




Fig. 1 Relation between frequencies of nasal and intestinal carriage for *S. aureus* (circles) and for MRSA (diamonds). Lines show the linear regression of *S. aureus* (dotted line) and MRSA (straight line), the slope of the linear regression lines are 0.55 and 0.53, and R^2 are 0.6012 and 0.7381 for *S. aureus* and MRSA respectively. The plotted data for *S. aureus* are extracted from: [3, 10, 17–19, 24, 44, 48, 77, 90] and for MRSA from: [10, 18–21, 23, 24, 27, 31, 33, 36, 38, 39, 51, 91]

Table 3 The MRSA intestinal carriage and nasal carriage detection frequencies in hospitalized adult patients at high risk

Reference	Total patients screened	Intestinal MRSA, percentage of total (<i>n</i>)	Nasal MRSA percentage of total (<i>n</i> /total ^a)	Method used
[18]	354	8 (29)	15 (52)	Stools
[38]	114	5 (6)	14 (16)	Rectal swabs
[51]	327	11 (36)	24 (117/484)	Stools
[20]	10	20 (2)	57 (16/28)	Perineal swabs
[33]	411	5 (22)	7 (31)	Perineal swabs
[91]	105	10 (10)	13 (34/256)	Perineal swabs
[24]	231	5 (11)	10 (23)	Perineal swabs
[27]	1,250	5 (57)	7 (90)	Rectal swabs
[23]	192	10 (20)	13 (25)	Groin swabs
[31]	1,181	13 (151)	13 (155)	Groin swabs
[39]	845	6 (54)	6 (54)	Rectal swabs
[19]	84	24 (20)	37 (17/46)	Perineal swabs
[21]	52	33 (17)	52 (53/101)	Groin swabs
[36]	758	8 (60)	8 (57)	Perineal swabs
[10]	213	16 (35)	26 (56)	Perineal swabs
Totals	6,127	9 (530/6,127)	12 (796/6,464)	

^aTotals are indicated when frequencies are based on different numbers of total patients screened.

Interestingly, when frequencies of nasal and intestinal MRSA carriage are plotted and compared with those reported for *S. aureus* (Fig. 1) it seems that the incidence of carriage of MRSA is lower than that for *S. aureus* for both the nares and the intestines, which makes sense, since MRSA has been emerging and disseminating as a colonizing bacterium much more recently than *S. aureus* has. However, the slope of the regression of 0.53 for MRSA is the same as that determined for *S. aureus* (0.55), which indicates a similar relation between both sites of carriage for MRSA and suggests that antibiotic resistance does not influence the relative colonization ability for these two sites.

Distribution of MRSA detection sites

Direct comparison of the distribution of intestinal carriers versus nasal carriers amongst MRSA-colonized patient groups could be deduced from 22 reports (Table 4) [9, 10, 13, 18, 19, 22–29, 31–39]. Table 4 surveys the frequency of patients with intestinal carriage and nasal carriage expressed as a percentage of all MRSA-colonized patients found in the various studies. In general, patients were considered colonized when at least two consecutive cultures from any place in the body grew MRSA, but also single colonization criteria were applied. The frequencies reported ranged from 5 to 76% and from 34 to 84% for intestinal and nasal carriage respectively. When the distribution of intestinal and nasal carriage in MRSA-colonized patients is calculated from all of these studies comprising more than 2,000 MRSA-colonized/infected patients, an overall contribution of 45% is found for intestinal carriage and 58% for nasal carriage for all MRSA-colonized patients.

In Table 4 the reported frequencies of intestinal MRSA carriage in the absence of nasal carriage are also summarized. For instance, in a study of an MRSA outbreak involving a total of 975 MRSA patients, perineal screening identified an additional 15% of culture-positive individuals compared with nasal screening only [25]. In this study, the intestines were reported to be the only positive site out of five culture sites in 10% of the cases. Klotz and coworkers reported that in addition to a high frequency of 24% MRSA-positive stool cultures, 13% of the MRSA strains were first observed in the stool before detecting MRSA in other material from these patients [30]. Similarly, in a large Canadian surveillance study [40], comprising more than 10,000 adult patients, the perineal area was found to be the initial site of MRSA colonization or infection in 41% of the cases. In patients older than 65 years of age, the perineum or rectum was the only positive site in 13%, indicating that nasal screening alone would be sub-optimal in elderly patients. More recently, Zhang et al. [39] reported that nasal screening alone would have detected 76% of MRSA carriers and that the inclusion of rectal screening increased the detection sensitivity to 96%, indicating 20% rectal carriers without nasal carriage. Reyes et al. [36] reported an intestinal carriage frequency without nasal carriage in 27% of their MRSA-positive patients, whereas Mody et al. [10] showed that 23% of their MRSA carriers were only colonized in the perianal area. Altogether, comparisons of rates of nasal and intestinal carriage of MRSA were recorded in 16 of the studies mentioned (Table 4). It can be calculated from these studies that of all MRSA colonized patients, 58% were colonized in the nares, whereas 45% were colonized in the intestines. Of the individuals with gastro-intestinal MRSA, 1 out of 3 (35%) did not carry

Table 4 Distribution of intestinal and nasal carriage in all MRSA-colonized patients detected in the studies

References	Total MRSA cases	Intestinal MRSA, percentage of total (n)	Nasal MRSA, percentage of total (n/total ^a)	Intestinal without nasal, percentage of total (n)	Intestinal without nasal, percentage of intestinal carriers	Method used	Patient group
[22]	63	5 (3)	34 (24/79)	Not specified	Not specified	Perineal swabs	General hospital patients, MRSA outbreak
[13]	117	61 (70)	53 (62)	Not specified	Not specified	Rectal swabs	General hospital patients, MRSA outbreak
[26]	11	18 (2)	55 (6)	0 (0)	0	Anal swabs	SCI patients
[32]	67	25 (17)	58 (37)	Not specified	Not specified	Perineal swabs	SCI patients
[25]	723	40 (289)	47 (369/789)	15 (145)	50	Perineal swabs	Hospital MRSA outbreak
[18]	62	47 (29)	84 (52)	16 (10)	35	Stools	Private skilled nursing facility patients
[35]	23	17 (4)	65 (15)	9 (2)	50	Perineal swabs	Acute care patients
[34]	19	42 (8)	68 (13)	5 (1)	13	Perineal swabs	Acute rehabilitation unit SCI patients
[38]	24	25 (6)	67 (16)	8 (2)	33	Rectal swabs	Skilled care unit patients
[33]	51	54 (22)	61 (31)	18 (9)	41	Perianal swabs	Acute rehabilitation unit
[37]	36	50 (18)	75 (27)	25 (9)	50	Groin swabs	General hospital, at risk patients
[28]	203	26 (52)	44 (89)	Not specified	Not specified	Perineal swabs	Neurosurgery unit patients
[29]	35	20 (7)	75 (21)	Not specified	Not specified	Perineal swabs	General hospital, at risk patients
[24]	30	37 (11)	77 (23)	17 (5)	46	Perineal swabs	ICU patients
[9]	96	49 (47)	72 (72)	Not specified	Not specified	Perineal swabs	University hospital, at risk patients
[27]	123	46 (57)	73 (90)	19 (23)	40	Rectal swabs	ICU patients
[23]	31	65 (20)	80 (25)	19 (6)	30	Groin swabs	General hospital, at risk patients
[31]	224	67 (151)	69 (155)	12 (26)	17	Groin swabs	ICU patients
[39]	71	76 (54)	76 (54)	20 (14)	26	Rectal swabs	General hospital, at risk patients
[19]	22	73 (16)	71 (12/17)	30 (5)	31	Perineal swabs	Community-dwelling SCD patients
[36]	78	77 (60)	73 (57)	27 (21)	35	Perineal swabs	ICU patients and other risk factors
[10]	86	41 (35)	65 (56)	11 (9)	11	Perianal swabs	Nursing home residents
Totals	2,195	45 (978/2,195)	58 (1,306/2,268)	18 (287/1,614)	35 (287/833)		

SCI, spinal cord injury; SCD, spinal cord dysfunction

^aTotals are indicated when frequencies reported are based on different total numbers of cases

MRSA in the nose. Of all MRSA-colonized individuals detected in these studies more than 1 out of 6 (18%) presented with intestinal carriage in the absence of nasal carriage. Such frequencies suggest that the intestines might be clinically relevant reservoirs of MRSA that should be taken into account during screening for such carriage during outbreak situations or when screening patients at risk.

Intestinal or perineal carriage as a risk factor for dissemination and infections

The perineum, as an area where *S. aureus* can colonize and multiply, was first recognized by Hare and Ridley [41], and

heavy contamination from the perineum to the groin and upper parts of the thighs is often observed [42]. Both intestinal and perineal carriage have been implicated as important contributors to environmental dissemination of *S. aureus*. Patients with intestinal colonization of *S. aureus* may serve as an important source of transmission, since they often contaminate the adjacent environment. Masaki et al. [43] performed a prospective culture survey to investigate a possible relationship between *S. aureus* types colonizing the rectum and respiratory tract and *S. aureus* types isolated from the environment. They simultaneously detected in both patients and the environment several *S. aureus* types. This indicates a potential route of contamination of the hospital environment. It was also reported that

patients with intestinal and nasal colonization of *S. aureus* had higher frequencies of incontinence or diarrhea than patients without *S. aureus* colonization. This may significantly contribute to the observed trend toward increased contamination of environmental surfaces [44, 45]. The latter studies also substantiated that patients who have diarrheal stools and heavy gastro-intestinal colonization with MRSA are associated with significantly greater environmental MRSA contamination than patients without MRSA in their stools.

Data from early studies suggested an association between intestinal colonization and the occurrence of infections. For example, a study on recurrent furunculosis showed that 56% of the patients sampled had positive perineal cultures [46]. This suggested intestinal *S. aureus* carriage to be an infection risk factor, although nasal carriage was not precisely assessed for those patients. Hospitalized patients who reported developing *S. aureus* lesions on the buttocks and lower half of the abdomen and back often had the causative strain isolated from the perineum as well [47]. It was found that intensive care and liver transplant unit patients with both nasal and intestinal colonization had significantly increased rates of *S. aureus* infection of 40% compared with an infection rate of 18% in patients with nasal carriage without intestinal carriage [48]. On the basis of these data it was concluded that rectal carriage represents a potential reservoir and simultaneous nasal and rectal carriage portended a greater risk of *S. aureus* infections than nasal carriage alone in ICU and liver transplant unit patients. The same authors also proposed that intestinal colonization could be associated with an increased frequency of colonization of skin sites, which was confirmed in later studies [44]. It was found that patients with nasal and intestinal colonization were significantly more likely than those with nasal colonization only to have positive skin cultures. This group performed a prospective study involving 71 patients. The patients enrolled were divided into three groups: those without nasal or intestinal colonization, those with nasal colonization only, and those with both nasal and intestinal colonization. The development of *S. aureus* infections was significantly different among the three groups. Only 1 out of 32 patients without nasal or intestinal colonization developed an *S. aureus* infection. *S. aureus* infections developed more often in patients with stool colonization (8 out of 26) than in patients with only nasal colonization (2 out of 13). However, due to the small number of infected patients the power of this study was too limited to reach statistical significance. Significant differences in staphylococcal infection between patients with *S. aureus* in the stools and patients with negative stool cultures for *S. aureus* were documented [49]. In an 8-month prospective study of inpatients known to have vancomycin-resistant enterococci (VRE) colonization, none of the 14

patients with stool cultures negative for *S. aureus* developed an *S. aureus* infection in the hospital. In contrast, more than half of *S. aureus*-colonized patients had an *S. aureus* infection documented.

Eradication of intestinal carriage

Given its relatively high incidence, prevention of or therapy for intestinal carriage should be clinically beneficial. However, most studies are focused on the elimination of nasal rather than intestinal carriage. Nasal carriage eradication with mupirocin ointment has been studied frequently (recently reviewed by Van Rijen et al. [50]) and is generally considered to be highly effective, at least in the short term. However, reacquisition of carriage may occur from extranasal sites. Dupeyron et al. [51] monitored mupirocin treatment of 86 patients and found treatment failure in 22. For this group, the stool carriage rate was significantly higher and stool carriage upon admission was independently associated with reacquisition of nasal carriage. Conflicting results regarding the effects of nasal carriage eradication on the prevalence of MSSA/MRSA infections have been reported. Some authors showed significantly decreased rates of nosocomial infections [52, 53], whereas others did not [54, 55].

Mupirocin treatment may only be marginally effective in the eradication of multi-site carriage and, therefore, therapies based upon combinations of nasal, skin, and intestinal carriage eradication methods have more recently been exploited. The use of oral rifampin, for treatment of intestinal carriage of *S. aureus* for various patient populations and healthy people was reviewed by Falagas et al. [56, 57]. Rifampin, however, showed limited success in MRSA elimination. More novel treatment modalities for intestinal MRSA elimination to control transmission or subsequent infections, e.g., using oral vancomycin, have been described merely in uncontrolled or observational studies. Oral vancomycin treatment results were reported that showed that the eradication of MRSA intestinal carriage by enteral vancomycin in subsets of adult ICU patients [58–60] as well as in pediatric patients [61, 62] was effective, but had limited effect on the prevention of transmission. The results from prospective controlled studies of intestinal MRSA decolonization are urgently awaited.

Colonization of the intestinal tract by *S. aureus* may have another important clinical implication. The co-existence of *S. aureus* and VRE was reported in more than 50% of the American patients studied [49]. In this study, stool specimens were tested for VRE and *S. aureus* at enrollment (baseline) and weekly thereafter. Patients with at least three stool cultures were included in the study. Of the 37 patients who completed the study, all were colonized with VRE; 62%

were colonized with *S. aureus* on at least one occasion, and 60% were colonized persistently. Patients with stool cultures positive for *S. aureus* were colonized with MRSA strains in 87% of all cases. Warren et al. [63] screened stools and rectal samples from 878 ICU patients for VRE and MRSA. Of 485 VRE-positive patients, 83 (17%) also had intestinal carriage of MRSA. Furuno et al. [64] reported that out of 57 patients with both nasal MRSA carriage and intestinal VRE, 23 (40%) also had intestinal carriage of MRSA. These studies suggest that the intestinal tract could provide an important reservoir for the emergence of vancomycin-resistant *S. aureus* (VRSA) isolates

The emergence of the rectal carriage of CA-MRSA has been documented in a study in which *S. aureus* was detected in 507 of 2,963 vaginal/rectal cultures from late pregnancy cases (17%) [65]. Interestingly, 3% of the pregnant women had vaginal–rectal colonization of an epidemic CA-MRSA strain. A significant association between *S. aureus* colonization and Group B streptococcus (GBS) colonization was found. In a subsequent study this association was analyzed in more detail [66] and the significant association between MSSA and GBS was confirmed. Pregnant women with MSSA carriage were significantly more likely to have postpartum fever than those who were *S. aureus*-negative. Surprisingly, a significant negative association between CA-MRSA carriage and GBS carriage was observed. Apparently, when a certain bacterial species inhabits a given niche other species may not be able to colonize. In the case of the vaginal carriage of MRSA or MSSA such bacterial interference may lead to novel modes of intervention based on interference therapy of GBS once the microbial molecules involved have been identified.

Potential role of *S. aureus* in intestinal disease

How *S. aureus* causes intestinal infections is still ill-defined. The basic mechanisms are quite enigmatic and the etiological processes are just beginning to be identified. For instance, Froberg et al. [67] presented histo-pathological evidence for the existence of a specific *S. aureus*-induced pseudomembranous intestinal disease, distinct from that seen during *C. difficile* infection in an unusual case of simultaneous infection with *C. difficile* and MRSA. Whereas *C. difficile* induces colonic pseudomembranes, the MRSA infection induced loosely adherent pseudomembranes in the small bowel.

Intestinal carriage of *S. aureus* may impose a risk factor for intestinal infection. Antibiotic treatment can lead to the overgrowth of bacteria in the intestine and induce enteritis or AAD. The role of intestinal *S. aureus* as a causative agent for enteritis or AAD and as a risk factor for other

infections gained renewed interest with the spreading of MRSA. MRSA has been suggested as a cause of AAD in hospitalized patients [68]. In this study *S. aureus* was the only identified pathogenic micro-organism to cause AAD in 47 patients and the presence of staphylococcal enterotoxin A was strongly associated with the development of diarrhea. A German study reported intestinal carriage of *S. aureus*, in the absence of *C. difficile* in 8% of patients with AAD [69]. Compelling evidence for the etiological role of MRSA in AAD was provided by excluding the involvement of *C. difficile*, numerous other bacterial pathogens and parasites, but also several enteric viruses in 11 patients with enterotoxin-producing MRSA intestinal carriage [70].

An extensive analysis of the prevalence of *C. difficile* and *S. aureus* in 2,727 stool samples of patients with diarrhea was performed. *C. difficile* grew from 148 specimens and 184 were positive for *C. difficile* toxin A/B analysis and altogether, a total of 252 stool samples were positively diagnosed with *C. difficile*. *S. aureus* was grown out of 198 fecal samples, of which 29 were identified as having MRSA [71]. In another study 10 MRSA-positive samples were detected out of 4,659 fecal samples [72]. Table 5 conclusively defines the interrelatedness between intestinal colonization by *S. aureus* and *C. difficile*.

Intestinal *S. aureus* colonization and disease development in infants and young children

In infants very high frequencies for intestinal *S. aureus* carriage were reported in early studies (reviewed by Williams [4], Table 6) and some of these suggested that acquisition of *S. aureus* occurred very early in life and probably as a consequence of nasal acquisition. Intestinal colonization in children was also studied more recently [73]. One hundred patients, below 16 years of age and attending the emergency department of a university hospital, who were analyzed for nasal and perineal carriage, included 20 *S. aureus*-colonized patients, of whom 2 (10%) had intestinal carriage in the absence of nasal carriage. Of this group, 17 patients (85%) had nasal carriage. Others studied *S. aureus* carriage in a child care center [74]. Of

Table 5 Frequency of detection of *C. difficile* and *S. aureus* in antibiotic-associated diarrhea (AAD) patients

Reference	Total	<i>C. difficile</i> , percentage (n)	<i>S. aureus</i> , percentage (n)
[68]	3,437	13 (460)	2 (60)
[70]	1,543	10 (159)	10 (151)
[69]	89	44 (39)	28 (25)
[72]	4,659	13 (591)	0.2 (10)
[71]	2,727	9 (252)	7 (198)

Table 6 *S. aureus* intestinal carriage in infants

References	Cases	Percentage	Method	Age
[92]	83	56	Stools	1–10 days
	22	100		2–6 months
	32	50		6–12 months
[76]	62	61	Stools	2 years
[73]	100	19	Perineal swabs	Hospitalized children up to 16 years
[78]	49	16	Rectal/stools	3 days
		57		1 week
		65		2 weeks
		65		4 weeks
		73		8 weeks
		73		6 months
		53		1 year
[98]	44	59	Stools	1 week
		61		1 month
		50		3 months
		39		6 months
		32		12 months
[77]	50	20	Rectal/stools	3 days
		40		1 week
		52		2 weeks
		60		4 weeks
		64		8 weeks
[79]	53	62	Rectal swabs	2 weeks
		70		4 weeks
[80]	324	13	Rectal swabs	3 days
		39		7 days
		52		14 days
		63		28 days
		72		2 months
		79		6 months

128 children who had swabs taken from nose, perineum, and throat, 8 (24%) had perineal carriage. Nasal and throat carriage frequency were higher with 15 (46%) and 22 (67%) respectively. An African study investigated the incidence of *S. aureus* in children aged 5 years and below suffering from sporadic diarrhea in Nigeria [75]. Out of 1,761 diarrheic fecal specimens collected, only 72 (4%) were positive for *S. aureus*.

Also, in more recent Swedish studies [76–79], high frequencies of intestinal *S. aureus* carriage during the first year of life were reported, and co-colonization of intestine and anterior nares with the same *S. aureus* strains, which were also found on the parents, suggested mother-to-child transmission. These studies were performed to test the hypothesis that development of allergic disease among children may be associated with differences in intestinal colonization patterns of *S. aureus*. Two-year-old allergic children from both Sweden and Estonia were reported to have significantly higher counts of *S. aureus* in the

intestines than non-allergic children [76]. In a prospective follow-up study, significantly increased *S. aureus* intestinal prevalence at 6 months of age was found in a group of allergic children compared with the non-allergic group. An interesting correlation between intestinal colonization with *S. aureus* at 2 and 4 weeks of age and the development of food allergy was observed by Lundell et al. [79]. They also found a correlation between perinatal intestinal *S. aureus* colonization and expression levels of the soluble immune modulator CD14, but not the levels of CD83, and concluded that colonization with *S. aureus* might modulate the development of the neonatal immune system. This indicates an interrelatedness between factors involved in the host's immune response, early colonization, and development of allergic disease. Adlerberth et al. [80] studied the hypothesis that infantile intestinal colonization patterns may influence sensitization to food allergens and atopic eczema. They analyzed a birth cohort of more than 300 infants from three European countries with regard to relations between intestinal colonization patterns during the first year and the development of atopic eczema and sensitization at 18 months of age. They reported a nearly significant association ($p=0.06$) between early intestinal colonization with *S. aureus* and increased risk of atopic eczema. Kalliomaki et al. [81] recently observed that higher numbers of fecal *S. aureus* carriage at 6 and 12 months of age were associated with obesity in children.

Bisgaard et al. [82], who studied a Danish birth cohort of 411 infants, did not observe any correlation between neonatal airway colonization with *S. aureus* at 1 month of age and the development of childhood asthma. In contrast, for colonization with the classical otitis media bacteria *S. pneumoniae*, *M. catarrhalis* and *H. influenzae* a significant association with increased persistent or acute wheezing and hospitalization for wheezing was observed. This seems to indicate a lack of influence of *S. aureus* colonization. However, since these studies were performed in a high-risk cohort, an alternative plausible interpretation of these data is that *S. aureus* to a greater extent than the other bacteria mentioned may modulate the neonatal immune system, and that the lower rates of wheezing associated with *S. aureus* colonization may actually reflect relative protection. This may be in line with observed differences in immune induction between gram-positive and gram-negative bacteria and the risk of childhood asthma [83]. The potential involvement of *S. aureus* enterotoxins (SE) in allergic diseases in early childhood by following 510 children from birth to 5 years of age was studied by Semic-Jusufagic et al. [84]. SE-mix-specific IgE, (SE-A, SE-C, and TSST-1) were measured to determine SE sensitization and correlated with atopic disease. Atopic children were nearly 4 times more SE-mix-sensitive than non-atopic children. Children with eczema were significantly more frequently SE-mix sensi-

tized than children without and the SE-mix sensitization rate increased significantly with increasing eczema severity. SE-mix sensitization was also significantly associated with current wheezing. Furthermore, SE-mix-sensitized children with wheezing had significantly higher airway reactivity than wheezing children who were not sensitized to SE-mix, which suggests that enterotoxins from *S. aureus* might be potential modifiers of childhood wheezing and eczema.

Intestinal MRSA in infants and young children at risk

Paired analysis of nasal and intestinal colonization during separate outbreaks of MRSA in children and newborns has been reported in a limited number of studies. After identification of a single case of MRSA infection, 128 children from a child care center were assessed for MSSA/MRSA carriage by perianal, nasal, and throat swabs [74]. This analysis identified an additional MRSA carrier only from the perianal swab, whereas the two other sites were negative. Singh et al. [85] found that during an MRSA outbreak at the neonatal intensive care unit of two hospitals, out of 373 infants analyzed, 24 were positive for MRSA. Of these, 7 (29%) had positive rectal cultures, of whom 1 (0.4%) was negative for nasal carriage. In 17 infants (71%) only the nasal culture was positive. Investigations during two MRSA outbreaks (one with a HA-MRSA and one with a CA-MRSA strain) were performed using paired nasal and intestinal screening [86]. Altogether, 1,792 newborns were screened and 50 were MRSA-positive. In the first hospital, out of 25 infants positively screened, 17 (71%) had nasal carriage and 5 (21%) had carriage in the rectum, all of whom also had nasal carriage. In the second hospital, 18 out of 25 positively screened (72%) had nasal carriage and 15 out of 25 infants (60%) had rectal carriage, of whom 3 (12%) did not have nasal carriage. The use of surveillance cultures of throat and rectal swabs in a pediatric intensive care unit is important [62]. Among 1,241 patients analyzed there were 29 MRSA carriers, of whom 14 (48%) had rectal carriage. Gustafsson et al. [87] described a study on MRSA carriage in 23 children adopted by Swedish families. Multiple swabs were

taken from the perineum, nose, and other sites. MRSA was detected in 13 of the children. The perineum culture was positive at least once in 9 children (69%) and the nose was positive in 9 children (69%). Interestingly, 4 perineal MRSA carriers (31%) did not have nasal carriage.

Altogether, from these studies it can be deduced that of all the newborns or young children in whom MRSA was detected, 74% had nasal or throat carriage and 44% had intestinal carriage (Table 7). Moreover, 10% had intestinal carriage without nasal or throat carriage, which indicates that, in addition to nasal or throat carriage, the perineum or intestine also constitutes a colonization site with potential infection risk in newborns and young children.

Concluding remarks

Intestinal carriage of *S. aureus* occurs in a significant fraction of both healthy and diseased human individuals. For healthy adults and hospitalized adults at risk the incidence figure is 20%. For MRSA the average fraction of intestinal carriers amongst adult patients at risk is 9%. In young children the colonization of the intestines with *S. aureus* occurs at a very high frequency within the first 6 months of life, after which the frequency drops. For newborns and young children intestinal colonization with MRSA was detected in 1–2% of patients screened. This shows that the clinical impact of this phenomenon may be significant, which was corroborated in a number of studies that associated carriage and intestinal infection. Both MSSA and MRSA seem to successfully colonize the human intestines. This further emphasizes that care has to be taken: not only MSSA infections but also MRSA infection can result from intestinal carriage. Since MSSA and MRSA nasal carriage has been implicated to be a highly important risk factor for infections, the same will probably apply to intestinal carriage. Therefore, prospective controlled studies of intestinal carriage as a predictor of clinical infection need to be performed to determine the significance of this site of colonization. Prolonged intestinal carriage, also among personnel, can be an important factor

Table 7 Intestinal MRSA carriage in young children at risk

Reference	Total patients screened	Total positive carriers	Rectal MRSA	Nasal MRSA	Rectal without nasal
[74]	128	1	1	0	1
[85]	373	24	7	17	1
[86]	1,792	50	20	35	3
[62]	1,241	29	14	26 ^a	3 ^a
[87]	23	13	9	9	4
Totals	3,557	117	51	87	12
Percentage of total positive carriers			44%	74%	10%

^a Throat swab analyzed instead of nasal swab

in the persistence of MRSA outbreaks in hospitals and can be a source of environmental contamination. The development of rapid molecular detection methods for MRSA carriage facilitates prophylaxis [88]. Nasally applied mupirocin very effectively eliminates nasal carriage and results in significant reductions in *S. aureus* infection rates. Mupirocin seems less effective for the elimination of intestinal carriage or other extra-nasal sites. Therefore, application of combined strategies, involving mupirocin in combination with oral antibiotics or other selective intestinal decontamination regimens, such as novel therapies based on polyclonal antibodies [89], may be more effective in cases of proven nasal and intestinal carriage. Because of the significant frequencies of intestinal carriage reported in a wide variety of patient groups and in healthy people it is recommended, in addition to nasal screening, to also include intestinal or perianal screening during surveillance.

Little is known on the age-related kinetics of intestinal carriage, but *S. aureus* seems to colonize the intestine of healthy newborns from very early in life onwards, and can be involved in the development of neonatal infectious disease. Later in life, intestinal carriage frequencies seem to drop. In concordance with nasal carriage, non-, intermittent and persistent intestinal carriage need to be defined in more detail. It is, therefore, of importance to determine whether specific variants of *S. aureus* are more proficient colonizers than others and which adhesion and virulence factors play a role in the transition from intestinal colonization to infection. Even more importantly, little is known on the environmental, bacteriological and physiological determinants of intestinal carriage. Preliminary data identify intestinal carriage as an infection risk factor and results from cohort studies suggest that there is interrelatedness among early intestinal colonization of *S. aureus*, the host's immune system, and the development of disease later in life. Additional research, however, will be needed to fully appreciate the importance of intestinal *S. aureus* colonization in association with infection.

Open Access This article is distributed under the terms of the Creative Commons Attribution Noncommercial License which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

References

1. Kluytmans J, van Belkum A, Verbrugh H (1997) Nasal carriage of *Staphylococcus aureus*: epidemiology, underlying mechanisms, and associated risks. *Clin Microbiol Rev* 10:505–520
2. Weinstein HJ (1959) The relation between the nasal-staphylococcal-carrier state and the incidence of postoperative complications. *N Engl J Med* 260:1303–1308
3. Wertheim HF, Melles DC, Vos MC, van Leeuwen W, van Belkum A et al (2005) The role of nasal carriage in *Staphylococcus aureus* infections. *Lancet Infect Dis* 5:751–762
4. Williams RE (1963) Healthy carriage of *Staphylococcus aureus*: its prevalence and importance. *Bacteriol Rev* 27:56–71
5. Larson HE, Price AB, Honour P, Borriello SP (1978) Clostridium difficile and the aetiology of pseudomembranous colitis. *Lancet* 311:1063–1066
6. Metan G, Zarakolu P, Unal S (2005) Rapid detection of antibacterial resistance in emerging Gram-positive cocci. *J Hosp Infect* 61:93–99
7. Crossley K, Solliday J (1980) Comparison of rectal swabs and stool cultures for the detection of gastrointestinal carriage of *Staphylococcus aureus*. *J Clin Microbiol* 11:433–434
8. Lee DC, Barlas D, Ryan JG, Ward MF, Sama AE et al (2002) Methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant enterococci: prevalence and predictors of colonization in patients presenting to the emergency department from nursing homes. *J Am Geriatr Soc* 50:1463–1465
9. Meurman O, Routamaa M, Peltonen R (2005) Screening for methicillin-resistant *Staphylococcus aureus*: which anatomical sites to culture. *J Hosp Infect* 61:351–353
10. Mody L, Kauffman CA, Donabedian S, Zervos M, Bradley SF (2008) Epidemiology of *Staphylococcus aureus* colonization in nursing home residents. *Clin Infect Dis* 46:1368–1373
11. Drews SJ, Willey BM, Kreiswirth N, Wang M, Ianes T et al (2006) Verification of the IDI-MRSA assay for detecting methicillin-resistant *Staphylococcus aureus* in diverse specimen types in a core clinical laboratory setting. *J Clin Microbiol* 44:3794–3796
12. Buehlmann M, Frei R, Fenner L, Dangel M, Fluckiger U et al (2008) Highly effective regimen for decolonization of methicillin-resistant *Staphylococcus aureus* carriers. *Infect Control Hosp Epidemiol* 29:510–516
13. Rimland D, Roberson B (1986) Gastrointestinal carriage of methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 24:137–138
14. Matthias JQ, Shooter RA, Williams RE (1957) *Staphylococcus aureus* in the faeces of hospital patients. *Lancet* 272:1172–1173
15. Boe J, Solberg CO, Vogelsang TM, Wormnes A (1964) Perineal carriers of staphylococci. *BMJ* 2:280–281
16. Ridley M (1959) Perineal carriage of *Staphylococcus aureus*. *BMJ* 1:270–273
17. Dancer SJ, Noble WC (1991) Nasal, axillary, and perineal carriage of *Staphylococcus aureus* among women: identification of strains producing epidermolytic toxin. *J Clin Pathol* 44:681–684
18. Lee YL, Cesario T, Gupta G, Flionis L, Tran C et al (1997) Surveillance of colonization and infection with *Staphylococcus aureus* susceptible or resistant to methicillin in a community skilled-nursing facility. *Am J Infect Control* 25:312–321
19. Roghmann MC, Gorman PH, Wallin MT, Kreisel K, Shurland S et al (2007) *Staphylococcus aureus* colonization in community-dwelling people with spinal cord dysfunction. *Arch Phys Med Rehabil* 88:979–983
20. Borer A, Gilad J, Yagupsky P, Peled N, Porat N et al (2002) Community-acquired methicillin-resistant *Staphylococcus aureus* in institutionalized adults with developmental disabilities. *Emerg Infect Dis* 8:966–970
21. Van Hal SJ, Stark D, Lockwood B, Marriott D, Harkness J (2007) Methicillin-resistant *Staphylococcus aureus* (MRSA) detection: comparison of two molecular methods (IDI-MRSA PCR assay and GenoType MRSA Direct PCR assay) with three selective MRSA agars (MRSA ID, MRSASelect, and CHROMagar MRSA) for use with infection-control swabs. *J Clin Microbiol* 45:2486–2490
22. Aeilts GD, Sapico FL, Canawati HN, Malik GM, Montgomerie JZ (1982) Methicillin-resistant *Staphylococcus aureus* colonization and infection in a rehabilitation facility. *J Clin Microbiol* 16:218–223

23. Bishop EJ, Grabsch EA, Ballard SA, Mayall B, Xie S et al (2006) Concurrent analysis of nose and groin swab specimens by the IDI-MRSA PCR assay is comparable to analysis by individual-specimen PCR and routine culture assays for detection of colonization by methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 44:2904–2908
24. Cavalcanti SM, Franca ER, Cabral C, Vilela MA, Montenegro F et al (2005) Prevalence of *Staphylococcus aureus* introduced into intensive care units of a University Hospital. *Braz J Infect Dis* 9:56–63
25. Coello R, Jimenez J, Garcia M, Arroyo P, Minguez D et al (1994) Prospective study of infection, colonization and carriage of methicillin-resistant *Staphylococcus aureus* in an outbreak affecting 990 patients. *Eur J Clin Microbiol Infect Dis* 13:74–81
26. Darouiche R, Wright C, Hamill R, Koza M, Lewis D et al (1991) Eradication of colonization by methicillin-resistant *Staphylococcus aureus* by using oral minocycline-rifampin and topical mupirocin. *Antimicrob Agents Chemother* 35:1612–1615
27. Eveillard M, de Lassence A, Lancien E, Barnaud G, Ricard JD et al (2006) Evaluation of a strategy of screening multiple anatomical sites for methicillin-resistant *Staphylococcus aureus* at admission to a teaching hospital. *Infect Control Hosp Epidemiol* 27:181–184
28. Gnanalingham KK, Elsaghier A, Kibbler C, Shieff C (2003) The impact of methicillin-resistant *Staphylococcus aureus* in a neurosurgical unit: a growing problem. *J Neurosurg* 98:8–13
29. Kampf G, Kramer A (2004) Eradication of methicillin-resistant *Staphylococcus aureus* with an antiseptic soap and nasal mupirocin among colonized patients—an open uncontrolled clinical trial. *Ann Clin Microbiol Antimicrob* 3:9
30. Klotz M, Zimmermann S, Opper S, Heeg K, Mutters R (2005) Possible risk for re-colonization with methicillin-resistant *Staphylococcus aureus* (MRSA) by faecal transmission. *Int J Hyg Environ Health* 208:401–405
31. Lim MS, Marshall CL, Spelman D (2006) Carriage of multiple subtypes of methicillin-resistant *Staphylococcus aureus* by intensive care unit patients. *Infect Control Hosp Epidemiol* 27:1063–1067
32. Maeder K, Ginunas VJ, Montgomerie JZ, Canawati HN (1993) Methicillin-resistant *Staphylococcus aureus* (MRSA) colonization in patients with spinal cord injury. *Paraplegia* 31:639–644
33. Manian FA, Senkel D, Zack J, Meyer L (2002) Routine screening for methicillin-resistant *Staphylococcus aureus* among patients newly admitted to an acute rehabilitation unit. *Infect Control Hosp Epidemiol* 23:516–519
34. Mylotte JM, Kahler L, Graham R, Young L, Goodnough S (2000) Prospective surveillance for antibiotic-resistant organisms in patients with spinal cord injury admitted to an acute rehabilitation unit. *Am J Infect Control* 28:291–297
35. Papia G, Louie M, Tralla A, Johnson C, Collins V et al (1999) Screening high-risk patients for methicillin-resistant *Staphylococcus aureus* on admission to the hospital: is it cost effective. *Infect Control Hosp Epidemiol* 20:473–477
36. Reyes RC, Stoakes L, Milburn S, Lennox G, Daniel J et al (2008) Evaluation of a new chromogenic medium for the detection of methicillin-resistant *Staphylococcus aureus* carriage on nasal and perianal specimens. *Diagn Microbiol Infect Dis* 60:225–227
37. Roberts S, Young H, Faulkner S, Bilkey M, Eyres S et al (2002) Value of broth cultures in detecting methicillin-resistant *Staphylococcus aureus*. *NZ Med J* 115:U191
38. Trick WE, Weinstein RA, DeMarais PL, Kuehnert MJ, Tomaska W et al (2001) Colonization of skilled-care facility residents with antimicrobial-resistant pathogens. *J Am Geriatr Soc* 49:270–276
39. Zhang SX, Drews SJ, Tomassi J, Katz KC (2007) Comparison of two versions of the IDI-MRSA assay using charcoal swabs for prospective nasal and nonnasal surveillance samples. *J Clin Microbiol* 45:2278–2280
40. Simor AE, Ofner-Agostini M, Paton S, McGeer A, Loeb M et al (2005) Clinical and epidemiologic features of methicillin-resistant *Staphylococcus aureus* in elderly hospitalized patients. *Infect Control Hosp Epidemiol* 26:838–841
41. Hare R, Ridley M (1958) Further studies on the transmission of *Staphylococcus aureus*. *BMJ* 1:69–73
42. Solberg CO (2000) Spread of *Staphylococcus aureus* in hospitals: causes and prevention. *Scand J Infect Dis* 32:587–595
43. Masaki H, Asoh N, Watanabe H, Tao M, Watanabe K et al (2003) Possible relationship between *Staphylococcus aureus* colonizing the respiratory tract and rectum and *S. aureus* isolated in a geriatric hospital environment. *Intern Med* 42:281–282
44. Bhalla A, Aron DC, Donskey CJ (2007) *Staphylococcus aureus* intestinal colonization is associated with increased frequency of *S. aureus* on skin of hospitalized patients. *BMC Infect Dis* 7:105
45. Boyce JM, Havill NL, Otter JA, Adams NM (2007) Widespread environmental contamination associated with patients with diarrhea and methicillin-resistant *Staphylococcus aureus* colonization of the gastrointestinal tract. *Infect Control Hosp Epidemiol* 28:1142–1147
46. Tulloch LG, Alder VG, Gillespie WA (1960) Treatment of chronic furunculosis. *BMJ* 2:354–356
47. Solberg CO (1965) A study of carriers of *Staphylococcus aureus* with special regard to quantitative bacterial estimations. *Acta Med Scand Suppl* 436:1–96
48. Squier C, Rihs JD, Risa KJ, Sagnimeni A, Wagener MM et al (2002) *Staphylococcus aureus* rectal carriage and its association with infections in patients in a surgical intensive care unit and a liver transplant unit. *Infect Control Hosp Epidemiol* 23:495–501
49. Ray AJ, Pultz NJ, Bhalla A, Aron DC, Donskey CJ (2003) Coexistence of vancomycin-resistant enterococci and *Staphylococcus aureus* in the intestinal tracts of hospitalized patients. *Clin Infect Dis* 37:875–881
50. Van Rijen MM, Bonten M, Wenzel RP, Kluytmans JA (2008) Intranasal mupirocin for reduction of *Staphylococcus aureus* infections in surgical patients with nasal carriage: a systematic review. *J Antimicrob Chemother* 61:254–261
51. Dupeyron C, Campillo B, Bordes M, Faubert E, Richardet JP et al (2002) A clinical trial of mupirocin in the eradication of methicillin-resistant *Staphylococcus aureus* nasal carriage in a digestive disease unit. *J Hosp Infect* 52:281–287
52. Dupeyron C, Campillo B, Richardet JP, Soussy CJ (2006) Long-term efficacy of mupirocin in the prevention of infections with methicillin-resistant *Staphylococcus aureus* in a gastroenterology unit. *J Hosp Infect* 63:385–392
53. Perl TM, Cullen JJ, Wenzel RP, Zimmerman MB, Pfaller MA et al (2002) Intranasal mupirocin to prevent postoperative *Staphylococcus aureus* infections. *N Engl J Med* 346:1871–1877
54. Harbarth S, Dharan S, Liassine N, Herrault P, Auckenthaler R et al (1999) Randomized, placebo-controlled, double-blind trial to evaluate the efficacy of mupirocin for eradicating carriage of methicillin-resistant *Staphylococcus aureus*. *Antimicrob Agents Chemother* 43:1412–1416
55. Wertheim HF, Verveer J, Boelens HA, van Belkum A, Verbrugh HA et al (2005) Effect of mupirocin treatment on nasal, pharyngeal, and perineal carriage of *Staphylococcus aureus* in healthy adults. *Antimicrob Agents Chemother* 49:1465–1467
56. Falagas ME, Bliziotis IA, Fragoulis KN (2007) Oral rifampin for eradication of *Staphylococcus aureus* carriage from healthy and sick populations: a systematic review of the evidence from comparative trials. *Am J Infect Control* 35:106–114
57. Falagas ME, Fragoulis KN, Bliziotis IA (2006) Oral rifampin for prevention of *Staphylococcus aureus* carriage-related infections in patients with renal failure—a meta-analysis of randomized controlled trials. *Nephrol Dial Transplant* 21:2536–2542
58. Cerda E, Abella A, de la Cal MA, Lorente JA, Garcia-Hierro P et al (2007) Enteral vancomycin controls methicillin-resistant

- Staphylococcus aureus* endemicity in an intensive care burn unit: a 9-year prospective study. *Ann Surg* 245:397–407
59. De la Cal MA, Cerda E, van Saene HK, Garcia-Hierro P, Negro E et al (2004) Effectiveness and safety of enteral vancomycin to control endemicity of methicillin-resistant *Staphylococcus aureus* in a medical/surgical intensive care unit. *J Hosp Infect* 56:175–183
 60. Silvestri L, van Saene HK, Milanese M, Fontana F, Gregori D et al (2004) Prevention of MRSA pneumonia by oral vancomycin decontamination: a randomised trial. *Eur Respir J* 23:921–926
 61. Solis A, Brown D, Hughes J, Van Saene HK, Heaf DP (2003) Methicillin-resistant *Staphylococcus aureus* in children with cystic fibrosis: an eradication protocol. *Pediatr Pulmonol* 36:189–195
 62. Thorburn K, Taylor N, Saladi SM, van Saene HK (2006) Use of surveillance cultures and enteral vancomycin to control methicillin-resistant *Staphylococcus aureus* in a paediatric intensive care unit. *Clin Microbiol Infect* 12:35–42
 63. Warren DK, Liao RS, Merz LR, Eveland M, Dunne WM Jr (2004) Detection of methicillin-resistant *Staphylococcus aureus* directly from nasal swab specimens by a real-time PCR assay. *J Clin Microbiol* 42:5578–5581
 64. Furuno JP, Perencevich EN, Johnson JA, Wright MO, McGregor JC et al (2005) Methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant Enterococci co-colonization. *Emerg Infect Dis* 11:1539–1544
 65. Chen KT, Huard RC, Della-Latta P, Saiman L (2006) Prevalence of methicillin-sensitive and methicillin-resistant *Staphylococcus aureus* in pregnant women. *Obstet Gynecol* 108:482–487
 66. Chen KT, Campbell H, Borrell LN, Huard RC, Saiman L et al (2007) Predictors and outcomes for pregnant women with vaginal-rectal carriage of community-associated methicillin-resistant *Staphylococcus aureus*. *Am J Perinatol* 24:235–240
 67. Froberg MK, Palavecino E, Dykoski R, Gerding DN, Peterson LR et al (2004) *Staphylococcus aureus* and *Clostridium difficile* cause distinct pseudomembranous intestinal diseases. *Clin Infect Dis* 39:747–750
 68. Gravet A, Rondeau M, Harf-Monteil C, Grunenberger F, Monteil H et al (1999) Predominant *Staphylococcus aureus* isolated from antibiotic-associated diarrhea is clinically relevant and produces enterotoxin A and the bicomponent toxin LukE-lukD. *J Clin Microbiol* 37:4012–4019
 69. Ackermann G, Thomalla S, Ackermann F, Schaumann R, Rodloff AC et al (2005) Prevalence and characteristics of bacteria and host factors in an outbreak situation of antibiotic-associated diarrhoea. *J Med Microbiol* 54:149–153
 70. Boyce JM, Havill NL, Maria B (2005) Frequency and possible infection control implications of gastrointestinal colonization with methicillin-resistant *Staphylococcus aureus*. *J Clin Microbiol* 43:5992–5995
 71. Flemming K, Ackermann G (2007) Prevalence of enterotoxin producing *Staphylococcus aureus* in stools of patients with nosocomial diarrhea. *Infection* 35:356–358
 72. Asha NJ, Tompkins D, Wilcox MH (2006) Comparative analysis of prevalence, risk factors, and molecular epidemiology of antibiotic-associated diarrhea due to *Clostridium difficile*, *Clostridium perfringens*, and *Staphylococcus aureus*. *J Clin Microbiol* 44:2785–2791
 73. Suggs AH, Maranan MC, Boyle-Vavra S, Daum RS (1999) Methicillin-resistant and borderline methicillin-resistant asymptomatic *Staphylococcus aureus* colonization in children without identifiable risk factors. *Pediatr Infect Dis J* 18:410–414
 74. Shahin R, Johnson IL, Jamieson F, McGeer A, Tolkin J et al (1999) Methicillin-resistant *Staphylococcus aureus* carriage in a child care center following a case of disease. Toronto Child Care Center Study Group. *Arch Pediatr Adolesc Med* 153:864–868
 75. Efuntoye MO, Adetosoye AI (2003) Enterotoxigenicity and drug sensitivity of staphylococci from children aged five years and below with sporadic diarrhoea. *East Afr Med J* 80:656–659
 76. Bjorksten B, Naaber P, Sepp E, Mikelsaar M (1999) The intestinal microflora in allergic Estonian and Swedish 2-year-old children. *Clin Exp Allergy* 29:342–346
 77. Lindberg E, Adlerberth I, Hesselmar B, Saalman R, Strannegard IL et al (2004) High rate of transfer of *Staphylococcus aureus* from parental skin to infant gut flora. *J Clin Microbiol* 42:530–534
 78. Lindberg E, Nowrouzian F, Adlerberth I, Wold AE (2000) Long-time persistence of superantigen-producing *Staphylococcus aureus* strains in the intestinal microflora of healthy infants. *Pediatr Res* 48:741–747
 79. Lundell AC, Adlerberth I, Lindberg E, Karlsson H, Ekberg S et al (2007) Increased levels of circulating soluble CD14 but not CD83 in infants are associated with early intestinal colonization with *Staphylococcus aureus*. *Clin Exp Allergy* 37:62–71
 80. Adlerberth I, Strachan DP, Matricardi PM, Ahme S, Orfei L et al (2007) Gut microbiota and development of atopic eczema in 3 European birth cohorts. *J Allergy Clin Immunol* 120:343–350
 81. Kalliomaki M, Collado MC, Salminen S, Isolauri E (2008) Early differences in fecal microbiota composition in children may predict overweight. *Am J Clin Nutr* 87:534–538
 82. Bisgaard H, Hermansen MN, Buchvald F, Loland L, Halkjaer LB et al (2007) Childhood asthma after bacterial colonization of the airway in neonates. *N Engl J Med* 357:1487–1495
 83. Michelow IC, Fracchia MS, Kinane TB (2008) Asthma and neonatal airway colonization. *N Engl J Med* 358:423–425; author reply 424–425
 84. Semic-Jusufagic A, Bachert C, Gevaert P, Holtappels G, Lowe L et al (2007) *Staphylococcus aureus* sensitization and allergic disease in early childhood: population-based birth cohort study. *J Allergy Clin Immunol* 119:930–936
 85. Singh K, Gavin PJ, Vescio T, Thomson RB Jr, Deddish RB et al (2003) Microbiologic surveillance using nasal cultures alone is sufficient for detection of methicillin-resistant *Staphylococcus aureus* isolates in neonates. *J Clin Microbiol* 41:2755–2757
 86. Rosenthal A, White D, Churilla S, Brodie S, Katz KC (2006) Optimal surveillance culture sites for detection of methicillin-resistant *Staphylococcus aureus* in newborns. *J Clin Microbiol* 44:4234–4236
 87. Gustafsson EB, Ringberg H, Johansson PJ (2007) MRSA in children from foreign countries adopted to Swedish families. *Acta Paediatr* 96:105–108
 88. Carroll KC (2008) Rapid diagnostics for methicillin-resistant *Staphylococcus aureus*: current status. *Mol Diagn Ther* 12:15–24
 89. Numan SC, Veldkamp P, Kuijper EJ, van den Berg RJ, van Dissel JT (2007) *Clostridium difficile*-associated diarrhoea: bovine anti-*Clostridium difficile* whey protein to help aid the prevention of relapses. *Gut* 56:888–889
 90. Singh N, Squier C, Wannstedt C, Keyes L, Wagener MM et al (2006) Impact of an aggressive infection control strategy on endemic *Staphylococcus aureus* infection in liver transplant recipients. *Infect Control Hosp Epidemiol* 27:122–126
 91. Grmek-Kosnik I, Ihan A, Dermota U, Rems M, Kosnik M et al (2005) Evaluation of separate vs pooled swab cultures, different media, broth enrichment and anatomical sites of screening for the detection of methicillin-resistant *Staphylococcus aureus* from clinical specimens. *J Hosp Infect* 61:155–161
 92. Buttiaux R, Pierret J (1949) Les staphylocoques pathogènes dans les selles des nourissons normaux. *Ann Inst Pasteur (Paris)* 76:480–482
 93. Brodie J, Sommerville T, Wilson SGF (1956) Coagulase positive staphylococci. A seral survey during the first six months of nursing training. *BMJ* 1:667–669
 94. Grun L (1958) Studies on intestinal staphylococci. *Arch Hyg Bakteriol* 142:3–7

95. Greendyke RM, Constantine HP, Magruder GB, Dean DC, Gardner JH et al (1958) Staphylococci on a medical ward, with special reference to fecal carriers. *Am J Clin Pathol* 30:318–322
96. Hofstad T, Wormnes A (1961) The effect of broad spectrum antibiotics on the faecal staphylococcal and monilial flora in man. *Acta Pathol Microbiol Scand* 51:275–279
97. Polakoff S, Richards ID, Parker MT, Lidwell OM (1967) Nasal and skin carriage of *Staphylococcus aureus* by patients undergoing surgical operation. *J Hyg (Lond)* 65:559–566
98. Bjorksten B, Sepp E, Julge K, Voor T, Mikelsaar M (2001) Allergy development and the intestinal microflora during the first year of life. *J Allergy Clin Immunol* 108:516–520