

# Intra-aortic balloon counterpulsation and infarct size in patients with acute anterior myocardial infarction without shock: The CRISP AMI Randomized Trial



*Manesh R. Patel, MD, Richard W. Smalling, MD, PhD,  
Holger Thiele, Prof Dr med, Huiman X. Barnhart, PhD, Yi Zhou, PhD,  
Praveen Chandra, MD, Derek Chew, MD, Marc Cohen, MD,  
John French, MB CHB, PhD, Divaka Perera, MD, E. Magnus Ohman, MD  
European Society of Cardiology – Hotline Presentation August 30<sup>th</sup> 2011*

- Despite improvements in STEMI care
  - The 6 month mortality remains high ~10%<sup>1</sup>
- Intra-aortic balloon counterpulsation
  - ↑Diastolic arterial pressure (coronary perfusion pressure)
  - ↓Simultaneously decrease afterload and left ventricular end diastolic pressure (LVEDP) - both work to decrease oxygen consumption
  - Decreases infarct expansion when placed prior to reperfusion in animal studies <sup>2,3</sup>

<sup>1</sup>Heart disease and stroke statistics--2009 update. Circulation 2009;119:e21-181.

<sup>2</sup>LeDoux JF et. al.. Catheterization & Cardiovascular Interventions 2008;72:513-21.

<sup>3</sup>Azevedo CF et. al. European Heart Journal 2005;26:1235-41.

# Primary Objective



To determine whether routine initiation of intra-aortic balloon counterpulsation (IABC) before mechanical reperfusion compared to standard of care (SOC) primary PCI decreases infarct size in patients with anterior ST-segment elevation myocardial infarction (STEMI) without cardiogenic shock

## Anterior STEMI without Shock

### Inclusion Criteria

- Anterior STEMI  
2 mm in 2 contiguous leads or  
at least 4 mm in the anterior  
leads
- Planned Primary PCI within 6 hrs
- Adult able to consent

Intra-aortic Balloon  
Counterpulsation prior to PCI

Randomize  
Open Label  
(n ~ 300)

Standard of Care Primary PCI

At least 12 hours of IABC post PCI

Routine Post PCI care

Cardiac MRI performed day 3-5 post PCI

Primary Endpoint: Infarct Size on CMR

1. All Patients with CMR data
2. Patients with Prox LAD occlusion TIMI 0/1 flow

Clinical Events – 6 months

# Exclusion Criteria


- Known Contraindication to MRI
- Prior Thrombolytic Therapy for STEMI
- Cardiogenic Shock
- Prior MI, CABG, or ESRD
- Contraindications to IABC
  - Known Severe AI, AAA, or severe peripheral artery disease
  - >400 lbs or < 4 feet

- **Sample Size**
  - Estimated Infarct size
    - All patients (25.3 -26.6% LV)<sup>1,2</sup> and (19.9 - 28.8% LV)<sup>1,2</sup> prox. LAD TIMI 0/1
  - 25% reduction (270 patients) 10% CMR data missing
  - >80% power, Type 1 error 0.025 (2-sided)
  - ~ **300 patients**
- **Primary Endpoint Evaluation: Infarct Size on CMR**
  - Modified ITT – all patients with CMR data
  - All CMR patients with proximal LAD occlusion TIMI 0/1
- **Primary Safety Evaluation: Major vascular complications and Major bleeding**
- **Clinical Outcomes: 6-month rate all cause mortality, MACE**

<sup>1</sup> Patel et al. Jacc: Cardiovascular Imaging 2010;3:52-60

<sup>2</sup> Thiele et al. Circulation 2008 Jul 1;118(1):49-57 Epub 2008 Jun 16

# CMR Protocol



Survey

<b>Function</b> 4-chamber 2-chamber	<b>Edema</b> 3 short axes	<b>Early enhancement</b> Short axes Apex-Base	<b>Function</b> Short axes Apex-Base	<b>Delayed enhancement</b> Short axes Apex-Base	<b>Delayed enhancement</b> 4-chamber 2-chamber
---	------------------------------	---	--	---	--

SSFP sequence  
(TR/TE/flip =  
3.2ms/1.2ms/60°)

T2 STIR sequence  
(TR/TE/flip =  
2 heart  
beats/80ms/90°)  
slice thickness:  
8-10 mm

Inversion recovery  
gradient echo  
sequence  
(TR/TE/flip  
2.8ms/1.1ms/15°)  
slice thickness:  
8-10 mm, no gap

SSFP sequence  
(TR/TE/flip =  
3.2ms/1.2ms/60°)  
slice thickness:  
8-10 mm, no gap


Inversion recovery  
gradient echo sequence  
(TR/TE/flip  
2.8ms/1.1ms/15°)  
slice thickness:  
8-10 mm, no gap



# Enrollment

9 countries, 30 sites, 337 patients

**CRISP**  
**ami**  
counterpulsation  
reduces infarct  
size pre pci in  
acute myocardial  
infarction



PI Name	Site Name	N
Sreenivas	CARE Hospital	58
Singh	Baroda Heart Institute and Research Centre	54
Blaxill	Leeds General Infirmary	32
Pijls	Catherina Hospital	28
Mills	Duke Univ. Med Center	23



# Study Conduct

	Randomized* N=337	
	IABC N=161	SOC N=176
Received intervention	153 (95.03%)	161 (91.48%)
Withdrew	4	2
Lost to follow-up	<b>Crossing over to IABC 15</b>	
<b>MRI not performed</b>	Sustained hypotension/Cardiogenic shock 12	
Died	To prevent event post <input type="checkbox"/> vessel dissection 1	
Unstable	Failed PCI of IR vessel 1	
	Continued chest pain 1	
Metallic contraindication	3	1
Unable to tolerate	11	18
Other	6	0
<b>MRI performed, not evaluable</b>	<b>5</b>	<b>7</b>

# Baseline Demographics

	<b>All (N=337)</b>	<b>IABC (N=161)</b>	<b>SOC (N=176)</b>
Age, median (25th, 75th), yrs	56.6 (48.4, 65.6)	56.1 (48.3, 64.3)	57.7 (48.6, 66.4)
Male, %	81.9	82.0	81.8
Race, %			
White	47.8	50.3	45.5
Asian	45.1	46.6	43.8
Black or African American	4.7	1.9	7.4
Other	2.1	1.2	2.8
Medical history, %			
Hypertension on drug tx.	29.4	24.2	34.1
Current nicotine use	31.8	33.1	30.7
Dyslipidemia on drug tx.	12.5	12.5	12.5
Diabetes mellitus	18.7	16.8	20.5

# Baseline Demographics (cont.)


	<b>All (N=337)</b>	<b>IABC (N=161)</b>	<b>SOC (N=176)</b>
SBP, median (25th, 75th), mm Hg	131.0 (118.0, 150.0)	130.0 (113.0, 150.0)	135.0 (120.0, 151.0)
DBP, median (25th, 75th), mm Hg	80.0 (70.0, 92.0)	80.0 (70.0, 92.0)	80.0 (71.5, 92.0)
HR, median (25th, 75th), bpm	81.0 (71.0, 94.0)	81.0 (71.0, 93.0)	80.0 (70.0, 94.0)
ST ↑ in anterior leads, no. (%)			
0–<2 mm	0 (0.0)	0 (0.0)	0 (0.0)
2–<4 mm	1 (0.3)	0 (0.0)	1 (0.6)
4–<6 mm	135 (40.1)	61 (37.9)	74 (42.0)
≥6 mm	201 (59.6)	100 (62.1)	101 (57.4)

# PCI Procedure

	All N=337	IABC N=161	SOC N=176
PCI			
PCI performed, %	94.3	96.3	92.6
Infarct-related artery			
Left anterior descending, %	97.6	99.4	96.0
Infarct-related artery stenosis location			
Proximal, %	62.9	64.8	61.2
Infarct-related artery TIMI flow pre-intervention			
Grade 0, %	65.3	66.0	64.7
Grade 1, %	10.3	11.3	9.4
Infarct-related artery final TIMI flow post-intervention			
Grade 3, %	94.2	92.9	95.3

# Time to Treatment

- First medical contact to first device\*
- Symptom onset to 1st device



# Primary outcome

	All (N=337)	IABC (N=161)	SOC (N=176)	P Value
<b>Primary endpoint</b>				
<b>Infarct size (% LV), modified ITT all patients with CMR data</b>				<b>0.060</b>
N	275	133	142	
Mean	39.8	42.1	37.5	
Median	38.8	42.8	36.2	
<b>Infarct size (% LV), modified ITT patients prox. LAD and TIMI flow 0/1</b>				<b>0.110</b>
N	192	93	99	
Mean	44.4	46.7	42.3	
Median	42.1	45.1	38.6	

Co-primary endpoint: 2-sided p=0.025


# 30-day Clinical Events

	<b>IABC (N=161)</b>	<b>SOC (N=176)</b>	<b>P Value</b>
Death, %	1.9*	4.0*	0.26*
Stroke, %	1.9	0.6	0.35
Major bleed per GUSTO 1 definition or transfusion, %	3.1	1.7	0.49
Vascular complications, (n) %	7(4.3)	2 (1.1)	0.09
Major limb ischemia requiring operative intervention (n)	0	0	
Distal embolization (n)	0	0	
Major dissection (n)	2	0	
Pseudoaneurysm or AV fistula (n)	3	2	
Hematoma >5 cm (n)	3	0	

\*From KM curves and log-rank test.



# All Cause Death – 6 months



	IABC (N=161)	SOC (N=176)	P Value
Death, %	1.9*	5.2*	0.12*
Death/recurrent MI/new or worsening CHF, %	6.3*	10.9*	0.15*
Death/shock/new or worsening CHF, %†	5.0*	12.0*	0.03*

\*From KM curves and log-rank test. †Exploratory analysis.

Among Patients with Acute Anterior STEMI without cardiogenic shock use of Intra-aortic counterpulsation prior to PCI compared to standard of care PCI:

1. Does not reduce infarct size
2. All cause mortality at 6 months was not different
3. Exploratory composite clinical endpoint favored of IABC

- These findings do not support the routine use of IABC prior to PCI in Anterior STEMI patients without cardiogenic shock,
- Clinicians should continue to be vigilant about identifying patients who are at risk for rapid deterioration or hypotension that may benefit from support, as seen with the cross-over in this trial (8.5%)
- Acute STEMI studies are feasible without significant increases in door-to-device times

# Acknowledgements



## **CRISP Steering Committee**

Manesh R. Patel, Holger Thiele, Richard W. Smalling, Praveen Chandra, Marc Cohen, Divaka Perera, Derek Chew, John French, E. Magnus Ohman

## **CRISP AMI investigators**

Sreenivas Kumar A., Singh, Blaxill, Pijls, Mills, Thomas, Henriksen, Smalling, Passey, Bashir, McCann, Weintraub, Cohen, Vranckx, Thiele, Reddy, Schwab, Ling, Garg, Chandra, Sinhal, Casale, Banerjee, Khanna, Hillegass, Varghese, Satler, Strasser, Biederman, Shavelle, Valente, Lefevre, Kaluski, Carozza Jr., Weeks, Bush, Saligrama, Bingi, Talwar, Diebele

## **DSMB**

Eric Bates, David Holmes, Richard Trout

## **Global Coordinating Center**

Duke Clinical Research Institute

- Pam Monds, Project Lead
- Dorothy J Wagstaff, Lead Clinical Data Specialist
- Joey Zhou, Huiman Barnhardt - Statistician
- Karen Ramsey, Lead CRA

## **MRI Core Lab**

The Heart Center Leipzig -University Hospital

- Matthias Gutberlet – Director, Maren Redlich
- Fabian Juhrich,

## **Regional Centers groups**

- Meredith Cooney, Flinders Coordinating Centre, Australia/NZ Lead
- Tanya Fawcett, MAQUET CV, Europe Lead
- Vaibhav S. Pawar, Jubilant Clinsys Ltd., India Lead

# CRISP ami

counterpulsation  
reduces infarct  
size pre pci in  
acute myocardial  
infarction