



Casetta, F., Coltorti, M., Ickert, R. B. , Bonadiman, C., Giacomoni, P. P. and Ntaflou, T. (2018) Intrusion of shoshonitic magmas at shallow crustal depth: T–P path, H₂O estimates, and AFC modeling of the Middle Triassic Predazzo Intrusive Complex (Southern Alps, Italy). *Contributions to Mineralogy and Petrology*, 173, 57. (doi:[10.1007/s00410-018-1483-0](https://doi.org/10.1007/s00410-018-1483-0)).

This is the author's final accepted version.

There may be differences between this version and the published version. You are advised to consult the publisher's version if you wish to cite from it.

<http://eprints.gla.ac.uk/166325/>

Deposited on: 09 August 2018

Enlighten – Research publications by members of the University of Glasgow
<http://eprints.gla.ac.uk>

Contributions to Mineralogy and Petrology

Intrusion of shoshonitic magmas at shallow crustal depth: T-P path, H₂O estimates and AFC modelling of the Middle Triassic Predazzo Intrusive Complex (Southern Alps, Italy)

--Manuscript Draft--

Manuscript Number:	CTMP-D-18-00030R1
Full Title:	Intrusion of shoshonitic magmas at shallow crustal depth: T-P path, H ₂ O estimates and AFC modelling of the Middle Triassic Predazzo Intrusive Complex (Southern Alps, Italy)
Article Type:	Original Paper
Keywords:	Predazzo Intrusive Complex; Sr and Nd isotopes; P-T estimates and magma water content; Crustal assimilation; Solidification time; Shoshonitic intrusion.
Corresponding Author:	Federico Casetta Universita degli Studi di Ferrara Ferrara, Italy ITALY
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Universita degli Studi di Ferrara
Corresponding Author's Secondary Institution:	
First Author:	Federico Casetta
First Author Secondary Information:	
Order of Authors:	Federico Casetta Massimo Coltorti Ryan B. Ickert Costanza Bonadiman Pier Paolo Giacomoni Theodoros Ntaflou
Order of Authors Secondary Information:	
Funding Information:	
Abstract:	<p>The multi-pulse shoshonitic Predazzo Intrusive Complex represents an ideal igneous laboratory for investigating the chemical and physical conditions of magma emplacement in a crustal context, since numerical models can be constrained by field evidence. It constitutes the most intriguing remnant of the Middle Triassic magmatic systems of the Dolomitic Area (Southern Alps), preserved by the Alpine tectonics. Predazzo Intrusive Complex comprises silica saturated (pyroxenites/gabbros to syenites), silica undersaturated (gabbros to syenites) and silica oversaturated (granites and syenogranites) rock suites. In this paper we modelled its emplacement and evolution with a multiple thermo-/oxy-barometric, hygrometric and EC-AFC approach. At odds with what proposed in literature but according to the field evidence, the emplacement of the Predazzo Intrusive Complex occurred at shallow depth (< 6 km). In this context, the different pulses differed slightly in bulk water content, but shared a common thermal regime, with temperatures between 1000-1100°C and ~600°C at low to moderate oxidizing conditions (-0.1 to +0.7 ΔFMQ). The interaction between the intrusion and the shallow crustal rocks was minimal, with Sr and Nd isotopic compositions indicating an average of 5-6% assimilation of crust. A thermo- and oxy-barometric comparison with the nearby Mt. Monzoni enabled also to speculate about the solidification time of the intrusion, which we infer took place over about 700 ka.</p>
Response to Reviewers:	To the Editor of Contributions to Mineralogy and Petrology

[Click here to view linked References](#)

1 **Intrusion of shoshonitic magmas at shallow crustal depth: T-P path, H₂O estimates and**
2 **AFC modelling of the Middle Triassic Predazzo Intrusive Complex (Southern Alps,**
3 **Italy)**

4
5
6
7
8
9 **Federico Casetta***

10 Department of Physics and Earth Sciences, University of Ferrara
11
12 Via Saragat 1, 44121 Ferrara, Italy

13
14
15
16 **Massimo Coltorti**

17 Department of Physics and Earth Sciences, University of Ferrara
18
19 Via Saragat 1, 44121 Ferrara, Italy

20
21
22
23 **Ryan B. Ickert**

24 Scottish Universities Environmental Research Centre, Scottish Enterprise Technology Park,
25
26 Rankine Avenue, East Kilbride, G75 0QF, UK

27
28
29 **Costanza Bonadiman**

30 Department of Physics and Earth Sciences, University of Ferrara
31
32 Via Saragat 1, 44121 Ferrara, Italy

33
34
35 **Pier Paolo Giacomoni**

36 Department of Physics and Earth Sciences, University of Ferrara
37
38 Via Saragat 1, 44121 Ferrara, Italy

39
40
41 **Theodoros Ntaflos**

42 Department of Lithospheric Research, Universität Wien
43
44 Althanstraße 14 (UZA II), 1090 Wien

45
46
47
48
49
50
51
52 * Corresponding author. Phone +39 0532 974721. E-mail: cstfrc@unife.it

30 **Intrusion of shoshonitic magmas at shallow crustal depth: T-P path, H₂O estimates and**
1
231 **AFC modelling of the Middle Triassic Predazzo Intrusive Complex (Southern Alps,**
3
4
32 **Italy)**

9
34 **Abstract**

10
11 The multi-pulse shoshonitic Predazzo Intrusive Complex represents an ideal igneous laboratory
1235
13 for investigating the chemical and physical conditions of magma emplacement in a crustal
1436
15 context, since numerical models can be constrained by field evidence. It constitutes the most
16
1737 intriguing remnant of the Middle Triassic magmatic systems of the Dolomitic Area (Southern
18
1938 Alps), preserved by the Alpine tectonics. Predazzo Intrusive Complex comprises silica
20
21 saturated (pyroxenites/gabbros to syenites), silica undersaturated (gabbros to syenites) and
2239
23 silica oversaturated (granites and syenogranites) rock suites. In this paper we modelled its
2440
25 emplacement and evolution with a multiple thermo-/oxy-barometric, hygrometric and EC-AFC
26
2741 approach. At odds with what proposed in literature but according to the field evidence, the
28
2942 emplacement of the Predazzo Intrusive Complex occurred at shallow depth (< 6 km). In this
30
3143 context, the different pulses differed slightly in bulk water content, but shared a common
32
33 thermal regime, with temperatures between 1000-1100°C and ~600°C at low to moderate
3444
35 oxydizing conditions (-0.1 to +0.7 ΔFMQ). The interaction between the intrusion and the
3645
37 shallow crustal rocks was minimal, with Sr and Nd isotopic compositions indicating an average
38
3946 of 5-6% assimilation of crust. [A thermo- and oxy-barometric](#) comparison with the nearby Mt.
40
4147 Monzoni enabled also to speculate about the solidification time of the intrusion, which we infer
42
43
4448 took place over about 700 ka.
45
46
47
48
49
50
5151

52
53
54
55
5653 **Keywords**

57
5854 Predazzo Intrusive Complex; Sr and Nd isotopes; P-T estimates and magma water content;
59
60
6155 Crustal assimilation; Solidification time; Shoshonitic intrusion.
62
63
64
65

56
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

The Middle Triassic magmatic event in the Southern Alps is expressed in volcano-plutonic sequences outcropping from the Brescian Alps, Alto Vicentino, Dolomites and Carnia areas (Italy) to the Karavanken region (Austria). Most of the igneous products are volcanic and volcanoclastic rocks, with subordinated dyke swarms. Intrusive bodies are instead rare, and mainly located in the Dolomites (Predazzo-Mt. Monzoni-Cima Pape) and Karavanken areas (Gasparotto and Simboli 1991; Gianolla 1992; Bonadiman et al. 1994; Visonà and Zanferrari 2000; Brack et al. 2005; Cassinis et al. 2008; Bellieni et al. 2010; Casetta et al. 2017). These intrusions represent snapshots of magmatic plumbing systems, emplaced during and/or immediately after the eruption of the overlying volcanic products. There are several outstanding geological problems regarding the nature of these complexes, mainly related to their emplacement conditions, and to the relationships between tectonics and magmatism. The Predazzo Intrusive Complex (PIC) represents the ideal “petrologic laboratory” for investigating and reconstructing the features of the Middle Triassic feeding systems, since it offers the possibility to constrain the theoretical and experimental modeling by means of field evidence. As in the case of the nearby Mt. Monzoni, the contact aureole between PIC and the surrounding sedimentary rocks enabled many authors to speculate about the geometry of the intrusive body and its field relationships to the shallow crust (Princivalle et al. 1999; Ferry et al. 2002; Povoden et al. 2002; Gallien et al. 2007). On the basis of petrological and field evidence, Casetta et al. (2017) identified three different magma batches constituting the PIC, interpreted as a multi-pulse body of shoshonitic affinity with variable alkalis and H₂O-content which led to differentiation from mafic to amphibole and biotite-bearing end-members. However, several issues are still unsolved, such as the (possible) interaction between PIC magmas and crust, and their chemical/physical conditions of emplacement. The latter resulting in a current uncertainty about the depth of the intrusion.

82 Few thermobarometric and oxybarometric data are reported in literature for the Middle Triassic
1
23 magmatic systems of the Dolomitic Area. According to Bonadiman et al. (1994), the oxygen
3
4
54 fugacity of the nearby Mt. Monzoni system was around the NNO buffer and the crystallization
6
75 temperature range was between 1044 and 589°C. Despite the clear evidence of a predominant
8
9
86 role of water during crystallization, as testified by the wide presence of hydrous phases in the
10
11
127 intrusive rocks, no estimates of the H₂O contents and *P-T* conditions of these Middle Triassic
13
14
38 plumbing systems have been performed so far. Taking into account the Al^{tot} content of
15
16
179 amphiboles, Menegazzo Vitturi et al. (1995) hypothesized a depth of 10-17 km for PIC
18
19
90 emplacement, but their values appeared in stark contrast with both the field evidence and the
20
21
2291 data reported by Visonà and Zanferrari (2000) for the similar and coeval Karawanken pluton
23
2492 (5-9 km).

25
26
93 Therefore, in this paper we made new geochemical and isotopic (⁸⁷Sr/⁸⁶Sr-¹⁴³Nd/¹⁴⁴Nd)
27
28
294 measurements on new samples from PIC that are representative of its main portions. A multiple
30
31
95 thermobarometric and hygrometric approach, based on the interaction between distinct single
32
33
3496 mineral, mineral pairs and mineral-melt equations, corroborated by appropriate simulations by
35
36
97 means of Rhyolite-MELTS software (Gualda et al. 2012), enabled us to: (i) provide for the first
37
38
398 time *P-T-fO₂* estimates and H₂O evaluation of PIC system over its entire evolution; (ii) verify
40
4199 if the depth of the intrusion obtained by the previous petrologic approaches well fit the field
42
43
44100 evidence. The resulting *P-T-fO₂* and H₂O values, together with the isotopic signatures of PIC
45
46
4701 rocks were then used as input to (iii) quantify the role of assimilation and fractional
47
48
102 crystallization processes in the generation of the main PIC magmatic suites.
49
50

5103 52 53 104 **2. Middle Triassic magmatism in the Southern Alps** 54

55 5605 **2.1. Geodynamic framework** 57

58
106 The Middle Triassic magmatic sequences of the Dolomitic Area are mainly composed of
59
60
6107 volcanites and volcanoclastites with subordinated and scattered intrusive bodies over an area of
62
63
64
65

108 about 2,000 km² (Vardabasso 1929, 1930; Castellarin et al. 1980, 1982; Lucchini et al. 1982;
1
209 Sloman 1989; Bonadiman et al. 1994; Coltorti et al. 1996; Gianolla et al. 2010; Casetta et al.
3
4
510 2017). The Middle Triassic geodynamic framework of Dolomitic Area and the whole Southern
6
711 Alps domain is still matter of debate, fostered by juxtaposition between the calc-
8
912 alkaline/shoshonitic orogenic affinity of the magmatic products and evidence for a concomitant
10
1113 extensional-transensional tectonic regime (Doglioni 1987; Stampfli and Borel 2002, 2004;
13
1414 Doglioni 2007). Several geodynamic models were thus proposed, invoking the presence of: (i)
15
1615 an aborted rift in a passive margin (Bernoulli and Lemoine 1980); (ii) a compression at the NW
18
1916 limb of the Paleo-Tethys (Castellarin et al. 1980); (iii) an “active” mantle upwelling (Stahle et
20
2117 al. 2001); (iv) a transition to back-arc conditions lasting since Carboniferous-Permian, triggered
23
2418 by the northward subduction of the Paleotethys remnants (Ziegler and Stampfli 2001; Stampfli
25
2619 and Borel 2002; 2004; Cassinis et al. 2008; Schmid et al. 2008 Zanetti et al. 2013); (v) an
28
2920 anorogenic rifting concomitant with **arc-like magmatism**, whose signature was inherited by a
30
31121 mantle source previously metasomatised by subduction-related components during the
32
33
3422 Hercynian orogenic cycle (Bonadiman et al. 1994). This last hypothesis was supported by the
35
3623 similar Sr-Nd isotope signatures of the Middle Triassic magmas in the Southern Alps and the
37
38
3924 Permian igneous products (260-290 Ma), whose expression were found along the Western and
40
4125 in the Eastern Alps sectors (Barth et al. 1993; Rottura et al. 1998; Monjoie et al. 2007;
42
43
4426 Schaltegger and Brack 2007; Marocchi et al. 2008; Quick et al. 2009; Willcock et al. 2015; Dal
45
4627 Piazz et al. 2015; Sinigoi et al. 2016; Manzotti et al. 2017). This kind of analogy led to speculate
47
48
4928 that similar mantle sources, and/or similar melting conditions, could have generated the Middle
50
5129 Triassic and the Permian magmatic episodes. In the Dolomitic Area, thick sequences of **basaltic**
52
53
5430 **andesitic to rhyolitic** ignimbrites associated to the Permian Atesina Volcanic District are also
55
5631 the main constituent of the crustal basement of the Dolomitic Area where magmas intruded
57
58
5932 during Middle Triassic.
60
6133

2.2. The Southern Alps intrusive bodies: geochemical and petrographic overview

In contrast to the large amount of basaltic/latitic volcanic rocks erupted in the Southern Alps during the Ladinian-Carnian, the intrusive bodies are more limited in volume. They are Cima Pape (< 3 km²), Mt. Monzoni (4.6 km²), Predazzo Intrusive Complex (25 km², [see below](#)) in the Dolomitic Area (Italy) and Karawanken (about 50 km²) in Carinthia (Austria).

The Cima Pape sill and the ca. 230 Ma Mt. Monzoni body (Borsi and Ferrara 1968) are composed of biotite and amphibole-bearing gabbroic to syenitic rocks with shoshonitic affinity, generated mainly by fractional crystallization from parental **basaltic to trachybasaltic magmas** (Gasparotto and Simboli 1991; Bonadiman et al. 1994; Della Lucia 1997). The emplacement of Mt. Monzoni body is thought to be controlled by a syn-genetic ESE-WNW transcurrent tectonics, which created the conditions for a shallow level intrusion and the subsequent magma differentiation. A shallow depth is also suggested for the Cima Pape sill, where a gradual textural transition to the overlying lavas, locally organized in columnar structures (Sarti and Ardizzoni 1984) is observable on field.

The Karawanken pluton (ca. 230 Ma, Lippoldt and Pidgeon 1974) comprises biotite and amphibole-bearing diorites/monzonites to granites/syenites, with rare aplites and pegmatites, showing an overall high-K calc-alkaline to shoshonitic affinity. To explain the production of Karawanken magmas, Visonà and Zanferrari (2000) invoked the occurrence of combined assimilation + **fractional crystallization** processes from a parental **basaltic** melt in a shallow magma chamber (5-9 km). The geochemical differences (i.e. La/Nb and Ba/Nb ratios) between **the** Karawanken pluton and the intrusions of the Dolomitic Area led the authors to suggest the origin of **Karawanken** magmas from an enriched mantle source not previously affected by subductive components, as proposed by Bonadiman et al. (1994) for the Mt. Monzoni body. Irrespective of the volumetric differences, common features of these intrusions are the high-K calc-alkaline to shoshonitic affinity, the modal occurrence of hydrated phases (biotite and amphibole) in the intercumulus assemblages and the shallow depth of emplacement.

2.3. The Predazzo Intrusive Complex

The Predazzo Intrusive Complex (PIC, Fig. 1) is a ring-like shaped multi-pulse intrusion with an overall volume of about 4.5 km³ cropping out in Trentino Alto Adige (NE Italy). Dated at 237.3 ± 1.0 Ma (U/Pb on zircon from granites, Mundil et al., 1996), it intruded the Permian Atesina Volcanic District rhyolitic ignimbrites, the Permo-Triassic sedimentary formations and the overlying volcanic sequences (Brack et al. 1996, 1997, 2005; Mietto et al. 2012), forming well-defined metamorphic aureoles (Princivalle et al. 1999; Ferry et al. 2002; Povoden et al. 2002; Gallien et al. 2007). The intrusion and its related volcanic products were almost completely preserved from the action of the Alpine tectonic event, and kept in their original position with respect to the surrounding rocks. According to Casetta et al. (2017), PIC rocks are grouped in three main units with different geochemical features (Fig. 1), named (in relative order of emplacement) Shoshonitic Silica Saturated (SS, 3.1 km³), Granitic Unit (GU, 1.1 km³) and Shoshonitic Silica Undersaturated (SU, 0.3 km³). The SS and SU suites are composed of pyroxenites/gabbros to syenites (Fig. 2), respectively quartz- and nepheline-bearing, formed as result of fractional crystallization (FC) processes in an almost closed system from two distinct starting monzogabbroic-like magmas (Petersen et al. 1980; Visonà 1997; Marrocchino et al. 2002; Casetta et al. 2017). The Granitic Unit is instead constituted by quartz and biotite-bearing granites/syenogranites (Fig. 2), and apparently seems to represent a different magma batch and/or the result of genetic processes independent from those that generated the SS and SU rocks (Menegazzo Vitturi et al. 1995; Visonà 1997; Casetta et al. 2017). However, the lack of isotopic studies made it impossible to definitively exclude for the three magma bodies a significant involvement of assimilated crustal rocks.

On the other side, the field relationships between the intrusive units, their related dykes and the host rocks suggest a shallow origin of this multi-pulse intrusion, as also testified by the presence of hypabyssal rocks at the hundred-meter-scale transition between the plutonic and the volcanic

186 sequences (Casetta et al. 2017). Nevertheless, the only thermobarometric studies present in
1 literature, based on the Al^{tot} content of amphiboles (Menegazzo Vitturi et al. 1995), proposed a
2 10-17 km depth for the PIC, giving rise to some uncertainty about its emplacement and
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3. Analytical methods

Whole rock major and trace element analyses were carried out at the Department of Physics and Earth Sciences (University of Ferrara, Italy) using ARL Advant-XP automated X-Ray fluorescence spectrometer. Full matrix correction procedure and intensities were elaborated following Traill and Lachance (1966). Accuracy and precision are better than 2-5% for major elements and 5-10% for trace elements. The detection limits are 0.01 wt% and 1-3 ppm for most of the major and trace element concentrations, respectively.

Rb, Sr, Y, Zr, Nb, Hf, Ta, Th, U, and Rare earth elements (REE) were analyzed by inductively coupled plasma-mass spectrometry (ICP-MS) using a Thermo Series X spectrometer with precision and accuracy better than 10% for all elements, well above the detection limit.

Mineral phases major element compositions were analyzed at the Department of Lithospheric Research, University of Wien (Austria), by using a CAMECA SX100 electron microprobe equipped with four WD and one ED spectrometers. The operating conditions were as follows: 15 kV accelerating voltage, 20 nA beam current, 20 s counting time on peak position. Natural and synthetic standards were used for calibration and PAP corrections were applied to the intensity data (Pouchou and Pichoir 1991).

Whole rock ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd isotopic analyses were performed at the Radiogenic Laboratory of the Scottish Universities Environmental Research Centre (SUERC) of Glasgow by means of a Sector-54 TIMS instrument. Strontium was loaded onto outgassed single Re filaments with a Ta-activator solution; Nd was loaded onto the side of a triple Ta-Re-Ta

212 filament assembly in H₂O and ran as Nd⁺. The sample preparation procedures for Sr-Nd isotopic
1
213 analyses have not been described previously, so we articulate them in full, together with
3
4
214 standards and instrumental analytical performance, in the Supplementary Material_1.
5
6

216 4. Whole rock geochemistry

217 4.1. Major and trace element

218 Shoshonitic Silica Saturated (SS) suite (Fig. 2, Table 1) is mainly constituted by monzogabbros
15
16 and monzodiorites, with Mg# (calculated as Mg/[Mg+Fe²⁺] mol%) of 65-45, and by
17
18 subordinated monzonites and syenites (Mg# of 53-26 and 45-20 respectively) randomly
19
20 distributed within the unit (Fig. 1). Volumetrically limited pyroxenitic (Mg# 64-47) and
21
22 gabbroic (Mg# 65-45) cumulates crop out in the southwestern sector of the intrusion.
23
24

225 Shoshonitic Silica Undersaturated (SU) suite (Fig. 2, Table 1) is on average more evolved than
26
27 the SS one, being dominated by the presence of nepheline-normative monzonites (Mg# 40-39)
28
29 and syenites (Mg# 36-15), over a minority composed of cumulitic gabbros (Mg# 56-50),
30
31 monzogabbros (Mg# 59-47) and monzodiorites (Mg# 55-27). Granitic Unit (GU, Fig. 2) is
32
33 composed of highly evolved granites and syenogranites (69-77 SiO₂ wt%, Table 1) that,
34
35 according to Chappell and White (2001) discriminating criteria, are ascribable to the I-type
36
37 granitoids (Al/[Na + K + Ca] < 1.1; low P₂O₅ wt%; decreasing Zr, Sr and Al₂O₃ wt% with
38
39 increasing SiO₂ wt%). Few samples show instead intermediate behaviour between I- and S-type
40
41 rocks.
42
43

44 All SS, SU and GU rocks have K-affinity (Fig. 2), and their K₂O and SiO₂ contents led to
45
46 classify them as belonging to shoshonitic (SS and SU) and High-K calc-alkaline (GU) series.
47
48

49 Despite the general similarities, the major and trace element distribution in the SS, SU and GU
50
51 rocks made it possible to pinpoint their origin from independent magmatic batches. At
52
53 comparable differentiation degrees, main discriminating features between the SS and SU rocks
54
55 are the relative enrichment in HFSE (Th, U, Pb), LREE and the higher Na₂O/K₂O ratio of the
56
57
58
59
60
61
62
63
64
65

latters. Such differences are in accordance with the predominant presence of amphibole and other Na and REE-rich phases in the SU rocks (Casetta et al. 2017).

4.2. $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ isotopes

Whole rock $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ isotopic ratios were analysed on representative samples of the SS, SU and GU suites (Table 2). Shoshonitic Silica Saturated samples have initial Sr isotope composition from 0.7039 to 0.7052, whereas their initial Nd isotope compositions range from 0.512191 to 0.512247. Shoshonitic Silica Undersaturated rocks are characterized by a generally higher $^{143}\text{Nd}/^{144}\text{Nd}_i$ varying from 0.512261 to 0.512289 and by $^{87}\text{Sr}/^{86}\text{Sr}_i$ comparable with the SS samples (0.7047 to 0.7063). $^{143}\text{Nd}/^{144}\text{Nd}_i$ range of GU rocks is between 0.512206 and 0.512304, thus comparable to those of both SS and SU suites. High Rb/Sr of the GU rocks resulted in imprecise initial calculated $^{87}\text{Sr}/^{86}\text{Sr}$ preventing any correlation between the GU and the SS/SU bodies.

5. Crystallization sequences and mineral chemistry

To better focus on the P - T - $f\text{O}_2$ conditions and on the H_2O contents of the PIC system, some remarks on the mineral phase compositions and crystallization sequences of the various units is hereafter summarized. A more detailed description of the petrographic and mineral chemistry features of these rocks was reported by Casetta et al. (2017). The new mineral phases analyses are reported in the [Supplementary Material_2](#).

Shoshonitic Silica Saturated rocks are dominated by the presence of clinopyroxene and plagioclase as cumulus phases; minor olivine and orthopyroxene can be found in gabbros and monzogabbros. The intercumulus assemblage is characterized by the ubiquitous presence of plagioclase and biotite, followed by the **appearance of** amphibole in gabbros to monzodiorites (Fig. 3). Magnetite and Ti-magnetite are often reported in association with biotite, whereas K-feldspar and accessory phases (quartz, apatite, ilmenite, sphene, zircon) modally increase in

264 more evolved rocks. The main alteration features of the SS rocks consist of sericite formation
1
265 at the expenses of feldspars, chlorite growth over clinopyroxene, amphibole and biotite, as well
3
4
266 as epidote formation at the expenses of clinopyroxene and plagioclase. The following
6
267 crystallization sequence can be deduced for SS rocks: *olivine* → *clinopyroxene*
8
268 (*±orthopyroxene*) → *Ti-magnetite/magnetite* → *plagioclase* → *biotite* → *amphibole* → *K-*
10
11 *feldspar* → *quartz* (*±accessories*).

14
270 The crystallization sequence of the SU magmatic suite mirrors its silica undersaturation and
15
16
271 higher Na₂O content. Orthopyroxene and quartz are in fact absent, and an earlier appearance of
18
19
272 amphibole at the expenses of biotite characterises the intercumulus assemblage of the SU rocks
20
21
273 (Fig. 3). Other Na-rich minerals like ferrosalitic clinopyroxene, abitic plagioclase and nepheline
23
24
274 (Visonà 1997) are variably present. Accessory phases are ugrandite group garnets, epidote,
25
26
275 apatite and titanite. As already highlighted by Visonà (1997), many portions of the SU body are
28
29
276 hydrothermalized, showing the formation of kaoline, sericite and scapolite. Where this
30
31
277 secondary assemblage does not occur, the primary differences between the SU and SS
32
33
34
278 magmatic suites are evident. The likely crystallization sequence of the SU rocks is: *olivine* →
35
36
279 *clinopyroxene* → *Ti-magnetite/magnetite* → *plagioclase* → *amphibole* → *biotite* → *K-feldspar*
37
38
39
380 → *nepheline* (*±accessories*).

40
41
281 Syenogranites and granites of the GU are quite homogeneous in composition, being constituted
42
43
282 by K-feldspar, plagioclase and quartz, locally associated to several other minerals, among
45
46
283 which Fe-rich biotite is the most common. Accessory phases are tourmaline, fluorite, sphene,
47
48
284 magnetite, apatite, ilmenite and zircon. Muscovite and chlorite are rare and always grow at the
49
50
51
285 expenses of biotite and amphibole in late hydrothermal stages, whereas the sericitization of
52
53
286 feldspars is pervasive. Less common phases are ematite, allanite, scheelite, xenotime, gummite,
54
55
56
287 thorite, uranium micas, molibdenite and other sulphides, irregularly disseminated and generated
57
58
288 in the later pneumatolitic and hydrothermal stages (Marzocchi 1987; Visonà 1997). The main
59
60
61
62
63
64
65

289 GU paragenesis is composed of: *K-feldspar + plagioclase + quartz + fluorite + biotite (chlorite*
1
290 *and/or muscovite) ± amphibole + accessories.*

291 **In the less altered samples, at comparable whole rock differentiation degree, the composition**
292 **of the most representative minerals clearly discriminate between the SS and SU magmatic**
293 **suites. For instance, biotites** and amphiboles are Al- and Na-enriched in the SU rocks with
294 respect to the SS ones. Amphibole in fact varies from the hornblendic and actinolitic
295 composition in the SS rocks ($\text{Na}_2\text{O} < 1.5 \text{ wt\%}$; $\text{Al}_2\text{O}_3 < 8 \text{ wt\%}$) to the hastingsitic/pargasitic
296 **composition** (rare actinolitic hornblende, Leake et al. 1997) in the SU ones (Na_2O up to ~ 2.5
297 wt\% ; Al_2O_3 up to $\sim 18 \text{ wt\%}$). Even the Na content of plagioclase differs between the two
298 magmatic suites: it ranges from An_{84} to An_{34} in SS samples, and from An_{50} to $\text{An}_{<23}$ in SU
299 rocks. Another discriminating feature is represented by the clinopyroxene composition,
300 augitic/salitic (En_{43-33}) in the SS and salitic/ferrosalitic (En_{37-24}) in the SU rocks (**Morimoto,**
301 **1988**). Furthermore, the manganese content of several phases is quite higher in the SU rocks:
302 Ti-magnetite (0.2-1.9 wt% SU; 0.1-0.4 wt% SS), biotite (0.1- 0.7 wt% SU; 0-0.4 wt% SS),
303 amphibole ($\sim 0.7 \text{ wt\%}$ SU; $\sim 0.5 \text{ wt\%}$ SS) and clinopyroxene (0.5-1.5 wt% SU; 0.2-0.8 wt% SS).

305 **6. P, T and H₂O estimates: thermobarometric evolution of the intrusion**

306 6.1. Recover the initial equilibrium conditions

307 In this section, we apply several methods for estimating the *P*, *T* and H₂O contents of PIC rocks,
308 taking into account the coexistence of minerals used as thermometers, barometers and/or
309 hygrometers in both the SS and SU crystallization sequences. Apart from the Holland and
310 Blundy (1994), Anderson and Smith (1995), Anderson (1996) and Henry et al. (2005) methods
311 (see below) applied to amphibole-plagioclase pairs and biotite, and specific for intrusive rocks,
312 the other equations used in this study are designed for volcanic samples. As often happens in
313 intrusive contexts, the identification of the **parental** “melt” from which crystals formed is
314 challenging (e.g. **Skaergaard Intrusion; Nielsen 2004; Namur and Humphreys 2018**). Intrusive

315 rocks, in fact, rarely correspond to bulk melt composition, representing a variable mixture of
1
316 cumulus and intercumulus minerals, removed by the crystallizing melts at various stages of
3
4
317 fractionation, thus in equilibrium with different melts in different moments. Casetta et al. (2017)
6
318 suggested that the cumulitic gabbros and pyroxenites, as well as most of the intermediately
8
319 evolved rocks of the complex, were generated by various extent of fractional crystallization
10
11
120 from a starting trachybasaltic magma in an almost closed system.

14
321 It is modelled that the mineral assemblages constituting the PIC rocks (e.g. clinopyroxenes of
15
16
322 SS gabbro) showed partial equilibrium with a more evolved melt (e.g. SS trachybasalt), from
18
19
323 which then segregated to form the cumulates. The cumulitic nature of a rock itself incorporates
20
21
324 the concept that its composition (e.g. Mg# of clinopyroxene) represents an independent physico-
23
24
325 chemical system with respect to the deriving melt. Consequently, if we try to consider the
25
26
326 intrusive (cumulitic) rock of PIC as a bulk representative of a melt (e.g. in terms of Mg# or
27
28
327 CaO/Al₂O₃), it is reasonable to find that its mineral constituents (e.g. clinopyroxene or
30
31
328 plagioclase), are in evident disequilibrium with the bulk composition (Fig. 4 and 5). They are
32
33
329 instead compositionally coherent with a segregation process from a more evolved melt. On the
35
36
330 other side, by taking into account more evolved rocks (e.g. monzogabbros to syenites), the
37
38
331 amount of crystal disequilibrium progressively decreases (Fig. 4 and 5). This because the
40
41
332 ultimate products of the SS/SU fractional crystallization processes, syenitic in composition,
42
43
333 progressively approach the eutectic of the system and likely resemble a melt composition: this
44
45
334 condition has been also proposed by Morse and Brady (2017) for the syenites of the 1300 Ma
47
48
335 old Kiglapait Intrusion (Labrador).

50
336 Evidences of the reliability of this genetic model are the observed textural and mineral
52
53
337 homogeneities of the SS/SU rocks: any kind of significant zoned texture would result from the
54
55
338 introduction of additional magma chamber processes (i.e. mixing) during the formation of the
57
58
339 PIC lithotypes, invalidating our assumptions. The unzoned texture, coupled with the crystal size
59
60
340 of PIC rocks (e.g. clinopyroxenes up to 6-7 mm in pyroxenites and gabbros) and with the small

341 volume of the intrusive body (4.5 km³), speaks also against a considerable effect of syn- to post-
1
342 crystallization diffusion processes, which can be a rate-limiting process for thermobarometric
3
4
343 estimates. On the other side, thermometers involving clinopyroxene in upper crustal context,
6
344 commencing cooling from relatively low *T*, hold the potential to record the peak temperature
8
345 conditions, especially in large-sized grains (Müller et al. 2013). If we couple the euhedral
10
11
346 unzoned texture of PIC clinopyroxenes with the diffusion rates proposed by several authors
13
14
347 (e.g. at about 1000°C $\log D^{(Ca)} = -21$ m²/s; $\log D^{(Ti)} = -22$ m²/s; $\log D^{(Fe)} = -21$ m²/s, where *D*
15
16
348 indicates the diffusion coefficient; see Brady and McCallister 1983; Dimanov et al. 1996;
18
19
349 Cherniak and Lyiang 2012; Müller et al. 2013 and references therein), we can argue that the
20
21
350 crystal compositions were nearly unmodified by significant diffusion processes. Similar
23
24
351 remarks can be made for plagioclase, whose crystallization in plutonic rocks at relatively low
25
26
352 *T* limit the efficiency of CaAl-NaSi diffusion to submicron length scales (Grove et al. 1984).
27
28
353 The mineral compositions were therefore considered as representative of the various stages of
30
31
354 fractional crystallization of the magmas inside PIC, and were used to constrain the physical
32
33
355 parameters of the magma chamber.
35
36
356 On the basis of these assumptions, we matched each mineral (e.g. clinopyroxene of SS
37
38
357 cumulates) with an estimated “melt” following the crystal/melt equilibrium partitioning (e.g.
40
41
358 $K_{Cpx-Liq}^{Fe-Mg}$, see below), to trace the P-T path of an hypothetical differentiation trend (e.g. SS
42
43
359 suite, see Casetta et al. 2017). This operation enabled us to compute the “melt” composition, to
45
46
360 skip the apparent disequilibrium between minerals and cumulitic rocks as a bulk and thus to
47
48
361 constrain the *P*, *T* and H₂O parameters of the less evolved melts in the PIC feeding system.
49
50
362 Afterwards, the application of the “traditional” thermobarometric equations for intrusive
52
53
363 contexts (i.e. to amphibole-plagioclase pairs and biotite, see below) enabled us to verify the
54
55
364 convergence between the various results as well as to unravel the *T-P-H₂O* conditions in the
57
58
365 later stages of crystallization. A complete list of the applied equations, results, and
59
60
366 corresponding errors is reported in Table 4 and discussed in detail in the following sections.
62
63
64
65

6.2. Clinopyroxene-melt thermobarometer and water content of primary magmas

Clinopyroxene is the dominant and ubiquitous phase in the SS and SU rocks, thus clinopyroxene-melt thermobarometry was considered as a valuable starting point to estimate the T - P intensive variables and H₂O content of PIC magmatic system since early stage of fractionation. Crystal-melt equilibrium conditions were tested by taking into account the experimentally determine range of $K_{\text{Fe-Mg}}^{\text{Cpx-Liq}} = 0.24\text{-}0.30$ at $T > 1050^\circ\text{C}$ by Putirka et al. (2003). First tests highlighted that all clinopyroxene crystals in the SS and SU units have $K_{\text{Fe-Mg}}^{\text{Cpx-Liq}} = 0.36\text{-}0.68$ (hereinafter the superscript “Liq” indicates the composition of the bulk rock), indicating a disequilibrium towards composition more evolved than their host rock (Fig. 4). The amount of disequilibrium, as said before, decreases with increasing the differentiation degree of the rock (Fig. 4). Such a decrease could be explained by the fact that more evolved lithotypes better approach the eutectic of the system and more likely simulate melt compositions.

Following the FC model of Casetta et al. (2017), clinopyroxenes of each lithotype were thus related to a calculated “melt” composition having the same chemical affinity of their host rock and a more evolved nature (Table 3), by which crystals retrieved their equilibrium conditions.

As shown in the flow chart of Fig. 6a, once chosen the clinopyroxene-melt couple of the SS suite, T - P pairs were estrapolated by means of the Putirka (2008) equations, assuming variable H₂O wt% contents of the melt. The water-dependant equations 32b and 33 (Putirka 2008), derived from the P -independent thermometer and the T -dependent barometer of Putirka (1996), were firstly applied to constrain the crystallization conditions of clinopyroxene from the melt, obtaining several T - P -H₂O triplets. Since these triplets are strongly dependant on the chosen melt and on the inferred H₂O content, they were checked, together with the melt composition, by means of the Rhyolite-MELTS calibration (Gualda et al. 2012). Iterated procedures (Fig. 6a) were developed until a clinopyroxene composition analogous to the starting one was reproduced

393 by Rhyolite-MELTS. In this way, the two independent approaches reinforce one to another,
1
394 giving a more robust framework for the thermobarometric and water results. Since the only left
3
4
395 starting parameter required by Rhyolite-MELTS was oxygen fugacity, we chose $f_{O_2} = NNO$ as
6
396 proposed by Bonadiman et al. (1994) for the Mt. Monzoni intrusion.

397 However, by increasing the differentiation degree of SS rocks, and/or considering the more
10
11
1398 alkaline SU magmas, the reliability of Rhyolite-MELTS decreases, as a function of the alkali-
13
14
1399 enrichment of the system and the related crystallization of amphibole and biotite, for whom
15
16
1400 thermodynamic parameters are not well constrained (Gualda et al. 2012). Therefore, the use of
18
19
1401 Masotta et al. (2013) thermobarometer was preferred for the SU suite and for the more evolved
20
21
1402 SS rocks, and was cross-checked with Putirka (2008) equations (Fig. 6b). As for SS rocks, once
22
23
1403 assessed the equilibrium between clinopyroxene and melt, T - P pairs at variable H_2O contents
25
26
1404 of the melt were extrapolated by means of the Putirka (2008) method. The resulting T - P - H_2O
27
28
1405 triplets were then inputed in Masotta et al. (2013) equation until the resulting T and P were
30
31
1406 comparable to those obtained by Putirka (2008) method at similar water contents.

33
1407 Some issues for handling this combination of modeling tools should be remarked: (i) since
35
1408 Rhyolite-MELTS is not suitable for amphibole and biotite-dominated rocks, it was only used
37
38
1409 to simulate the composition of clinopyroxene in equilibrium with the least evolved SS magmas,
40
1410 at the P , T and H_2O conditions determined by the previous calculations; (ii) the alkaline nature
42
43
1411 of PIC (especially SU) differentiated rocks implies that clinopyroxene-melt behavior during
45
1412 crystallization and cooling depends not only on Fe and Mg, but also on Ca, Na and Al, so
47
48
1413 Masotta et al. (2013) equations became progressively more reliable; (iii) iterated and cross-
49
50
1414 checked calculations using Putirka (2008), Rhyolite-MELTS and Masotta et al. (2013) methods
52
53
1415 (Fig. 6a-b) were proposed to unravel P - T - H_2O crystallization conditions avoiding circular
54
55
1416 relationships between the equations.

57
1417 The errors proposed for the thermobarometers, oxybarometers and hygrometers are reported in
59
60
1418 Table 4: while temperatures could be defined within a narrow error range (± 10 to $\pm 20^\circ C$) with
61
62
63
64
65

419 both Putirka et al. (2008) and Masotta et al. (2013) equations, the P estimates were affected by
1
420 high uncertainties, varying from ± 1.2 (Masotta et al. 2013) to ± 2.6 kbar (Putirka et al. 2008).
3

422 6.2.1. Clinopyroxene crystallization conditions of Shoshonitic Silica Saturated rocks 8

423 The most magnesian clinopyroxene in SS rocks were found within pyroxenites ($Mg\#_{Cpx}$ 76-78),
10
11 whereas the less magnesian compositions ($Mg\#_{Cpx}$ 63-71) were in the monzodiorites and
12
13 monzonites. The early T - P - H_2O crystallization conditions were likely represented by the
14
15 former, which attained equilibrium with a trachybasaltic melt ($Mg\#$ 51; Fig. 4a, Table 3), further
16
17 confirming the first FC step identified by Casetta et al. (2017). Clinopyroxene-melt
18
19 thermobarometers indicated a P - T range of 1.6-2.3 kbar and 1070-1050°C, for water contents
20
21 of 2.0-2.5 wt% (Table 4). The best fitting data in by Rhyolite-MELTS simulations were
22
23 obtained at pressure of 1.5 kbar, temperature of 1060°C and 2.5 H_2O wt%. Consequently, 2.0-
24
25 2.5 H_2O wt% could be assumed as the water contents of the first SS magmas intruding the PIC.
26
27 Clinopyroxenes in the cumulitic gabbros were slightly less homogeneous and less magnesian
28
29 ($Mg\#_{Cpx}$ 69-71 cores; 65-66 rims), and resulted in equilibrium with melts varying in $Mg\#$ from
30
31 43 to 40 (cores) to about $Mg\#$ 36 (rims), as shown in Fig. 4a and Table 3. Pressure-temperature
32
33 paths indicated that cores attained equilibrium in a P range of 0.5-1.9 kbar and T of 910-980°C,
34
35 while rims formed at 0.3-1.2 kbar and 900-920°C, considering H_2O contents of 3.0-4.0 wt%.
36
37 Clinopyroxenes in monzogabbros and monzodiorites were nearly unzoned although
38
39 compositionally comparable to those in gabbros, showing $Mg\#_{Cpx}$ of 68-72. As for gabbroic
40
41 rocks, equilibrium was reached with intermediately evolved magmas ($Mg\#$ 40, Fig. 4a, Table
42
43 3), by which thermobarometric equations indicated a T range of 940-990°C at P of 1.4-1.6 kbar,
44
45 for a H_2O contents in the melt of 3.0-4.0 wt%. All results are reported in Table 4 and Fig. 7a.
46
47
48
49
50
51
52
53
54
55
56
57

58 6.2.2. Clinopyroxene crystallization conditions of Shoshonitic Silica Undersaturated rocks 59

444 Clinopyroxenes from the SU suite are more alkaline than those of the SS rocks, ranging from
1
445 salitic-aegirinaugitic to ferrosalitic compositions. The most magnesian compositions in gabbros
3
446 and monzogabbros (Mg#_{Cpx} 67-63) were in equilibrium with Mg# 36 melts (Fig. 4b, Table 3).
6
447 Pressure-temperature paths indicate a *P* range of 0.2-1.6 kbar and *T* of 920-940°C at water
8
448 contents of about 1.0-1.5 wt%. Clinopyroxenes in syenites (Mg#_{Cpx} 60-48) show composition
10
1449 close to the equilibrium with the whole rock (Fig. 4b, Table 3), which represent a theoretical
13
1450 slightly evolved trachytic melt. Thermobarometers yielded a *P* of 0.2-1.6 kbar at *T* of 810-
15
1451 860°C at water contents between 4.0 and 5.0 wt%. Results are reported in Table 4 and shown
18
1452 in Fig. 7a.
20
21
22
23

454 6.3. Amphibole and amphibole-plagioclase thermobarometer and hygrometer 25

455 Pressure, temperature and H₂O conditions for amphibole crystallization were estimated using
27
28
456 amphibole-plagioclase Al-exchange thermobarometers (Holland and Blundy 1994; Anderson
30
457 and Smith 1995; Anderson 1996) and a single amphibole thermobarometers (Ridolfi et al. 2010;
32
33
458 Ridolfi and Renzulli 2012). Amphibole-plagioclase equations were used for mineral pairs in
35
459 both SS and SU suites, whereas the single amphibole thermobarometers were applied to all
37
38
460 amphibole compositions in the SS rocks, but only for amphibole in gabbros in the SU suite.
40
461 This because of the higher alkali content of SU differentiated magmas, well above the
42
43
462 compositional ranges admitted for the calibration of the empirical geothermometers of Ridolfi
45
463 et al. (2010) and Ridolfi and Renzulli (2012). For the SS and SU amphiboles, the single
47
48
464 amphibole equations enabled also the calculation of the water content of the melt from which
49
50
465 they crystallized. Contrary to the clinopyroxene-melt equations, the error range for the
52
53
466 barometric estimates based on amphibole-plagioclase and single amphibole was lower, ranging
54
55
467 between ±0.03 and ±0.6 kbar, whereas errors on *T* estimates were slightly higher (±23 to ±40°C,
57
58
468 Table 4).
59
60
469
61
62
63
64
65

470 6.3.1. Amphibole crystallization conditions of Shoshonitic Silica Saturated and Shoshonitic
1
471 Silica Undersaturated rocks
3

472 Pressure-temperature paths obtained from amphibole-plagioclase thermobarometers yielded a
6
473 T - P range of 750-845°C and 0.1-1.2 kbar in SS gabbros to monzodiorites (Fig. 7a, Table 4).
8

474 Single amphibole calculations for the same lithotypes confirmed these values, indicating P =
10
11
1475 0.7-1.4 kbar and T = 720-810°C. The estimated H₂O contents of melts in equilibrium with
13
14
476 amphiboles of SS gabbros to monzodiorites was between 4.8 and 6.0 wt% (Fig. 7b, Table 4).
15

16
1477 Amphibole-plagioclase estimates for the SU suite yielded similar P intervals (0.4-1.0 kbar) but
18
19
478 at lower T (620-660°C) for gabbros, and P - T ranges of 1.6-1.9 kbar and 870-880°C taking into
20
21
479 account the mineral pairs in the monzogabbroic rocks (Fig. 7a, Table 4). The single amphibole
22
23
480 method, applied only to the actinolitic hornblendes in SU gabbros, provided comparable
25
26
481 pressures (0.5-0.6 kbar) and temperatures (700-720°C) with respect to the SS rocks (Fig. 7a,
27
28
482 Table 4). The calculated water contents of the melts were similar to those obtained for the SS
30
31
483 suite (5.1-5.7 wt%, Fig. 7b, Table 4).
32

33
484 As expected, the water contents calculated by these models were higher than those obtained
35
36
485 from the clinopyroxene-melt calculations. Amphibole crystallization, in fact, occurred later
37
38
486 than clinopyroxene, thus the melt from which amphibole precipitated should have undergone
40
41
487 differentiation in some extent, and its water content should have increased. If we consider the
42
43
488 water enrichment linked to the fractional crystallization of the SS/SU magmas, together with
44
45
489 the concomitant precipitation of volatile-bearing phases, the H₂O contents resulted from the
47
48
490 single amphibole equations appear in line with the calculations proposed by the FC model of
49
50
491 Casetta et al. (2017).
52

53
492
54
55
493 6.4. Biotite thermometer
56
57

58
494 Biotite is a ubiquitous phase occurring in the late inter-cumulus mineral assemblages of almost
59
60
495 all SS, SU and GU rocks. Its presence enabled us to provide some estimates on the temperatures
61
62
63
64
65

496 of these assemblages, by using of the empirical single-mineral thermometer of Henry et al.
1
497 (2005), based on the Ti content of biotite.
3

4
498
5
6
499 *6.4.1. Biotite crystallization conditions of Shoshonitic Silica Saturated, Shoshonitic Silica*
8
500 *Undersaturated and Granitic Unit rocks*
10

11
501 Temperature of biotite crystallization in SS pyroxenitic cumulates resulted between 690 and
13
14
502 740°C, whereas a *T* interval of 600-660°C was considered representative of SS gabbros to
15
16
503 monzodiorites. Biotites from more differentiated SS monzonites yielded a *T* of about 540-
18
19
504 580°C (Fig. 7a, Table 4). Biotite in SU gabbros and monzogabbros indicated comparable
20
21
505 temperatures, ranging between 640 and 660°C (Fig. 7a, Table 4).
22

23
24
506 The thermometer was also applied to GU syenogranites, resulting in a *T* range of 420-570°C:
25
26
507 the biotites in GU rocks are however iron-rich, with $Mg/(Mg+Fe_{tot}) < 0.275$, just out of the
27
28
508 compositional field for which the thermometer was calibrated. Consequently, the resulting
30
31
509 temperatures were not used for futher modeling purposes.
32

33
34
510
35
36
511 **6.5. Plagioclase-melt thermobarometer and hygrometer**
37

38
512 Plagioclase appears in almost all PIC rocks, from the cumulitic pyroxenites, where it is part of
40
41
513 the intercumulus assemblages, to the syenitic rocks, where, together with K-feldspar, dominates
42
43
514 the paragenesis. Its ubiquitous presence made it an additional tool for investigating the *T* and
44
45
515 H₂O content of PIC magmas along the entire SS and SU differentiation trends by means of
47
48
516 Putirka (2008) and Lange et al. (2009) equations. Plagioclase from GU granites and
49
50
517 syenogranites are compositionally close to the pure albitic end-member, probably because of
52
53
518 hydrothermal/alteration processes. This made them unsuitable for the hygrometer application,
54
55
519 being its calibration limited to An₃₇ plagioclase (Lange et al. 2009).
57

58
520 As for clinopyroxene, plagioclase equilibrium with its host rock was constrained by means of
59
60
521 the experimentally determined values of $Kd_{An-Ab}^{Pl-Liq} = 0.10 \pm 0.05$ at *T* < 1050 °C and 0.27 ± 0.11
62
63
64
65

522 at $T \geq 1050$ °C (Putirka 2008). However, Mollo et al. (2011) demonstrated that $K_{\text{An-Ab}}^{\text{Plag-Liq}}$
1
523 is highly sensitive to the cooling rate of the melts, being thus variable between 0.2 ± 0.02
3
524 (cooling rate of $0.5^\circ\text{C}/\text{min}$) and 0.35 ± 0.03 (cooling rate of $15^\circ\text{C}/\text{min}$). Thus, all plagioclase-
6
525 melt equilibria (even at $T < 1050^\circ\text{C}$) were considered following a $K_{\text{An-Ab}}^{\text{Plag-Liq}}$ in the range of
8
526 0.27 ± 0.11 (Putirka 2008), whose interval is also in accordance with the results obtained by
10
11 Mollo et al. (2011). Cooling rate is in fact an essential factor for the crystallization dynamics of
12
13 intrusive bodies, as testified by the plagioclase morphological variations in PIC rocks, where it
14
15 appears as both cumulus and intercumulus phase throughout the entire SS/SU fractionation
16
17 trends. According to crystallization sequences, plagioclase appeared after clinopyroxene and
18
19 Fe-Ti oxides, just before or even contemporary to biotite and amphibole in the intercumulus
20
21 assemblages. Consequently, we considered the thermobarometric values obtained by
22
23 clinopyroxene, amphibole and biotite calculations for each magmatic suite (see above) to
24
25 identify the P - T - H_2O interval in which plagioclase crystallized from the melt. As for
26
27 clinopyroxene, plagioclase often appeared in disequilibrium with its whole rock composition
28
29 (Fig. 5), and the amount of disequilibrium gradually decreases with increasing the
30
31 differentiation degree of the corresponding whole rock. Plagioclase in monzodiorites were in
32
33 fact closer to the equilibrium with respect to those of pyroxenites and gabbros, whereas most
34
35 of the plagioclase from monzonites and syenites were in equilibrium with their host rock (Fig.
36
37 5). As previously mentioned for clinopyroxene, such a trend could be justified by considering
38
39 that the more evolved rocks likely resemble melt compositions, being close to the eutectic of
40
41 the system. Thus, plagioclase compositions were related to estimated melt compositions,
42
43 according to the differentiation trends proposed by Casetta et al. (2017), to retrieve the
44
45 equilibrium conditions. Once equilibrated, plagioclase and melt compositions were used as
46
47 input for the Lange et al. (2009) hygrometer to calculate the amount of H_2O dissolved in the
48
49 melt (see flow chart of Fig. 6). For each sample, input T and P required in Lange et al. (2009)
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

547 equation were chosen in the T - P interval between those estimated by clinopyroxene-melt
548 calculations and those resulted from amphibole and biotite.

549 Pressure, temperature and H_2O were calculated by iterating Lange et al. (2009) method and
550 equation 24a of Putirka (2008) until a matching T was found. In any case, small variations of P
551 scarcely affect **the results of the** hygrometers: ± 1 kbar corresponds to ± 0.1 wt% H_2O calculated
552 in the melt (Giacomoni et al. 2014). Since the P interval obtained by the previous
553 (clinopyroxene and amphibole) barometers was roughly between 0.2 and 1.9 kbar for all PIC
554 rocks, the H_2O estimates were almost entirely constrained by temperature changes. As a
555 consequence, the match between T obtained with Lange et al. (2009) and equation 24a of
556 Putirka (2008) was considered a reliable cross-check and a robust constraint on our geothermal
557 modelling (Fig. 6).

558 *6.5.1. Plagioclase crystallization conditions of Shoshonitic Silica Saturated rocks*

559 Data from clinopyroxene-melt simulations on pyroxenites indicated that SS primary magmas
560 began crystallizing at about 1.5 kbar, 1060°C and 2.5 H_2O wt%. The more anorthitic plagioclase
561 (An_{84-73}) were analyzed in the pyroxenites intercumulus assemblage, following clinopyroxene
562 and preceding biotite, thus the T range of plagioclase crystallization is constrained by the
563 temperatures of crystallization of these two phases. Since pressure does not sensitively affect
564 the calculations, a pressure of 1.5 kbar was used according to the results from clinopyroxene-
565 melt calculations (Fig. 6 and 7). Results indicated that the most anorthitic plagioclase (An_{84-73})
566 crystallized in equilibrium with a **trachybasaltic melt (Mg# 51; Fig. 5, Table 3)**: at $P = 1.5$ kbar,
567 T of crystallization resulted of 1060-1081°C for a H_2O content **of 3.0 wt%**.

568 Plagioclase in SS gabbros were in compositional continuity with those of pyroxenites, varying
569 between An_{73} and An_{51} in both the cumulus and intercumulus assemblages. As for plagioclase
570 inside pyroxenites, the marked disequilibrium (Fig. 5) was probably related to the cumulitic
571 nature of gabbros and/or a variable cooling rate of the magma. The crystallization sequence and
572

573 the comparison with the T ranges obtained from clinopyroxene, amphibole and biotite
1 suggested that plagioclase crystallized in a temperature interval of 980 to 740°C. Plagioclase
574 with An₇₃ reached the equilibrium with a Mg# 49 melt (Fig. 5): at P of 1.5 kbar, T and H₂O
3 content resulted of 1016-1053°C and 3.1 wt% respectively. An₆₃₋₅₁ plagioclases were
575 equilibrated with a **basaltic trachyandesitic** melt (Fig. 5, **Table 3**), yielding T of 920-1021°C
6 and a water content of 3.2-3.4 wt% at pressures of 1.2-1.5 kbar (Fig. 7, Table 4). It is worth
576 noting that for these temperatures, Putirka (2008) proposed a potential partitioning $Kd_{An-Ab}^{Plag-Liq}$
8 = 0.05-0.15 to attest equilibrium. However, after several iterations, the modelled P - T -H₂O
577 terms for PIC magmas were attained only within a $Kd_{An-Ab}^{Plag-Liq}$ of 0.27±0.11. Such apparent
10 discrepancies could be explained considering the dependency of An-Ab equilibrium coefficient
11 with the cooling rate (Mollo et al. 2011).
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

584 Plagioclase from monzogabbros to monzonites were in a compositional range from An₅₃ to
27 An₃₄, resulting in equilibrium with **basaltic trachyandesitic** to **trachyandesitic** melts (Fig. 5,
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

6.5.2. Plagioclase crystallization conditions of Shoshonitic Silica Undersaturated rocks

490 Plagioclase in SU rocks range from An₅₀₋₄₃ in gabbros to An₃₈₋₂₃ in monzogabbros and
42 monzodiorites, and reach the more albitic compositions in syenites, where anorthite content is
43 low (An₂₃-An₂). Plagioclase in gabbros were equilibrated by an intermediately evolved melt,
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

598 Apart from the nearly pure albitic compositions (likely effects of secondary processes),
1
599 plagioclase in monzogabbros to syenites vary from An₃₈ to An₂₃. They were equilibrated with
3
600 an evolved **trachytic** melt (Fig. 5, **Table 3**), thought to be the final product of the SU
6
601 differentiation trend (Casetta et al. 2017). At $T - P$ space of 920-936°C and 1.2 kbar, the
8
602 estimated water contents of melt in equilibrium with An₃₈ plagioclases was about 4.4 wt%,
10
11 progressively increasing with the differentiation of the samples (Fig. 7, Table 4).
13

14 15 16 1605 6.6. K-Feldspar-melt hygrometer

18
19
606 Except for pyroxenites, K-feldspar is present in all PIC rocks, where it occurs as intercumulus
20
21 (gabbros to monzodiorites) and cumulus (monzonites/syenites) phase. Its composition was used
22
23 to constrain the amount of H₂O dissolved in the co-existing melt by means of the equation
24
25 proposed by Mollo et al. (2015). Such method, based on the Or-Ab exchange between K-
26
609 feldspar and melt **is calibrated for alkaline differentiated magmas, thus proper for the SS, SU**
27
28 **and GU rocks compositions**. Together with the data obtained by the previous equations, such
29
30 estimates enabled to “track” the H₂O contents in the progressively differentiating melts.
31
32 According to Mollo et al. (2015), this method can be applied only to K-feldspar with Or₄₄₋₈₆,
33
34 whereas no reliable results were produced for Or_{>86} as it extended outside the range of calibrated
35
36 compositions. K-feldspar and whole rock equilibrium evaluation was attained by minimizing
37
38 the difference between predicted and measured $K^{K-Feld-Liq}Kd_{Or-Ab}$, following equation 2 of the
39
40 Mollo et al. (2015) model. These conditions were satisfied only by considering K-feldspar
41
42 compositions in equilibrium with **trachyandesitic to trachytic melts (Table 3)**, further
43
616 confirming the correspondence between the crystallization sequence and the progressive
44
45 differentiation model proposed by Casetta et al. (2017).
46
47 According to the equilibration temperatures recorded by clinopyroxene, amphibole, biotite and
48
618 plagioclase, a T of 900-800°C was used as input for all samples of the SS and SU suites, whereas
49
50 lower temperatures (800-700°C) were considered for GU syenogranites.
51
52
53
54
55
56
57
58
622
59
60
623
62
63
64
65

624
1
625
3
4
626
6
627
8
9
628
10
11
629
13
14
630
15
16
631
18
19
632
20
21
633
22
23
634
25
26
635
27
28
636
30
31
637
32
33
638
35
639
37
38
640
40
641
42
642
43
44
643
47
644
49
50
645
52
646
54
55
647
57
648
59
60
61
62
63
64
65

6.6.1. K-Feldspar crystallization conditions of Shoshonitic Silica Saturated, Shoshonitic Silica Undersaturated and Granitic Unit rocks

Water concentration of melts in equilibrium with K-feldspar in SS gabbros to monzogabbros were in the range of 4.4-5.3 wt% and 5.6-6.5 wt% respectively, whereas higher H₂O contents (6.1-7.3 wt%) were recorded in monzodiorites. The highest water values were calculated in SS syenites, that in turn show a larger variability of values (H₂O = 4.6-7.3 wt%; Fig. 7, Table 4).

K-feldspar analysed in SU monzogabbros and monzonites indicated H₂O contents of the crystallizing melts between 5.5-6.3 wt% and 6.7-7.8 wt%, respectively. H₂O estimates for GU syenogranites range between 8 and 11 wt%, being however meaningless in the physical system compatible with the body emplacement. The solubility of water at $P < 2.5$ kbar for rhyolitic-trachytic melts in fact do not exceed 7 wt%, as experimentally and theoretically determined in the T range of 500-1000°C (Di Matteo et al. 2004; Liu et al. 2005). These values were therefore discharged in the discussion.

7. Oxygen fugacity

The two main magmatic suites (SS and SU) are formed by the differentiation of melts which began crystallizing at pressures lower than 2.0 kbar and at similar T ranges (1050-1000°C). Main differences between the SS and SU suites could be found by taking into account the total amount of water of the crystallizing melts: SS primitive melts in fact were characterized by an higher water contents with respect to that of SU ones (2.0-2.5 wt% vs. 1.0-1.5 wt%, respectively). Classically, it is considered that dissolved molecular H₂O reacts with oxygens of the silicate network producing two OH⁻ groups (e.g. Stolper 1982; Silver and Stolper 1985; Kohn 2000). The overall reaction reading as $H_2O + O^{2-} = 2OH^-$ may thus be considered as the counterpart of the water solubility in evaluating the redox conditions.

649 **To effectively quantify such differences, we** estimated the $f\text{O}_2$ conditions of each portion of PIC
1
650 by means of the T - $f\text{O}_2$ model of Burkhard (1991), based on the biotite/K-feldspar/magnetite
3
4
651 equilibrium in biotite-bearing intrusive rocks. Burkhard's (1991) empirical equation
6
652 complements that of Kress and Carmichael (1988), enabling to calculate oxygen fugacity and
8
9
653 temperature by two independent formulas. Water fugacity required in Burkhard's (1991)
10
11
654 equation was calculated at a fixed P of 1.5 kbar following Burnham et al. (1969). Sanidine
13
14
655 activity was considered according to Waldbaum and Thompson (1969), whereas magnetite
15
16
656 activity was approximated at unity (Burkhard 1991). Since this equation required as input the
18
19
657 amount of Fe^{2+} in biotite, $\text{Fe}^{2+}/\text{Fe}^{3+}$ ratios of SS, SU and GU biotites were calculated following
20
21
658 Dymek (1983). Because the compositional spectrum of biotite and sanidine in each rock is
23
24
659 slightly variable, we calculated two T - $f\text{O}_2$ pairs for each lithotype (Fig. 8, Table 4), to account
25
26
660 for the possible oxygen fugacity ranges. According to Burkhard (1991), the error on each
27
28
661 calculation was of ± 0.3 log units.
30

31
662 As shown in Fig. 8, oxygen fugacity in SS rocks resulted in a range from -14.1 to -10.7 $\log f\text{O}_2$,
32
33
663 at temperatures comprised between 790 and 1000°C (Table 4). Such values plot between the
35
36
664 FMQ and NNO buffers (+0.2 to +0.7 ΔFMQ , Fig. 8), and are well comparable to the values
37
38
665 chosen as input for the Rhyolite-MELTS model above developed (-9.34 $\log f\text{O}_2$ at 1060°C),
40
41
666 further confirming the validity of the calculated parameters. Rocks from the SU and GU suites
42
43
667 were characterized by values of -11.9 to -10.2 $\log f\text{O}_2$ at T of 920-1050°C (-0.1 to +0.3 ΔFMQ ,
44
45
668 Fig. 8), and -12.9 to -11.8 $\log f\text{O}_2$ at T of 850-920°C (around +0.4 ΔFMQ , Fig. 8), respectively
47
48
669 (Table 4). Syenogranites of the GU suite were characterized by a behavior comparable to that
49
50
670 of SS rocks, whereas SU rocks record slightly more reduced conditions.
52

53 671 54 55 672 **8. EC-AFC processes** 57

58
673 As shown by the thermobarometric and hygrometric models, the two main magmatic suites (SS
59
60
674 and SU) emplaced and crystallized at comparable pressures and temperatures conditions. The
62
63
64
65

675 only differences can be identified when water content and oxygen fugacity are considered,
1
676 being the SU system characterized by lower H₂O content and by more reduced conditions.
3
677 Anyway, such slight divergences are not able to justify the marked geochemical variations
6
678 between SS and SU magmas (see also Casetta et al. 2017). To **account for** the geochemical
8
679 features of the SS and SU batches, we developed several Energy-Constrained Assimilation and
10
680 Fractional Crystallization (EC-AFC) models (Bohrson and Spera 2001; Spera and Bohrson
13
681 2001), based on ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd. The purpose of this calculation was to discriminate
15
682 between the original (i.e. mantle-derived) signature of SS/SU magmas and **the effects** of
18
683 assimilation (and/or contamination) during the emplacement and differentiation of PIC
20
684 magmas. Specifically, the main target of the models were: (i) verifying if the isotopic signature
23
685 of SU rocks was generated an interaction between SS magmas and crust; (ii) quantify the
25
686 eventual (if any) crustal assimilation during the differentiation of the SS/SU magmatic suites;
28
687 (iii) discriminate between assimilation and contamination processes.

688 A wide spectrum of Permo-Triassic crustal rock isotopic signatures from the literature were
32
689 considered in the EC-AFC model as potential assimilants. These are Triassic carbonates (Martin
35
690 and Macdougall 1995; Blendiger et al. 2015), Permian intrusives **of Mt. Croce (central-eastern**
37
691 **Southern Alps; Rottura et al. 1997) and Serie dei Laghi (southwestern Alps; Sinigoi et al. 2016,**
40
692 **and references therein)**, Permian rhyolitic ignimbrites of the Atesina Volcanic District (Barth
42
693 et al. 1993), and the Kinzigite formation (**southwestern Alps; Voshage et al. 1990).**
45
694 Geothermobarometric and hygrometric results (Table 4) were used to to calculate assimilants
47
695 and starting magmas' specific heat [J/(KgK)], heat of crystallization (J/Kg) and heat of fusion
49
696 (J/Kg) according to Spera (2000), Bohrson and Spera (2001) and Spera and Bohrson (2001,
52
697 Table 5). Specific heat [J/(KgK)] and liquidus T (°C) of carbonate assimilant (Table 5) were
54
698 considered accordingly to Dallai et al. (2011) and Eppelbaum et al. (2014).

699 The absence of ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd **variations** with increasing silica content for both SS
59
700 and SU suites and the lack of **overlap** between SS and SU compositional fields suggested that
62
63
64
65

701 the two suites have a different origin. A slight $^{87}\text{Sr}/^{86}\text{Sr}$ increase was however noted for both
1
702 SS and SU samples while approaching the intrusion borders (Table 2). On the other side, the
3
703 $^{143}\text{Nd}/^{144}\text{Nd}$ of both SS and SU rocks were not sensitive to the distance from intrusion edges,
4
6
704 ruling out the occurrence of contamination processes during magma emplacement. Therefore,
8
705 the Nd isotopic enrichment of the SU suite can be considered as a primary feature, directly
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

709 The EC-AFC model from a starting SS composition (Fig. 9) showed that none of the chosen
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

51 A second EC-AFC model was attempted to explain the isotopic variations of SU samples, but
52 their scattered distribution prevented to retrace the assimilation path. Nevertheless, this model
53 was used just to quantify the crustal contribution in the Nd isotopic variations of the SU
54 samples. As a result, the $^{143}\text{Nd}/^{144}\text{Nd}$ range of the SU suite was achieved by a 2-7% assimilation

of the crustal components (carbonates, Atesina Volcanic District rhyolitic ignimbrites, Mt. Croce granodiorites, Serie dei Laghi granites and amphibolitic kinzigites).

On the whole, the Nd isotopic ranges of both SS and SU suites could be explained by low amounts of assimilation, whereas Sr isotopic variability is more likely derived from contamination processes at the intrusion borders during magma emplacement, where fluid mobilization and local metamorphic reactions occurred (Ferry et al. 2002; Gallien et al. 2007).

As reported in Table 2, the Sr isotopic signature of PIC rocks appears to be more sensitive to their distance to the intrusion edges rather than to their degree of differentiation, being thus the $^{143}\text{Nd}/^{144}\text{Nd}$ ratio the best proxy to discriminate the two magmatic suites. Given this, the Sr variations of the SU suite at near constant $^{143}\text{Nd}/^{144}\text{Nd}$ could be also affected by i) slight alteration effects, unavoidable for such a small body and/or ii) the contact halo between PIC and the wall rocks.

9. Discussion

9.1. Emplacement of the Shoshonitic Silica Saturated and Shoshonitic Silica Undersaturated magmas

The SS and SU intrusive bodies formed as result of fractional crystallization of two distinct trachybasaltic melts, which differentiated along two independent trends over a short time span.

The result of this process was the generation of a large spectrum of intrusive rocks whose cumulus+intercumulus assemblages progressively vary in composition, reaching the most evolved nature in the syenitic rocks. The textural/compositional homogeneity of the main cumulus phases of PIC rocks, together with the scarce efficiency of syn- to post-crystallization diffusion processes on such large-sized crystals, led us to hypothesize that their composition is directly function of the physical conditions at which they formed and segregated from the melt.

In this scenario, clinopyroxene and plagioclase were considered as “snapshots” of the fractional

751 crystallization processes, able to provide informations on the *T-P* conditions of the melts from
1
752 which they generated.

3
4
753 If we compare the clinopyroxene and plagioclase compositions to that of their host rock, in fact,
5
6
754 the disequilibrium is evident (Figg. 4 and 5). Clinopyroxene of SS gabbros (Mg# 55), for
7
8
9
755 example, are characterized by Mg# of 65-71, ideally in equilibrium with melt with Mg# < 43,
10
11
12
756 quite far from the bulk rock composition of the gabbro. The most magnesian ones (Mg#_{Cpx} 76-
13
14
757 78), in pyroxenites, are theoretically in equilibrium with a trachybasaltic melt (Mg# 51), further
15
16
17
758 confirming that the mafic rocks of the complex are cumulates formed during the initial stages
18
19
20
759 of fractionation of the trachybasaltic melt. On the other hand, the less magnesian clinopyroxene
21
22
760 crystals tracked in syenites were almost in equilibrium with their host rocks, corroborating the
23
24
25
761 assumption that the more differentiated rocks of the complex resemble the composition of melts
26
27
762 and approach the eutectic of the system. On the basis of the results obtained by the proposed
28
29
763 “equilibration model”, in turn supporting the differentiation model of Casetta et al. (2017), PIC
30
31
764 rocks were formed during progressive stages of fractional crystallization from starting
32
33
765 trachybasaltic melts.

34
35
36
766 Once evaluated the disequilibrium conditions and reconduted the clinopyroxene/plagioclase
37
38
39
767 compositions to their ideal equilibrium conditions, thermobarometric and hygrometric
40
41
42
768 calculations enabled us to retrieve the *T-P-H₂O* parameters of the less evolved melts in the
43
44
769 feeding system. The “classical” thermobarometric approach for intrusive rocks (amphibole-
45
46
770 plagioclase and biotite equations) was then used to verify the previous results. These
47
48
771 thermobarometers and hygrometers provided *T-P-H₂O* informations for the later crystallization
49
50
772 stages.

51
52
53
773 According to our estimates, some differences between the emplacement conditions of SS and
54
55
56
774 SU magmas can be highlighted. Oxygen fugacities of the PIC plumbing system are between -
57
58
775 0.1 and +0.7 Δ FMQ (Fig. 8), confirming the oxidation conditions proposed by Bonadiman et
59
60
776 al. (1994) for Mt. Monzoni intrusion at comparable temperature intervals, further highlighting

777 the low to moderate oxidizing components that characterized the magmatic systems of the
1 Dolomitic Area during Triassic.

778
3
4
779 On the basis of clinopyroxene-melt thermometric and hygrometric calculations, SS magmas
6
780 result slightly water-enriched ($H_2O = 2.0-2.5$ wt%) with respect to SU ones ($H_2O = 1.0- 1.5$
8
781 wt%), during the first crystallization stages ($T \sim 1000-1100^\circ C$, Fig. 7, Table 4). With increasing
10
11
1782 differentiation, the water content of PIC rocks increased up to 5-8 wt%, value compatible with
13
14
783 the amount of fractionation (79-94%) proposed by Casetta et al. (2017), by taking into account
15
16
1784 the role of amphibole and biotite during differentiation.

18
19
785 The $T-P$ estimates (Fig. 7, Table 4) indicated that both magma batches cooled from a starting T
20
21
786 of $\sim 1000-1100^\circ C$ (cumulus assemblage) down to $\sim 600^\circ C$ (intercumulus assemblage), sharing
22
23
24
787 a common thermal regime. The pressure obtained by applying various mineral-mineral and
25
26
788 mineral-melt equilibria were subsequently used to estimate the emplacement depth of the PIC
27
28
2789 plumbing system, and all the pressures were filtered to extrapolate the best-fit from all models,
30
31
790 taking into account the errors of the different methods (Table 4). The resulting pressure values
32
33
791 were between 0.4 and 1.7 kbar for both SS and SU intrusive bodies, yielding a depth of about
35
36
792 1.4-5.6 km considering a $\Delta P/\Delta z$ of 0.29 kbar/km. With respect to the depth (10-17 km) proposed
37
38
793 by Menegazzo Vitturi et al. (1995), our estimates suggested that PIC represents an intrusion in
40
41
794 the shallow crust, with SS and SU magma batches emplaced within the same thermal regime.

43
795 Despite the common geothermobarometric evolution and slightly different water contents, SS
45
46
796 and SU bodies can be distinguished in terms of $^{87}Sr/^{86}Sr$ and $^{143}Nd/^{144}Nd$, reinforcing the
47
48
797 preliminary PIC model that, on the basis of mineral and bulk rock elemental geochemistry
49
50
51
798 (Casetta et al. 2017) suggested that SS and SU series represent two independent magma pulses.
52
53
799 EC-AFC models (Bohrson and Spera 2001; Spera and Bohrson 2001) show that assimilation of
54
55
800 various types of Permo-Triassic crustal components by a starting SS magma could not account
57
58
801 for the higher $^{143}Nd/^{144}Nd$ of the SU batch, ruling out any possible derivation of the SU suite
59
60
802 from the SS one. Therefore, the Nd isotopic enrichment of the SU suite could be interpreted as

803 an original mantle signature or, alternatively, could be ascribed to assimilation processes
1
804 occurring within the crust. We prefer the first hypothesis, since the second one would imply the
3
4
805 existence of a common mantle-segregated magma with a $^{143}\text{Nd}/^{144}\text{Nd}$ ratio high enough to be
6
806 able to “split” in the SS and SU trends by varying the nature of the assimilant, and such a magma
8
9
807 has never been documented in the Dolomitic Area.

11 The same EC-AFC models demonstrated that both suites assimilated a **small** proportion of
13
14
809 crustal components (5-6% on average, Fig. 9). That amount of assimilation is, however, able to
15
16
810 explain the $^{143}\text{Nd}/^{144}\text{Nd}$ variation throughout SS/SU samples, being that the $^{87}\text{Sr}/^{86}\text{Sr}$ was
18
19
811 affected by late contamination that occurred at the intrusion edges. The fit between PIC isotopic
20
21
812 data and the crustal end-members of the Triassic carbonates, the Permian magmatic rocks and
23
24
813 the metamorphic basement of the Ivrea Zone is consistent with a geochemical influence on the
25
26
814 PIC of a basement that could be attributed to the Ivrea Zone. This hypothesis was also suggested
27
28
815 by Barth et al. (1993) to explain the Atesina Volcanic District-Cima d’Asta magma genesis.

31 In summary, both SS and SU batches **are characterized by a strong contribution of an enriched**
32
33
817 **mantle component, even if the lack of primitive products cannot directly quantify its role in the**
35
36
818 **original source. Anyway, if examined in the light of their temporal relationships, the major,**
37
38
819 **trace element and Sr-Nd isotopic signature of the SS and SU bodies enable to define the**
40
41
820 **progressive evolution of the PIC magmatic suites. In turn, this geochemical evolution could be**
42
43
821 **linked to a time-related (slight) variation of the mantle source, even if an accurate dating of the**
45
46
822 **two single bodies would help in better deciphering the time scale of such process. On the other**
47
48
823 **side, the magmatic evolution in the shallow crust for both the SS and SU batches was confined**
49
50
824 **in the same thermal regime.**

9.2. Origin of the Granitic Unit

58 The GU rocks (granites and syenogranites) differ from the other PIC magmatic suites (SS and
59
60
828 SU), mainly in terms of silica saturation, FeO, MgO, K₂O and Rb contents (Casetta et al. 2017).

829 Moreover the estimated volume of the GU portion of the complex (1.1 km³) is unrealistically
1
830 larger than that of the SS/SU syenites (≤ 0.25 km³) if we consider the already high (> 90%)
3
831 fractionation degrees from which these latter generated. **This means that, if the GU portion**
6
832 **was the result of ~95% fractionation of a SS/SU magma, ~20 km³ of SS/SU rocks should be**
8
833 **intruded somewhere. This fact is in stark contrast to what is observable on field.**

11
834 The GU magmas evolved at redox states comparable to SS/SU ones (about +0.4 Δ FMQ) and
13
835 record temperatures down to 800-850°C (Fig. 8). Put this all together, it is reasonable to
15
836 hypothesize that **the GU granites/syenogranites were generated by magmatic differentiation**
18
837 **from a calc-alkaline parental melt, the primitive/intermediate products of which are not exposed**
20
838 **in the PIC area.** It should be emphasized that GU rocks are similar to the Middle Triassic calc-
23
839 alkaline/**high-K calc-alkaline rhyolites** found in Carnia (**Tarvisio area, Julian Alps**; Gianolla
25
840 1992) and Alto Vicentino regions (**Recoaro-Schio area, Southern Alps**; De Vecchi et al. 1974;
27
841 Barbieri et al. 1982). In these latter, the entire **high-K calc-alkaline trend** is documented, further
30
842 highlighting the existence of a similar differentiation suite during the Triassic magmatic event.
32
843 In such a context, GU rocks could represent the corresponding intrusives of the rhyolites of
35
844 these areas, even if further studies are needed to better investigate this comparison.
37
38
39
40

846 9.3 Solidification time of PIC

42
847 According to Bonadiman et al. (1994) the solidification time of Mt. Monzoni intrusion,
45
848 emplaced at *P-T-z* conditions similar to those of PIC, was about 300 ka. The thermobarometric
47
849 results obtained for PIC magmas and their subsequent application to the Spera (2000), Bohrsen
49
50
850 and Spera (2001) and Spera and Bohrsen (2001) equations, enabled to model the cooling time
52
851 of PIC, expected to be slightly higher than that of Mt. Monzoni, due to its larger volume (4.5
54
852 km³ vs. 1.0-1.5 km³).

57
853 In order to make an accurate comparison with Bonadiman et al. (1994), Spera (1980) equation
59
60
854 was applied to PIC by assuming the thermal estimates already used for the EC-AFC calculations
62
63
64
65

855 and/or derived from Ryolite-MELTS modelling (see Table 5). **By taking into account the whole**
1
856 **volume of PIC, the solidification time resulted about 700 ka. Even if this time value was**
3
4
857 **obtained without discriminating the single volumes of the SS, SU and GU bodies, it was**
6
858 **considered reliable taking into account the similar thermal regimes calculated for the SS, SU**
8
9
859 **and GU suites and the limited re-heating effects provided by the subsequent magma pulses.**
10

10. Conclusions

11
1860 A coupled thermobarometric and isotopic study of the main magmatic suites of the Predazzo
13
14
861 Intrusive Complex enabled us to set out some remarkable points on the emplacement conditions
15
16
1862 of the Middle Triassic magmas in the Dolomitic Area. They can be summarized as:
18
19
863
20
21
864
22

23
24
865 1. Predazzo Intrusive Complex emplacement occurred at a depth of 1.4-5.6 km (0.40-1.65 kbar),
25
26
866 consistent with the field relationships suggesting consanguinity with the overlying effusive
27
28
867 rocks. Its solidification time was about 700 ka, slightly higher but comparable to that proposed
30
31
868 by Bonadiman et al. (1994) for the near Mt. Monzoni intrusion.
32

33
869 2. Despite similar *T-P* of magma emplacement of the three unites, H₂O and *f*O₂ data indicated
35
36
870 that SU primary magmas were characterized by lower water contents and oxidizing conditions
37
38
871 than the SS ones (1.0-1.5 vs. 2.0-2.5 H₂O wt%; -0.1/+0.33 vs. +0.2/+0.7 ΔFMQ, respectively).
40
41
872 Plagioclase-melt, amphibole and K-feldspar-melt hygrometers indicated highly hydrated
42
43
873 conditions of magmas and progressive H₂O enrichment during differentiation.
44

45
874 3. The ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd of PIC magmas plot in the enriched mantle source field. This
47
48
875 feature speaks in favour of the presence of a subduction-signature in the mantle beneath
49
50
876 Southern Alps during Triassic, as already proposed by Bonadiman et al. (1994) and Zanetti et
52
53
877 al. (2013). **Even if the absence of primary products in the Dolomitic Area prevents to exactly**
54
55
878 **quantify the role of the enriched mantle component in the magma genesis, its contribution was**
57
879 **undoubtedly high.**
59

- 880 4. The EC-AFC models cannot account for the genesis of SU rocks via crustal assimilation
1
881 starting from a SS magma, thus indicating two distinct geochemical signatures for SS and SU
3
882 magmas. In turn, this difference could be ascribed to a time-related change in the mantle source,
6
883 slightly moving towards a more $^{143}\text{Nd}/^{144}\text{Nd}$ enriched component.
8
884 5. A low degree of crustal assimilation (5-6%) was able to explain the Nd isotopic variability
10
885 of both SS and SU magmas, whereas the Sr isotopic variations were more likely the results of
13
886 contamination processes at PIC edges. Fractional crystallization was the major (and almost
15
887 only) process acting during the differentiation of SS and SU magmas, confirming what
18
888 hypothesized by Petersen et al. (1980) and Casetta et al. (2017). The signature of the Triassic
20
889 magmas is consistent with an Ivrea-like basement beneath the Dolomitic Area.
23
890 6. Despite petrography, whole-rock and mineral chemistry suggested an I-type affinity for GU
25
891 granites/syenogranites, the wide range and the uncertainty on their isotopic signature leaves the
28
892 question on the origin of GU suite still unsolved. However, the geochemical and petrographic
30
893 similarities with the coeval calc-alkaline/high-K calc-alkaline rhyolites found in Carnia
32
894 (Gianolla 1992) and Alto Vicentino (De Vecchi et al. 1974; Barbieri et al. 1982; Bellieni et al.
35
895 2010) suggest the presence of a high-K calc-alkaline series also in the Dolomitic Area.
37
38
39
40

897 **Acknowledgments**

42

898 We are grateful to Andrea Marzoli and Matteo Masotta for their careful and constructive
45
899 reviews that significantly improved the quality of the paper, and we would also acknowledge
47
900 Timothy L. Grove for his editorial guidance. The IUSS Mobility Research Programme of the
49
50
901 University of Ferrara, grant n. 570 for Long Period, 2016 (FC) and The Italian National
52
902 Research Program PRIN_2015/prot. 20158A9 (CB) supported this research. Anne Kelly and
54
55
903 Vincent Gallagher are also thanked for making the Sr and Nd isotopic analyses at SUERC.
57
58
904
59
60

905 **References**

62
63
64
65

- 906 Anderson JL (1996) Status of thermobarometry in granitic batholiths. *Geol Soc Am Spec Pap*
1
907 315: 125-138. <https://doi.org/10.1017/S0263593300006544>
3
- 908 Anderson JL, Smith DR (1995) The effects of temperature and fO_2 on the Al-in-hornblende
6
909 barometer. *Am Min* 80(5-6): 549-559.
8
- 910 Barbieri G, De Vecchi GP, De Zanche V, Mietto P, Sedeà R (1982) Stratigrafia e petrologia del
10
911 magmatismo triassico nell'area di Recoaro. Guida alla geologia del Sudalpino centro-
13
912 orientale. *Soc Geol It Guide Geologiche Regionali*: 179-187.
15
- 913 Barth S, Oberli F, Meier M, Blattner P, Bargossi GM, Di Battistini G (1993) The evolution of
18
914 a calc-alkaline basic to silicic magma system: geochemical and Rb–Sr, Sm–Nd, and $^{18}O/^{16}O$
20
915 isotopic evidence from the Late Hercynian Atesina-Cima d'Asta volcanoplutonic complex,
23
916 northern Italy. *Geochim Cosmochim Acta* 57: 4285-4300. <https://doi.org/10.1016/0016->
25
917 7037(93)90323-O
27
- 918 Bellieni G, Fioretti AM, Marzoli A, Visonà D (2010) Permo–Paleogene magmatism in the
30
919 eastern Alps. *Rend Lincei* 21: S51-S71. <https://doi.org/10.1007/s12210-010-0095-z>
32
- 920 Bernoulli D, Lemoine M (1980) Birth and Early Evolution of the Tethys: the Overall Situation.
35
921 *Mem Bur Rech Géol Minières* 115: 168-179.
37
- 922 Blendinger W, Lohmeier S, Bertini A, Meißner E, Sattler CD (2015) A new model for the
40
923 formation of Dolomite in the Triassic dolomites, Northern Italy. *J Pet Geo* 38(1): 5-36.
42
924 10.1111/jpg.12596
44
- 925 Bohron WA, Spera FJ (2001) Energy-constrained open-system magmatic processes II:
47
926 application of energy-constrained assimilation–fractional crystallization (EC-AFC) model
49
50
927 to magmatic systems. *J Petrol* 42(5): 1019-1041.
52
928 <https://doi.org/10.1093/petrology/42.5.1019>
54
- 929 Bonadiman C, Coltorti M, Siena F (1994) Petrogenesis and T- fO_2 estimates of Mt. Monzoni
57
930 complex (Central Dolomites, Southern Alps): a Triassic shoshonitic intrusion in a trascurrent
59
931 geodynamic setting. *Eur J Mineral* 6: 943-966. 10.1127/ejm/6/6/0943
62
63
64
65

- 932 Borsi S, Ferrara G (1968) Isotopic age measurements of the M. Monzoni intrusive complex.
1
933 Miner Petrogr Acta 14: 171-183.
3
4
934 Brack P, Mundil R, Oberli F, Meier M, Rieber H (1996) Biostratigraphic and radiometric age
6
935 data question the Milankovitch characteristics of the Latemar cycles (Southern Alps, Italy).
8
936 Geology 24(4): 371-375. [https://doi.org/10.1130/0091-
12
13 7613\(1996\)024<0371:BARADQ>2.3.CO;2](https://doi.org/10.1130/0091-
10
11 7613(1996)024<0371:BARADQ>2.3.CO;2)
14
938 Brack P, Mundil R, Oberli F, Meier M, Rieber H (1997) Biostratigraphic and radiometric age
15
16 data question the Milankovitch characteristics of the Latemar cycles (Southern Alps, Italy).
17
18 Reply: Geology 25(5): 471-472.
19
20
21
22 941 Brack P, Rieber H, Nicora A, Mundil R (2005) The Global boundary Stratotype Section and
23
24 Point (GSSP) of the Ladinian Stage (Middle Triassic) at Bagolino (Southern Alps, Northern
25
26 Italy) and its implications for the Triassic time scale. Episodes 28 (4): 233-244.
27
28
29 944 **Brady JB, McCallister RH (1983) Diffusion data for clinopyroxenes from homogenization and
30
31 self-diffusion experiments. Am Min 68: 95-105**
32
33
34 946 Burkhard DJM (1991) Temperature and redox path of biotite-bearing intrusives: a method of
35
36 estimation applied to S- and I-type granites from Australia. Earth Planet Sci Lett 104: 89-
37
38 98. [https://doi.org/10.1016/0012-821X\(91\)90240-I](https://doi.org/10.1016/0012-821X(91)90240-I)
39
40
41 949 Burnham DJM, Holloway JR, Davis NF (1969) Thermodynamic properties of water to 1000°C
42
43 and 10,000 bars. Geol Soc Am Spec Pap 132: 96.
44
45
46 951 Casetta F, Coltorti M, Marrocchino E (2017) Petrological evolution of the Middle Triassic
47
48 Predazzo Intrusive Complex, Italian Alps. Int Geol Rev 60(8): 977-997.
49
50
51 953 <https://doi.org/10.1080/00206814.2017.1363676>
52
53
54 954 Cassinis G, Cortesogno L, Gaggero L, Perotti CR, Buzzi L (2008) Permian to Triassic
55
56 geodynamic and magmatic evolution of the Brescian Prealps (eastern Lombardy, Italy). B
57
58 Soc Geol Ital 127(3): 501-518.
59
60
61
62
63
64
65

- 957 Castellarin A, Lucchini F, Rossi PL, Sartori L, Simboli G, Somnavilla E (1982) Note
1
958 geologiche sulle intrusioni di Predazzo e dei M. Monzoni. Guida alla geologia del Sudalpino
3
4
959 centro-orientale: Guide geologiche regionali SGI: 211-219.
5
6
960 Castellarin A, Lucchini F, Rossi PL, Simboli G, Bosellini A, Somnavilla E (1980) Middle
8
9
961 Triassic magmatism in Southern Alps II: A geodynamic model. Riv Ital Paleontol S 85(3-
10
11
962 4): 1111-1124.
13
14
963 Chappell BW, White AJR (2001) Two contrasting granite types: 25 years later. Aust J Earth
15
16
964 Sci 48(4): 489-499. 10.1046/j.1440-0952.2001.00882.x
18
19
965 **Cherniak DJ, Liang Y (2012) Ti diffusion in natural pyroxene. *Geochim Cosmochim Acta* 98:**
20
21
966 **31-47. <https://doi.org/10.1016/j.gca.2012.09.021>**
22
23
967 Coltorti M, Siena F, Visonà D (1996) Aspetti petrologici del magmatismo Triassico dell'area
25
26
968 di Predazzo. 78° Riunione estiva Soc. Geol. It: Geologia delle Dolomiti. San Cassiano
27
28
969 (1996), Conference Abstract.
30
31
970 Dallai L, Cioni R, Boschi C, D'Oriano C (2011) Carbonate-derived CO₂ purging magma at
32
33
971 depth: influence on the eruptive activity of Somma-Vesuvius, Italy. Earth Planet Sci Lett
35
36
972 310(1): 84-95. <https://doi.org/10.1016/j.epsl.2011.07.013>
37
38
973 Dal Piaz G, Bistacchi A, Gianotti F et al. (2015) Carta Geologica d'Italia-Foglio 070 Monte
40
41
974 Cervino. Carta Geologica d'Italia, 1:50.000 scale: 70.
42
43
975 De Vecchi G, De Zanche V, Sedeo R (1974) Osservazioni preliminari sulle manifestazioni
44
45
976 magmatiche triassiche nelle Prealpi Vicentine (area di Recoaro-Schio-Posina). B Soc Geol
47
48
977 Ital 93(2): 397-409.
49
50
978 Della Lucia A (1997) Aspetti vulcanologici e petrologici del complesso vulcano-plutonico
52
53
979 Triassico di Cima Pape (Pale di San Lucano, BL). Unpublished master thesis, University of
54
55
980 Ferrara.
57
58
59
60
61
62
63
64
65

- 981 Dimanov A, Jaoul O, Sautter V. (1996) Calcium self-diffusion in natural diopside single
1
982 crystals. *Geochim Cosmochim Acta* 60(21): 4095-4106. [https://doi.org/10.1016/S0016-
4 7037\(96\)00250-5](https://doi.org/10.1016/S0016-
3 7037(96)00250-5)
5
6
984 Di Matteo V, Carroll MR, Behrens H, Vetere F, Brooker RA (2004) Water solubility in
8
985 trachytic melts. *Chem Geol* 213(1-3): 187-196.
10
11 <https://doi.org/10.1016/j.chemgeo.2004.08.042>
13
14
987 Doglioni C (1987) Tectonics of the Dolomites (Southern Alps, Northern Italy). *J Struct Geol* 9:
15
16 181-193. [https://doi.org/10.1016/0191-8141\(87\)90024-1](https://doi.org/10.1016/0191-8141(87)90024-1)
18
19
989 Doglioni C (2007) Tectonics of the Dolomites. *Bull Angew Geol* 12(2): 11-15.
20
21
990 Dymek FR (1983) Titanium, aluminium and interlayer cation substitution in biotite from high-
22
23 grade gneisses, West Greenland. *Am Mineral* 68: 880-899.
24
25
992 Eppelbaum L, Kutasov I, Pilchin A (2014) Thermal properties of rocks and density of fluids.
26
27 In: *Applied geothermics*. Springer Berlin Heidelberg, pp. 99-149.
28
29
993 Ferry JM, Wing BA, Penniston-Dorland SC, Rumble D (2002) The direction of fluid flow
30
31 during contact metamorphism of siliceous carbonate rocks: new data for the Monzoni and
32
33 Predazzo aureoles, northern Italy, and a global review. *Contrib Mineral Petrol* 142(6): 679-
34
35 699. <https://doi.org/10.1007/s00410-001-0316-7>
36
37
996 Gallien F, Abart R, Wyhlidal S (2007) Contact metamorphism and selective metasomatism of
38
39 the layered Bellerophon Formation in the eastern Monzoni contact aureole, northern Italy.
40
41 *Mineral Petrol* 91(1): 25-53. <https://doi.org/10.1007/s00710-007-0184-6>
42
43
1000 Gasparotto G, Simboli G (1991) Mineralogia, petrografia e schemi evolutivi delle magmatiti
44
45 triassiche del complesso di Cima Pape (Dolomiti Orientali). *Mineral Petrogr Acta* 34: 205-
46
47 234.
48
49
1001 Giacomoni PP, Ferlito C, Coltorti M, Bonadiman C, Lanzafame G (2014) Plagioclase as archive
50
51 of magma ascent dynamics on “open conduit” volcanoes: The 2001–2006 eruptive period at
52
53 Mt. Etna. *Earth Sci Rev* 138: 371-393. <https://doi.org/10.1016/j.earscirev.2014.06.009>
54
55
1002
56
57
1003
58
59
1004
60
61
1005
62
63
1006
64
65

- 1007 Gianolla P (1992) Evoluzione mediotriassica del vulcanismo di Rio Freddo (Api Giulie, Italia).
1
1008 Mem Sci Geol 44: 193-209.
3
4
1009 Gianolla P, Avanzini M, Breda A et al. (2010) Dolomites, 7th International Triassic Field
6
1010 Workshop, Field trip to the World Heritage Site of the Tethyan Triassic. September 5-10
8
1011 2010, Dolomites, Southern Alps, Italy. With the adesion of Fondazione Dolomiti-
10
1012 Dolomiten-Dolomites-Dolomitis Unesco.
13
14
1013 **Grove TL, Baker MB, Kinzler RJ (1984) Coupled CaAl-NaSi diffusion in plagioclase feldspar:**
15
1014 **Experiments and applications to cooling rate speedometry. Geochim Cosmochim Acta 48:**
18
1015 **2113-2121.**
20
21
1016 Gualda GAR, Ghiorso MS, Lemons RV, Carley TL (2012) Rhyolite-MELTS: a Modified
22
1017 Calibration of MELTS Optimized for Silica-rich, Fluid-bearing Magmatic Systems. J Petrol
25
1018 53: 875-890. <https://doi.org/10.1093/petrology/egr080>
26
27
28
1019 Henry DJ, Guidotti CV, Thomson JA (2005) The Ti-saturation surface for low-to-medium
30
1020 pressure metapelitic biotites: Implications for geothermometry and Ti-substitution
31
1021 mechanisms. Am Mineral 90(2-3): 316-328. <https://doi.org/10.2138/am.2005.1498>
32
33
1022 Holland T, Blundy J (1994) Non-ideal interactions in calcic amphiboles and their bearing on
36
1023 amphibole-plagioclase thermometry. Contrib Mineral Petrol 116(4): 433-447.
37
1024 <https://doi.org/10.1007/BF00310910>
42
43
1025 Huebner JS, Sato M (1970) The oxygen fugacity-temperature relationships of manganese oxide
44
1026 and nickel oxide buffers. Am Mineral 55: 934-952.
45
46
47
1027 Ickert RB (2013) Algorithms for estimating uncertainties in initial radiogenic isotope ratios and
48
1028 model ages. Chem Geol 340: 131-138. <https://doi.org/10.1016/j.chemgeo.2013.01.001>
49
50
1029 **Kohn SC (2000) The dissolution mechanisms of water in silicate melts; a synthesis of recent**
53
1030 **data. Mineral Mag 64(3): 389-408. <https://doi.org/10.1180/002646100549463>**
54
55
1031 Kress VC, Carmichael IS (1988) Stoichiometry of the iron oxidation reactions in silicate melts.
58
1032 Am Mineral 73: 1267-1274.
59
60
61
62
63
64
65

1033 Kress VC, Carmichael IS (1991) The compressibility of silicate liquids containing Fe₂O₃ and
1 the effect of composition, temperature, oxygen fugacity and pressure on their redox states.
1034 Contrib Mineral Petrol 108(1-2): 82-92. 10.1007/BF00307328
1035
1036 Lange RA, Frey HM, Hektor J (2009) A thermodynamic model for the plagioclase-liquid
1037 hygrometer/thermometer. Am Mineral 94(4): 494-506.
1038 <https://doi.org/10.2138/am.2009.3011>
1039
1040 Laurenzi MA, Visonà D, (1996) ⁴⁰Ar/³⁹Ar Chronology of Predazzo magmatic complex
1041 (Southern Alps, Italy). 78° Riunione estiva Soc. Geol. It: Geologia delle Dolomiti. San
1042 Cassiano (1996): 16-18.
1043
1044 Le Maitre RW (2002) A classification of igneous rocks and glossary of terms. Cambridge
1045 University Press.
1046
1047 Leake BE, Arps CES, Birch WD (1997) Nomenclature of amphiboles: Report of the
1048 Subcommittee on Amphiboles of the International Mineralogical Association, Commission
1049 on New Minerals and Mineral Names. Am Mineral 82: 1019-1037.
1050 <https://doi.org/10.1180/minmag.1997.061.405.13>
1051
1052 Lippolt H, Pidgeon R (1974) Isotopic mineral ages of a diorite from the Eisenkappel intrusion,
1053 Austria. Zeitschrift für Naturforschung 29a.
1054
1055 Liu Y, Zhang Y, Behrens H (2005) Solubility of H₂O in rhyolitic melts at low pressures and a
1056 new empirical model for mixed H₂O-CO₂ solubility in rhyolitic melts. J Volcanol Geotherm
1057 Res 143(1-3): 219-235. <https://doi.org/10.1016/j.jvolgeores.2004.09.019>
1058
1059 Lucchini F, Rossi PL, Simboli G (1982) Il magmatismo triassico dell'area di Predazzo (Alpi
1060 Meridionali, Italia). In Castellarin A, Vai GB (eds) Guida alla Geologia del Sudalpino
1061 centro-orientale: Guide Geologiche Regionali Società Geologica Italiana: pp. 221-230.
1062
1063 Lugmair GW, Marti K (1978) Lunar initial ¹⁴³Nd/¹⁴⁴Nd: differential evolution of Lunar crust
1064 and mantle. Earth Planet Sci Lett 39: 349-357. [https://doi.org/10.1016/0012-821X\(78\)90021-3](https://doi.org/10.1016/0012-821X(78)90021-3)

- 1059 Manzotti P, Ballèvre M, Dal Piaz GV (2017) Continental gabbros in the Dent Blanche Tectonic
1 System (Western Alps): from the pre-Alpine crustal structure of the Adriatic palaeo-margin
1060 2 to the geometry of an alleged subduction interface. *J Geol Soc* 174(3): 541-556.
3
4
1061 5
6
1062 7 Marocchi M, Morelli C, Mair V, Klötzli U, Bargossi GM (2008) Evolution of large silicic
8 magma systems: new U–Pb zircon data on the NW Permian Athesian Volcanic Group
1063 9 (Southern Alps, Italy). *J Geol* 116: 480-498.
10
11
1064 12
13
1065 14 Marrocchino E, Coltorti M, Visonà D, Thirwall, MF (2002) Petrology of Predazzo magmatic
15 complex (Trento, Italy). *Geochim Cosmochim Acta* 66(15°): A486-A486.
16
1066 17
18
1067 19 Martin EE, Macdougall JD (1995) Sr and Nd isotopes at the Permian/Triassic boundary: A record
20 of climate change. *Chem Geol* 125(1-2): 73-99. [https://doi.org/10.1016/0009-](https://doi.org/10.1016/0009-2541(95)00081-V)
21
1068 22
23
1069 24
25
1070 26 Marzocchi C (1987) Petrologia e metallogenesi del complesso eruttivo di Predazzo (TN): con
27 particolare riguardo al settore meridionale. University of Ferrara, unpublished master thesis.
28
1071 29
30
1072 31 Masotta M, Mollo S, Freda C, Gaeta M, Moore G (2013) Clinopyroxene–liquid thermometers
32 and barometers specific to alkaline differentiated magmas. *Contrib Mineral Petrol* 166(6):
33
1073 34
35
1074 36
37
1075 38 Menegazzo Vitturi L, Visonà D, Zantedeschi C (1995) Amphibole composition in rocks from
39 Predazzo volcano-plutonic complex (Southern Alps, Italy). *Mem Sci Geol* 47: 87-94.
40
1076 41
42
1077 43 Mietto P, Manfrin S, Preto N et al. (2012) The Global Boundary Stratotype Section and Point
44 (GSSP) of the Carnian Stage (Late Triassic) at Prati di Stuores/ Stuores Wiesen Section
45
1078 46
47
1079 48
49
50
1080 51 Mollo S, Masotta M, Forni F, Bachmann O, De Astis G, Moore G, Scarlato P (2015) A K-
52
1081 53
54
1082 55
56
57
58
59
60
61
62
63
64
65

- 1083 Mollo S, Putirka K, Iezzi G, Del Gaudio P, Scarlato P (2011) Plagioclase–melt (dis) equilibrium
1
1084 due to cooling dynamics: implications for thermometry, barometry and hygrometry. *Lithos*
2
3
4
1085 125(1): 221-235. <https://doi.org/10.1016/j.lithos.2011.02.008>
5
6
1086 Monjoie P, Bussy F, Schaltegger U, Mulch A, Lapierre H, Pfeifer HR (2007) Contrasting
7
8
9
1087 magma types and timing of intrusion in the Permian layered mafic complex of Mont Collon
10
11
1088 (Western Alps, Valais, Switzerland): evidence from U/Pb zircon and $^{40}\text{Ar}/^{39}\text{Ar}$ amphibole
12
13
14
1089 dating. *Swiss J Geosci* 100(1): 125-135. <https://doi.org/10.1007/s00015-007-1210-8>
15
16
1090 **Morimoto N (1988) Nomenclature of pyroxenes. *Mineral Petrol* 39(1): 55-76.**
17
18
1091 <https://doi.org/10.1007/BF01226262>
19
20
1092 **Morse SA, Brady JB (2017) Thermal History of the Upper Zone of the Kiglapait Intrusion. *J*
21
22
23
1093 *Petrol* 58(7): 1319-1332. 10.1093/petrology/egx055**
24
25
1094 **Müller T, Dohmen R, Becker HW, Ter Heege JH, Chakraborty S (2013) Fe-Mg interdiffusion
26
27
28
1095 rates in clinopyroxene: experimental data and implications for Fe–Mg exchange
29
30
31
1096 geothermometers. *Contrib Mineral Petrol* 166(6): 1563-1576. 10.1007/s00410-013-0941-y**
32
33
1097 Mundil R, Brack P, Laurenzi MA (1996) High resolution U/Pb single zircon age
34
35
36
1098 determinations: new constraints on the timing of Middle Triassic magmatism in the Southern
37
38
1099 Alps. 78° Riunione estiva Soc. Geol. It: Geologia delle Dolomiti. San Cassiano (1996),
39
40
1100 Conference Abstract.
41
42
1101 Myers JT, Eugster HP (1983) The system Fe-Si-O: Oxygen buffer calibrations to 1500 K.
43
44
45
1102 *Contrib Mineral Petrol* 82(1): 75-90. <https://doi.org/10.1007/BF00371177>
46
47
1103 **Namur O, Humphreys MCS (2018) Trace Element Constraints on the Differentiation and
48
49
50
1104 Crystal Mush Solidification in the Skaergaard Intrusion, Greenland. *J Petrol* 1-31. doi:
51
52
1105 [10.1093/petrology/egy032](https://doi.org/10.1093/petrology/egy032)**
53
54
1106 **Nielsen TFD (2004) The Shape and Volume of the Skaergaard Intrusion, Greenland:
55
56
57
1107 Implications for Mass Balance and Bulk Composition. *J Petrol* 45(3): 507-530.**
58
59
1108 [10.1093/petrology/egg092](https://doi.org/10.1093/petrology/egg092)
60
61
62
63
64
65

- 1109 Petersen JS, Morten L, Simboli G, Lucchini F (1980) REE abundances in the Predazzo-
1
1110 Monzoni intrusive complex, Dolomites, North Italy. Riv Ital Paleontol S 85: 1065-1080.
2
3
4
1111 Pin C, Sills JD (1986) Petrogenesis of layered gabbros and ultramafic rocks from Val Sesia, the
5
6
1112 Ivrea Zone, NW Italy: trace element and isotope geochemistry. Geol Soc Spec Publ 24: 231-
7
8
1113 249. <https://doi.org/10.1144/GSL.SP.1986.024.01.21>
9
10
1114 Pouchou JL, Pichoir F (1991) Quantitative analysis of homogeneous or stratified microvolumes
11
12
1115 applying the model "PAP". In Electron probe quantitation. Springer US, pp. 31-75.
13
14
1116 https://doi.org/10.1007/978-1-4899-2617-3_4
15
16
1117 Povoden E, Horacek M, Abart R (2002) Contact metamorphism of siliceous dolomite and
17
18
1118 impure limestones from the Werfen formation in the eastern Monzoni contact aureole.
19
20
1119 Mineral Petrol 76(1): 99-120. <https://doi.org/10.1007/s007100200034>
21
22
23
1120 Princivalle F, Della Giusta A, De Min A, Piccirillo EM (1999) Crystal chemistry and
24
25
1121 significance of cation ordering in Mg-Al rich spinels from high-grade hornfels (Predazzo-
26
27
1122 Monzoni, NE Italy). Mineral Mag 63(2): 257-257.
28
29
1123 <https://doi.org/10.1180/002646199548367>
30
31
32
33
1124 Putirka KD (2008) Thermometers and barometers for volcanic systems. Rev Mineral Geochem
34
35
1125 69(1): 61-120. <https://doi.org/10.2138/rmg.2008.69.3>
36
37
38
39
1126 Putirka K, Johnson M, Kinzler R, Longhi J, Walker D (1996) Thermobarometry of mafic
40
41
1127 igneous rocks based on clinopyroxene-liquid equilibria, 0–30 kbar. Contrib Mineral Petrol
42
43
1128 123(1): 92-108. <https://doi.org/10.1007/s004100050145>
44
45
46
47
1129 Putirka K, Mikaelian H, Ryerson F, Shaw H (2003) New clinopyroxene-liquid
48
49
1130 thermobarometers for mafic, evolved, and volatile-bearing lava compositions, with
50
51
1131 applications to lavas from Tibet and the Snake River Plain, Idaho. Am Mineral 88: 1542-
52
53
1132 1554. <https://doi.org/10.2138/am-2003-1017>
54
55
56
57
58
59
60
61
62
63
64
65

- 1133 Quick JE, Sinigoi S, Peressini G, Demarchi G, Wooden JL, Sbisà A (2009) Magmatic plumbing
1 of a large Permian caldera exposed to a depth of 25 km. *Geology* 37: 603-606.
1134 <https://doi.org/10.1130/G30003A.1>
1135
- 1136 Ridolfi F, Renzulli A (2012) Calcic amphiboles in calc-alkaline and alkaline magmas:
8 thermobarometric and chemometric empirical equations valid up to 1,130° C and 2.2 GPa.
1137 *Contrib Mineral Petrol* 163(5): 877-895. <https://doi.org/10.1007/s00410-011-0704-6>
1138
- 1139 Ridolfi F, Renzulli A, Puerini M (2010) Stability and chemical equilibrium of amphibole in
14 calc-alkaline magmas: an overview, new thermobarometric formulations and application to
1140 subduction-related volcanoes. *Contrib Mineral Petrol* 160(1): 45-66.
1141 <https://doi.org/10.1007/s00410-009-0465-7>
1142
- 1143 Rotenberg E, Davis DW, Amelin Y, Ghosh S, Bergquist BA (2012) Determination of the decay-
25 constant of ⁸⁷Rb by laboratory accumulation of ⁸⁷Sr. *Geochim Cosmochim Acta* 85: 41-57.
1144 <https://doi.org/10.1016/j.gca.2012.01.016>
1145
- 1146 Rottura A, Bargossi GM, Caggianelli A, Del Moro A, Visonà D, Tranne CA (1998) Origin and
31 significance of the Permian high-K calc-alkaline magmatism in the central-eastern Southern
1147 Alps, Italy. *Lithos* 45: 329-348. [https://doi.org/10.1016/S0024-4937\(98\)00038-3](https://doi.org/10.1016/S0024-4937(98)00038-3)
1148
- 1149 Rottura A, Del Moro A, Caggianelli A, Bargossi GM, Gasparotto G (1997) Petrogenesis of the
40 Monte Croce granitoids in the context of Permian magmatism in the Southern Alps, Italy.
1150 *Eur J Mineral* 9(6): 1293-1310. [10.1127/ejm/9/6/1293](https://doi.org/10.1127/ejm/9/6/1293)
1151
- 1152 Sarti M, Ardizzoni F (1984) Tettonica Triassica nel gruppo di Cima Pape-Pale di Sanson
47 (Dolomiti Bellunesi). *Mem Sci Geol* 36: 353-370.
1153
- 1154 Schaltegger U, Brack P (2007) Crustal-scale magmatic systems during intracontinental strike-
52 slip tectonics: U, Pb and Hf isotopic constraints from Permian magmatic rocks of the
1155 Southern Alps. *Int J Earth Sci* 96(6): 1131-1151. <https://doi.org/10.1007/s00531-006-0165-8>
1156
- 1157 8
59
60
61
62
63
64
65

- 1158 Schmid SM, Bernoulli D, Fügenschuh B et al. (2008) The Alpine-Carpathian-Dinaridic
1
1159 orogenic system: correlation and evolution of tectonic units. *Swiss J Geosci* 101(1): 139-
2
3
4
1160 183. <https://doi.org/10.1007/s00015-008-1247-3>
5
6
1161 **Silver L, Stolper E (1985) A thermodynamic model for hydrous silicate melts. *J Geol* 93(2):**
7
8
1162 **161-177.**
9
10
1163 Sinigoi S, Quick JE, Demarchi G, Klotzli U (2016) Production of hybrid granitic magma at the
11
12
13
14
1164 advancing front of basaltic underplating: Inferences from the Sesia Magmatic System (south
15
16
17
1165 - western Alps, Italy). *Lithos* 252-253: 109-122. <https://doi.org/10.1016/j.lithos.2016.02.018>
18
19
1166 Sloman LE (1989) Triassic shoshonites from the dolomites, northern Italy: Alkaline arc rocks
20
21
1167 in a strike-slip setting. *J Geophys Res: Solid Earth* 94(B4): 4655-4666.
22
23
1168 10.1029/JB094iB04p04655
24
25
1169 Spera FJ (1980) Thermal evolution of plutons: a parameterized approach. *Science* 207: 299-301.
26
27
1170 10.1126/science.207.4428.299
28
29
1171 Spera FJ (2000) Physical properties of magmas. In Sigurdsson H (ed.) *Encyclopedia of*
30
31
1172 *Volcanoes*. New York: Academic Press, pp. 171-190.
32
33
1173 Spera FJ, Bohron WA (2001) Energy-constrained open-system magmatic processes I: General
34
35
1174 model and energy-constrained assimilation and fractional crystallization (EC-AFC)
36
37
1175 formulation. *J Petrol* 42(5): 999-1018. <https://doi.org/10.1093/petrology/42.5.999>
38
39
40
1176 Stahle V, Frenzel G, Hess JC, Saupé F, Schmidt ST, Schneider W (2001) Permian metabasalt
41
42
1177 and Triassic alkaline dykes in the northern Ivrea zone: clues to the post-Variscan
43
44
1178 geodynamic evolution of the Southern Alps. *Schweiz Mineral Petrogr Mitt* 81(1): 1-21.
45
46
47
48
1179 Stampfli GM, Borel GD (2002) A plate tectonic model for the Paleozoic and Mesozoic
49
50
1180 constrained by dynamic plate boundaries and restored synthetic oceanic isochrones. *Earth*
51
52
1181 *Planet Sci Lett* 196(1): 17-33. [https://doi.org/10.1016/S0012-821X\(01\)00588-X](https://doi.org/10.1016/S0012-821X(01)00588-X)
53
54
55
1182 Stampfli GM, Borel GD (2004) The TRANSMED transects in space and time: constraints on
56
57
1183 the paleotectonic evolution of the Mediterranean domain. In *The TRANSMED Atlas: The*
58
59
60
61
62
63
64
65

1184 Mediterranean region from crust to mantle, pp. 53-80. <https://doi.org/10.1007/978-3-642->
1
1185 18919-7_3
2
3
4
1186 **Stolper E (1982) The speciation of water in silicate melts. *Geochim Cosmochim Acta* 46(12):**
5
6
1187 **2609-2620. [https://doi.org/10.1016/0016-7037\(82\)90381-7](https://doi.org/10.1016/0016-7037(82)90381-7)**
7
8
9
1188 Traill RJ, Lachance GR (1966) A practical solution to the matrix problem in X-ray analysis. II.
10
11 Application to a multicomponent alloy system. *Can Spectrosc* 11(3): 63-71.
12
13
14
1189 Vardabasso S (1929) Rapporti tra attività magmatica e vicende tettoniche nella provincia
15
16 petrografica di Predazzo: *Studi Trentini di Scienze Naturali* 11.
17
18
19
1190 Vardabasso S (1930) Carta geologica del territorio eruttivo di Predazzo e Monzoni. Ufficio
20
21 Idrografico del Magistrato alle Acque di Venezia, 1:25000 scale.
22
23
24
1191 Visonà D (1997) The Predazzo multipulse intrusive body (Western Dolomites, Italy). Field and
25
26 mineralogical studies. *Mem Sci Geol* 49: 117-125.
27
28
29
1192 Visonà D, Zanferrari A (2000) Some constraints on geochemical features in the Triassic mantle
30
31 of the easternmost Austroalpine–Southalpine domain: evidence from the Karawanken pluton
32
33 (Carinthia, Austria). *Int J Earth Sci* 89(1): 40-51. <https://doi.org/10.1007/s005310050316>
34
35
36
1193 Voshage H, Hofmann AW, Mazzucchelli M, Rivalenti G, Sinigoi S, Raczek I, Demarchi G
37
38 (1990) Isotopic evidence from the Ivrea Zone for a hybrid lower crust formed by magmatic
39
40 underplating. *Nature* 347(6295): 731-736. [10.1038/347731a0](https://doi.org/10.1038/347731a0)
41
42
43
1194 Waldbaum DR, Thompson JB (1969) Mixing properties of sanidine crystalline solutions: IV.
44
45 Phase diagrams from equations of state. *Am Mineral* 54: 1274-1298.
46
47
48
1195 Willcock MAW, Bargossi GM, Weinberg RF, Gasparotto G, Cas RAF, Giordano G, Marocchi
49
50 M (2015) A complex magma reservoir system for a large volume intra-to extra-caldera
51
52 ignimbrite: Mineralogical and chemical architecture of the VEI8, Permian Ora ignimbrite
53
54 (Italy). *J Volcanol Geotherm Res* 306: 17-40.
55
56
57
1196 <https://doi.org/10.1016/j.jvolgeores.2015.09.015>
58
59
60
61
62
63
64
65

- 1209 Zanetti A, Mazzucchelli M, Sinigoi S, Giovanardi T, Peressini G, Fanning M (2013) SHRIMP
1
1210 U–Pb Zircon Triassic intrusion age of the Finero mafic complex (Ivrea–Verbano zone,
2
3
4
1211 Western Alps) and its geodynamic implications. *J Petrol* 54(11): 2235-2265.
5
6
1212 <https://doi.org/10.1093/petrology/egt046>
7
8
1213 Ziegler PA, Stampfli GM (2001) Late Palaeozoic-Early Mesozoic plate boundary
9
10
1214 reorganization: collapse of the Variscan orogen and opening of Neotethys. *Nat Brescia* 25:
11
12
1215 17-34.
13
14
1216 Zindler A, Hart SR (1986) Chemical geodynamics. *Annu Rev Earth Planet Sci* 14: 493-571.
15
16
17
18
19
20
21

1218 **Figure Captions**

1219 **Fig. 1 (colour online)**

1220 Simplified geological map of the Predazzo Intrusive Complex (PIC). On the left bottom corner,
22
23
1221 the study area location with respect to the NE sector of Italy is shown
24
25

1223 **Fig. 2 (colour online)**

1224 (a) Total Alkali vs. Silica and (b) K₂O vs. Na₂O diagrams for Predazzo Intrusive Complex rocks
26
27
1225 (Le Maitre 2002). SS = Shoshonitic Silica Saturated; SU = Shoshonitic Silica Undersaturated;
28
29
1226 GU = Granitic Unit
30
31
32
33

1228 **Fig. 3 (colour online)**

1229 Textural sketches of Predazzo Intrusive Complex rocks and corresponding photomicrographs
34
35
1230 showing the crystallization relationships between biotite and amphibole in the intercumulus
36
37
1231 assemblage of (a, b) Shoshonitic Silica Saturated monzodiorites and (c, d) Shoshonitic Silica
38
39
1232 Undersaturated monzogabbros. Each labelled mineral phase in the right photomicrographs is
40
41
1233 contraddistinct by a different colour in the left sketches. Amph = amphibole; Ap = apatite; Bt
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1234 = biotite; Cpx = clinopyroxene; K-Feld = K-feldspar; Ox = Fe-Ti oxide; Plag = plagioclase; Qz
1
1235 = quartz; Ser = sericite
2
3

4
1236
5

6

1237 **Fig. 4 (colour online)**
8

9

1238 Mg# vs. $Kd_{Fe-Mg}^{Cpx-Liq}$ of clinopyroxene from the (a) Shoshonitic Silica Saturated and (b)
10 Shoshonitic Silica Undersaturated rocks. The white symbols represent the disequilibrium
11 between crystals and their host rock composition, whose whole rock Mg# (wr Mg#) is also
1239 reported. The filled symbols represent the attained equilibrium conditions [after the calculation](#)
13 [of melt composition in equilibrium](#). “Liq” indicates both the composition of the crystals host
14 rocks (white symbols, prior to equilibration process) and that of the calculated “melt” (filled
15 symbols, after equilibration process). The length of the dashed arrows is directly function of
16 the amount of disequilibrium between the crystals and their host rock. The equilibrium range
1241 of $Kd_{Fe-Mg}^{Cpx-Liq} = 0.24-0.30$ (black lines) is from Putirka et al. (2003)
17
18
19
20
21
22
23
24
25
26
27
28
29
30

31

1247
32

33

1248 **Fig. 5 (colour online)**
35

36

1249 Anorthite (X_{An}) vs. $Kd_{An-Ab}^{Plag-Liq}$ of plagioclase from the (a) Shoshonitic Silica Saturated and
37 (b) Shoshonitic Silica Undersaturated rocks. As in Fig. 4, white symbols represent the
38 disequilibrium between crystals and their host rock composition (whose Mg# is reported as “wr
39 Mg#”), whereas filled symbols represent the attained equilibrium conditions after the
40 equilibration processes. “Liq” indicates both the composition of the crystals host rocks (white
41 symbols, prior to equilibration process) and that of the calculated “melt” (filled symbols, after
42 equilibration process). The length of the dashed arrows is directly function of the amount of
43 disequilibrium between the crystals and their host rock. The chosen equilibrium range of Kd_{An-Ab}^{Plag-}
44 $Liq = 0.27 \pm 0.11$ (black lines) is from Putirka (2008)
45
46
47
48
49
50
51
52
53
54
55
56
57

58

1258
59

60

1259 **Fig. 6 (colour online)**
62

63

64

65

1260 Flow chart representing the rationality of our models for clinopyroxene (Cpx) and plagioclase
1
1261 (Plag) thermobarometry and hygrometry, from the starting mineral compositions (black
2
3
4
1262 squares) to the final P - T - H_2O triplets. (a) Model for clinopyroxenes in the less differentiated
5
6
1263 Shoshonitic Silica Saturated rocks; (b) Model for clinopyroxenes in the evolved Shoshonitic
7
8
1264 Silica Saturated and in the Shoshonitic Silica Undersaturated rocks; (c) Model for plagioclases
9
10
1265 in both Shoshonitic Silica Saturated and Shoshonitic Silica Undersaturated rocks
11
12
13

14
15
16
1267 **Fig. 7 (colour online)**
17
18

19
1268 (a) Temperature vs. pressure and (b) temperature vs. water content diagrams for Predazzo
20
21
1269 Intrusive Complex rocks. SS = Shoshonitic Silica Saturated; SU = Shoshonitic Silica
22
23
1270 Undersaturated. The reported values were obtained from the several thermobarometric
24
25
1271 approaches explained in text. Cpx = clinopyroxene; Amph = amphibole; Plag = plagioclase; K-
26
27
1272 Feld = K-feldspar. Dashed interval in (a) represent the best-fit pressure range between the
28
29
1273 various estimates, whereas the shaded field represents the temperature interval resulted from
30
31
1274 biotite thermometry. Depth was calculated by considering a $\Delta P/\Delta z$ of 0.29 kbar/km
32
33
34
35

36
37
38
1276 **Fig. 8 (colour online)**
39
40

41
1277 Temperature vs. oxygen fugacity (expressed in $\log fO_2$) diagram for Predazzo Intrusive
42
43
1278 Complex rocks. SS = Shoshonitic Silica Saturated; SU = Shoshonitic Silica Undersaturated.
44
45
1279 FMQ and NNO buffers are from Myers and Eugster (1983) and Huebner and Sato (1970),
46
47
1280 respectively. Up on the right, insert with the same oxygen fugacity data expressed in terms of
48
49
1281 ΔFMQ to facilitate the comparison between the various magmatic suites
50
51
52

53
54
55
1283 **Fig. 9 (Colour online)**
56
57

58
1284 Diagrams showing the $^{87}Sr/^{86}Sr$ vs. $^{143}Nd/^{144}Nd$ isotopic ratios of Predazzo Intrusive Complex
59
60
1285 rocks compared to those of Permo-Triassic crustal components (all values corrected to 234 Ma).
61
62
63
64
65

1286 SS = Shoshonitic Silica Saturated; SU = Shoshonitic Silica Undersaturated. Triassic carbonates
1
1287 field from Martin and Macdougall (1995) and Blendiger et al. (2015); Permian intrusives (Mt.
2
3
4
1288 Croce and Serie dei Laghi) field from Rottura et al. (1997) and Sinigoi et al. (2016, and
5
6
1289 references therein); Athesina Volcanic District (AVD) Permian ignimbrites field from Barth et
7
8
1290 al. (1993); Val Sesia Permian mafic rocks field from Voshage et al. (1990), Pin and Sills (1996)
9
10
1291 and Sinigoi et al. (2016); Kinzigite formation field from Voshage et al. (1990). (a) Sr-Nd
11
12
1292 isotopic ratios of Predazzo Intrusive Complex rocks plotted against those of the Permo-Triassic
13
14
15
1293 crustal components, the enriched 1 (EMI), high- μ (HIMU, where $\mu = {}^{238}\text{U}/{}^{204}\text{Pb}$) and depleted
16
17
1294 (DMM) mantle end-members (Zindler and Hart 1986); the insert corresponds to the area
18
19
20
21
1295 reported in diagram (b). (b) EC-AFC trends (Bohrson and Spera 2001; Spera and Bohrson 2001)
22
23
24
1296 for the Shoshonitic Silica Saturated suite by considering variable crustal assimilants. 1 =
25
26
1297 assimilation of Triassic carbonates; 2 = assimilation of AVD ignimbrites; 3 = assimilation of
27
28
1298 Mt. Croce granodiorites; 4 = assimilation of Serie dei Laghi granites; 5 = assimilation of
29
30
1299 Kinzigite formation amphibolites
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

[Click here to view linked References](#)

1 **Intrusion of shoshonitic magmas at shallow crustal depth: T-P path, H₂O estimates and**
2 **AFC modelling of the Middle Triassic Predazzo Intrusive Complex (Southern Alps,**
3 **Italy)**

4
5 **Federico Casetta***

6 Department of Physics and Earth Sciences, University of Ferrara

7 Via Saragat 1, 44121 Ferrara, Italy

8
9 **Massimo Coltorti**

10 Department of Physics and Earth Sciences, University of Ferrara

11 Via Saragat 1, 44121 Ferrara, Italy

12
13 **Ryan B. Ickert**

14 Scottish Universities Environmental Research Centre, Scottish Enterprise Technology Park,

15 Rankine Avenue, East Kilbride, G75 0QF, UK

16
17 **Costanza Bonadiman**

18 Department of Physics and Earth Sciences, University of Ferrara

19 Via Saragat 1, 44121 Ferrara, Italy

20
21 **Pier Paolo Giacomoni**

22 Department of Physics and Earth Sciences, University of Ferrara

23 Via Saragat 1, 44121 Ferrara, Italy

24
25 **Theodoros Ntaflos**

26 Department of Lithospheric Research, Universität Wien

27 Althanstraße 14 (UZA II), 1090 Wien

28
29 * Corresponding author. Phone +39 0532 974721. E-mail: cstfrc@unife.it

30 **Intrusion of shoshonitic magmas at shallow crustal depth: T-P path, H₂O estimates and**
1
231 **AFC modelling of the Middle Triassic Predazzo Intrusive Complex (Southern Alps,**
3
4
32 **Italy)**
6
733
8

9
34 **Abstract**
10

11 The multi-pulse shoshonitic Predazzo Intrusive Complex represents an ideal igneous laboratory
1235
13 for investigating the chemical and physical conditions of magma emplacement in a crustal
1436
15 context, since numerical models can be constrained by field evidence. It constitutes the most
16
1737 intriguing remnant of the Middle Triassic magmatic systems of the Dolomitic Area (Southern
18
1938 Alps), preserved by the Alpine tectonics. Predazzo Intrusive Complex comprises silica
20
21 saturated (pyroxenites/gabbros to syenites), silica undersaturated (gabbros to syenites) and
2239
23 silica oversaturated (granites and syenogranites) rock suites. In this paper we modelled its
2440
25 emplacement and evolution with a multiple thermo-/oxy-barometric, hygrometric and EC-AFC
26
2741 approach. At odds with what proposed in literature but according to the field evidence, the
28
2942 emplacement of the Predazzo Intrusive Complex occurred at shallow depth (< 6 km). In this
30
3143 context, the different pulses differed slightly in bulk water content, but shared a common
32
33 thermal regime, with temperatures between 1000-1100°C and ~600°C at low to moderate
3444
35 oxydizing conditions (-0.1 to +0.7 ΔFMQ). The interaction between the intrusion and the
3645
37 shallow crustal rocks was minimal, with Sr and Nd isotopic compositions indicating an average
38
3946 of 5-6% assimilation of crust. A thermo- and oxy-barometric comparison with the nearby Mt.
40
4147 Monzoni enabled also to speculate about the solidification time of the intrusion, which we infer
42
43 took place over about 700 ka.
4448
45
46
47
48
49
50
5151
52
5352
54
55

563 **Keywords**
57

5854 Predazzo Intrusive Complex; Sr and Nd isotopes; P-T estimates and magma water content;
59
60 Crustal assimilation; Solidification time; Shoshonitic intrusion.
6155
62
63
64
65

56
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1. Introduction

The Middle Triassic magmatic event in the Southern Alps is expressed in volcano-plutonic sequences outcropping from the Brescian Alps, Alto Vicentino, Dolomites and Carnia areas (Italy) to the Karavanken region (Austria). Most of the igneous products are volcanic and volcanoclastic rocks, with subordinated dyke swarms. Intrusive bodies are instead rare, and mainly located in the Dolomites (Predazzo-Mt. Monzoni-Cima Pape) and Karavanken areas (Gasparotto and Simboli 1991; Gianolla 1992; Bonadiman et al. 1994; Visonà and Zanferrari 2000; Brack et al. 2005; Cassinis et al. 2008; Bellieni et al. 2010; Casetta et al. 2017). These intrusions represent snapshots of magmatic plumbing systems, emplaced during and/or immediately after the eruption of the overlying volcanic products. There are several outstanding geological problems regarding the nature of these complexes, mainly related to their emplacement conditions, and to the relationships between tectonics and magmatism. The Predazzo Intrusive Complex (PIC) represents the ideal “petrologic laboratory” for investigating and reconstructing the features of the Middle Triassic feeding systems, since it offers the possibility to constrain the theoretical and experimental modeling by means of field evidence. As in the case of the nearby Mt. Monzoni, the contact aureole between PIC and the surrounding sedimentary rocks enabled many authors to speculate about the geometry of the intrusive body and its field relationships to the shallow crust (Princivalle et al. 1999; Ferry et al. 2002; Povoden et al. 2002; Gallien et al. 2007). On the basis of petrological and field evidence, Casetta et al. (2017) identified three different magma batches constituting the PIC, interpreted as a multi-pulse body of shoshonitic affinity with variable alkalis and H₂O-content which led to differentiation from mafic to amphibole and biotite-bearing end-members. However, several issues are still unsolved, such as the (possible) interaction between PIC magmas and crust, and their chemical/physical conditions of emplacement. The latter resulting in a current uncertainty about the depth of the intrusion.

82 Few thermobarometric and oxybarometric data are reported in literature for the Middle Triassic
1
23 magmatic systems of the Dolomitic Area. According to Bonadiman et al. (1994), the oxygen
3
4
54 fugacity of the nearby Mt. Monzoni system was around the NNO buffer and the crystallization
6
75 temperature range was between 1044 and 589°C. Despite the clear evidence of a predominant
8
9
86 role of water during crystallization, as testified by the wide presence of hydrous phases in the
10
11
127 intrusive rocks, no estimates of the H₂O contents and *P-T* conditions of these Middle Triassic
13
14
38 plumbing systems have been performed so far. Taking into account the Al^{tot} content of
15
16
179 amphiboles, Menegazzo Vitturi et al. (1995) hypothesized a depth of 10-17 km for PIC
18
19
90 emplacement, but their values appeared in stark contrast with both the field evidence and the
20
21
2291 data reported by Visonà and Zanferrari (2000) for the similar and coeval Karawanken pluton
23
2492 (5-9 km).

25
26
93 Therefore, in this paper we made new geochemical and isotopic (⁸⁷Sr/⁸⁶Sr-¹⁴³Nd/¹⁴⁴Nd)
27
28
294 measurements on new samples from PIC that are representative of its main portions. A multiple
30
31
95 thermobarometric and hygrometric approach, based on the interaction between distinct single
32
33
3496 mineral, mineral pairs and mineral-melt equations, corroborated by appropriate simulations by
35
36
97 means of Rhyolite-MELTS software (Gualda et al. 2012), enabled us to: (i) provide for the first
37
38
398 time *P-T-fO₂* estimates and H₂O evaluation of PIC system over its entire evolution; (ii) verify
40
4199 if the depth of the intrusion obtained by the previous petrologic approaches well fit the field
42
43
44100 evidence. The resulting *P-T-fO₂* and H₂O values, together with the isotopic signatures of PIC
45
46
4701 rocks were then used as input to (iii) quantify the role of assimilation and fractional
47
48
102 crystallization processes in the generation of the main PIC magmatic suites.
49
50

5103 52 53 104 **2. Middle Triassic magmatism in the Southern Alps** 54

55 5605 **2.1. Geodynamic framework** 57

58
106 The Middle Triassic magmatic sequences of the Dolomitic Area are mainly composed of
59
60
6107 volcanites and volcanoclastites with subordinated and scattered intrusive bodies over an area of
62
63
64
65

108 about 2,000 km² (Vardabasso 1929, 1930; Castellarin et al. 1980, 1982; Lucchini et al. 1982;
1
209 Sloman 1989; Bonadiman et al. 1994; Coltorti et al. 1996; Gianolla et al. 2010; Casetta et al.
3
4
510 2017). The Middle Triassic geodynamic framework of Dolomitic Area and the whole Southern
6
711 Alps domain is still matter of debate, fostered by juxtaposition between the calc-
8
912 alkaline/shoshonitic orogenic affinity of the magmatic products and evidence for a concomitant
10
1113 extensional-transensional tectonic regime (Doglioni 1987; Stampfli and Borel 2002, 2004;
13
14
1514 Doglioni 2007). Several geodynamic models were thus proposed, invoking the presence of: (i)
15
16
1715 an aborted rift in a passive margin (Bernoulli and Lemoine 1980); (ii) a compression at the NW
18
1916 limb of the Paleo-Tethys (Castellarin et al. 1980); (iii) an “active” mantle upwelling (Stahle et
20
21
2217 al. 2001); (iv) a transition to back-arc conditions lasting since Carboniferous-Permian, triggered
23
2418 by the northward subduction of the Paleotethys remnants (Ziegler and Stampfli 2001; Stampfli
25
26
2719 and Borel 2002; 2004; Cassinis et al. 2008; Schmid et al. 2008 Zanetti et al. 2013); (v) an
28
2920 anorogenic rifting concomitant with arc-like magmatism, whose signature was inherited by a
30
31
3221 mantle source previously metasomatised by subduction-related components during the
32
33
3422 Hercynian orogenic cycle (Bonadiman et al. 1994). This last hypothesis was supported by the
35
36
3723 similar Sr-Nd isotope signatures of the Middle Triassic magmas in the Southern Alps and the
38
3924 Permian igneous products (260-290 Ma), whose expression were found along the Western and
40
4125 in the Eastern Alps sectors (Barth et al. 1993; Rottura et al. 1998; Monjoie et al. 2007;
42
43
4426 Schaltegger and Brack 2007; Marocchi et al. 2008; Quick et al. 2009; Willcock et al. 2015; Dal
45
4627 Piazz et al. 2015; Sinigoi et al. 2016; Manzotti et al. 2017). This kind of analogy led to speculate
47
48
4928 that similar mantle sources, and/or similar melting conditions, could have generated the Middle
50
5129 Triassic and the Permian magmatic episodes. In the Dolomitic Area, thick sequences of basaltic
52
53
5430 andesitic to rhyolitic ignimbrites associated to the Permian Atesina Volcanic District are also
55
5631 the main constituent of the crustal basement of the Dolomitic Area where magmas intruded
57
58
5932 during Middle Triassic.
60
6133

2.2. The Southern Alps intrusive bodies: geochemical and petrographic overview

In contrast to the large amount of basaltic/latitic volcanic rocks erupted in the Southern Alps during the Ladinian-Carnian, the intrusive bodies are more limited in volume. They are Cima Pape (< 3 km²), Mt. Monzoni (4.6 km²), Predazzo Intrusive Complex (25 km², see below) in the Dolomitic Area (Italy) and Karawanken (about 50 km²) in Carinthia (Austria).

The Cima Pape sill and the ca. 230 Ma Mt. Monzoni body (Borsi and Ferrara 1968) are composed of biotite and amphibole-bearing gabbroic to syenitic rocks with shoshonitic affinity, generated mainly by fractional crystallization from parental basaltic to trachybasaltic magmas (Gasparotto and Simboli 1991; Bonadiman et al. 1994; Della Lucia 1997). The emplacement of Mt. Monzoni body is thought to be controlled by a syn-genetic ESE-WNW transcurrent tectonics, which created the conditions for a shallow level intrusion and the subsequent magma differentiation. A shallow depth is also suggested for the Cima Pape sill, where a gradual textural transition to the overlying lavas, locally organized in columnar structures (Sarti and Ardizzoni 1984) is observable on field.

The Karawanken pluton (ca. 230 Ma, Lippoldt and Pidgeon 1974) comprises biotite and amphibole-bearing diorites/monzonites to granites/syenites, with rare aplites and pegmatites, showing an overall high-K calc-alkaline to shoshonitic affinity. To explain the production of Karawanken magmas, Visonà and Zanferrari (2000) invoked the occurrence of combined assimilation + fractional crystallization processes from a parental basaltic melt in a shallow magma chamber (5-9 km). The geochemical differences (i.e. La/Nb and Ba/Nb ratios) between the Karawanken pluton and the intrusions of the Dolomitic Area led the authors to suggest the origin of Karawanken magmas from an enriched mantle source not previously affected by subductive components, as proposed by Bonadiman et al. (1994) for the Mt. Monzoni body. Irrespective of the volumetric differences, common features of these intrusions are the high-K calc-alkaline to shoshonitic affinity, the modal occurrence of hydrated phases (biotite and amphibole) in the intercumulus assemblages and the shallow depth of emplacement.

2.3. The Predazzo Intrusive Complex

The Predazzo Intrusive Complex (PIC, Fig. 1) is a ring-like shaped multi-pulse intrusion with an overall volume of about 4.5 km³ cropping out in Trentino Alto Adige (NE Italy). Dated at 237.3 ± 1.0 Ma (U/Pb on zircon from granites, Mundil et al., 1996), it intruded the Permian Atesina Volcanic District rhyolitic ignimbrites, the Permo-Triassic sedimentary formations and the overlying volcanic sequences (Brack et al. 1996, 1997, 2005; Mietto et al. 2012), forming well-defined metamorphic aureoles (Princivalle et al. 1999; Ferry et al. 2002; Povoden et al. 2002; Gallien et al. 2007). The intrusion and its related volcanic products were almost completely preserved from the action of the Alpine tectonic event, and kept in their original position with respect to the surrounding rocks. According to Casetta et al. (2017), PIC rocks are grouped in three main units with different geochemical features (Fig. 1), named (in relative order of emplacement) Shoshonitic Silica Saturated (SS, 3.1 km³), Granitic Unit (GU, 1.1 km³) and Shoshonitic Silica Undersaturated (SU, 0.3 km³). The SS and SU suites are composed of pyroxenites/gabbros to syenites (Fig. 2), respectively quartz- and nepheline-bearing, formed as result of fractional crystallization (FC) processes in an almost closed system from two distinct starting monzogabbroic-like magmas (Petersen et al. 1980; Visonà 1997; Marrocchino et al. 2002; Casetta et al. 2017). The Granitic Unit is instead constituted by quartz and biotite-bearing granites/syenogranites (Fig. 2), and apparently seems to represent a different magma batch and/or the result of genetic processes independent from those that generated the SS and SU rocks (Menegazzo Vitturi et al. 1995; Visonà 1997; Casetta et al. 2017). However, the lack of isotopic studies made it impossible to definitively exclude for the three magma bodies a significant involvement of assimilated crustal rocks.

On the other side, the field relationships between the intrusive units, their related dykes and the host rocks suggest a shallow origin of this multi-pulse intrusion, as also testified by the presence of hypabyssal rocks at the hundred-meter-scale transition between the plutonic and the volcanic

186 sequences (Casetta et al. 2017). Nevertheless, the only thermobarometric studies present in
1 literature, based on the Al^{tot} content of amphiboles (Menegazzo Vitturi et al. 1995), proposed a
2 10-17 km depth for the PIC, giving rise to some uncertainty about its emplacement and
3
4 188 crystallization conditions due to the discrepancy between the barometric- and field-based
5
6
7 189 estimates.
8
9
10 190

14 192 **3. Analytical methods**

16 Whole rock major and trace element analyses were carried out at the Department of Physics
17 and Earth Sciences (University of Ferrara, Italy) using ARL Advant-XP automated X-Ray
18
19 194 fluorescence spectrometer. Full matrix correction procedure and intensities were elaborated
20
21 195 following Traill and Lachance (1966). Accuracy and precision are better than 2-5% for major
22
23
24 196 elements and 5-10% for trace elements. The detection limits are 0.01 wt% and 1-3 ppm for most
25
26
27 197 of the major and trace element concentrations, respectively.
28
29 198

30
31 199 Rb, Sr, Y, Zr, Nb, Hf, Ta, Th, U, and Rare earth elements (REE) were analyzed by inductively
32
33
34 200 coupled plasma-mass spectrometry (ICP-MS) using a Thermo Series X spectrometer with
35
36 201 precision and accuracy better than 10% for all elements, well above the detection limit.
37

38
39 202 Mineral phases major element compositions were analyzed at the Department of Lithospheric
40
41 203 Research, University of Wien (Austria), by using a CAMECA SX100 electron microprobe
42
43 204 equipped with four WD and one ED spectrometers. The operating conditions were as follows:
44
45
46 205 15 kV accelerating voltage, 20 nA beam current, 20 s counting time on peak position. Natural
47
48
49 206 and synthetic standards were used for calibration and PAP corrections were applied to the
50
51 207 intensity data (Pouchou and Pichoir 1991).
52

53 208 Whole rock ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd isotopic analyses were performed at the Radiogenic
54
55
56 209 Laboratory of the Scottish Universities Environmental Research Centre (SUERC) of Glasgow
57
58 210 by means of a Sector-54 TIMS instrument. Strontium was loaded onto outgassed single Re
59
60
61 211 filaments with a Ta-activator solution; Nd was loaded onto the side of a triple Ta-Re-Ta
62
63
64
65

212 filament assembly in H₂O and ran as Nd⁺. The sample preparation procedures for Sr-Nd isotopic
1 analyses have not been described previously, so we articulate them in full, together with
213 standards and instrumental analytical performance, in the Supplementary Material_1.
2
3
4
214
5
6
215
7
8

216 **4. Whole rock geochemistry**

217 4.1. Major and trace element

218 Shoshonitic Silica Saturated (SS) suite (Fig. 2, Table 1) is mainly constituted by monzogabbros
19 and monzodiorites, with Mg# (calculated as Mg/[Mg+Fe²⁺] mol%) of 65-45, and by
20 subordinated monzonites and syenites (Mg# of 53-26 and 45-20 respectively) randomly
21 distributed within the unit (Fig. 1). Volumetrically limited pyroxenitic (Mg# 64-47) and
22 gabbroic (Mg# 65-45) cumulates crop out in the southwestern sector of the intrusion.
23

224 Shoshonitic Silica Undersaturated (SU) suite (Fig. 2, Table 1) is on average more evolved than
25 the SS one, being dominated by the presence of nepheline-normative monzonites (Mg# 40-39)
26 and syenites (Mg# 36-15), over a minority composed of cumulitic gabbros (Mg# 56-50),
27 monzogabbros (Mg# 59-47) and monzodiorites (Mg# 55-27). Granitic Unit (GU, Fig. 2) is
28 composed of highly evolved granites and syenogranites (69-77 SiO₂ wt%, Table 1) that,
29 according to Chappell and White (2001) discriminating criteria, are ascribable to the I-type
30 granitoids (Al/[Na + K + Ca] < 1.1; low P₂O₅ wt%; decreasing Zr, Sr and Al₂O₃ wt% with
31 increasing SiO₂ wt%). Few samples show instead intermediate behaviour between I- and S-type
32 rocks.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

48 All SS, SU and GU rocks have K-affinity (Fig. 2), and their K₂O and SiO₂ contents led to
49 classify them as belonging to shoshonitic (SS and SU) and High-K calc-alkaline (GU) series.
50

51 Despite the general similarities, the major and trace element distribution in the SS, SU and GU
52 rocks made it possible to pinpoint their origin from independent magmatic batches. At
53 comparable differentiation degrees, main discriminating features between the SS and SU rocks
54 are the relative enrichment in HFSE (Th, U, Pb), LREE and the higher Na₂O/K₂O ratio of the
55
56
57
58
59
60
61
62
63
64
65

latters. Such differences are in accordance with the predominant presence of amphibole and other Na and REE-rich phases in the SU rocks (Casetta et al. 2017).

4.2. $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ isotopes

Whole rock $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ isotopic ratios were analysed on representative samples of the SS, SU and GU suites (Table 2). Shoshonitic Silica Saturated samples have initial Sr isotope composition from 0.7039 to 0.7052, whereas their initial Nd isotope compositions range from 0.512191 to 0.512247. Shoshonitic Silica Undersaturated rocks are characterized by a generally higher $^{143}\text{Nd}/^{144}\text{Nd}_i$ varying from 0.512261 to 0.512289 and by $^{87}\text{Sr}/^{86}\text{Sr}_i$ comparable with the SS samples (0.7047 to 0.7063). $^{143}\text{Nd}/^{144}\text{Nd}_i$ range of GU rocks is between 0.512206 and 0.512304, thus comparable to those of both SS and SU suites. High Rb/Sr of the GU rocks resulted in imprecise initial calculated $^{87}\text{Sr}/^{86}\text{Sr}$ preventing any correlation between the GU and the SS/SU bodies.

5. Crystallization sequences and mineral chemistry

To better focus on the P - T - $f\text{O}_2$ conditions and on the H_2O contents of the PIC system, some remarks on the mineral phase compositions and crystallization sequences of the various units is hereafter summarized. A more detailed description of the petrographic and mineral chemistry features of these rocks was reported by Casetta et al. (2017). The new mineral phases analyses are reported in the Supplementary Material_2.

Shoshonitic Silica Saturated rocks are dominated by the presence of clinopyroxene and plagioclase as cumulus phases; minor olivine and orthopyroxene can be found in gabbros and monzogabbros. The intercumulus assemblage is characterized by the ubiquitous presence of plagioclase and biotite, followed by the appearance of amphibole in gabbros to monzodiorites (Fig. 3). Magnetite and Ti-magnetite are often reported in association with biotite, whereas K-feldspar and accessory phases (quartz, apatite, ilmenite, sphene, zircon) modally increase in

264 more evolved rocks. The main alteration features of the SS rocks consist of sericite formation
1
265 at the expenses of feldspars, chlorite growth over clinopyroxene, amphibole and biotite, as well
3
4
266 as epidote formation at the expenses of clinopyroxene and plagioclase. The following
6
267 crystallization sequence can be deduced for SS rocks: *olivine* → *clinopyroxene*
8
268 (*±orthopyroxene*) → *Ti-magnetite/magnetite* → *plagioclase* → *biotite* → *amphibole* → *K-*
10
11 *feldspar* → *quartz* (*±accessories*).

14
15 The crystallization sequence of the SU magmatic suite mirrors its silica undersaturation and
16
17 higher Na₂O content. Orthopyroxene and quartz are in fact absent, and an earlier appearance of
18
19 amphibole at the expenses of biotite characterises the intercumulus assemblage of the SU rocks
20
21 (Fig. 3). Other Na-rich minerals like ferrosalitic clinopyroxene, abitic plagioclase and nepheline
22
23 (Visonà 1997) are variably present. Accessory phases are ugrandite group garnets, epidote,
24
25 apatite and titanite. As already highlighted by Visonà (1997), many portions of the SU body are
26
27 hydrothermalized, showing the formation of kaoline, sericite and scapolite. Where this
28
29 secondary assemblage does not occur, the primary differences between the SU and SS
30
31 magmatic suites are evident. The likely crystallization sequence of the SU rocks is: *olivine* →
32
33 *clinopyroxene* → *Ti-magnetite/magnetite* → *plagioclase* → *amphibole* → *biotite* → *K-feldspar*
34
35 → *nepheline* (*±accessories*).

38
39
40
41 Syenogranites and granites of the GU are quite homogeneous in composition, being constituted
42
43 by K-feldspar, plagioclase and quartz, locally associated to several other minerals, among
44
45 which Fe-rich biotite is the most common. Accessory phases are tourmaline, fluorite, sphene,
46
47 magnetite, apatite, ilmenite and zircon. Muscovite and chlorite are rare and always grow at the
48
49 expenses of biotite and amphibole in late hydrothermal stages, whereas the sericitization of
50
51 feldspars is pervasive. Less common phases are ematite, allanite, scheelite, xenotime, gummite,
52
53 thorite, uranium micas, molibdenite and other sulphides, irregularly disseminated and generated
54
55 in the later pneumatolitic and hydrothermal stages (Marzocchi 1987; Visonà 1997). The main
56
57
58
59
60
61
62
63
64
65

289 GU paragenesis is composed of: *K-feldspar + plagioclase + quartz + fluorite + biotite (chlorite*
1
290 *and/or muscovite) ± amphibole + accessories.*

291 In the less altered samples, at comparable whole rock differentiation degree, the composition
2
3
4
5
6
7
292 of the most representative minerals clearly discriminate between the SS and SU magmatic
8
9
10
293 suites. For instance, biotites and amphiboles are Al- and Na-enriched in the SU rocks with
11
12
13
14
15
294 respect to the SS ones. Amphibole in fact varies from the hornblendic and actinolitic
16
17
18
19
20
295 composition in the SS rocks ($\text{Na}_2\text{O} < 1.5 \text{ wt\%}$; $\text{Al}_2\text{O}_3 < 8 \text{ wt\%}$) to the hastingsitic/pargasitic
21
22
23
296 composition (rare actinolitic hornblende, Leake et al. 1997) in the SU ones (Na_2O up to ~ 2.5
24
25
26
27
28
297 wt\% ; Al_2O_3 up to $\sim 18 \text{ wt\%}$). Even the Na content of plagioclase differs between the two
29
30
31
32
33
298 magmatic suites: it ranges from An_{84} to An_{34} in SS samples, and from An_{50} to $\text{An}_{<23}$ in SU
34
35
36
299 rocks. Another discriminating feature is represented by the clinopyroxene composition,
37
38
39
300 augitic/salitic (En_{43-33}) in the SS and salitic/ferrosalitic (En_{37-24}) in the SU rocks (Morimoto,
40
41
42
301 1988). Furthermore, the manganese content of several phases is quite higher in the SU rocks:
43
44
45
302 Ti-magnetite (0.2-1.9 wt% SU; 0.1-0.4 wt% SS), biotite (0.1- 0.7 wt% SU; 0-0.4 wt% SS),
46
47
48
303 amphibole ($\sim 0.7 \text{ wt\%}$ SU; $\sim 0.5 \text{ wt\%}$ SS) and clinopyroxene (0.5-1.5 wt% SU; 0.2-0.8 wt% SS).
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

305 **6. P, T and H₂O estimates: thermobarometric evolution of the intrusion**

306 6.1. Recover the initial equilibrium conditions

307 In this section, we apply several methods for estimating the *P*, *T* and H₂O contents of PIC rocks,
308 taking into account the coexistence of minerals used as thermometers, barometers and/or
309 hygrometers in both the SS and SU crystallization sequences. Apart from the Holland and
310 Blundy (1994), Anderson and Smith (1995), Anderson (1996) and Henry et al. (2005) methods
311 (see below) applied to amphibole-plagioclase pairs and biotite, and specific for intrusive rocks,
312 the other equations used in this study are designed for volcanic samples. As often happens in
313 intrusive contexts, the identification of the parental “melt” from which crystals formed is
314 challenging (e.g. Skaergaard Intrusion; Nielsen 2004; Namur and Humphreys 2018). Intrusive

315 rocks, in fact, rarely correspond to bulk melt composition, representing a variable mixture of
1
316 cumulus and intercumulus minerals, removed by the crystallizing melts at various stages of
3
4
317 fractionation, thus in equilibrium with different melts in different moments. Casetta et al. (2017)
6
318 suggested that the cumulitic gabbros and pyroxenites, as well as most of the intermediately
8
319 evolved rocks of the complex, were generated by various extent of fractional crystallization
10
11
320 from a starting trachybasaltic magma in an almost closed system.
13

14
321 It is modelled that the mineral assemblages constituting the PIC rocks (e.g. clinopyroxenes of
15
16
322 SS gabbro) showed partial equilibrium with a more evolved melt (e.g. SS trachybasalt), from
18
19
323 which then segregated to form the cumulates. The cumulitic nature of a rock itself incorporates
20
21
324 the concept that its composition (e.g. Mg# of clinopyroxene) represents an independent physico-
23
24
325 chemical system with respect to the deriving melt. Consequently, if we try to consider the
25
26
326 intrusive (cumulitic) rock of PIC as a bulk representative of a melt (e.g. in terms of Mg# or
27
28
327 CaO/Al₂O₃), it is reasonable to find that its mineral constituents (e.g. clinopyroxene or
30
31
328 plagioclase), are in evident disequilibrium with the bulk composition (Fig. 4 and 5). They are
32
33
329 instead compositionally coherent with a segregation process from a more evolved melt. On the
35
36
330 other side, by taking into account more evolved rocks (e.g. monzogabbros to syenites), the
37
38
331 amount of crystal disequilibrium progressively decreases (Fig. 4 and 5). This because the
40
41
332 ultimate products of the SS/SU fractional crystallization processes, syenitic in composition,
42
43
333 progressively approach the eutectic of the system and likely resemble a melt composition: this
45
46
334 condition has been also proposed by Morse and Brady (2017) for the syenites of the 1300 Ma
47
48
335 old Kiglapait Intrusion (Labrador).
49

50
336 Evidences of the reliability of this genetic model are the observed textural and mineral
52
53
337 homogeneities of the SS/SU rocks: any kind of significant zoned texture would result from the
54
55
338 introduction of additional magma chamber processes (i.e. mixing) during the formation of the
57
58
339 PIC lithotypes, invalidating our assumptions. The unzoned texture, coupled with the crystal size
59
60
340 of PIC rocks (e.g. clinopyroxenes up to 6-7 mm in pyroxenites and gabbros) and with the small
62
63
64
65

341 volume of the intrusive body (4.5 km³), speaks also against a considerable effect of syn- to post-
1
342 crystallization diffusion processes, which can be a rate-limiting process for thermobarometric
3
4
343 estimates. On the other side, thermometers involving clinopyroxene in upper crustal context,
6
344 commencing cooling from relatively low T , hold the potential to record the peak temperature
8
345 conditions, especially in large-sized grains (Müller et al. 2013). If we couple the euhedral
10
11
346 unzoned texture of PIC clinopyroxenes with the diffusion rates proposed by several authors
13
14
347 (e.g. at about 1000°C $\log D^{(Ca)} = -21$ m²/s; $\log D^{(Ti)} = -22$ m²/s; $\log D^{(Fe)} = -21$ m²/s, where D
15
16
348 indicates the diffusion coefficient; see Brady and McCallister 1983; Dimanov et al. 1996;
18
19
349 Cherniak and Lyiang 2012; Müller et al. 2013 and references therein), we can argue that the
20
21
350 crystal compositions were nearly unmodified by significant diffusion processes. Similar
23
24
351 remarks can be made for plagioclase, whose crystallization in plutonic rocks at relatively low
25
26
352 T limit the efficiency of CaAl-NaSi diffusion to submicron length scales (Grove et al. 1984).
27
28
353 The mineral compositions were therefore considered as representative of the various stages of
30
31
354 fractional crystallization of the magmas inside PIC, and were used to constrain the physical
32
33
355 parameters of the magma chamber.
35
36
356 On the basis of these assumptions, we matched each mineral (e.g. clinopyroxene of SS
37
38
357 cumulates) with an estimated “melt” following the crystal/melt equilibrium partitioning (e.g.
40
41
358 $K_{Cpx-Liq}^{Fe-Mg}$, see below), to trace the P-T path of an hypothetical differentiation trend (e.g. SS
42
43
359 suite, see Casetta et al. 2017). This operation enabled us to compute the “melt” composition, to
45
46
360 skip the apparent disequilibrium between minerals and cumulitic rocks as a bulk and thus to
47
48
361 constrain the P , T and H₂O parameters of the less evolved melts in the PIC feeding system.
49
50
362 Afterwards, the application of the “traditional” thermobarometric equations for intrusive
52
53
363 contexts (i.e. to amphibole-plagioclase pairs and biotite, see below) enabled us to verify the
54
55
364 convergence between the various results as well as to unravel the T - P -H₂O conditions in the
57
58
365 later stages of crystallization. A complete list of the applied equations, results, and
59
60
366 corresponding errors is reported in Table 4 and discussed in detail in the following sections.
62
63
64
65

6.2. Clinopyroxene-melt thermobarometer and water content of primary magmas

Clinopyroxene is the dominant and ubiquitous phase in the SS and SU rocks, thus clinopyroxene-melt thermobarometry was considered as a valuable starting point to estimate the T - P intensive variables and H_2O content of PIC magmatic system since early stage of fractionation. Crystal-melt equilibrium conditions were tested by taking into account the experimentally determine range of $Kd_{Fe-Mg}^{Cpx-Liq} = 0.24-0.30$ at $T > 1050^\circ C$ by Putirka et al. (2003). First tests highlighted that all clinopyroxene crystals in the SS and SU units have $Kd_{Fe-Mg}^{Cpx-Liq} = 0.36-0.68$ (hereinafter the superscript “Liq” indicates the composition of the bulk rock), indicating a disequilibrium towards composition more evolved than their host rock (Fig. 4). The amount of disequilibrium, as said before, decreases with increasing the differentiation degree of the rock (Fig. 4). Such a decrease could be explained by the fact that more evolved lithotypes better approach the eutectic of the system and more likely simulate melt compositions.

Following the FC model of Casetta et al. (2017), clinopyroxenes of each lithotype were thus related to a calculated “melt” composition having the same chemical affinity of their host rock and a more evolved nature (Table 3), by which crystals retrieved their equilibrium conditions. As shown in the flow chart of Fig. 6a, once chosen the clinopyroxene-melt couple of the SS suite, T - P pairs were estrapolated by means of the Putirka (2008) equations, assuming variable H_2O wt% contents of the melt. The water-dependant equations 32b and 33 (Putirka 2008), derived from the P -independent thermometer and the T -dependent barometer of Putirka (1996), were firstly applied to constrain the crystallization conditions of clinopyroxene from the melt, obtaining several T - P - H_2O triplets. Since these triplets are strongly dependant on the chosen melt and on the inferred H_2O content, they were checked, together with the melt composition, by means of the Rhyolite-MELTS calibration (Gualda et al. 2012). Iterated procedures (Fig. 6a) were developed until a clinopyroxene composition analogous to the starting one was reproduced

393 by Rhyolite-MELTS. In this way, the two independent approaches reinforce one to another,
1
394 giving a more robust framework for the thermobarometric and water results. Since the only left
3
4
395 starting parameter required by Rhyolite-MELTS was oxygen fugacity, we chose $fO_2 = NNO$ as
6
396 proposed by Bonadiman et al. (1994) for the Mt. Monzoni intrusion.

397 However, by increasing the differentiation degree of SS rocks, and/or considering the more
10
11
398 alkaline SU magmas, the reliability of Rhyolite-MELTS decreases, as a function of the alkali-
13
14
399 enrichment of the system and the related crystallization of amphibole and biotite, for whom
15
16
400 thermodynamic parameters are not well constrained (Gualda et al. 2012). Therefore, the use of
18
19
401 Masotta et al. (2013) thermobarometer was preferred for the SU suite and for the more evolved
20
21
402 SS rocks, and was cross-checked with Putirka (2008) equations (Fig. 6b). As for SS rocks, once
23
24
403 assessed the equilibrium between clinopyroxene and melt, T - P pairs at variable H_2O contents
25
26
404 of the melt were extrapolated by means of the Putirka (2008) method. The resulting T - P - H_2O
27
28
405 triplets were then inputed in Masotta et al. (2013) equation until the resulting T and P were
30
31
406 comparable to those obtained by Putirka (2008) method at similar water contents.
32

33
407 Some issues for handling this combination of modeling tools should be remarked: (i) since
35
36
408 Rhyolite-MELTS is not suitable for amphibole and biotite-dominated rocks, it was only used
37
38
409 to simulate the composition of clinopyroxene in equilibrium with the least evolved SS magmas,
40
41
410 at the P , T and H_2O conditions determined by the previous calculations; (ii) the alkaline nature
42
43
411 of PIC (especially SU) differentiated rocks implies that clinopyroxene-melt behavior during
44
45
412 crystallization and cooling depends not only on Fe and Mg, but also on Ca, Na and Al, so
47
48
413 Masotta et al. (2013) equations became progressively more reliable; (iii) iterated and cross-
49
50
414 checked calculations using Putirka (2008), Rhyolite-MELTS and Masotta et al. (2013) methods
52
53
415 (Fig. 6a-b) were proposed to unravel P - T - H_2O crystallization conditions avoiding circular
54
55
416 relationships between the equations.

57
58
417 The errors proposed for the thermobarometers, oxybarometers and hygrometers are reported in
59
60
418 Table 4: while temperatures could be defined within a narrow error range (± 10 to $\pm 20^\circ C$) with
61
62
63
64
65

both Putirka et al. (2008) and Masotta et al. (2013) equations, the P estimates were affected by high uncertainties, varying from ± 1.2 (Masotta et al. 2013) to ± 2.6 kbar (Putirka et al. 2008).

6.2.1. Clinopyroxene crystallization conditions of Shoshonitic Silica Saturated rocks

The most magnesian clinopyroxene in SS rocks were found within pyroxenites ($Mg\#_{Cpx}$ 76-78), whereas the less magnesian compositions ($Mg\#_{Cpx}$ 63-71) were in the monzodiorites and monzonites. The early T - P - H_2O crystallization conditions were likely represented by the former, which attained equilibrium with a trachybasaltic melt ($Mg\#$ 51; Fig. 4a, Table 3), further confirming the first FC step identified by Casetta et al. (2017). Clinopyroxene-melt thermobarometers indicated a P - T range of 1.6-2.3 kbar and 1070-1050°C, for water contents of 2.0-2.5 wt% (Table 4). The best fitting data in by Rhyolite-MELTS simulations were obtained at pressure of 1.5 kbar, temperature of 1060°C and 2.5 H_2O wt%. Consequently, 2.0-2.5 H_2O wt% could be assumed as the water contents of the first SS magmas intruding the PIC. Clinopyroxenes in the cumulitic gabbros were slightly less homogeneous and less magnesian ($Mg\#_{Cpx}$ 69-71 cores; 65-66 rims), and resulted in equilibrium with melts varying in $Mg\#$ from 43 to 40 (cores) to about $Mg\#$ 36 (rims), as shown in Fig. 4a and Table 3. Pressure-temperature paths indicated that cores attained equilibrium in a P range of 0.5-1.9 kbar and T of 910-980°C, while rims formed at 0.3-1.2 kbar and 900-920°C, considering H_2O contents of 3.0-4.0 wt%. Clinopyroxenes in monzogabbros and monzodiorites were nearly unzoned although compositionally comparable to those in gabbros, showing $Mg\#_{Cpx}$ of 68-72. As for gabbroic rocks, equilibrium was reached with intermediately evolved magmas ($Mg\#$ 40, Fig. 4a, Table 3), by which thermobarometric equations indicated a T range of 940-990°C at P of 1.4-1.6 kbar, for a H_2O contents in the melt of 3.0-4.0 wt%. All results are reported in Table 4 and Fig. 7a.

6.2.2. Clinopyroxene crystallization conditions of Shoshonitic Silica Undersaturated rocks

444 Clinopyroxenes from the SU suite are more alkaline than those of the SS rocks, ranging from
1
445 salitic-aegirinaugitic to ferrosalitic compositions. The most magnesian compositions in gabbros
3
446 and monzogabbros ($Mg\#_{Cpx}$ 67-63) were in equilibrium with $Mg\#$ 36 melts (Fig. 4b, Table 3).
5
6
447 Pressure-temperature paths indicate a P range of 0.2-1.6 kbar and T of 920-940°C at water
8
9
448 contents of about 1.0-1.5 wt%. Clinopyroxenes in syenites ($Mg\#_{Cpx}$ 60-48) show composition
10
11
449 close to the equilibrium with the whole rock (Fig. 4b, Table 3), which represent a theoretical
13
14
450 slightly evolved trachytic melt. Thermobarometers yielded a P of 0.2-1.6 kbar at T of 810-
15
16
451 860°C at water contents between 4.0 and 5.0 wt%. Results are reported in Table 4 and shown
18
19
452 in Fig. 7a.
20
21
22
23

454 6.3. Amphibole and amphibole-plagioclase thermobarometer and hygrometer 25

455 Pressure, temperature and H₂O conditions for amphibole crystallization were estimated using
26
27
456 amphibole-plagioclase Al-exchange thermobarometers (Holland and Blundy 1994; Anderson
28
29
30
457 and Smith 1995; Anderson 1996) and a single amphibole thermobarometers (Ridolfi et al. 2010;
31
32
33
458 Ridolfi and Renzulli 2012). Amphibole-plagioclase equations were used for mineral pairs in
34
35
36
459 both SS and SU suites, whereas the single amphibole thermobarometers were applied to all
37
38
39
460 amphibole compositions in the SS rocks, but only for amphibole in gabbros in the SU suite.
40
41
461 This because of the higher alkali content of SU differentiated magmas, well above the
42
43
462 compositional ranges admitted for the calibration of the empirical geothermometers of Ridolfi
44
45
463 et al. (2010) and Ridolfi and Renzulli (2012). For the SS and SU amphiboles, the single
46
47
48
464 amphibole equations enabled also the calculation of the water content of the melt from which
49
50
51
465 they crystallized. Contrary to the clinopyroxene-melt equations, the error range for the
52
53
466 barometric estimates based on amphibole-plagioclase and single amphibole was lower, ranging
54
55
467 between ± 0.03 and ± 0.6 kbar, whereas errors on T estimates were slightly higher (± 23 to $\pm 40^\circ\text{C}$,
56
57
58
468 Table 4).
59
60
61
62
63
64
65

470 6.3.1. Amphibole crystallization conditions of Shoshonitic Silica Saturated and Shoshonitic
1
471 Silica Undersaturated rocks
3

472 Pressure-temperature paths obtained from amphibole-plagioclase thermobarometers yielded a
6
473 T - P range of 750-845°C and 0.1-1.2 kbar in SS gabbros to monzodiorites (Fig. 7a, Table 4).
8

474 Single amphibole calculations for the same lithotypes confirmed these values, indicating $P =$
10
11
1475 0.7-1.4 kbar and $T = 720$ -810°C. The estimated H₂O contents of melts in equilibrium with
13
14
476 amphiboles of SS gabbros to monzodiorites was between 4.8 and 6.0 wt% (Fig. 7b, Table 4).
15

16
1477 Amphibole-plagioclase estimates for the SU suite yielded similar P intervals (0.4-1.0 kbar) but
18
19
478 at lower T (620-660°C) for gabbros, and P - T ranges of 1.6-1.9 kbar and 870-880°C taking into
20
21
479 account the mineral pairs in the monzogabbroic rocks (Fig. 7a, Table 4). The single amphibole
22
23
480 method, applied only to the actinolitic hornblendes in SU gabbros, provided comparable
25
26
481 pressures (0.5-0.6 kbar) and temperatures (700-720°C) with respect to the SS rocks (Fig. 7a,
27
28
482 Table 4). The calculated water contents of the melts were similar to those obtained for the SS
30
31
483 suite (5.1-5.7 wt%, Fig. 7b, Table 4).
32

33
484 As expected, the water contents calculated by these models were higher than those obtained
35
36
485 from the clinopyroxene-melt calculations. Amphibole crystallization, in fact, occurred later
37
38
486 than clinopyroxene, thus the melt from which amphibole precipitated should have undergone
40
41
487 differentiation in some extent, and its water content should have increased. If we consider the
42
43
488 water enrichment linked to the fractional crystallization of the SS/SU magmas, together with
44
45
489 the concomitant precipitation of volatile-bearing phases, the H₂O contents resulted from the
47
48
490 single amphibole equations appear in line with the calculations proposed by the FC model of
49
50
491 Casetta et al. (2017).
52

53
492
54
55
493 6.4. Biotite thermometer
56
57

58
494 Biotite is a ubiquitous phase occurring in the late inter-cumulus mineral assemblages of almost
59
60
495 all SS, SU and GU rocks. Its presence enabled us to provide some estimates on the temperatures
61
62
63
64
65

496 of these assemblages, by using of the empirical single-mineral thermometer of Henry et al.
1
497 (2005), based on the Ti content of biotite.
3

4
498
5
6
499 *6.4.1. Biotite crystallization conditions of Shoshonitic Silica Saturated, Shoshonitic Silica*
8
500 *Undersaturated and Granitic Unit rocks*
10

11
501 Temperature of biotite crystallization in SS pyroxenitic cumulates resulted between 690 and
13
14
502 740°C, whereas a T interval of 600-660°C was considered representative of SS gabbros to
15
16
503 monzodiorites. Biotites from more differentiated SS monzonites yielded a T of about 540-
18
19
504 580°C (Fig. 7a, Table 4). Biotite in SU gabbros and monzogabbros indicated comparable
20
21
505 temperatures, ranging between 640 and 660°C (Fig. 7a, Table 4).
22

23
24
506 The thermometer was also applied to GU syenogranites, resulting in a T range of 420-570°C:
25
26
507 the biotites in GU rocks are however iron-rich, with $Mg/(Mg+Fe_{tot}) < 0.275$, just out of the
27
28
508 compositional field for which the thermometer was calibrated. Consequently, the resulting
30
31
509 temperatures were not used for further modeling purposes.
32

33
34
510
35
36
511 **6.5. Plagioclase-melt thermobarometer and hygrometer**
37

38
512 Plagioclase appears in almost all PIC rocks, from the cumulitic pyroxenites, where it is part of
40
41
513 the intercumulus assemblages, to the syenitic rocks, where, together with K-feldspar, dominates
42
43
514 the paragenesis. Its ubiquitous presence made it an additional tool for investigating the T and
44
45
515 H_2O content of PIC magmas along the entire SS and SU differentiation trends by means of
47
48
516 Putirka (2008) and Lange et al. (2009) equations. Plagioclase from GU granites and
49
50
517 syenogranites are compositionally close to the pure albitic end-member, probably because of
52
53
518 hydrothermal/alteration processes. This made them unsuitable for the hygrometer application,
54
55
519 being its calibration limited to An_{37} plagioclase (Lange et al. 2009).
56
57

58
520 As for clinopyroxene, plagioclase equilibrium with its host rock was constrained by means of
59
60
521 the experimentally determined values of $^{Pl-Liq}Kd_{An-Ab} = 0.10 \pm 0.05$ at $T < 1050$ °C and 0.27 ± 0.11
62
63
64
65

522 at $T \geq 1050$ °C (Putirka 2008). However, Mollo et al. (2011) demonstrated that $K_{\text{An-Ab}}^{\text{Plag-Liq}}$
1
523 is highly sensitive to the cooling rate of the melts, being thus variable between 0.2 ± 0.02
3
524 (cooling rate of $0.5^\circ\text{C}/\text{min}$) and 0.35 ± 0.03 (cooling rate of $15^\circ\text{C}/\text{min}$). Thus, all plagioclase-
6
525 melt equilibria (even at $T < 1050^\circ\text{C}$) were considered following a $K_{\text{An-Ab}}^{\text{Plag-Liq}}$ in the range of
8
526 0.27 ± 0.11 (Putirka 2008), whose interval is also in accordance with the results obtained by
10
11 Mollo et al. (2011). Cooling rate is in fact an essential factor for the crystallization dynamics of
13
14 intrusive bodies, as testified by the plagioclase morphological variations in PIC rocks, where it
15
16 appears as both cumulus and intercumulus phase throughout the entire SS/SU fractionation
18
19 trends. According to crystallization sequences, plagioclase appeared after clinopyroxene and
20
21 Fe-Ti oxides, just before or even contemporary to biotite and amphibole in the intercumulus
22
23 assemblages. Consequently, we considered the thermobarometric values obtained by
25
26 clinopyroxene, amphibole and biotite calculations for each magmatic suite (see above) to
27
28 identify the P - T - H_2O interval in which plagioclase crystallized from the melt. As for
29
30 clinopyroxene, plagioclase often appeared in disequilibrium with its whole rock composition
31
32 (Fig. 5), and the amount of disequilibrium gradually decreases with increasing the
33
34 differentiation degree of the corresponding whole rock. Plagioclase in monzodiorites were in
35
36 fact closer to the equilibrium with respect to those of pyroxenites and gabbros, whereas most
37
38 of the plagioclase from monzonites and syenites were in equilibrium with their host rock (Fig.
39
40 5). As previously mentioned for clinopyroxene, such a trend could be justified by considering
41
42 that the more evolved rocks likely resemble melt compositions, being close to the eutectic of
43
44 the system. Thus, plagioclase compositions were related to estimated melt compositions,
45
46 according to the differentiation trends proposed by Casetta et al. (2017), to retrieve the
47
48 equilibrium conditions. Once equilibrated, plagioclase and melt compositions were used as
49
50 input for the Lange et al. (2009) hygrometer to calculate the amount of H_2O dissolved in the
51
52 melt (see flow chart of Fig. 6). For each sample, input T and P required in Lange et al. (2009)
53
54
55
56
57
58
59
60
61
62
63
64
65

equation were chosen in the T - P interval between those estimated by clinopyroxene-melt calculations and those resulted from amphibole and biotite.

Pressure, temperature and H_2O were calculated by iterating Lange et al. (2009) method and equation 24a of Putirka (2008) until a matching T was found. In any case, small variations of P scarcely affect the results of the hygrometers: ± 1 kbar corresponds to ± 0.1 wt% H_2O calculated in the melt (Giacomoni et al. 2014). Since the P interval obtained by the previous (clinopyroxene and amphibole) barometers was roughly between 0.2 and 1.9 kbar for all PIC rocks, the H_2O estimates were almost entirely constrained by temperature changes. As a consequence, the match between T obtained with Lange et al. (2009) and equation 24a of Putirka (2008) was considered a reliable cross-check and a robust constraint on our geothermal modelling (Fig. 6).

6.5.1. Plagioclase crystallization conditions of Shoshonitic Silica Saturated rocks

Data from clinopyroxene-melt simulations on pyroxenites indicated that SS primary magmas began crystallizing at about 1.5 kbar, 1060°C and 2.5 H_2O wt%. The more anorthitic plagioclase (An_{84-73}) were analyzed in the pyroxenites intercumulus assemblage, following clinopyroxene and preceding biotite, thus the T range of plagioclase crystallization is constrained by the temperatures of crystallization of these two phases. Since pressure does not sensitively affect the calculations, a pressure of 1.5 kbar was used according to the results from clinopyroxene-melt calculations (Fig. 6 and 7). Results indicated that the most anorthitic plagioclase (An_{84-73}) crystallized in equilibrium with a trachybasaltic melt (Mg# 51; Fig. 5, Table 3): at $P = 1.5$ kbar, T of crystallization resulted of 1060-1081°C for a H_2O content of 3.0 wt%.

Plagioclase in SS gabbros were in compositional continuity with those of pyroxenites, varying between An_{73} and An_{51} in both the cumulus and intercumulus assemblages. As for plagioclase inside pyroxenites, the marked disequilibrium (Fig. 5) was probably related to the cumulitic nature of gabbros and/or a variable cooling rate of the magma. The crystallization sequence and

573 the comparison with the T ranges obtained from clinopyroxene, amphibole and biotite
1 suggested that plagioclase crystallized in a temperature interval of 980 to 740°C. Plagioclase
574 with An₇₃ reached the equilibrium with a Mg# 49 melt (Fig. 5): at P of 1.5 kbar, T and H₂O
3 content resulted of 1016-1053°C and 3.1 wt% respectively. An₆₃₋₅₁ plagioclases were
575 equilibrated with a basaltic trachyandesitic melt (Fig. 5, Table 3), yielding T of 920-1021°C
6 and a water content of 3.2-3.4 wt% at pressures of 1.2-1.5 kbar (Fig. 7, Table 4). It is worth
576 noting that for these temperatures, Putirka (2008) proposed a potential partitioning $K_{\text{D}}^{\text{Plag-Liq}}_{\text{An-Ab}}$
8 = 0.05-0.15 to attest equilibrium. However, after several iterations, the modelled P - T -H₂O
9 terms for PIC magmas were attained only within a $K_{\text{D}}^{\text{Plag-Liq}}_{\text{An-Ab}}$ of 0.27±0.11. Such apparent
577 discrepancies could be explained considering the dependency of An-Ab equilibrium coefficient
10 and a water content of 3.2-3.4 wt% at pressures of 1.2-1.5 kbar (Fig. 7, Table 4). It is worth
11 noting that for these temperatures, Putirka (2008) proposed a potential partitioning $K_{\text{D}}^{\text{Plag-Liq}}_{\text{An-Ab}}$
12 = 0.05-0.15 to attest equilibrium. However, after several iterations, the modelled P - T -H₂O
13 terms for PIC magmas were attained only within a $K_{\text{D}}^{\text{Plag-Liq}}_{\text{An-Ab}}$ of 0.27±0.11. Such apparent
14 discrepancies could be explained considering the dependency of An-Ab equilibrium coefficient
15 with the cooling rate (Mollo et al. 2011).
16

583 Plagioclase from monzogabbros to monzonites were in a compositional range from An₅₃ to
25 An₃₄, resulting in equilibrium with basaltic trachyandesitic to trachyandesitic melts (Fig. 5,
26 Table 3). At P of 1.2 kbar, T resulted between 917 and 989°C, for water contents of 3.9-4.8
584 wt%, progressively increasing with the differentiation degree of the samples (Fig. 7, Table 4).
27

588 6.5.2. Plagioclase crystallization conditions of Shoshonitic Silica Undersaturated rocks

589 Plagioclase in SU rocks range from An₅₀₋₄₃ in gabbros to An₃₈₋₂₃ in monzogabbros and
40 monzodiorites, and reach the more albitic compositions in syenites, where anorthite content is
41 low (An₂₃-An₂). Plagioclase in gabbros were equilibrated by an intermediately evolved melt,
42 trachyandesitic in composition (Fig. 5, Table 3). According to the themobarometric data
43 obtained by multiple geothermometers (clinopyroxene, amphibole and biotite), the temperature
591 crystallization interval of plagioclase ranges between 940 and 720°C. Plagioclase-melt
44 thermometers and hygrometers yielded a T range of 975-990°C at P of 1.5 kbar, for a water
45 content of 4.1 wt%.
46
47
48
49
50
594
51
52
53
595
54
55
596
56
57
58
597
59
60
61
62
63
64
65

598 Apart from the nearly pure albitic compositions (likely effects of secondary processes),
1
599 plagioclase in monzogabbros to syenites vary from An₃₈ to An₂₃. They were equilibrated with
3
600 an evolved trachytic melt (Fig. 5, Table 3), thought to be the final product of the SU
6
601 differentiation trend (Casetta et al. 2017). At $T - P$ space of 920-936°C and 1.2 kbar, the
8
602 estimated water contents of melt in equilibrium with An₃₈ plagioclases was about 4.4 wt%,
10
11 progressively increasing with the differentiation of the samples (Fig. 7, Table 4).
13

14 15 16 1605 6.6. K-Feldspar-melt hygrometer

18
19
606 Except for pyroxenites, K-feldspar is present in all PIC rocks, where it occurs as intercumulus
20
21 (gabbros to monzodiorites) and cumulus (monzonites/syenites) phase. Its composition was used
22
23 to constrain the amount of H₂O dissolved in the co-existing melt by means of the equation
24
25 proposed by Mollo et al. (2015). Such method, based on the Or-Ab exchange between K-
26
609 feldspar and melt is calibrated for alkaline differentiated magmas, thus proper for the SS, SU
27
28 and GU rocks compositions. Together with the data obtained by the previous equations, such
29
30 estimates enabled to “track” the H₂O contents in the progressively differentiating melts.
31
32 According to Mollo et al. (2015), this method can be applied only to K-feldspar with Or₄₄₋₈₆,
33
34 whereas no reliable results were produced for Or_{>86} as it extended outside the range of calibrated
35
36 compositions. K-feldspar and whole rock equilibrium evaluation was attained by minimizing
37
38 the difference between predicted and measured $K\text{-Feld-Liq}Kd_{\text{Or-Ab}}$, following equation 2 of the
39
40 Mollo et al. (2015) model. These conditions were satisfied only by considering K-feldspar
41
42 compositions in equilibrium with trachyandesitic to trachytic melts (Table 3), further
43
616 confirming the correspondence between the crystallization sequence and the progressive
44
45 differentiation model proposed by Casetta et al. (2017).
46
47 According to the equilibration temperatures recorded by clinopyroxene, amphibole, biotite and
48
618 plagioclase, a T of 900-800°C was used as input for all samples of the SS and SU suites, whereas
49
50 lower temperatures (800-700°C) were considered for GU syenogranites.
51
52
53
54
55
56
57
58
622
59
60
623
62
63
64
65

624
1
625
3
4
626
6
627
8
9
628
10
11
629
13
14
630
15
16
631
18
19
632
20
21
633
22
23
634
25
26
635
27
28
636
30
31
637
32
33
638
35
639
37
38
640
40
641
42
642
44
45
643
47
644
49
50
645
52
646
54
55
647
57
648
59
60
61
62
63
64
65

6.6.1. *K.Feldspar crystallization conditions of Shoshonitic Silica Saturated, Shoshonitic Silica Undersaturated and Granitic Unit rocks*

Water concentration of melts in equilibrium with K-feldspar in SS gabbros to monzogabbros were in the range of 4.4-5.3 wt% and 5.6-6.5 wt% respectively, whereas higher H₂O contents (6.1-7.3 wt%) were recorded in monzodiorites. The highest water values were calculated in SS syenites, that in turn show a larger variability of values (H₂O = 4.6-7.3 wt%; Fig. 7, Table 4). K-feldspar analysed in SU monzogabbros and monzonites indicated H₂O contents of the crystallizing melts between 5.5-6.3 wt% and 6.7-7.8 wt%, respectively. H₂O estimates for GU syenogranites range between 8 and 11 wt%, being however meaningless in the physical system compatible with the body emplacement. The solubility of water at $P < 2.5$ kbar for rhyolitic-trachytic melts in fact do not exceed 7 wt%, as experimentally and theoretically determined in the T range of 500-1000°C (Di Matteo et al. 2004; Liu et al. 2005). These values were therefore discharged in the discussion.

7. Oxygen fugacity

The two main magmatic suites (SS and SU) are formed by the differentiation of melts which began crystallizing at pressures lower than 2.0 kbar and at similar T ranges (1050-1000°C). Main differences between the SS and SU suites could be found by taking into account the total amount of water of the crystallizing melts: SS primitive melts in fact were characterized by an higher water contents with respect to that of SU ones (2.0-2.5 wt% vs. 1.0-1.5 wt%, respectively). Classically, it is considered that dissolved molecular H₂O reacts with oxygens of the silicate network producing two OH⁻ groups (e.g. Stolper 1982; Silver and Stolper 1985; Kohn 2000). The overall reaction reading as $H_2O + O^{2-} = 2OH^-$ may thus be considered as the counterpart of the water solubility in evaluating the redox conditions.

649 To effectively quantify such differences, we estimated the $f\text{O}_2$ conditions of each portion of PIC
1
650 by means of the T - $f\text{O}_2$ model of Burkhard (1991), based on the biotite/K-feldspar/magnetite
3
4
651 equilibrium in biotite-bearing intrusive rocks. Burkhard's (1991) empirical equation
6
652 complements that of Kress and Carmichael (1988), enabling to calculate oxygen fugacity and
8
9
653 temperature by two independent formulas. Water fugacity required in Burkhard's (1991)
10
11
654 equation was calculated at a fixed P of 1.5 kbar following Burnham et al. (1969). Sanidine
13
14
655 activity was considered according to Waldbaum and Thompson (1969), whereas magnetite
15
16
656 activity was approximated at unity (Burkhard 1991). Since this equation required as input the
18
19
657 amount of Fe^{2+} in biotite, $\text{Fe}^{2+}/\text{Fe}^{3+}$ ratios of SS, SU and GU biotites were calculated following
20
21
658 Dymek (1983). Because the compositional spectrum of biotite and sanidine in each rock is
23
24
659 slightly variable, we calculated two T - $f\text{O}_2$ pairs for each lithotype (Fig. 8, Table 4), to account
25
26
660 for the possible oxygen fugacity ranges. According to Burkhard (1991), the error on each
27
28
661 calculation was of ± 0.3 log units.
30

31
662 As shown in Fig. 8, oxygen fugacity in SS rocks resulted in a range from -14.1 to -10.7 $\log f\text{O}_2$,
32
33
663 at temperatures comprised between 790 and 1000°C (Table 4). Such values plot between the
35
36
664 FMQ and NNO buffers (+0.2 to +0.7 ΔFMQ , Fig. 8), and are well comparable to the values
37
38
665 chosen as input for the Rhyolite-MELTS model above developed (-9.34 $\log f\text{O}_2$ at 1060°C),
40
41
666 further confirming the validity of the calculated parameters. Rocks from the SU and GU suites
42
43
667 were characterized by values of -11.9 to -10.2 $\log f\text{O}_2$ at T of 920-1050°C (-0.1 to +0.3 ΔFMQ ,
44
45
668 Fig. 8), and -12.9 to -11.8 $\log f\text{O}_2$ at T of 850-920°C (around +0.4 ΔFMQ , Fig. 8), respectively
47
48
669 (Table 4). Syenogranites of the GU suite were characterized by a behavior comparable to that
49
50
670 of SS rocks, whereas SU rocks record slightly more reduced conditions.
52

53 671 54 55 672 **8. EC-AFC processes** 57

58
673 As shown by the thermobarometric and hygrometric models, the two main magmatic suites (SS
59
60
674 and SU) emplaced and crystallized at comparable pressures and temperatures conditions. The
62
63
64
65

675 only differences can be identified when water content and oxygen fugacity are considered,
1
676 being the SU system characterized by lower H₂O content and by more reduced conditions.
3
4
677 Anyway, such slight divergences are not able to justify the marked geochemical variations
6
678 between SS and SU magmas (see also Casetta et al. 2017). To account for the geochemical
8
9
679 features of the SS and SU batches, we developed several Energy-Constrained Assimilation and
10
11
680 Fractional Crystallization (EC-AFC) models (Bohrson and Spera 2001; Spera and Bohrson
13
14
681 2001), based on ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd. The purpose of this calculation was to discriminate
15
16
682 between the original (i.e. mantle-derived) signature of SS/SU magmas and the effects of
18
19
683 assimilation (and/or contamination) during the emplacement and differentiation of PIC
20
21
684 magmas. Specifically, the main target of the models were: (i) verifying if the isotopic signature
23
24
685 of SU rocks was generated an interaction between SS magmas and crust; (ii) quantify the
25
26
686 eventual (if any) crustal assimilation during the differentiation of the SS/SU magmatic suites;
27
28
687 (iii) discriminate between assimilation and contamination processes.
30

31
688 A wide spectrum of Permo-Triassic crustal rock isotopic signatures from the literature were
32
33
689 considered in the EC-AFC model as potential assimilants. These are Triassic carbonates (Martin
35
36
690 and Macdougall 1995; Blendiger et al. 2015), Permian intrusives of Mt. Croce (central-eastern
37
38
691 Southern Alps; Rottura et al. 1997) and Serie dei Laghi (southwestern Alps; Sinigoi et al. 2016,
40
41
692 and references therein), Permian rhyolitic ignimbrites of the Atesina Volcanic District (Barth
42
43
693 et al. 1993), and the Kinzigite formation (southwestern Alps; Voshage et al. 1990).
45
46
694 Geothermobarometric and hygrometric results (Table 4) were used to to calculate assimilants
47
48
695 and starting magmas' specific heat [J/(KgK)], heat of crystallization (J/Kg) and heat of fusion
49
50
696 (J/Kg) according to Spera (2000), Bohrson and Spera (2001) and Spera and Bohrson (2001,
52
53
697 Table 5). Specific heat [J/(KgK)] and liquidus T (°C) of carbonate assimilant (Table 5) were
54
55
698 considered accordingly to Dallai et al. (2011) and Eppelbaum et al. (2014).
57

58
699 The absence of ⁸⁷Sr/⁸⁶Sr and ¹⁴³Nd/¹⁴⁴Nd variations with increasing silica content for both SS
59
60
700 and SU suites and the lack of overlap between SS and SU compositional fields suggested that
62
63
64
65

701 the two suites have a different origin. A slight $^{87}\text{Sr}/^{86}\text{Sr}$ increase was however noted for both
1
702 SS and SU samples while approaching the intrusion borders (Table 2). On the other side, the
3
703 $^{143}\text{Nd}/^{144}\text{Nd}$ of both SS and SU rocks were not sensitive to the distance from intrusion edges,
4
6
704 ruling out the occurrence of contamination processes during magma emplacement. Therefore,
8
705 the Nd isotopic enrichment of the SU suite can be considered as a primary feature, directly
9
10
11 function of its mantle source or alternatively derived from assimilation of crustal components.
12
13
14 In this view, the EC-AFC equations were applied to the less contaminated compositions to
15
16
1708 better constrain the Sr and (especially) Nd isotopic variations among the magmatic suites.
18
19
709 The EC-AFC model from a starting SS composition (Fig. 9) showed that none of the chosen
20
21
220 crustal components was able to drive the initial SS $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ ratios towards the
23
24
25 SU field. This simulation reinforced the primary nature of the Nd isotopes enrichment of the
26
27
712 SU body, ruling out the genesis of the SU rocks via crustal assimilation by an SS starting
28
29
30 magma. The model was instead able to explain the isotopic variability of the SS samples.
31
32
714 Almost all assimilation models well fitted the isotopic trend of SS rocks, suggesting that small
33
34
35 amounts of crustal components were assimilated during magma storage. As shown in Fig. 9,
36
37
716 the less depleted $^{143}\text{Nd}/^{144}\text{Nd}$ end-members were achieved by an interaction between magma
38
39
40 and 2-10% of crust, represented by kinzigites (Voshage et al. 1990), Mt. Croce granodiorites
41
42
43 (Rottura et al. 1997) and/or Atesina Volcanic District rhyolitic ignimbrites (Barth et al. 1993).
44
45
719 A 5% assimilation of carbonates (Martin and Macdougall 1995; Blendiger et al. 2015) and/or
46
47
48 Serie dei Laghi-like granites (Sinigoi et al. 2016) instead reproduced the more depleted
49
50
51 $^{143}\text{Nd}/^{144}\text{Nd}$ ratios.
52
53
54
55
56
57
58
725 A second EC-AFC model was attempted to explain the isotopic variations of SU samples, but
59
60
61 their scattered distribution prevented to retrace the assimilation path. Nevertheless, this model
62
63
64 was used just to quantify the crustal contribution in the Nd isotopic variations of the SU
65
66
67 samples. As a result, the $^{143}\text{Nd}/^{144}\text{Nd}$ range of the SU suite was achieved by a 2-7% assimilation

726 of the crustal components (carbonates, Atesina Volcanic District rhyolitic ignimbrites, Mt.
1 Croce granodiorites, Serie dei Laghi granites and amphibolitic kinzigites).

727
3
4
728 On the whole, the Nd isotopic ranges of both SS and SU suites could be explained by low
6
729 amounts of assimilation, whereas Sr isotopic variability is more likely derived from
8
730 contamination processes at the intrusion borders during magma emplacement, where fluid
10
11 mobilization and local metamorphic reactions occurred (Ferry et al. 2002; Gallien et al. 2007).
12
13
14
732 As reported in Table 2, the Sr isotopic signature of PIC rocks appears to be more sensitive to
15
16
733 their distance to the intrusion edges rather than to their degree of differentiation, being thus the
18
19
734 $^{143}\text{Nd}/^{144}\text{Nd}$ ratio the best proxy to discriminate the two magmatic suites. Given this, the Sr
20
21
735 variations of the SU suite at near constant $^{143}\text{Nd}/^{144}\text{Nd}$ could be also affected by i) slight
22
23
736 alteration effects, unavoidable for such a small body and/or ii) the contact halo between PIC
25
26
737 and the wall rocks.
27
28
29
30

739 **9. Discussion**

740 9.1. Emplacement of the Shoshonitic Silica Saturated and Shoshonitic Silica Undersaturated
35
741 magmas
37

742 The SS and SU intrusive bodies formed as result of fractional crystallization of two distinct
40
743 trachybasaltic melts, which differentiated along two independent trends over a short time span.
42

744 The result of this process was the generation of a large spectrum of intrusive rocks whose
45
745 cumulus+intercumulus assemblages progressively vary in composition, reaching the most
47
48
746 evolved nature in the syenitic rocks. The textural/compositional homogeneity of the main
49
50
747 cumulus phases of PIC rocks, together with the scarce efficiency of syn- to post-crystallization
52
53
748 diffusion processes on such large-sized crystals, led us to hypothesize that their composition is
54
55
749 directly function of the physical conditions at which they formed and segregated from the melt.
57

750 In this scenario, clinopyroxene and plagioclase were considered as “snapshots” of the fractional
59
60
61
62
63
64
65

751 crystallization processes, able to provide informations on the *T-P* conditions of the melts from
1
252 which they generated.
3

4
753 If we compare the clinopyroxene and plagioclase compositions to that of their host rock, in fact,
5
6
754 the disequilibrium is evident (Figg. 4 and 5). Clinopyroxene of SS gabbros (Mg# 55), for
7
8
755 example, are characterized by Mg# of 65-71, ideally in equilibrium with melt with Mg# < 43,
9
10
11
1756 quite far from the bulk rock composition of the gabbro. The most magnesian ones (Mg#_{Cpx} 76-
12
13
14
1757 78), in pyroxenites, are theoretically in equilibrium with a trachybasaltic melt (Mg# 51), further
15
16
1758 confirming that the mafic rocks of the complex are cumulates formed during the initial stages
17
18
19
1759 of fractionation of the trachybasaltic melt. On the other hand, the less magnesian clinopyroxene
20
21
22
2760 crystals tracked in syenites were almost in equilibrium with their host rocks, corroborating the
23
24
2761 assumption that the more differentiated rocks of the complex resemble the composition of melts
25
26
2762 and approach the eutectic of the system. On the basis of the results obtained by the proposed
27
28
2763 “equilibration model”, in turn supporting the differentiation model of Casetta et al. (2017), PIC
29
30
31
3764 rocks were formed during progressive stages of fractional crystallization from starting
32
33
3765 trachybasaltic melts.
34
35

36
3766 Once evaluated the disequilibrium conditions and reconduted the clinopyroxene/plagioclase
37
38
3767 compositions to their ideal equilibrium conditions, thermobarometric and hygrometric
39
40
4768 calculations enabled us to retrieve the *T-P-H₂O* parameters of the less evolved melts in the
41
42
43
4769 feeding system. The “classical” thermobarometric approach for intrusive rocks (amphibole-
44
45
4770 plagioclase and biotite equations) was then used to verify the previous results. These
46
47
48
4771 thermobarometers and hygrometers provided *T-P-H₂O* informations for the later crystallization
49
50
5772 stages.
51
52

53
5773 According to our estimates, some differences between the emplacement conditions of SS and
54
55
5774 SU magmas can be highlighted. Oxygen fugacities of the PIC plumbing system are between -
56
57
58
5775 0.1 and +0.7 Δ FMQ (Fig. 8), confirming the oxidation conditions proposed by Bonadiman et
59
60
6776 al. (1994) for Mt. Monzoni intrusion at comparable temperature intervals, further highlighting
61
62
63
64
65

777 the low to moderate oxidizing components that characterized the magmatic systems of the
1 Dolomitic Area during Triassic.

778
3
4
779 On the basis of clinopyroxene-melt thermometric and hygrometric calculations, SS magmas
6
780 result slightly water-enriched ($H_2O = 2.0-2.5$ wt%) with respect to SU ones ($H_2O = 1.0- 1.5$
8
781 wt%), during the first crystallization stages ($T \sim 1000-1100^\circ C$, Fig. 7, Table 4). With increasing
10
782 differentiation, the water content of PIC rocks increased up to 5-8 wt%, value compatible with
13
783 the amount of fractionation (79-94%) proposed by Casetta et al. (2017), by taking into account
15
784 the role of amphibole and biotite during differentiation.

18
19
785 The $T-P$ estimates (Fig. 7, Table 4) indicated that both magma batches cooled from a starting T
20
21
786 of $\sim 1000-1100^\circ C$ (cumulus assemblage) down to $\sim 600^\circ C$ (intercumulus assemblage), sharing
23
24
787 a common thermal regime. The pressure obtained by applying various mineral-mineral and
25
26
788 mineral-melt equilibria were subsequently used to estimate the emplacement depth of the PIC
27
28
789 plumbing system, and all the pressures were filtered to extrapolate the best-fit from all models,
30
31
790 taking into account the errors of the different methods (Table 4). The resulting pressure values
32
33
791 were between 0.4 and 1.7 kbar for both SS and SU intrusive bodies, yielding a depth of about
35
36
792 1.4-5.6 km considering a $\Delta P/\Delta z$ of 0.29 kbar/km. With respect to the depth (10-17 km) proposed
37
38
793 by Menegazzo Vitturi et al. (1995), our estimates suggested that PIC represents an intrusion in
40
41
794 the shallow crust, with SS and SU magma batches emplaced within the same thermal regime.
42
43
795 Despite the common geothermobarometric evolution and slightly different water contents, SS
45
46
796 and SU bodies can be distinguished in terms of $^{87}Sr/^{86}Sr$ and $^{143}Nd/^{144}Nd$, reinforcing the
47
48
797 preliminary PIC model that, on the basis of mineral and bulk rock elemental geochemistry
49
50
798 (Casetta et al. 2017) suggested that SS and SU series represent two independent magma pulses.
52
53
799 EC-AFC models (Bohrson and Spera 2001; Spera and Bohrson 2001) show that assimilation of
54
55
800 various types of Permo-Triassic crustal components by a starting SS magma could not account
57
58
801 for the higher $^{143}Nd/^{144}Nd$ of the SU batch, ruling out any possible derivation of the SU suite
59
60
802 from the SS one. Therefore, the Nd isotopic enrichment of the SU suite could be interpreted as
62
63
64
65

803 an original mantle signature or, alternatively, could be ascribed to assimilation processes
1
804 occurring within the crust. We prefer the first hypothesis, since the second one would imply the
3
4
805 existence of a common mantle-segregated magma with a $^{143}\text{Nd}/^{144}\text{Nd}$ ratio high enough to be
6
806 able to “split” in the SS and SU trends by varying the nature of the assimilant, and such a magma
8
9
807 has never been documented in the Dolomitic Area.

11
808 The same EC-AFC models demonstrated that both suites assimilated a small proportion of
13
14
809 crustal components (5-6% on average, Fig. 9). That amount of assimilation is, however, able to
15
16
810 explain the $^{143}\text{Nd}/^{144}\text{Nd}$ variation throughout SS/SU samples, being that the $^{87}\text{Sr}/^{86}\text{Sr}$ was
18
19
811 affected by late contamination that occurred at the intrusion edges. The fit between PIC isotopic
20
21
812 data and the crustal end-members of the Triassic carbonates, the Permian magmatic rocks and
23
24
813 the metamorphic basement of the Ivrea Zone is consistent with a geochemical influence on the
25
26
814 PIC of a basement that could be attributed to the Ivrea Zone. This hypothesis was also suggested
27
28
815 by Barth et al. (1993) to explain the Atesina Volcanic District-Cima d’Asta magma genesis.

30
31
816 In summary, both SS and SU batches are characterized by a strong contribution of an enriched
32
33
817 mantle component, even if the lack of primitive products cannot directly quantify its role in the
35
36
818 original source. Anyway, if examined in the light of their temporal relationships, the major,
37
38
819 trace element and Sr-Nd isotopic signature of the SS and SU bodies enable to define the
40
41
820 progressive evolution of the PIC magmatic suites. In turn, this geochemical evolution could be
42
43
821 linked to a time-related (slight) variation of the mantle source, even if an accurate dating of the
45
46
822 two single bodies would help in better deciphering the time scale of such process. On the other
47
48
823 side, the magmatic evolution in the shallow crust for both the SS and SU batches was confined
49
50
824 in the same thermal regime.

52 53 825 54 55 826 9.2. Origin of the Granitic Unit

57
58
827 The GU rocks (granites and syenogranites) differ from the other PIC magmatic suites (SS and
59
60
828 SU), mainly in terms of silica saturation, FeO, MgO, K₂O and Rb contents (Casetta et al. 2017).

829 Moreover the estimated volume of the GU portion of the complex (1.1 km^3) is unrealistically
larger than that of the SS/SU syenites ($\leq 0.25 \text{ km}^3$) if we consider the already high ($> 90\%$)
fractionation degrees from which these latter generated. This means that, if the GU portion
was the result of $\sim 95\%$ fractionation of a SS/SU magma, $\sim 20 \text{ km}^3$ of SS/SU rocks should be
intruded somewhere. This fact is in stark contrast to what is observable on field.

The GU magmas evolved at redox states comparable to SS/SU ones (about $+0.4 \Delta\text{FMQ}$) and
record temperatures down to $800\text{-}850^\circ\text{C}$ (Fig. 8). Put this all together, it is reasonable to
hypothesize that the GU granites/syenogranites were generated by magmatic differentiation
from a calc-alkaline parental melt, the primitive/intermediate products of which are not exposed
in the PIC area. It should be emphasized that GU rocks are similar to the Middle Triassic calc-
alkaline/high-K calc-alkaline rhyolites found in Carnia (Tarvisio area, Julian Alps; Gianolla
1992) and Alto Vicentino regions (Recoaro-Schio area, Southern Alps; De Vecchi et al. 1974;
Barbieri et al. 1982). In these latter, the entire high-K calc-alkaline trend is documented, further
highlighting the existence of a similar differentiation suite during the Triassic magmatic event.
In such a context, GU rocks could represent the corresponding intrusives of the rhyolites of
these areas, even if further studies are needed to better investigate this comparison.

9.3 Solidification time of PIC

According to Bonadiman et al. (1994) the solidification time of Mt. Monzoni intrusion,
emplaced at P - T - z conditions similar to those of PIC, was about 300 ka. The thermobarometric
results obtained for PIC magmas and their subsequent application to the Spera (2000), Bohrsen
and Spera (2001) and Spera and Bohrsen (2001) equations, enabled to model the cooling time
of PIC, expected to be slightly higher than that of Mt. Monzoni, due to its larger volume (4.5
 km^3 vs. $1.0\text{-}1.5 \text{ km}^3$).

In order to make an accurate comparison with Bonadiman et al. (1994), Spera (1980) equation
was applied to PIC by assuming the thermal estimates already used for the EC-AFC calculations

855 and/or derived from Ryolite-MELTS modelling (see Table 5). By taking into account the whole
1
856 volume of PIC, the solidification time resulted about 700 ka. Even if this time value was
3
4
857 obtained without discriminating the single volumes of the SS, SU and GU bodies, it was
5
6
858 considered reliable taking into account the similar thermal regimes calculated for the SS, SU
8
9
859 and GU suites and the limited re-heating effects provided by the subsequent magma pulses.
10

10. Conclusions

11
1860 A coupled thermobarometric and isotopic study of the main magmatic suites of the Predazzo
13
14
861 Intrusive Complex enabled us to set out some remarkable points on the emplacement conditions
15
16
1862 of the Middle Triassic magmas in the Dolomitic Area. They can be summarized as:
18
19
863
20
21
864
22

23
24
865 1. Predazzo Intrusive Complex emplacement occurred at a depth of 1.4-5.6 km (0.40-1.65 kbar),
25
26
866 consistent with the field relationships suggesting consanguinity with the overlying effusive
27
28
867 rocks. Its solidification time was about 700 ka, slightly higher but comparable to that proposed
30
31
868 by Bonadiman et al. (1994) for the near Mt. Monzoni intrusion.
32

33
869 2. Despite similar T - P of magma emplacement of the three unites, H_2O and fO_2 data indicated
35
36
870 that SU primary magmas were characterized by lower water contents and oxidizing conditions
37
38
871 than the SS ones (1.0-1.5 vs. 2.0-2.5 H_2O wt%; -0.1/+0.33 vs. +0.2/+0.7 ΔFMQ , respectively).
40
41
872 Plagioclase-melt, amphibole and K-feldspar-melt hygrometers indicated highly hydrated
42
43
873 conditions of magmas and progressive H_2O enrichment during differentiation.
44

45
874 3. The $^{87}Sr/^{86}Sr$ and $^{143}Nd/^{144}Nd$ of PIC magmas plot in the enriched mantle source field. This
47
48
875 feature speaks in favour of the presence of a subduction-signature in the mantle beneath
49
50
876 Southern Alps during Triassic, as already proposed by Bonadiman et al. (1994) and Zanetti et
52
53
877 al. (2013). Even if the absence of primary products in the Dolomitic Area prevents to exactly
54
55
878 quantify the role of the enriched mantle component in the magma genesis, its contribution was
57
879 undoubtedly high.
59
60
61
62
63
64
65

880 4. The EC-AFC models cannot account for the genesis of SU rocks via crustal assimilation
1
881 starting from a SS magma, thus indicating two distinct geochemical signatures for SS and SU
3
882 magmas. In turn, this difference could be ascribed to a time-related change in the mantle source,
4
6
883 slightly moving towards a more $^{143}\text{Nd}/^{144}\text{Nd}$ enriched component.
8

884 5. A low degree of crustal assimilation (5-6%) was able to explain the Nd isotopic variability
9
10
11
885 of both SS and SU magmas, whereas the Sr isotopic variations were more likely the results of
13
14
886 contamination processes at PIC edges. Fractional crystallization was the major (and almost
15
16
887 only) process acting during the differentiation of SS and SU magmas, confirming what
18
19
888 hypothesized by Petersen et al. (1980) and Casetta et al. (2017). The signature of the Triassic
20
21
889 magmas is consistent with an Ivrea-like basement beneath the Dolomitic Area.
22
23

890 6. Despite petrography, whole-rock and mineral chemistry suggested an I-type affinity for GU
25
26
891 granites/syenogranites, the wide range and the uncertainty on their isotopic signature leaves the
27
28
892 question on the origin of GU suite still unsolved. However, the geochemical and petrographic
30
31
893 similarities with the coheval calc-alkaline/high-K calc-alkaline rhyolites found in Carnia
32
33
894 (Gianolla 1992) and Alto Vicentino (De Vecchi et al. 1974; Barbieri et al. 1982; Bellieni et al.
35
36
895 2010) suggest the presence of a high-K calc-alkaline series also in the Dolomitic Area.
37
38

896 40 897 **Acknowledgments** 42

898 We are grateful to Andrea Marzoli and Matteo Masotta for their careful and constructive
45
899 reviews that significantly improved the quality of the paper, and we would also acknowledge
47
48
900 Timothy L. Grove for his editorial guidance. The IUSS Mobility Research Programme of the
49
50
901 University of Ferrara, grant n. 570 for Long Period, 2016 (FC) and The Italian National
52
53
902 Research Program PRIN_2015/prot. 20158A9 (CB) supported this research. Anne Kelly and
54
55
903 Vincent Gallagher are also thanked for making the Sr and Nd isotopic analyses at SUERC.
57

904 58 905 **References** 60 61

- 906 Anderson JL (1996) Status of thermobarometry in granitic batholiths. *Geol Soc Am Spec Pap*
1
907 315: 125-138. <https://doi.org/10.1017/S0263593300006544>
3
- 908 Anderson JL, Smith DR (1995) The effects of temperature and fO_2 on the Al-in-hornblende
6
909 barometer. *Am Min* 80(5-6): 549-559.
8
- 910 Barbieri G, De Vecchi GP, De Zanche V, Mietto P, Sedeà R (1982) Stratigrafia e petrologia del
10
911 magmatismo triassico nell'area di Recoaro. Guida alla geologia del Sudalpino centro-
13
912 orientale. *Soc Geol It Guide Geologiche Regionali*: 179-187.
15
- 913 Barth S, Oberli F, Meier M, Blattner P, Bargossi GM, Di Battistini G (1993) The evolution of
18
914 a calc-alkaline basic to silicic magma system: geochemical and Rb–Sr, Sm–Nd, and $^{18}O/^{16}O$
20
915 isotopic evidence from the Late Hercynian Atesina-Cima d'Asta volcanoplutonic complex,
23
916 northern Italy. *Geochim Cosmochim Acta* 57: 4285-4300. <https://doi.org/10.1016/0016->
25
917 7037(93)90323-O
27
- 918 Bellieni G, Fioretti AM, Marzoli A, Visonà D (2010) Permo–Paleogene magmatism in the
30
919 eastern Alps. *Rend Lincei* 21: S51-S71. <https://doi.org/10.1007/s12210-010-0095-z>
32
- 920 Bernoulli D, Lemoine M (1980) Birth and Early Evolution of the Tethys: the Overall Situation.
35
921 *Mem Bur Rech Géol Minières* 115: 168-179.
37
- 922 Blendinger W, Lohmeier S, Bertini A, Meißner E, Sattler CD (2015) A new model for the
40
923 formation of Dolomite in the Triassic dolomites, Northern Italy. *J Pet Geo* 38(1): 5-36.
42
924 10.1111/jpg.12596
44
- 925 Bohron WA, Spera FJ (2001) Energy-constrained open-system magmatic processes II:
47
926 application of energy-constrained assimilation–fractional crystallization (EC-AFC) model
49
50
927 to magmatic systems. *J Petrol* 42(5): 1019-1041.
52
928 <https://doi.org/10.1093/petrology/42.5.1019>
54
- 929 Bonadiman C, Coltorti M, Siena F (1994) Petrogenesis and T- fO_2 estimates of Mt. Monzoni
57
930 complex (Central Dolomites, Southern Alps): a Triassic shoshonitic intrusion in a transcurrent
59
931 geodynamic setting. *Eur J Mineral* 6: 943-966. 10.1127/ejm/6/6/0943
62
63
64
65

- 932 Borsi S, Ferrara G (1968) Isotopic age measurements of the M. Monzoni intrusive complex.
1
933 Miner Petrogr Acta 14: 171-183.
3
4
934 Brack P, Mundil R, Oberli F, Meier M, Rieber H (1996) Biostratigraphic and radiometric age
6
935 data question the Milankovitch characteristics of the Latemar cycles (Southern Alps, Italy).
8
936 Geology 24(4): 371-375. [https://doi.org/10.1130/0091-
12
13 7613\(1996\)024<0371:BARADQ>2.3.CO;2](https://doi.org/10.1130/0091-
10
11 7613(1996)024<0371:BARADQ>2.3.CO;2)
14
938 Brack P, Mundil R, Oberli F, Meier M, Rieber H (1997) Biostratigraphic and radiometric age
15
16 data question the Milankovitch characteristics of the Latemar cycles (Southern Alps, Italy).
17
18 Reply: Geology 25(5): 471-472.
19
20
21
22 941 Brack P, Rieber H, Nicora A, Mundil R (2005) The Global boundary Stratotype Section and
23
24 Point (GSSP) of the Ladinian Stage (Middle Triassic) at Bagolino (Southern Alps, Northern
25
26 Italy) and its implications for the Triassic time scale. Episodes 28 (4): 233-244.
27
28
29 944 Brady JB, McCallister RH (1983) Diffusion data for clinopyroxenes from homogenization and
30
31 self-diffusion experiments. Am Min 68: 95-105
32
33
34 946 Burkhard DJM (1991) Temperature and redox path of biotite-bearing intrusives: a method of
35
36 estimation applied to S- and I-type granites from Australia. Earth Planet Sci Lett 104: 89-
37
38 98. [https://doi.org/10.1016/0012-821X\(91\)90240-I](https://doi.org/10.1016/0012-821X(91)90240-I)
39
40
41 949 Burnham DJM, Holloway JR, Davis NF (1969) Thermodynamic properties of water to 1000°C
42
43 and 10,000 bars. Geol Soc Am Spec Pap 132: 96.
44
45
46 951 Casetta F, Coltorti M, Marrocchino E (2017) Petrological evolution of the Middle Triassic
47
48 Predazzo Intrusive Complex, Italian Alps. Int Geol Rev 60(8): 977-997.
49
50
51 953 <https://doi.org/10.1080/00206814.2017.1363676>
52
53
54 954 Cassinis G, Cortesogno L, Gaggero L, Perotti CR, Buzzi L (2008) Permian to Triassic
55
56 geodynamic and magmatic evolution of the Brescian Prealps (eastern Lombardy, Italy). B
57
58 Soc Geol Ital 127(3): 501-518.
59
60
61
62
63
64
65

- 957 Castellarin A, Lucchini F, Rossi PL, Sartori L, Simboli G, Sommavilla E (1982) Note
1
958 geologiche sulle intrusioni di Predazzo e dei M. Monzoni. Guida alla geologia del Sudalpino
3
4
959 centro-orientale: Guide geologiche regionali SGI: 211-219.
5
6
960 Castellarin A, Lucchini F, Rossi PL, Simboli G, Bosellini A, Sommavilla E (1980) Middle
8
9
961 Triassic magmatism in Southern Alps II: A geodynamic model. Riv Ital Paleontol S 85(3-
10
11
962 4): 1111-1124.
13
14
963 Chappell BW, White AJR (2001) Two contrasting granite types: 25 years later. Aust J Earth
15
16
964 Sci 48(4): 489-499. 10.1046/j.1440-0952.2001.00882.x
18
19
965 Cherniak DJ, Liang Y (2012) Ti diffusion in natural pyroxene. Geochim Cosmochim Acta 98:
20
21
966 31-47. <https://doi.org/10.1016/j.gca.2012.09.021>
22
23
967 Coltorti M, Siena F, Visonà D (1996) Aspetti petrologici del magmatismo Triassico dell'area
25
26
968 di Predazzo. 78° Riunione estiva Soc. Geol. It: Geologia delle Dolomiti. San Cassiano
27
28
969 (1996), Conference Abstract.
30
31
970 Dallai L, Cioni R, Boschi C, D'Oriano C (2011) Carbonate-derived CO₂ purging magma at
32
33
971 depth: influence on the eruptive activity of Somma-Vesuvius, Italy. Earth Planet Sci Lett
35
36
972 310(1): 84-95. <https://doi.org/10.1016/j.epsl.2011.07.013>
37
38
973 Dal Piaz G, Bistacchi A, Gianotti F et al. (2015) Carta Geologica d'Italia-Foglio 070 Monte
40
41
974 Cervino. Carta Geologica d'Italia, 1:50.000 scale: 70.
42
43
975 De Vecchi G, De Zanche V, Sedeà R (1974) Osservazioni preliminari sulle manifestazioni
44
45
976 magmatiche triassiche nelle Prealpi Vicentine (area di Recoaro-Schio-Posina). B Soc Geol
47
48
977 Ital 93(2): 397-409.
49
50
978 Della Lucia A (1997) Aspetti vulcanologici e petrologici del complesso vulcano-plutonico
52
53
979 Triassico di Cima Pape (Pale di San Lucano, BL). Unpublished master thesis, University of
54
55
980 Ferrara.
57
58
59
60
61
62
63
64
65

- 981 Dimanov A, Jaoul O, Sautter V. (1996) Calcium self-diffusion in natural diopside single
1 crystals. *Geochim Cosmochim Acta* 60(21): 4095-4106. [https://doi.org/10.1016/S0016-
3 7037\(96\)00250-5](https://doi.org/10.1016/S0016-
982 7037(96)00250-5)
4
983
5
6
984 Di Matteo V, Carroll MR, Behrens H, Vetere F, Brooker RA (2004) Water solubility in
8 trachytic melts. *Chem Geol* 213(1-3): 187-196.
985 <https://doi.org/10.1016/j.chemgeo.2004.08.042>
10
11
986
12
13
14
987 Doglioni C (1987) Tectonics of the Dolomites (Southern Alps, Northern Italy). *J Struct Geol* 9:
15 181-193. [https://doi.org/10.1016/0191-8141\(87\)90024-1](https://doi.org/10.1016/0191-8141(87)90024-1)
16
988
17
18
989 Doglioni C (2007) Tectonics of the Dolomites. *Bull Angew Geol* 12(2): 11-15.
19
20
990
21
991 Dymek FR (1983) Titanium, aluminium and interlayer cation substitution in biotite from high-
22 grade gneisses, West Greenland. *Am Mineral* 68: 880-899.
23
992
24
993 Eppelbaum L, Kutasov I, Pilchin A (2014) Thermal properties of rocks and density of fluids.
25 In: *Applied geothermics*. Springer Berlin Heidelberg, pp. 99-149.
26
994
27
995 Ferry JM, Wing BA, Penniston-Dorland SC, Rumble D (2002) The direction of fluid flow
28 during contact metamorphism of siliceous carbonate rocks: new data for the Monzoni and
29 Predazzo aureoles, northern Italy, and a global review. *Contrib Mineral Petrol* 142(6): 679-
30 699. <https://doi.org/10.1007/s00410-001-0316-7>
31
996
32
997
33
998 Gallien F, Abart R, Wyhlidal S (2007) Contact metamorphism and selective metasomatism of
34 the layered Bellerophon Formation in the eastern Monzoni contact aureole, northern Italy.
35 *Mineral Petrol* 91(1): 25-53. <https://doi.org/10.1007/s00710-007-0184-6>
36
999
37
1000
38
1001 Gasparotto G, Simboli G (1991) Mineralogia, petrografia e schemi evolutivi delle magmatiti
39 triassiche del complesso di Cima Pape (Dolomiti Orientali). *Mineral Petrogr Acta* 34: 205-
40 234.
41
1002
42
1003
43
1004
44
1005
45
1006
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1007 Gianolla P (1992) Evoluzione mediotriassica del vulcanismo di Rio Freddo (Api Giulie, Italia).
1
1008 Mem Sci Geol 44: 193-209.
3
4
1009 Gianolla P, Avanzini M, Breda A et al. (2010) Dolomites, 7th International Triassic Field
6
1010 Workshop, Field trip to the World Heritage Site of the Tethyan Triassic. September 5-10
8
1011 2010, Dolomites, Southern Alps, Italy. With the adesion of Fondazione Dolomiti-
10
1012 Dolomiten-Dolomites-Dolomitis Unesco.
13
14
1013 Grove TL, Baker MB, Kinzler RJ (1984) Coupled CaAl-NaSi diffusion in plagioclase feldspar:
15
1014 Experiments and applications to cooling rate speedometry. *Geochim Cosmochim Acta* 48:
18
1015 2113-2121.
20
21
1016 Gualda GAR, Ghiorso MS, Lemons RV, Carley TL (2012) Rhyolite-MELTS: a Modified
22
1017 Calibration of MELTS Optimized for Silica-rich, Fluid-bearing Magmatic Systems. *J Petrol*
25
1018 53: 875-890. <https://doi.org/10.1093/petrology/egr080>
26
27
28
1019 Henry DJ, Guidotti CV, Thomson JA (2005) The Ti-saturation surface for low-to-medium
30
1020 pressure metapelitic biotites: Implications for geothermometry and Ti-substitution
31
1021 mechanisms. *Am Mineral* 90(2-3): 316-328. <https://doi.org/10.2138/am.2005.1498>
32
33
1022 Holland T, Blundy J (1994) Non-ideal interactions in calcic amphiboles and their bearing on
36
1023 amphibole-plagioclase thermometry. *Contrib Mineral Petrol* 116(4): 433-447.
37
1024 <https://doi.org/10.1007/BF00310910>
38
39
40
1025 Huebner JS, Sato M (1970) The oxygen fugacity-temperature relationships of manganese oxide
43
1026 and nickel oxide buffers. *Am Mineral* 55: 934-952.
44
45
46
1027 Ickert RB (2013) Algorithms for estimating uncertainties in initial radiogenic isotope ratios and
48
1028 model ages. *Chem Geol* 340: 131-138. <https://doi.org/10.1016/j.chemgeo.2013.01.001>
49
50
1029 Kohn SC (2000) The dissolution mechanisms of water in silicate melts; a synthesis of recent
53
1030 data. *Mineral Mag* 64(3): 389-408. <https://doi.org/10.1180/002646100549463>
54
55
56
1031 Kress VC, Carmichael IS (1988) Stoichiometry of the iron oxidation reactions in silicate melts.
58
1032 *Am Mineral* 73: 1267-1274.
59
60
61
62
63
64
65

- 1033 Kress VC, Carmichael IS (1991) The compressibility of silicate liquids containing Fe₂O₃ and
1
1034 the effect of composition, temperature, oxygen fugacity and pressure on their redox states.
2
3
4
1035 Contrib Mineral Petrol 108(1-2): 82-92. 10.1007/BF00307328
5
6
1036 Lange RA, Frey HM, Hektor J (2009) A thermodynamic model for the plagioclase-liquid
7
8
1037 hygrometer/thermometer. Am Mineral 94(4): 494-506.
9
10
1038 <https://doi.org/10.2138/am.2009.3011>
11
12
1039 Laurenzi MA, Visonà D, (1996) ⁴⁰Ar/³⁹Ar Chronology of Predazzo magmatic complex
13
14
1040 (Southern Alps, Italy). 78° Riunione estiva Soc. Geol. It: Geologia delle Dolomiti. San
15
16
1041 Cassiano (1996): 16-18.
17
18
1042 Le Maitre RW (2002) A classification of igneous rocks and glossary of terms. Cambridge
19
20
21
1043 University Press.
22
23
1044 Leake BE, Arps CES, Birch WD (1997) Nomenclature of amphiboles: Report of the
24
25
1045 Subcommittee on Amphiboles of the International Mineralogical Association, Commission
26
27
1046 on New Minerals and Mineral Names. Am Mineral 82: 1019-1037.
28
29
1047 <https://doi.org/10.1180/minmag.1997.061.405.13>
30
31
32
1048 Lippolt H, Pidgeon R (1974) Isotopic mineral ages of a diorite from the Eisenkappel intrusion,
33
34
1049 Austria. Zeitschrift für Naturforschung 29a.
35
36
37
1050 Liu Y, Zhang Y, Behrens H (2005) Solubility of H₂O in rhyolitic melts at low pressures and a
38
39
1051 new empirical model for mixed H₂O-CO₂ solubility in rhyolitic melts. J Volcanol Geotherm
40
41
1052 Res 143(1-3): 219-235. <https://doi.org/10.1016/j.jvolgeores.2004.09.019>
42
43
44
1053 Lucchini F, Rossi PL, Simboli G (1982) Il magmatismo triassico dell'area di Predazzo (Alpi
45
46
1054 Meridionali, Italia). In Castellarin A, Vai GB (eds) Guida alla Geologia del Sudalpino
47
48
1055 centro-orientale: Guide Geologiche Regionali Società Geologica Italiana: pp. 221-230.
49
50
51
1056 Lugmair GW, Marti K (1978) Lunar initial ¹⁴³Nd/¹⁴⁴Nd: differential evolution of Lunar crust
52
53
1057 and mantle. Earth Planet Sci Lett 39: 349-357. <https://doi.org/10.1016/0012->
54
55
1058 821X(78)90021-3
56
57
58
59
60
61
62
63
64
65

- 1059 Manzotti P, Ballèvre M, Dal Piaz GV (2017) Continental gabbros in the Dent Blanche Tectonic
1 System (Western Alps): from the pre-Alpine crustal structure of the Adriatic palaeo-margin
1060 2 to the geometry of an alleged subduction interface. *J Geol Soc* 174(3): 541-556.
3
4
1061 5
6
1062 Marocchi M, Morelli C, Mair V, Klötzli U, Bargossi GM (2008) Evolution of large silicic
8 magma systems: new U–Pb zircon data on the NW Permian Athesian Volcanic Group
1063 9 (Southern Alps, Italy). *J Geol* 116: 480-498.
11
1064 12
13
1065 Marrocchino E, Coltorti M, Visonà D, Thirwall, MF (2002) Petrology of Predazzo magmatic
14 complex (Trento, Italy). *Geochim Cosmochim Acta* 66(15°): A486-A486.
15
16
1066 17
18
1067 Martin EE, Macdougall JD (1995) Sr and Nd isotopes at the Permian/Triassic boundary: A record
19 of climate change. *Chem Geol* 125(1-2): 73-99. [https://doi.org/10.1016/0009-](https://doi.org/10.1016/0009-2541(95)00081-V)
20
21
1068 22
23
1069 24
25
1070 Marzocchi C (1987) Petrologia e metallogenesi del complesso eruttivo di Predazzo (TN): con
26 particolare riguardo al settore meridionale. University of Ferrara, unpublished master thesis.
27
28
1071 29
30
1072 Masotta M, Mollo S, Freda C, Gaeta M, Moore G (2013) Clinopyroxene–liquid thermometers
31 and barometers specific to alkaline differentiated magmas. *Contrib Mineral Petrol* 166(6):
32
33
1073 34
35
1074 36
37
1075 38
39
1076 40
41
1077 42
43
1078 44
45
1079 46
47
48
1080 49
50
1081 51
52
1082 53
54
55
56
57
58
59
60
61
62
63
64
65

- 1083 Mollo S, Putirka K, Iezzi G, Del Gaudio P, Scarlato P (2011) Plagioclase–melt (dis) equilibrium
1 due to cooling dynamics: implications for thermometry, barometry and hygrometry. *Lithos*
1084 125(1): 221-235. <https://doi.org/10.1016/j.lithos.2011.02.008>
2
3
4
1085
5
6
1086 Monjoie P, Bussy F, Schaltegger U, Mulch A, Lapierre H, Pfeifer HR (2007) Contrasting
7
8
9
1087 magma types and timing of intrusion in the Permian layered mafic complex of Mont Collon
10
11
1088 (Western Alps, Valais, Switzerland): evidence from U/Pb zircon and $^{40}\text{Ar}/^{39}\text{Ar}$ amphibole
12
13
14
1089 dating. *Swiss J Geosci* 100(1): 125-135. <https://doi.org/10.1007/s00015-007-1210-8>
15
16
1090 Morimoto N (1988) Nomenclature of pyroxenes. *Mineral Petrol* 39(1): 55-76.
17
18
1091 <https://doi.org/10.1007/BF01226262>
19
20
21
1092 Morse SA, Brady JB (2017) Thermal History of the Upper Zone of the Kiglapait Intrusion. *J*
22
23
1093 *Petrol* 58(7): 1319-1332. 10.1093/petrology/egx055
24
25
26
1094 Müller T, Dohmen R, Becker HW, Ter Heege JH, Chakraborty S (2013) Fe-Mg interdiffusion
27
28
1095 rates in clinopyroxene: experimental data and implications for Fe–Mg exchange
29
30
31
1096 geothermometers. *Contrib Mineral Petrol* 166(6): 1563-1576. 10.1007/s00410-013-0941-y
32
33
1097 Mundil R, Brack P, Laurenzi MA (1996) High resolution U/Pb single zircon age
34
35
1098 determinations: new constraints on the timing of Middle Triassic magmatism in the Southern
36
37
1099 Alps. 78° Riunione estiva Soc. Geol. It: *Geologia delle Dolomiti*. San Cassiano (1996),
40
41
1100 Conference Abstract.
42
43
1101 Myers JT, Eugster HP (1983) The system Fe-Si-O: Oxygen buffer calibrations to 1500 K.
44
45
1102 *Contrib Mineral Petrol* 82(1): 75-90. <https://doi.org/10.1007/BF00371177>
46
47
48
1103 Namur O, Humphreys MCS (2018) Trace Element Constraints on the Differentiation and
49
50
1104 Crystal Mush Solidification in the Skaergaard Intrusion, Greenland. *J Petrol* 1-31. doi:
51
52
1105 10.1093/petrology/egy032
53
54
55
1106 Nielsen TFD (2004) The Shape and Volume of the Skaergaard Intrusion, Greenland:
56
57
1107 Implications for Mass Balance and Bulk Composition. *J Petrol* 45(3): 507-530.
58
59
1108 10.1093/petrology/egg092
60
61
62
63
64
65

- 1109 Petersen JS, Morten L, Simboli G, Lucchini F (1980) REE abundances in the Predazzo-
1
1110 Monzoni intrusive complex, Dolomites, North Italy. Riv Ital Paleontol S 85: 1065-1080.
2
3
4
1111 Pin C, Sills JD (1986) Petrogenesis of layered gabbros and ultramafic rocks from Val Sesia, the
5
6
1112 Ivrea Zone, NW Italy: trace element and isotope geochemistry. Geol Soc Spec Publ 24: 231-
7
8
1113 249. <https://doi.org/10.1144/GSL.SP.1986.024.01.21>
9
10
1114 Pouchou JL, Pichoir F (1991) Quantitative analysis of homogeneous or stratified microvolumes
12
13
1115 applying the model "PAP". In Electron probe quantitation. Springer US, pp. 31-75.
14
15
1116 https://doi.org/10.1007/978-1-4899-2617-3_4
16
17
1117 Povoden E, Horacek M, Abart R (2002) Contact metamorphism of siliceous dolomite and
18
19
1118 impure limestones from the Werfen formation in the eastern Monzoni contact aureole.
20
21
1119 Mineral Petrol 76(1): 99-120. <https://doi.org/10.1007/s007100200034>
22
23
24
25
1120 Princivalle F, Della Giusta A, De Min A, Piccirillo EM (1999) Crystal chemistry and
26
27
1121 significance of cation ordering in Mg-Al rich spinels from high-grade hornfels (Predazzo-
28
29
1122 Monzoni, NE Italy). Mineral Mag 63(2): 257-257.
30
31
1123 <https://doi.org/10.1180/002646199548367>
32
33
34
35
1124 Putirka KD (2008) Thermometers and barometers for volcanic systems. Rev Mineral Geochem
36
37
1125 69(1): 61-120. <https://doi.org/10.2138/rmg.2008.69.3>
38
39
40
1126 Putirka K, Johnson M, Kinzler R, Longhi J, Walker D (1996) Thermobarometry of mafic
41
42
1127 igneous rocks based on clinopyroxene-liquid equilibria, 0–30 kbar. Contrib Mineral Petrol
43
44
1128 123(1): 92-108. <https://doi.org/10.1007/s004100050145>
45
46
47
48
1129 Putirka K, Mikaelian H, Ryerson F, Shaw H (2003) New clinopyroxene-liquid
49
50
1130 thermobarometers for mafic, evolved, and volatile-bearing lava compositions, with
51
52
1131 applications to lavas from Tibet and the Snake River Plain, Idaho. Am Mineral 88: 1542-
53
54
1132 1554. <https://doi.org/10.2138/am-2003-1017>
55
56
57
58
59
60
61
62
63
64
65

- 1133 Quick JE, Sinigoi S, Peressini G, Demarchi G, Wooden JL, Sbisà A (2009) Magmatic plumbing
1 of a large Permian caldera exposed to a depth of 25 km. *Geology* 37: 603-606.
1134 <https://doi.org/10.1130/G30003A.1>
1135
- 1136 Ridolfi F, Renzulli A (2012) Calcic amphiboles in calc-alkaline and alkaline magmas:
8 thermobarometric and chemometric empirical equations valid up to 1,130° C and 2.2 GPa.
1137 *Contrib Mineral Petrol* 163(5): 877-895. <https://doi.org/10.1007/s00410-011-0704-6>
1138
- 1139 Ridolfi F, Renzulli A, Puerini M (2010) Stability and chemical equilibrium of amphibole in
14 calc-alkaline magmas: an overview, new thermobarometric formulations and application to
1140 subduction-related volcanoes. *Contrib Mineral Petrol* 160(1): 45-66.
1141 <https://doi.org/10.1007/s00410-009-0465-7>
1142
- 1143 Rotenberg E, Davis DW, Amelin Y, Ghosh S, Bergquist BA (2012) Determination of the decay-
25 constant of ⁸⁷Rb by laboratory accumulation of ⁸⁷Sr. *Geochim Cosmochim Acta* 85: 41-57.
1144 <https://doi.org/10.1016/j.gca.2012.01.016>
1145
- 1146 Rottura A, Bargossi GM, Caggianelli A, Del Moro A, Visonà D, Tranne CA (1998) Origin and
31 significance of the Permian high-K calc-alkaline magmatism in the central-eastern Southern
1147 Alps, Italy. *Lithos* 45: 329-348. [https://doi.org/10.1016/S0024-4937\(98\)00038-3](https://doi.org/10.1016/S0024-4937(98)00038-3)
1148
- 1149 Rottura A, Del Moro A, Caggianelli A, Bargossi GM, Gasparotto G (1997) Petrogenesis of the
37 Monte Croce granitoids in the context of Permian magmatism in the Southern Alps, Italy.
1150 *Eur J Mineral* 9(6): 1293-1310. [10.1127/ejm/9/6/1293](https://doi.org/10.1127/ejm/9/6/1293)
1151
- 1152 Sarti M, Ardizzoni F (1984) Tettonica Triassica nel gruppo di Cima Pape-Pale di Sanson
47 (Dolomiti Bellunesi). *Mem Sci Geol* 36: 353-370.
1153
- 1154 Schaltegger U, Brack P (2007) Crustal-scale magmatic systems during intracontinental strike-
52 slip tectonics: U, Pb and Hf isotopic constraints from Permian magmatic rocks of the
1155 Southern Alps. *Int J Earth Sci* 96(6): 1131-1151. <https://doi.org/10.1007/s00531-006-0165-8>
1156
1157 8

- 1158 Schmid SM, Bernoulli D, Fügenschuh B et al. (2008) The Alpine-Carpathian-Dinaridic
1
1159 orogenic system: correlation and evolution of tectonic units. *Swiss J Geosci* 101(1): 139-
2
3
4
1160 183. <https://doi.org/10.1007/s00015-008-1247-3>
5
6
1161 Silver L, Stolper E (1985) A thermodynamic model for hydrous silicate melts. *J Geol* 93(2):
7
8
1162 161-177.
9
10
1163 Sinigoi S, Quick JE, Demarchi G, Klotzli U (2016) Production of hybrid granitic magma at the
11
12
13
14
1164 advancing front of basaltic underplating: Inferences from the Sesia Magmatic System (south
15
16
1165 - western Alps, Italy). *Lithos* 252-253: 109-122. <https://doi.org/10.1016/j.lithos.2016.02.018>
17
18
1166 Sloman LE (1989) Triassic shoshonites from the dolomites, northern Italy: Alkaline arc rocks
19
20
21
1167 in a strike-slip setting. *J Geophys Res: Solid Earth* 94(B4): 4655-4666.
22
23
1168 10.1029/JB094iB04p04655
24
25
1169 Spera FJ (1980) Thermal evolution of plutons: a parameterized approach. *Science* 207: 299-301.
26
27
1170 10.1126/science.207.4428.299
28
29
1171 Spera FJ (2000) Physical properties of magmas. In Sigurdsson H (ed.) *Encyclopedia of*
30
31
32
1172 *Volcanoes*. New York: Academic Press, pp. 171-190.
33
34
35
1173 Spera FJ, Bohron WA (2001) Energy-constrained open-system magmatic processes I: General
36
37
38
1174 model and energy-constrained assimilation and fractional crystallization (EC-AFC)
39
40
1175 formulation. *J Petrol* 42(5): 999-1018. <https://doi.org/10.1093/petrology/42.5.999>
41
42
43
1176 Stahle V, Frenzel G, Hess JC, Saupé F, Schmidt ST, Schneider W (2001) Permian metabasalt
44
45
1177 and Triassic alkaline dykes in the northern Ivrea zone: clues to the post-Variscan
46
47
48
1178 geodynamic evolution of the Southern Alps. *Schweiz Mineral Petrogr Mitt* 81(1): 1-21.
49
50
1179 Stampfli GM, Borel GD (2002) A plate tectonic model for the Paleozoic and Mesozoic
51
52
53
1180 constrained by dynamic plate boundaries and restored synthetic oceanic isochrones. *Earth*
54
55
1181 *Planet Sci Lett* 196(1): 17-33. [https://doi.org/10.1016/S0012-821X\(01\)00588-X](https://doi.org/10.1016/S0012-821X(01)00588-X)
56
57
1182 Stampfli GM, Borel GD (2004) The TRANSMED transects in space and time: constraints on
58
59
60
1183 the paleotectonic evolution of the Mediterranean domain. In *The TRANSMED Atlas: The*
61
62
63
64
65

1184 Mediterranean region from crust to mantle, pp. 53-80. <https://doi.org/10.1007/978-3-642->
1
1185 18919-7_3
2
3
4
1186 Stolper E (1982) The speciation of water in silicate melts. *Geochim Cosmochim Acta* 46(12):
5
6
1187 2609-2620. [https://doi.org/10.1016/0016-7037\(82\)90381-7](https://doi.org/10.1016/0016-7037(82)90381-7)
7
8
1188 Traill RJ, Lachance GR (1966) A practical solution to the matrix problem in X-ray analysis. II.
9
10
1189 Application to a multicomponent alloy system. *Can Spectrosc* 11(3): 63-71.
11
12
1190 Vardabasso S (1929) Rapporti tra attività magmatica e vicende tettoniche nella provincia
13
14
1191 petrografica di Predazzo: *Studi Trentini di Scienze Naturali* 11.
15
16
1192 Vardabasso S (1930) Carta geologica del territorio eruttivo di Predazzo e Monzoni. Ufficio
17
18
1193 Idrografico del Magistrato alle Acque di Venezia, 1:25000 scale.
19
20
21
1194 Visonà D (1997) The Predazzo multipulse intrusive body (Western Dolomites, Italy). Field and
22
23
1195 mineralogical studies. *Mem Sci Geol* 49: 117-125.
24
25
1196 Visonà D, Zanferrari A (2000) Some constraints on geochemical features in the Triassic mantle
26
27
1197 of the easternmost Austroalpine–Southalpine domain: evidence from the Karawanken pluton
28
29
1198 (Carinthia, Austria). *Int J Earth Sci* 89(1): 40-51. <https://doi.org/10.1007/s005310050316>
30
31
1199 Voshage H, Hofmann AW, Mazzucchelli M, Rivalenti G, Sinigoi S, Raczek I, Demarchi G
32
33
1200 (1990) Isotopic evidence from the Ivrea Zone for a hybrid lower crust formed by magmatic
34
35
1201 underplating. *Nature* 347(6295): 731-736. [10.1038/347731a0](https://doi.org/10.1038/347731a0)
36
37
1202 Waldbaum DR, Thompson JB (1969) Mixing properties of sanidine crystalline solutions: IV.
38
39
1203 Phase diagrams from equations of state. *Am Mineral* 54: 1274-1298.
40
41
1204 Willcock MAW, Bargossi GM, Weinberg RF, Gasparotto G, Cas RAF, Giordano G, Marocchi
42
43
1205 M (2015) A complex magma reservoir system for a large volume intra-to extra-caldera
44
45
1206 ignimbrite: Mineralogical and chemical architecture of the VEI8, Permian Ora ignimbrite
46
47
1207 (Italy). *J Volcanol Geotherm Res* 306: 17-40.
48
49
1208 <https://doi.org/10.1016/j.jvolgeores.2015.09.015>
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1209 Zanetti A, Mazzucchelli M, Sinigoi S, Giovanardi T, Peressini G, Fanning M (2013) SHRIMP
 1 U–Pb Zircon Triassic intrusion age of the Finero mafic complex (Ivrea–Verbano zone,
 1210 Western Alps) and its geodynamic implications. *J Petrol* 54(11): 2235-2265.
 1211 <https://doi.org/10.1093/petrology/egt046>
 1212
 1213 Ziegler PA, Stampfli GM (2001) Late Palaeozoic-Early Mesozoic plate boundary
 1214 reorganization: collapse of the Variscan orogen and opening of Neotethys. *Nat Brescia* 25:
 1215 17-34.
 1216
 1217 Zindler A, Hart SR (1986) Chemical geodynamics. *Annu Rev Earth Planet Sci* 14: 493-571.

Figure Captions

Fig. 1 (colour online)

Simplified geological map of the Predazzo Intrusive Complex (PIC). On the left bottom corner, the study area location with respect to the NE sector of Italy is shown

Fig. 2 (colour online)

(a) Total Alkali vs. Silica and (b) K_2O vs. Na_2O diagrams for Predazzo Intrusive Complex rocks (Le Maitre 2002). SS = Shoshonitic Silica Saturated; SU = Shoshonitic Silica Undersaturated; GU = Granitic Unit

Fig. 3 (colour online)

Textural sketches of Predazzo Intrusive Complex rocks and corresponding photomicrographs showing the crystallization relationships between biotite and amphibole in the intercumulus assemblage of (a, b) Shoshonitic Silica Saturated monzodiorites and (c, d) Shoshonitic Silica Undersaturated monzogabbros. Each labelled mineral phase in the right photomicrographs is contraddistinct by a different colour in the left sketches. Amph = amphibole; Ap = apatite; Bt

1234 = biotite; Cpx = clinopyroxene; K-Feld = K-feldspar; Ox = Fe-Ti oxide; Plag = plagioclase; Qz
1
1235 = quartz; Ser = sericite
2
3

4
1236
5

6

1237 **Fig. 4 (colour online)**
8

9
1238 Mg# vs. $Kd_{Fe-Mg}^{Cpx-Liq}$ of clinopyroxene from the (a) Shoshonitic Silica Saturated and (b)
10
11 Shoshonitic Silica Undersaturated rocks. The white symbols represent the disequilibrium
1239 between crystals and their host rock composition, whose whole rock Mg# (wr Mg#) is also
13
14 reported. The filled symbols represent the attained equilibrium conditions after the calculation
1240 of melt composition in equilibrium. “Liq” indicates both the composition of the crystals host
15
16 rocks (white symbols, prior to equilibration process) and that of the calculated “melt” (filled
1241 symbols, after equilibration process). The length of the dashed arrows is directly function of
17
18 the amount of disequilibrium between the crystals and their host rock. The equilibrium range
19
1242 of $Kd_{Fe-Mg}^{Cpx-Liq} = 0.24-0.30$ (black lines) is from Putirka et al. (2003)
20
21
22
23
24
25
26
27
28
29
1246
30

31
1247
32

33

1248 **Fig. 5 (colour online)**
35

36
1249 Anorthite (X_{An}) vs. $Kd_{An-Ab}^{Plag-Liq}$ of plagioclase from the (a) Shoshonitic Silica Saturated and
37
38 (b) Shoshonitic Silica Undersaturated rocks. As in Fig. 4, white symbols represent the
1250 disequilibrium between crystals and their host rock composition (whose Mg# is reported as “wr
39
40 Mg#”), whereas filled symbols represent the attained equilibrium conditions after the
41
42 equilibration processes. “Liq” indicates both the composition of the crystals host rocks (white
43
1252 symbols, prior to equilibration process) and that of the calculated “melt” (filled symbols, after
44
45 equilibration process). The length of the dashed arrows is directly function of the amount of
46
47 disequilibrium between the crystals and their host rock. The chosen equilibrium range of Kd_{An-Ab}^{Plag-}
48
1254 $Kd_{An-Ab}^{Liq} = 0.27 \pm 0.11$ (black lines) is from Putirka (2008)
49
50
51
52
53
54
55
56
57

58
1258
59

60

1259 **Fig. 6 (colour online)**
61

62

63

64

65

1260 Flow chart representing the rationality of our models for clinopyroxene (Cpx) and plagioclase
1
1261 (Plag) thermobarometry and hygrometry, from the starting mineral compositions (black
2
3
4
1262 squares) to the final P - T - H_2O triplets. (a) Model for clinopyroxenes in the less differentiated
5
6
1263 Shoshonitic Silica Saturated rocks; (b) Model for clinopyroxenes in the evolved Shoshonitic
7
8
1264 Silica Saturated and in the Shoshonitic Silica Undersaturated rocks; (c) Model for plagioclases
9
10
1265 in both Shoshonitic Silica Saturated and Shoshonitic Silica Undersaturated rocks
11
12
13

14
15
16
1267 **Fig. 7 (colour online)**
17
18

19
1268 (a) Temperature vs. pressure and (b) temperature vs. water content diagrams for Predazzo
20
21
1269 Intrusive Complex rocks. SS = Shoshonitic Silica Saturated; SU = Shoshonitic Silica
22
23
1270 Undersaturated. The reported values were obtained from the several thermobarometric
24
25
1271 approaches explained in text. Cpx = clinopyroxene; Amph = amphibole; Plag = plagioclase; K-
26
27
1272 Feld = K-feldspar. Dashed interval in (a) represent the best-fit pressure range between the
28
29
1273 various estimates, whereas the shaded field represents the temperature interval resulted from
30
31
1274 biotite thermometry. Depth was calculated by considering a $\Delta P/\Delta z$ of 0.29 kbar/km
32
33
34
35


36
37
38
1276 **Fig. 8 (colour online)**
39
40


41
1277 Temperature vs. oxygen fugacity (expressed in $\log fO_2$) diagram for Predazzo Intrusive
42
43
1278 Complex rocks. SS = Shoshonitic Silica Saturated; SU = Shoshonitic Silica Undersaturated.
44
45
1279 FMQ and NNO buffers are from Myers and Eugster (1983) and Huebner and Sato (1970),
46
47
1280 respectively. Up on the right, insert with the same oxygen fugacity data expressed in terms of
48
49
1281 ΔFMQ to facilitate the comparison between the various magmatic suites
50
51
52


53
54
55
1283 **Fig. 9 (Colour online)**
56
57



58
1284 Diagrams showing the $^{87}Sr/^{86}Sr$ vs. $^{143}Nd/^{144}Nd$ isotopic ratios of Predazzo Intrusive Complex
59
60
1285 rocks compared to those of Permo-Triassic crustal components (all values corrected to 234 Ma).
61
62
63
64
65


1286 SS = Shoshonitic Silica Saturated; SU = Shoshonitic Silica Undersaturated. Triassic carbonates
1
1287 field from Martin and Macdougall (1995) and Blendiger et al. (2015); Permian intrusives (Mt.
2
3
4
1288 Croce and Serie dei Laghi) field from Rottura et al. (1997) and Sinigoi et al. (2016, and
5
6
1289 references therein); Athesina Volcanic District (AVD) Permian ignimbrites field from Barth et
7
8
1290 al. (1993); Val Sesia Permian mafic rocks field from Voshage et al. (1990), Pin and Sills (1996)
9
10
1291 and Sinigoi et al. (2016); Kinzigite formation field from Voshage et al. (1990). (a) Sr-Nd
11
12
1292 isotopic ratios of Predazzo Intrusive Complex rocks plotted against those of the Permo-Triassic
13
14
15
1293 crustal components, the enriched 1 (EMI), high- μ (HIMU, where $\mu = {}^{238}\text{U}/{}^{204}\text{Pb}$) and depleted
16
17
1294 (DMM) mantle end-members (Zindler and Hart 1986); the insert corresponds to the area
18
19
20
1295 reported in diagram (b). (b) EC-AFC trends (Bohrson and Spera 2001; Spera and Bohrson 2001)
21
22
23
1296 for the Shoshonitic Silica Saturated suite by considering variable crustal assimilants. 1 =
24
25
1297 assimilation of Triassic carbonates; 2 = assimilation of AVD ignimbrites; 3 = assimilation of
26
27
28
1298 Mt. Croce granodiorites; 4 = assimilation of Serie dei Laghi granites; 5 = assimilation of
29
30
1299 Kinzigite formation amphibolites
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

























Table 1. Major and trace element whole rock composition of selected Shoshonitic Silica Saturated (SS), Shoshonitic Silica Undersaturated (SU) and Granitic Unit (GU) rocks. Px: pyroxenite; Gb: gabbro; Mzgb: monzogabbro; Mzdr: monzodiorite; Mz: monzonite; Sy: syenite; Gr: granite; Sygr: syenogranite. Fe₂O₃ and FeO were calculated by considering a Fe₂O₃/FeO ratio of 0.15, in agreement with a fO₂ around FMQ buffer (Kress and Carmichael, 1991). Mg# = Mg/[Mg Fe²⁺] mol%; n.d. = not detected. All trace element (ppm) were analysed by ICP-MS except Pb, Zn, Ni, Co, Cr, V, Cu and Ba (XRF).

Suite	SS	SS	SS	SS	SS	SS	SU	SU	SU	SU	SU	SU	GU	GU
Lithology	Px	Gb	Mzgb	Mzdr	Mz	Sy	Gb	Mzgb	Mzdr	Mz	Sy	Sy	Gr	Sygr
Sample	FC31E	FC30C	FC43D	FC37B	EM79	EM64	EM15	FC2B	EM9	EM6	EM5	EM107	EM90B	EM31
Oxide (wt%)														
SiO ₂	35.48	45.64	52.56	55.43	61.84	67.91	49.83	52.39	55.48	56.86	60.22	62.44	76.41	70.21
TiO ₂	2.07	1.08	0.85	0.80	0.43	0.17	0.93	1.13	0.24	0.58	0.46	0.28	0.04	0.26
Al ₂ O ₃	6.20	16.78	18.92	17.30	19.82	16.54	17.91	17.08	23.80	21.71	19.58	18.68	13.79	15.58
Fe ₂ O ₃	3.74	1.55	1.03	1.01	0.47	0.22	1.34	1.06	0.37	0.59	0.49	0.21	0.11	0.31
FeO	24.90	10.34	6.88	6.74	3.11	1.51	8.91	7.05	2.44	3.94	3.31	1.44	0.74	2.07
MnO	0.25	0.18	0.19	0.20	0.07	0.07	0.18	0.41	0.18	0.19	0.16	0.11	0.02	0.03
MgO	12.54	6.97	4.03	3.73	1.14	0.22	4.97	4.80	0.51	1.50	0.84	0.45	0.28	0.24
CaO	13.57	14.16	8.68	6.71	3.34	0.90	9.56	8.28	3.94	4.81	3.80	3.20	1.65	1.01
Na ₂ O	0.51	1.09	2.79	3.24	3.62	4.27	2.81	4.80	5.29	5.16	5.70	3.54	5.44	3.99
K ₂ O	0.73	2.16	3.37	4.30	5.96	8.18	2.80	2.62	7.73	4.38	5.25	9.22	1.51	6.29
P ₂ O ₅	0.02	0.05	0.68	0.54	0.20	0.00	0.76	0.38	0.02	0.28	0.18	0.43	0.00	0.00
Tot.	100	100	100	100	100	100	100	100	100	100	100	100	100	100
LOI	0.00	2.57	1.33	1.09	1.72	0.99	1.21	1.57	4.33	1.13	0.60	1.20	1.53	1.23
Mg#	47.29	54.56	51.08	49.65	39.48	20.72	49.86	54.81	27.26	40.34	31.01	35.57	40.60	17.33
Trace element (ppm)														
Pb	14.8	14.2	17.8	27.5	21.0	23.0	13.0	44.1	45.0	37.0	28.0	27.0	n.d.	13.0
Zn	148	72.3	103	107	51.0	54.0	115	202	147	148	86.0	50.0	8.00	30.0
Ni	69.3	32.0	5.90	5.90	6.00	5.00	15.0	5.90	11.0	5.00	8.00	6.00	4.00	7.00
Co	62.9	52.7	27.2	24.4	9.00	6.00	40.0	29.5	4.00	9.00	7.00	3.00	n.d.	2.00
Cr	61.7	68.6	6.9	11.2	9.00	5.00	23.0	58.9	9.00	3.00	5.00	1.00	5.00	3.00
V	1600	565	153	144	32.0	15.0	239	165	36.0	80.0	67.0	29.0	13.0	9.00
Cu	13.0	17.3	85.7	79.4	n.d.	n.d.	n.d.	35.3	n.d.	n.d.	n.d.	n.d.	n.d.	n.d.
Rb	16.2	42.3	52.7	81.4	221	187	42.5	89.1	326	184	97.9	318	112	284
Ba	178	223	479	546	772	553	782	482	506	1129	498	616	121	190
Sr	95.4	440	676	556	755	284	849	621	818	1351	254	569	83.4	28.2
Nb	3.83	3.68	10.5	13.4	18.0	24.0	7.22	44.4	60.0	18.0	35.3	52.0	64.7	48.7
Zr	47.0	39.0	38.0	34.5	208	323	35.1	359	499	262	206	623	153	211
Hf	1.60	1.48	1.54	1.67	1.71	5.31	1.51	7.48	4.54	n.d.	5.66	6.95	5.05	5.81
U	0.22	0.48	0.93	1.59	2.46	3.89	1.08	2.92	11.8	7.39	5.26	11.6	11.0	2.55
Th	0.66	1.67	3.68	6.56	12.5	27.0	3.60	9.57	46.7	27.8	33.7	112	44.3	14.4
Y	14.1	11.1	11.9	13.7	21.9	24.5	13.6	23.8	13.7	26.0	14.8	49.9	27.1	5.0
Ta	0.26	0.30	0.52	0.68	n.d.	n.d.	0.51	1.57	n.d.	n.d.	1.34	n.d.	1.51	1.06
La	4.49	7.16	15.9	18.5	26.6	24.3	17.7	95.6	63.9	45.9	38.4	117	2.12	9.75
Ce	12.4	15.5	37.2	49.6	59.5	56.8	44.5	195	99.5	85.1	125	237	6.44	33.6
Pr	2.01	2.47	4.34	4.96	7.10	6.96	4.80	13.6	8.99	8.95	8.04	24.9	1.24	2.51
Nd	10.5	11.9	18.6	20.8	27.2	27.1	20.8	47.3	24.3	30.1	28.9	81.8	7.46	9.21
Sm	2.92	2.87	3.63	4.11	5.60	5.34	4.11	7.18	3.24	4.94	4.49	12.9	3.46	1.55
Eu	0.65	0.77	0.89	0.96	1.70	1.07	1.16	1.33	0.89	1.68	0.93	1.75	0.21	0.09
Gd	2.67	2.50	3.19	3.66	5.20	4.74	3.71	6.86	4.18	5.46	4.03	12.2	3.57	1.44
Tb	0.46	0.43	0.47	0.55	0.78	0.76	0.54	0.88	0.41	0.60	0.56	1.55	0.82	0.21
Dy	2.41	2.12	2.17	2.58	4.16	4.35	2.47	3.93	2.24	3.18	2.55	8.67	4.95	1.08
Ho	0.47	0.42	0.43	0.52	0.89	0.92	0.49	0.80	0.52	0.75	0.54	1.72	1.06	0.23
Er	1.22	1.08	1.12	1.38	2.25	2.57	1.27	2.22	1.59	2.09	1.53	5.34	3.04	0.73
Tm	0.18	0.17	0.18	0.22	0.38	0.44	0.20	0.37	0.26	0.35	0.27	0.84	0.52	0.13
Yb	1.06	0.96	0.95	1.25	2.51	3.08	1.08	2.16	1.78	2.30	1.61	5.78	3.17	0.89
Lu	0.15	0.14	0.15	0.19	0.35	0.47	0.17	0.34	0.27	0.35	0.25	0.83	0.45	0.14

Table 2. Whole rock $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ isotopes of Shoshonitic Silica Saturated (SS), Shoshonitic Silica Undersaturated (SU) and Granitic Unit (GU) rocks. Samples were corrected for 234 Ma (Laurenzi and Visonà 1996; Mundil et al. 1996) of radiogenic ingrowth using the trace element abundances determined by ICP-MS, the decay rates of Rotenberg et al. (2012) and Lugmair and Marti (1978). Initial ratios (i) and uncertainties (2σ) were propagated according to Ickert (2013).

Sample	Lithology	Suite	Age (Ma)	Rb (ppm)	Sr (ppm)	$^{87}\text{Sr}/^{86}\text{Sr}$	$^{87}\text{Rb}/^{86}\text{Sr}$	$^{87}\text{Sr}/^{86}\text{Sr}(i)$	2σ	Sm (ppm)	Nd (ppm)	$^{143}\text{Nd}/^{144}\text{Nd}$	$^{147}\text{Sm}/^{144}\text{Nd}$	$^{143}\text{Nd}/^{144}\text{Nd}(i)$	2σ	Distance to edge (m)
FC31E	Pyroxenite	SS	234	16.19	95.45	0.706630	0.4907	0.705024	0.000025	2.92	10.46	-	0.1483	-	-	170
FC30C	Gabbro	SS	234	42.27	440.27	0.706149	0.2777	0.705240	0.000020	2.87	11.86	0.512415	0.1463	0.512191	0.000007	420
FC43D	Monzogabbro	SS	234	52.73	675.82	0.705304	0.2257	0.704565	0.000022	3.63	18.60	0.512392	0.1180	0.512211	0.000007	565
EM79	Monzonite	SS	234	220.50	754.80	0.707524	0.8452	0.704757	0.000032	5.60	27.16	0.512389	0.1246	0.512198	-	740
EM64	Syenite	SS	234	186.70	284.10	0.710211	1.9018	0.703985	0.000065	5.34	27.12	0.512429	0.1189	0.512247	0.000006	1000
EM15	Gabbro	SU	234	42.55	849.41	0.705182	0.1449	0.704707	0.000018	4.11	20.83	0.512444	0.1192	0.512261	0.000008	385
FC2B	Monzogabbro	SU	234	89.14	620.85	0.707719	0.4154	0.706359	0.000028	7.18	47.25	0.512430	0.0918	0.512289	0.000005	135
EM107	Syenite	SU	234	318.30	569.10	0.710856	1.6187	0.705556	0.000055	12.94	81.78	0.512412	0.0956	0.512266	0.000006	300
EM5	Syenite	SU	234	97.87	253.65	0.709851	1.1166	0.706196	0.000042	4.49	28.87	-	0.0826	-	-	158
EM31	Syenogranite	GU	234	324.50	148.19	-	-	-	-	9.69	62.08	0.512350	0.0942	0.512206	0.000007	376
EM88	Syenogranite	GU	234	371.86	15.99	-	-	-	-	8.43	39.41	0.512502	0.1291	0.512304	0.000016	735

Table 3. Major element composition of the calculated melts used to retrieve the equilibrium conditions for clinopyroxene and plagioclase thermobarometry and for K-feldspar hygrometry (see text for explanation). SS = Shoshonitic Silica Saturated; SU = Shoshonitic Silica Undersaturated; Mg# = $Mg/[Mg Fe^{2+}]$ mol%.

Method	Suite	Crystal host rock	Estimated melt composition to attain equilibrium												Mg#	
			SiO ₂ (wt%)	TiO ₂	Al ₂ O ₃	Fe ₂ O ₃	FeO	MnO	MgO	CaO	Na ₂ O	K ₂ O	P ₂ O ₅			
Clinopyroxene-melt thermobarometer	SS	Pyroxenite	50.9	1.2	20.7	1.0	7.0	0.1	4.0	9.3	2.9	2.3	0.6	50.8	Trachybasalt	
	SS	Gabbro	54.9	0.6	21.0	0.8	5.4	0.1	2.1	8.2	3.2	3.2	0.4	41.4	Basaltic trachyandesite	
	SS	Gabbro	64.1	0.6	17.3	0.7	4.5	0.2	1.5	2.7	3.1	5.2	0.2	36.6	Trachyandesite	
	SS	Monzogabbro/monzodiorite	62.6	0.6	17.2	0.6	3.9	0.1	1.5	4.3	3.3	5.6	0.3	40.2	Trachyandesite	
	SU	Gabbro/monzogabbro	62.2	0.3	18.5	0.3	1.7	0.1	0.5	3.1	3.9	9.0	0.6	36.4	Trachyandesite	
	SU	Syenite	63.5	0.3	19.7	0.3	1.8	0.1	0.4	1.4	5.7	7.0	0.0	28.0	Trachyte	
Plagioclase-melt thermobarometer	SS	Pyroxenite	50.0	1.1	17.1	1.3	8.8	0.2	5.1	10.9	2.5	2.1	0.7	50.7	Trachybasalt	
	SS	Gabbro	53.6	0.8	17.4	1.1	7.6	0.2	4.1	8.4	2.8	3.4	0.5	49.3	Basaltic trachyandesite	
	SS	Monzogabbro	57.9	0.6	17.8	0.8	5.1	0.1	2.7	6.6	3.8	4.4	0.1	48.5	Basaltic trachyandesite	
	SS	Monzodiorite/monzonite	62.6	0.6	17.2	0.6	3.9	0.1	1.5	4.3	3.3	5.6	0.3	40.2	Trachyandesite	
	SU	Gabbro	57.8	0.7	20.2	0.6	4.0	0.2	1.5	4.9	4.7	5.1	0.4	39.3	Trachyandesite	
	SU	Monzogabbro to syenite	62.7	0.4	18.4	0.4	2.5	0.1	0.5	3.0	5.3	6.7	0.0	24.1	Trachyte	
K-Feldspar-melt hygrometer	SS	Gabbro/monzogabbro	62.6	0.6	17.2	0.6	3.9	0.1	1.5	4.3	3.3	5.6	0.3	40.2	Trachyandesite	
	SS	Monzodiorite to syenite	64.3	0.3	19.5	0.3	1.8	0.1	0.3	1.4	5.2	7.1	0.0	20.3	Trachyte	
	SU	Monzogabbro/monzodiorite	63.5	0.3	19.7	0.3	1.8	0.1	0.4	1.4	5.7	7.0	0.0	28.0	Trachyte	

Table 4. Pressure, temperature, fO_2 and H_2O parameters obtained by mineral pairs, mineral-melt and single mineral calculations on Shoshonitic Silica Saturated (SS), Shoshonitic Silica Undersaturated (SU) and Granitic Unit (GU) rocks. For each applied method, the corresponding reference, target and error on the single parameters are reported. Values used as input for each calculation are marked with (*); n.r. indicates a parameter that was not required as input by the chosen equation. Cpx = clinopyroxene; Amph = amphibole; Plag = plagioclase; Bt = biotite; K-Feld = K-feldspar; Mt = magnetite.

Suite and sample	Method	Target	Reference	T(°C)	Error (°C)	P (kbar)	Error (kbar)	H ₂ O (wt%)	Error (wt%)	log fO_2	Error (log fO_2)
SS Pyroxenite	Cpx-melt	P,T	Putirka (2008), eq. 32b, 33	1050-1070	±10-20	1.6-2.3	±2.6	2-2.5 (*)	-	n.r.	-
SS Pyroxenite	Cpx-melt	Cpx comp.	Rhyolite-MELTS (Gualda et al. 2012)	1060 (*)	-	1.5 (*)	-	2.5 (*)	-	-9.3 (*)	-
SS Gabbro	Cpx-melt	P,T	Putirka (2008), eq. 32b, 33	910-930°C	±10-20	1.0-1.9	±2.6	3-4 (*)	-	n.r.	-
SS Gabbro	Cpx-melt	P,T	Masotta et al. (2013), eq. Talk and Palk2012	910-980	±18	0.5-1.4	±1.2	3-4 (*)	-	n.r.	-
SS Monzogabbro/monzodiorite	Cpx-melt	P,T	Masotta et al. (2013), eq. Talk and Palk2012	940-990	±18	1.4-1.6	±1.2	3-4 (*)	-	n.r.	-
SU Gabbro/monzogabbro	Cpx-melt	P,T	Putirka (2008), eq. 32b, 33	930-940	±10-20	0.4-1.6	±2.6	1-1.5 (*)	-	n.r.	-
SU Gabbro/monzogabbro	Cpx-melt	P,T	Masotta et al. (2013), eq. Talk and Palk2012	920-940	±18	0.2-1.4	±1.2	1.5 (*)	-	n.r.	-
SU Syenite	Cpx-melt	P,T	Putirka (2008), eq. 32b, 33	810-820	±10-20	0.2-1.4	±2.6	4-5 (*)	-	n.r.	-
SU Syenite	Cpx-melt	P, T	Masotta et al. (2013), eq. Talk and Palk2012	850-860	±18	1.3-1.6	±1.2	4-5 (*)	-	n.r.	-
SS Gabbro to monzodiorite	Amph-Plag	T	Holland and Blundy (1994)	750-845	±40	n.r.	-	n.r.	-	n.r.	-
SS Gabbro to monzodiorite	Amph-Plag	P	Anderson and Smith (1995); Anderson (1996)	n.r.	-	0.1-1.2	±0.6	n.r.	-	n.r.	-
SS Gabbro to monzodiorite	Amph	P,T,H ₂ O	Ridolfi et al. (2010); Ridolfi and Renzulli (2012)	720-810	±23-24	0.7-1.4	±0.05-0.15	4.8-6.0	±0.8	-	-
SU Gabbro/monzogabbro	Amph-Plag	T	Holland and Blundy (1994)	620-880	±40	n.r.	-	n.r.	-	n.r.	-
SU Gabbro/monzogabbro	Amph-Plag	P	Anderson and Smith (1995); Anderson (1996)	n.r.	-	0.4-1.9	±0.6	n.r.	-	n.r.	-
SU Gabbro	Amph	P,T,H ₂ O	Ridolfi et al. (2010); Ridolfi and Renzulli (2012)	700-720	±23-24	0.5-0.6	±0.05-0.15	5.1-5.7	±0.8	-	-
SS Pyroxenite	Bt	T	Henry et al. (2005)	690-740	±12-24	n.r.	-	n.r.	-	n.r.	-
SS Gabbro to monzonite	Bt	T	Henry et al. (2005)	540-660	±12-24	n.r.	-	n.r.	-	n.r.	-
SU Gabbro/monzogabbro	Bt	T	Henry et al. (2005)	640-660	±12-24	n.r.	-	n.r.	-	n.r.	-
SS Pyroxenite	Plag-melt	H ₂ O	Lange et al. (2009)	1060 (*)	-	1.5 (*)	-	3.0	±0.7	n.r.	-
SS Pyroxenite	Plag-melt	P,T	Putirka (2008), eq. 24a	1076-1081	±36	1.5 (*)	-	3.0 (*)	-	n.r.	-
SS Gabbro	Plag-melt	H ₂ O	Lange et al. (2009)	920-1016 (*)	-	1.2-1.5 (*)	-	3.1-3.4	±0.7	n.r.	-
SS Gabbro	Plag-melt	P,T	Putirka (2008), eq. 24a	999-1053	±36	1.2-1.5 (*)	-	3.1-3.4 (*)	-	n.r.	-
SS Monzogabbro/monzodiorite	Plag-melt	H ₂ O	Lange et al. (2009)	920 (*)	-	1.2 (*)	-	3.9-4.8	±0.7	n.r.	-
SS Monzogabbro/monzodiorite	Plag-melt	P,T	Putirka (2008), eq. 24a	917-989	±36	1.2 (*)	-	3.9-4.8 (*)	-	n.r.	-

Table 3. (continued)

Suite and sample	Method	Target	Reference	T(°C)	Error (°C)	P (kbar)	Error (kbar)	H₂O (wt%)	Error (wt%)	logfO₂	Error (log fO₂)
SU Gabbro	Plag-melt	H ₂ O	Lange et al. (2009)	990 (*)	-	1.5 (*)	-	4.1	±0.7	n.r.	-
SU Gabbro	Plag-melt	P,T	Putirka (2008), eq. 24a	975-990	±36	1.5 (*)	-	4.1 (*)	-	n.r.	-
SU Monzogabbro	Plag-melt	H ₂ O	Lange et al. (2009)	920 (*)	-	1.2 (*)	-	4.4	±1.0	n.r.	-
SU Monzogabbro	Plag-melt	P,T	Putirka (2008), eq. 24a	936	±36	1.2 (*)	-	4.4 (*)	-	n.r.	-
SS Gabbro to monzogabbro	K-Feld-melt	H ₂ O	Mollo et al. (2015)	800-900 (*)	-	n.r.	-	4.4-6.5	±0.94	n.r.	-
SS Monzodiorite to syenite	K-Feld-melt	H ₂ O	Mollo et al. (2015)	800-900 (*)	-	n.r.	-	4.6-7.3	±0.94	n.r.	-
SU Monzogabbro to monzonite	K-Feld-melt	H ₂ O	Mollo et al. (2015)	800-900 (*)	-	n.r.	-	5.5-7.8	±0.94	n.r.	-
SS Gabbro	Bt/K-Feld/Mt	fO ₂	Burkhard (1991)	865-966	±7	1.5 (*)	-	n.r.	-	-11.2/-12.9	±0.3
SS Monzogabbro/monzodiorite	Bt/K-Feld/Mt	fO ₂	Burkhard (1991)	806-993	±7	1.5 (*)	-	n.r.	-	-10.7-13.8	±0.3
SS Monzonite	Bt/K-Feld/Mt	fO ₂	Burkhard (1991)	786-881	±7	1.5 (*)	-	n.r.	-	-12.4/-14.1	±0.3
SU Gabbro/monzogabbro	Bt/K-Feld/Mt	fO ₂	Burkhard (1991)	916-1054	±7	1.5 (*)	-	n.r.	-	-10.2/-11.9	±0.3
GU Syenogranite	Bt/K-Feld/Mt	fO ₃	Burkhard (1991)	853-922	±7	1.5 (*)	-	n.r.	-	-11.8/-12.9	±0.3

Table 5. Starting Shoshonitic Silica Saturated (SS) and Shoshonitic Silica Undersaturated (SU) magmas, assimilant end-members and input thermodynamic data used in the EC-AFC model. Calculated and absolute values are from Bohrson and Spera (2001) and Spera and Bohrson (2001) upper crustal case. Specific heat and liquidus T of carbonate assimilant were calculated according to Dallai et al. (2011) and Eppelbaum et al. (2014) respectively. References are also reported for all chosen assimilants compositions.

Starting magma	Liquidus T(°C)	Initial T (°C)	Solidus T(°C)	Specific Heat [J/(KgK)]	Heat of Crystallization (J/Kg)	Heat of Fusion (J/Kg)
SS monzogabbro	1212	1100	600	1567	396000	270000
SU monzogabbro	1251.65	1100	600	1515	396000	270000

Assimilant	Liquidus T(°C)	Initial T (°C)	Specific Heat [J/(KgK)]	Reference for isotopic $^{87}\text{Sr}/^{86}\text{Sr}$ and $^{143}\text{Nd}/^{144}\text{Nd}$ composition
Carbonate	650	300	1170	Martin and Macdougall (1995); Blendiger et al. (2015)
Rhyolitic Ignimbrite (AVD)	1077	300	1376	Barth et al. (1993)
Mt. Croce Granodiorites	1111	300	1398	Rottura et al. (1997)
Serie dei Laghi Granites	1026	300	1379	Sinigoï et al. (2016) and references therein
Kinzigite Amphibolite	1196	300	1447	Voshage et al. (1990)