

Invariant submanifolds of metric contact pairs

Amine Hadjar¹ · Paola Piu²

Received: 20 July 2015 / Accepted: 9 April 2016 / Published online: 25 April 2016
© Fondazione Annali di Matematica Pura ed Applicata and Springer-Verlag Berlin Heidelberg 2016

Abstract We show that ϕ -invariant submanifolds of metric contact pairs with orthogonal characteristic foliations make constant angles with the Reeb vector fields. Our main result is that for the normal case such submanifolds of dimension at least 2 are all minimal. We prove that an odd-dimensional ϕ -invariant submanifold of a metric contact pair with orthogonal characteristic foliations inherits a contact form with an almost contact metric structure, and this induced structure is contact metric if and only if the submanifold is tangent to one Reeb vector field and orthogonal to the other one. Furthermore, we show that the leaves of the two characteristic foliations of the differentials of the contact pair are minimal. We also prove that when one Reeb vector field is Killing and spans one characteristic foliation, the metric contact pair is locally the product of a contact metric manifold with \mathbb{R} .

Keywords Metric contact pair · Minimal invariant submanifold · Metric f -structure · Almost contact metric manifold

Mathematics Subject Classification 53C25 · 53B20 · 53D10 · 53B35 · 53C12

1 Introduction

On a Riemannian manifold endowed with a tensor field φ of type $(1, 1)$, a submanifold is said to be φ -invariant (or invariant) when its tangent bundle is preserved by φ . Under some compatibility conditions between φ and the metric, as is for example the case for almost

✉ Paola Piu
piu@unica.it
Amine Hadjar
mohamed.hadjar@uha.fr

¹ Laboratoire de Mathématiques, Informatique et Applications, Université de Haute Alsace, 4, Rue des Frères Lumière, 68093 Mulhouse Cedex, France

² Dipartimento di Matematica e Informatica, Università degli studi di Cagliari, Via Ospedale 72, 09124 Cagliari, Italy

Hermitian manifolds, the questions concerning the minimality of the submanifold and the nature of the induced structure are natural and interesting. It is well known that invariant submanifolds of a contact metric manifold are minimal, and the same holds for those of a Kähler manifold. However, Vaisman in [20] proved that a J -invariant submanifold of a Vaisman manifold (J being the complex structure) is minimal if and only if it inherits a Vaisman structure. These manifolds are a subclass of locally conformally Kähler (lcK) manifolds. Dragomir and Ornea in [13] generalized this result by showing that a J -invariant submanifold of an lcK manifold is minimal if and only if the submanifold is tangent to the Lee vector field (and therefore tangent to the anti-Lee vector field). By a statement of Bande and Kotschick [8], normal metric contact pairs of type $(h, 0)$ are nothing but non-Kähler Vaisman manifolds, so the results of Vaisman and Dragomir–Ornea apply to these manifolds.

A metric contact pair is a manifold of even dimension endowed with a special case of a metric f -structure with two complemented frames in the sense of Yano [21]. It carries an endomorphism field ϕ of corank 2 and two natural commuting almost complex structures J and T of opposite orientations. When J and T are both integrable, the structure is said to be normal. For J -invariant or T -invariant submanifolds on a normal metric contact pair with decomposable ϕ , the problem was solved by Bande and the first author [7] by proving that these submanifolds are minimal if and only if they are tangent to both the two Reeb vector fields of the structure, and they gave an example of such submanifold on which the contact pair of the ambient manifold does not induce a contact pair. Regarding the ϕ -invariant case, they gave partial results.

In this paper, we study ϕ -invariant submanifolds of a metric contact pair $(M, \alpha_1, \alpha_2, \phi, g)$ of any type (h, k) with decomposable ϕ (or equivalently with orthogonal characteristic foliations), by looking at the angles that Reeb vector fields make with these submanifolds. We prove that these two angles are constant; then, we solve completely the problem of minimality and that of induced structures. First we show the following.

A connected ϕ -invariant submanifold N of a metric contact pair with decomposable ϕ and Reeb vector fields Z_1 and Z_2 satisfies one of the following properties:

- N is even-dimensional and tangent to both Z_1 and Z_2 .
- N is one-dimensional and contained in one of the two-dimensional leaves of the vertical foliation spanned by Z_1 and Z_2 .
- N is of odd dimension ≥ 3 everywhere tangent to one Reeb vector field Z_1 and orthogonal to the other one Z_2 , or vice versa.
- N is of odd dimension ≥ 3 , nowhere tangent and nowhere orthogonal to Z_1 and Z_2 making two constant angles with them. In this case, the dimensions of the characteristic foliations are $\geq \dim N$.

For the minimality problem, as a consequence we prove the following.

Any ϕ -invariant submanifold of dimension ≥ 2 of a normal metric contact pair with decomposable ϕ is minimal.

The one-dimensional case is obvious since it concerns vertical geodesics, i.e., those which are integral curves of $c_1 Z_1 + c_2 Z_2$ with c_i constant functions.

Now let us return to the question concerning the induced structure on a ϕ -invariant submanifold N of a metric contact pair. When N is even-dimensional, it is tangent to the Reeb vector fields and then it is J -invariant (and T -invariant). Although it does not always inherit a contact pair [7], we observe that it carries at least a metric f -structure with two complemented frames. By normality on the ambient manifold, we get a \mathcal{K} -structure (see [10] for a definition) on N .

For the case N of odd dimension along which the Reeb vector fields are nowhere tangent and nowhere orthogonal, when ϕ is decomposable, we prove that N carries a contact form and an almost contact metric structure which is not contact metric, while for the other cases (N tangent to one Reeb vector field and orthogonal to the other one) it has been shown in [11] that N is a contact metric submanifold of one of the characteristic leaves of the contact pair. By normality on the ambient manifold, the induced structure on N will be normal.

Furthermore, we prove that the leaves of the characteristic foliations of $d\alpha_1$ and $d\alpha_2$ of a metric contact pair $(M, \alpha_1, \alpha_2, \phi, g)$ with decomposable ϕ are minimal. Their leaves are ϕ -invariant and tangent to both the two Reeb vector fields, but for our proof the integrability of J or T is not needed. We give an example where one of these two foliations is not totally geodesic. When the type numbers of the contact pair are $(h, 0)$, i.e., when the Reeb vector field Z_2 spans the characteristic distribution of α_1 , we prove the following.

If Z_2 is Killing, then the two characteristic foliations of the contact pair are totally geodesic, and then, the metric contact pair is locally product of a contact metric manifold with \mathbb{R} . By normality, the first factor will be Sasakian.

2 Preliminaries

Blair et al. [11] introduced in Hermitian geometry the notion of *bicontact structure*. This topic was formulated again as *contact pair* by Bande in his PhD thesis in 2000 and then together with the first author in [3]. A pair of 1-forms (α_1, α_2) on a manifold M is said to be a *contact pair* of type (h, k) if

$$\begin{aligned} \alpha_1 \wedge (d\alpha_1)^h \wedge \alpha_2 \wedge (d\alpha_2)^k & \text{ is a volume form,} \\ (d\alpha_1)^{h+1} = 0 \quad \text{and} \quad (d\alpha_2)^{k+1} = 0. \end{aligned}$$

The latter two conditions guarantee the integrability of the two subbundles of the tangent bundle

$$T\mathcal{F}_i = \{X : \alpha_i(X) = 0, d\alpha_i(X, Y) = 0 \forall Y\}, \quad i = 1, 2.$$

They determine the *characteristic foliations* \mathcal{F}_1 of α_1 and \mathcal{F}_2 of α_2 which are transverse and complementary. The leaves of \mathcal{F}_1 and \mathcal{F}_2 are contact manifolds of dimension $2k + 1$ and $2h + 1$, respectively, with contact forms induced by α_2 and α_1 (see [3]). We also define the $(2h + 2k)$ -dimensional *horizontal subbundle* \mathcal{H} to be the intersection of the kernels of α_1 and α_2 .

The equations

$$\begin{aligned} \alpha_1(Z_1) = \alpha_2(Z_2) = 1, \quad \alpha_1(Z_2) = \alpha_2(Z_1) = 0, \\ i_{Z_1}d\alpha_1 = i_{Z_1}d\alpha_2 = i_{Z_2}d\alpha_1 = i_{Z_2}d\alpha_2 = 0, \end{aligned}$$

where i_X is the contraction with the vector field X , determine uniquely the two vector fields Z_1 and Z_2 , called *Reeb vector fields*. Since they commute [3], they give rise to a locally free \mathbb{R}^2 -action, an integrable distribution called *Reeb distribution*, and then a foliation \mathcal{V} of M by surfaces. The subbundle $T\mathcal{V} = \mathbb{R}Z_1 \oplus \mathbb{R}Z_2$ is called the *vertical subbundle* and the tangent bundle of M splits as:

$$TM = T\mathcal{F}_1 \oplus T\mathcal{F}_2 = \mathcal{H} \oplus \mathbb{R}Z_1 \oplus \mathbb{R}Z_2.$$

A *contact pair structure* [4] on a manifold M is a triple $(\alpha_1, \alpha_2, \phi)$, where (α_1, α_2) is a contact pair and ϕ a tensor field of type $(1, 1)$ such that:

$$\phi^2 = -Id + \alpha_1 \otimes Z_1 + \alpha_2 \otimes Z_2, \quad \phi Z_1 = \phi Z_2 = 0.$$

The rank of ϕ is $\dim M - 2$ and $\alpha_i \circ \phi = 0$, for $i = 1, 2$.

The endomorphism ϕ is said to be *decomposable* if $\phi(T\mathcal{F}_i) \subset T\mathcal{F}_i$, for $i = 1, 2$. This is a natural condition that allows to have on each leaf of \mathcal{F}_1 and \mathcal{F}_2 an induced almost contact structure.

In [5], the notion of *normality* for a contact pair structure is defined as the integrability of both of the two natural commuting almost complex structures of opposite orientations $J = \phi - \alpha_2 \otimes Z_1 + \alpha_1 \otimes Z_2$ and $T = \phi + \alpha_2 \otimes Z_1 - \alpha_1 \otimes Z_2$ on M . This is equivalent to the vanishing of the tensor field

$$N^1(X, Y) = [\phi, \phi](X, Y) + 2d\alpha_1(X, Y)Z_1 + 2d\alpha_2(X, Y)Z_2$$

where $[\phi, \phi]$ is the Nijenhuis tensor of ϕ .

A Riemannian metric g on a manifold endowed with a contact pair structure is said to be *associated* [4] if

$$g(X, \phi Y) = (d\alpha_1 + d\alpha_2)(X, Y) \quad \text{and} \quad g(X, Z_i) = \alpha_i(X) \quad \text{for } i = 1, 2.$$

Such a metric is necessarily *compatible* with respect to the contact pair structure, which means that

$$g(\phi X, \phi Y) = g(X, Y) - \alpha_1(X)\alpha_1(Y) - \alpha_2(X)\alpha_2(Y).$$

Moreover, the subbundles $\mathbb{R}Z_1, \mathbb{R}Z_2$ and \mathcal{H} are pairwise orthogonal.

A *metric contact pair* on a manifold M is a four-tuple $(\alpha_1, \alpha_2, \phi, g)$ where $(\alpha_1, \alpha_2, \phi)$ is a contact pair structure and g an associated metric with respect to it. Such a manifold M will also be called a *metric contact pair* for short.

On a metric contact pair, the endomorphism field ϕ is decomposable if and only if the characteristic foliations $\mathcal{F}_1, \mathcal{F}_2$ are orthogonal [4]. In this case, the leaves of \mathcal{F}_j are minimal submanifolds [6]; they carry contact metric structures induced by (ϕ, Z_j, α_j, g) , for $j \neq i$ (see [4]), and by the normality they become Sasakian [5]. Of course, the product of two contact metric manifolds (or a contact metric manifold with \mathbb{R}) gives rise to a metric contact pair with decomposable endomorphism, and the structure is normal if and only if the two factors are Sasakian. It is important to note that there exist metric contact pairs with decomposable ϕ which are not locally products of contact metric manifolds as shown in the following example (see also [6] for a similar construction).

Example 1 Consider the simply connected six-dimensional nilpotent Lie group G^6 with structure equations:

$$\begin{aligned} d\alpha_1 &= \alpha_3 \wedge \alpha_5, & d\alpha_2 &= \alpha_4 \wedge \alpha_6, & d\alpha_6 &= \alpha_4 \wedge \alpha_5, \\ d\alpha_3 &= d\alpha_4 = d\alpha_5 = 0 \end{aligned}$$

The pair (α_1, α_2) is a contact pair of type $(1, 1)$ with Reeb vector fields (Y_1, Y_2) , the Y_i 's being dual to the α_i 's. The characteristic distribution of α_2 (respectively, α_1) is spanned by Y_4, Y_6 and Y_2 (respectively, Y_3, Y_5 and Y_1). Take the metric

$$g = \alpha_1^2 + \alpha_2^2 + \frac{1}{2} (\alpha_3^2 + \alpha_4^2 + \alpha_5^2 + \alpha_6^2)$$

and the decomposable endomorphism ϕ defined to be zero on Y_1, Y_2 , and

$$\phi Y_5 = Y_3, \quad \phi Y_3 = -Y_5, \quad \phi Y_6 = Y_4, \quad \phi Y_4 = -Y_6$$

Then $(\alpha_1, \alpha_2, \phi, g)$ is a left invariant metric contact pair on the Lie group G^6 . We can easily see that the leaves of the two characteristic foliations are Sasakian, though the metric contact pair is not normal because $N^1(Y_3, Y_4) = [\phi, \phi](Y_3, Y_4) = Y_4 \neq 0$. We also remark that the characteristic foliation of α_1 is totally geodesic, while from $g(\nabla_{Y_4} Y_6, Y_5) = -\frac{1}{2}$ follows that the characteristic foliation of α_2 is not totally geodesic.

Since the structure constants of the nilpotent Lie algebra of G^6 are rational, there exist cocompact lattices Γ of G^6 . Now the metric contact pair on G^6 descends to all these quotients G^6/Γ and we obtain closed nilmanifolds carrying the same type of structure. All the remarks concerning the structure we constructed on G^6 still remain valid on the metric contact pairs G^6/Γ .

For the normal case, we can give the following.

Example 2 As a manifold consider the product $H^6 = \mathbb{H}_3 \times \mathbb{H}_3$ where \mathbb{H}_3 is the three-dimensional Heisenberg group. Let $\{\alpha_1, \alpha_2, \alpha_3\}$ (respectively, $\{\beta_1, \beta_2, \beta_3\}$) be a basis of the cotangent space at the identity for the first (respectively, second) factor \mathbb{H}_3 satisfying

$$\begin{aligned} d\alpha_3 &= \alpha_1 \wedge \alpha_2, & d\alpha_1 &= d\alpha_2 = 0, \\ d\beta_3 &= \beta_1 \wedge \beta_2, & d\beta_1 &= d\beta_2 = 0. \end{aligned}$$

The pair (α_3, β_3) determines a contact pair of type $(1, 1)$ on H^6 with Reeb vector fields (X_3, Y_3) , the X_i 's (respectively, the Y_i 's) being dual to the α_i 's (respectively, the β_i 's). The left invariant metric

$$g = \alpha_3^2 + \beta_3^2 + \frac{1}{2}(\alpha_1^2 + \beta_1^2 + \alpha_2^2 + \beta_2^2)$$

is associated to the pair with decomposable endomorphism ϕ given by $\phi(X_2) = X_1$ and $\phi(Y_2) = Y_1$. The metric contact pair $(H^6, \alpha_3, \beta_3, \phi, g)$ is normal because it is the product of two Sasakian manifolds. Also here H^6 admits cocompact lattices Γ and the structure descends to the nilmanifolds H^6/Γ as normal metric contact pairs.

Some other interesting examples and properties of such structures were given in [2–7]. In the following remark, we describe how metric contact pairs relate to other well-known structures.

Remark 1 Normal metric contact pairs with decomposable endomorphism were already studied in [11] under the name bicontact Hermitian manifolds of bidegree $(1, 1)$. They were regarded as a generalization of the Calabi–Eckmann manifolds. A metric contact pair of type (h, k) is a special case of metric f -structure of rank $2h + 2k$ with two complemented frames in the sense of Yano [21]. The normality condition of metric contact pairs, i.e., the integrability of both almost complex structures J and T is equivalent to the normality condition as an f -structure which consists exactly on the vanishing of the tensor field N^1 described before. We can also observe that a normal metric contact pair is a special case of \mathcal{K} -structures in the sense of Blair, Ludden and Yano [10]. Normal metric contact pairs of type $(h, 0)$ are necessarily \mathcal{S} -structures (see [10] for a definition), but those of type (h, k) with $h > 0$ and $k > 0$ are not. Furthermore, it has been shown in [8] that normal metric contact pairs of type $(h, 0)$ are nothing but non-Kähler Vaisman manifolds (called \mathcal{PK} -manifolds in [19]). The \mathcal{P} -manifolds of Vaisman [19] are necessarily metric contact pairs of type $(h, 0)$ where the Reeb vector fields are Killing, and they include the subclass of \mathcal{PK} -manifolds.

In the course of our work, we will need the following lemmas.

Lemma 1 (Bande et al. [2]) *On a manifold endowed with a contact pair structure with decomposable ϕ and a compatible metric, for every X we have that $\nabla_X Z_1$ and $\nabla_X Z_2$ are horizontal.*

Lemma 2 *On a manifold endowed with a contact pair structure with decomposable ϕ and a compatible metric, for every X horizontal we have that $\nabla_{Z_1} X$ and $\nabla_{Z_2} X$ are horizontal.*

Proof By the previous lemma, $\alpha_j(\nabla_X Z_i) = 0$ for $i, j = 1, 2$, giving

$$\begin{aligned} \alpha_j(\nabla_{Z_i} X) &= \alpha_j([Z_i, X]) \\ &= Z_i \alpha_j(X) - X \alpha_j(Z_i) - 2d\alpha_j(Z_i, X) \\ &= 0. \end{aligned}$$

□

3 ϕ -Invariant submanifolds

A submanifold N of a metric contact pair is said to be ϕ -invariant if its tangent bundle TN is preserved by the endomorphism field ϕ . We will denote by Z_i^T (respectively, Z_i^\perp) the tangential (respectively, normal) component of the two Reeb vector fields Z_1 and Z_2 along N . In the following proposition, we recall some properties from [7] we need and concerning the positions of the Reeb vector fields along a ϕ -invariant submanifold N .

Proposition 1 (Bande et al. [7])

1. *Along the ϕ -invariant submanifold N the four sections Z_1^T, Z_2^T, Z_1^\perp and Z_2^\perp are vertical.*
2. *There is no point p of N such that the tangent vectors $(Z_1)_p$ and $(Z_2)_p$ are both orthogonal to the tangent space $T_p N$.*
3. *If at a point p of N one Reeb vector field is tangent to N and the second one is transverse, then the second one is orthogonal to N at p .*

To this, we can add the following.

Proposition 2 *If at a point p of the ϕ -invariant submanifold N one of the Reeb vector fields is orthogonal to N , then the second one is tangent to N at p .*

Proof Suppose that $(Z_1)_p$ is orthogonal to $T_p N$. First by Proposition 1-(2) we have $(Z_2^T)_p \neq 0$. Moreover, $(Z_2^T)_p$ is orthogonal to $(Z_1)_p$. Next by Proposition 1-(1) the vector $(Z_2^T)_p$ lies in the plane $((Z_1)_p, (Z_2)_p)$ and we get $(Z_2^T)_p = (Z_2)_p$. □

Now after these observations, we can state the following proposition.

Proposition 3 *A ϕ -invariant submanifold which is tangent to both the Reeb vector fields has even dimension. Otherwise, its dimension is odd and its tangent bundle intersects the Reeb distribution along a line bundle.*

Indeed, it is clear that when a submanifold is tangent to both Z_1 and Z_2 at a point, since its tangent space at that point is preserved by ϕ it is also preserved by J which is almost complex. Thus, the dimension is even.

Now let us describe a little more a ϕ -invariant submanifold N which is not tangent to the Reeb distribution. We have seen that the four sections Z_1^T, Z_1^\perp, Z_2^T and Z_2^\perp are vertical.

When at a point p of N one of the two Reeb vector fields is not tangent to N , the second one is not orthogonal to the submanifold N (Proposition 2), and this occurs on a whole open set of N . Moreover, at those points both families $\{Z_1^T, Z_2^T\}$ and $\{Z_1^\perp, Z_2^\perp\}$ have rank one. Any tangent vector of N at p which is orthogonal to $\{Z_1^T, Z_2^T\}$ is horizontal, since it is orthogonal to $\{Z_1^\perp, Z_2^\perp\}$ too and then orthogonal to $\{Z_1, Z_2\}$. Then by the ϕ -invariance of N , ϕ is almost complex when acting on horizontal vectors tangent to N , i.e., on the horizontal part $\mathcal{H} \cap TN$ of TN . This explains the odd dimension of N and the fact that the Reeb distribution still remains nowhere tangent to the submanifold. Notice that $\{Z_1^T, Z_2^T\}$ spans a line bundle which is the intersection of the tangent bundle TN with the Reeb distribution that can be called the vertical part of TN . Moreover, we have

$$TN = (T\mathcal{V} \cap TN) \oplus (\mathcal{H} \cap TN).$$

When N is a one-dimensional ϕ -invariant submanifold, then it is any one-dimensional submanifold of any two-dimensional leaf of the vertical foliation \mathcal{V} . So in the sequel, we will suppose that the dimension of N is at least 3.

For simplicity, suppose that Z_2 is nowhere tangent to N , so that Z_1 is nowhere orthogonal to N (by Proposition 2). Normalizing Z_1^T , we get on N a unit vector field spanning the vertical part $T\mathcal{V} \cap TN$ of TN :

$$\zeta = \frac{1}{\|Z_1^T\|} Z_1^T.$$

Now along N , the equation

$$\zeta = (\cos \theta_1)Z_1 + (\sin \theta_1)Z_2$$

defines a smooth function θ_1 on N taking values in $] -\pi/2, \pi/2[$. This function is the measure of the angle that Z_1 makes with the submanifold, i.e., the oriented angle (Z_1, Z_1^T) in the plane (Z_1, Z_2) oriented by the almost complex structure J . When Z_1 is nowhere tangent to the submanifold, similarly we get a function θ_2 measuring the oriented angle (Z_2, Z_2^T) that is the angle between Z_2 and the submanifold, and satisfying $Z_2^T / \|Z_2^T\| = (\cos \theta_2)Z_2 + (\sin \theta_2)(-Z_1)$ with $-\pi/2 < \theta_2 < \pi/2$.

We will say that the submanifold N is *leaning* when both Reeb vector fields are leaning along N , i.e., when they are nowhere tangent and nowhere orthogonal to N . This means that the functions θ_1 and θ_2 are well defined and take nonvanishing values.

Theorem 1 *Let $(M, \alpha_1, \alpha_2, \phi, g)$ be a metric contact pair of type (h, k) with decomposable ϕ . Suppose that M carries a leaning ϕ -invariant submanifold N of odd dimension $2n + 1 \geq 3$. Then,*

1. *for the angle (Z_1, Z_1^T) , we have $0 < \theta_1 < \pi/2$,*
2. *for the angle (Z_2, Z_2^T) , we have $\theta_2 = \theta_1 - \pi/2$,*
3. *each nonzero horizontal tangent vector X of N decomposes as $X = X_1 + X_2$, X_i being nonzero horizontal vector tangent to the characteristic foliation \mathcal{F}_j for $j \neq i$, with $\|X_2\|^2 / \|X_1\|^2 = \tan \theta_1$,*
4. *for the type numbers, we have $h \geq n$ and $k \geq n$, i.e., the dimensions of the two characteristic foliations are at least $2n + 1$.*

In this theorem, Property 3 means that X makes with $T\mathcal{F}_2$ (or more precisely with $\mathcal{H} \cap T\mathcal{F}_2$) an angle of absolute value $\theta'_1 = \arctan \sqrt{\tan \theta_1}$, and of course with $\mathcal{H} \cap T\mathcal{F}_1$ an angle $\theta'_2 = \pi/2 - \theta'_1$. Property 1 states that along N the vertical part of TN separates the vertical

plane (Z_1, Z_2) into two half-planes, each one containing one Reeb vector field, and for ζ we also have

$$\zeta = \frac{1}{\|Z_2^T\|} Z_2^T = (\cos \theta_2)Z_2 + (\sin \theta_2)(-Z_1).$$

Proof Take any local horizontal nonvanishing vector field X of N . Such a vector field always exists since the dimension of N is at least 3. Along N , it decomposes as $X = X_1 + X_2$ with X_i horizontal and tangent to \mathcal{F}_j , for $j \neq i$. By the ϕ -invariance of N , ϕX is also tangent to N , and ϕX_i is horizontal and tangent to \mathcal{F}_j , for $j \neq i$, by decomposability of ϕ . In order to compute the vertical part of $[X, \phi X]$, we have

$$g([X, \phi X], Z_1) = \alpha_1([X_1, \phi X_1]) + \alpha_1([X_1, \phi X_2]) + \alpha_1([X_2, \phi X_1]) + \alpha_1([X_2, \phi X_2]).$$

For the first term, we have $\alpha_1([X_1, \phi X_1]) = -2d\alpha_1(X_1, \phi X_1) = 2g(X_1, X_1)$, and the last three terms vanish. Thus, we have

$$g([X, \phi X], Z_1) = 2\|X_1\|^2$$

and similarly

$$g([X, \phi X], Z_2) = 2\|X_2\|^2$$

so that the vertical part of $[X, \phi X]$ is the vector field $2\|X_1\|^2 Z_1 + 2\|X_2\|^2 Z_2$.

Since $[X, \phi X]$ is also tangent to N , its vertical part is then collinear with $\zeta = (\cos \theta_1)Z_1 + (\sin \theta_1)Z_2$. Now from $\cos \theta_1 > 0$ and $\sin \theta_1 \neq 0$, we obtain that $\sin \theta_1 > 0$, i.e., $0 < \theta_1 < \pi/2$, and $\theta_2 = \theta_1 - \pi/2$. Moreover, X_1, X_2 do not vanish. For the measure of the angle (X, X_1) , we obtain its tangent which is $\|X_2\|/\|X_1\| = \sqrt{\tan \theta_1}$.

Regarding the dimensions of the characteristic foliations, take at any point p of N a basis $\{e_1, \dots, e_{2n}\}$ of the horizontal part $\mathcal{H} \cap T_p N$ of the tangent space $T_p N$. Each vector e_l being horizontal decomposes as $e_l = e_{l1} + e_{l2}$ with $e_{li} \in \mathcal{H} \cap T_p \mathcal{F}_j$ for $j \neq i$. Let $\lambda_1, \dots, \lambda_{2n}$ any real numbers such that $\sum_{l=1}^{2n} \lambda_l e_{l1} = 0$. Put $X = \sum_{l=1}^{2n} \lambda_l e_l$. Then $X = X_1 + X_2$ with $X_i = \sum_{l=1}^{2n} \lambda_l e_{li}$ lying in $T_p \mathcal{F}_j$ for $j \neq i$. Applying Property 3 of this theorem to the horizontal vector X , we get $X = 0$ since X_1 is supposed to be zero. Hence, from $\sum_{l=1}^{2n} \lambda_l e_l = 0$ we obtain that $\lambda_l = 0$ for all l . Finally the vectors e_{11}, \dots, e_{2n-1} are linearly independent in $\mathcal{H} \cap T_p \mathcal{F}_2$ and then $h \geq n$. In the same way, we get $k \geq n$, and this completes the proof. \square

Example 3 Consider the metric contact pairs on the nilpotent Lie group G^6 and its closed nilmanifolds G^6/Γ described in Example 1. For any two arbitrary nonzero real numbers a and b , the three vectors

$$X = aY_3 + bY_6, \quad \phi X = -aY_5 + bY_4 \quad \text{and} \quad Z = [X, \phi X] = a^2Y_1 + b^2Y_2$$

span a ϕ -invariant subalgebra of the Lie algebra of G^6 which determines a three-dimensional foliation \mathcal{N} in G^6 (and also in the nilmanifolds G^6/Γ). Each leaf N of \mathcal{N} is ϕ -invariant, leaning, minimal and non totally geodesic. The vertical part of TN is spanned by Z restricted to N , and the angle θ_1 that the Reeb vector field Y_1 makes with N satisfies

$$\cos \theta_1 = a^2/\sqrt{a^4 + b^4} \quad \text{and} \quad \sin \theta_1 = b^2/\sqrt{a^4 + b^4}.$$

By a suitable choice of a and b , we can see that the angle θ_1 can take any value in $]0, \pi/2[$.

In the same way, we can have ϕ -invariant submanifolds on the normal metric contact pairs H^6 and its nilmanifolds described in Example 2.

Example 4 Take again any nonzero real numbers a and b . The three vectors $X = aX_2 + bY_1$, $\phi X = aX_1 - bY_2$ and $Z = [X, \phi X] = a^2X_3 + b^2Y_3$ span a ϕ -invariant subalgebra of the Lie algebra of H^6 which determines a three-dimensional foliation \mathcal{N} in H^6 (and also in each nilmanifold H^6/Γ). Each leaf is ϕ -invariant, leaning and totally geodesic. Moreover, the angle that the Reeb vector field X_3 makes with the leaf has tangent equal to b^2/a^2 .

In order to obtain an example of a closed ϕ -invariant leaning submanifold, we have just to choose suitably a lattice of H^6 . Indeed let L_e be the leaf passing through the identity element of the Lie group H^6 . We can see that the Lie subgroup L_e is nothing but the Heisenberg group that admits cocompact lattices (see, e.g., [7] or [15] to get an explicit one). Take any of such lattices which we will call Γ . Because L_e is a subgroup of H^6 , we have that Γ is also a lattice of H^6 . Now the closed nilmanifold $N^3 = L_e/\Gamma$ of L_e is a submanifold of the nilmanifold $M^6 = H^6/\Gamma$ of H^6 . As explained before since the normal contact pair on H^6 is left invariant, it descends to the quotient M^6 as a normal metric contact pair $(\tilde{\alpha}_3, \tilde{\beta}_3, \tilde{\phi}, \tilde{g})$ of type $(1, 1)$ with decomposable endomorphism $\tilde{\phi}$. Finally we obtain a normal metric contact pair M^6 with a decomposable endomorphism, carrying a closed leaning $\tilde{\phi}$ -invariant submanifold N^3 .

Before stating our main theorem concerning minimality of ϕ -invariant submanifolds on normal metric contact pairs, we first give an important lemma which allows us to describe more the angles between odd-dimensional ϕ -invariant submanifold and the Reeb vector fields.

Lemma 3 *Let $(M, \alpha_1, \alpha_2, \phi, g)$ be a metric contact pair with decomposable ϕ , and N a connected ϕ -invariant submanifold of odd dimension ≥ 3 . If at a point of N the Reeb vector field Z_2 is not tangent to N , then Z_2 is everywhere transverse to N , the measure θ_1 of the oriented angle that Z_1 makes with N is constant and $0 \leq \theta_1 < \pi/2$.*

Actually for the leaning case the four angles $\theta_1, \theta_2, \theta'_1$ and θ'_2 described above are all constant. For $\theta_1 = 0$, at each point Z_1 is tangent and Z_2 is orthogonal to the submanifold. As a matter of fact, Z_2 must be transverse to N because of the odd dimension and then orthogonal to N by Proposition 1-(3). For example, this is the case for the leaves of the characteristic foliation \mathcal{F}_2 .

Proof The proof will be given in two stages. Let N' be the nonempty open set of points of N on which Z_2 is transverse to N . The function θ_1 is well defined on N' and we have obviously $0 \leq \theta_1 < \pi/2$. We will first prove that θ_1 is locally constant on N' (Stage 1). Next we will prove that N' is nothing but N (Stage 2).

Stage 1, step 1:

Let X be any (local) unit vector field of N' orthogonal to ζ . Then it is horizontal because it is also orthogonal to $J\zeta$. Since ζ is unit, $g(\nabla_X \zeta, \zeta) = 0$. We also have

$$g(\nabla_X \zeta, J\zeta) = g(\nabla_X X, J\zeta) + g([X, \zeta], J\zeta) = 0$$

since $\nabla_X X$ is horizontal by Lemma 2, and $[X, \zeta]$ is tangent to N' . Because ζ and $J\zeta$ span the vertical bundle, $\nabla_X \zeta$ is horizontal along N' .

Stage 1, step 2:

Differentiating $\zeta = (\cos \theta_1)Z_1 + (\sin \theta_1)Z_2$, we obtain

$$\nabla_X \zeta = X(\theta_1)J\zeta + \cos \theta_1 \nabla_X Z_1 + \sin \theta_1 \nabla_X Z_2. \tag{1}$$

By step 1 and Lemma 1, we have that $\nabla_X \zeta, \nabla_X Z_1$ and $\nabla_X Z_2$ are horizontal; then, (1) implies that along N'

$$X(\theta_1) = 0.$$

Stage 1, step 3:

Take an X as above. The dimension of N' being ≥ 3 such an X always exists. By the ϕ -invariance of N' , ϕX and $[X, \phi X]$ are also tangent to the submanifold. Using the fact that X and ϕX are horizontal, we have

$$g(Z_1 + Z_2, [X, \phi X]) = (\alpha_1 + \alpha_2)([X, \phi X]) = -2(d\alpha_1 + d\alpha_2)(X, \phi X) = 2g(X, X) = 2.$$

Replacing $Z_1 + Z_2$ by $(\cos \theta_1 + \sin \theta_1)\zeta + (\cos \theta_1 - \sin \theta_1)J\zeta$ and using $0 \leq \theta_1 < \pi/2$ we obtain

$$g([X, \phi X], \zeta) = \frac{2}{\cos \theta_1 + \sin \theta_1}$$

because $J\zeta$ is orthogonal to the submanifold. Then we have

$$[X, \phi X] = \frac{2}{\cos \theta_1 + \sin \theta_1} \zeta + Y \tag{2}$$

for some Y horizontal and tangent to the submanifold. By step 1, since $X, \phi X$ and Y are horizontal and tangent to N' , we get $X(\theta_1) = \phi X(\theta_1) = 0$, then $[X, \phi X](\theta_1) = 0$ and also $Y(\theta_1) = 0$. Hence, (2) implies that along N'

$$\zeta(\theta_1) = 0.$$

Finally by steps 2 and 3 the function θ_1 is locally constant on the open set N' .

Stage 2:

Let N'' be the complement set of the open set N' in N , that is, the set of the points of N where Z_2 is tangent to N . At these points, Z_1 is not tangent to N , and the function θ_2 is well defined on an open set of N containing N'' . By the same arguments as before, θ_2 is locally constant, giving that N'' is open because it consists on the vanishing points of θ_2 . By the connectedness of N , we have that N'' is empty and

$$N = N'.$$

Now the function θ_1 is well defined and constant on the whole N , and of course Z_2 is everywhere transverse to the submanifold, completing the proof. □

A first immediate consequence of our Lemma 3, using Proposition 3, is the following theorem which describes all possible relative positions of a ϕ -invariant submanifold with respect to both the Reeb vector fields.

Theorem 2 *Let N be a connected ϕ -invariant submanifold of a metric contact pair with decomposable ϕ . Then N satisfies one of the following properties:*

1. N is even-dimensional and tangent to both Reeb vector fields.
2. N is one-dimensional and contained in one of the two-dimensional leaves of the vertical foliation.
3. N is of odd dimension ≥ 3 everywhere tangent to one Reeb vector field Z_1 and orthogonal to the other one Z_2 , or vice versa.
4. N is of odd dimension ≥ 3 , nowhere tangent and nowhere orthogonal to the Reeb vector fields making two constant angles with them.

Remark 2 Except the one-dimensional case, a ϕ -invariant submanifold of a metric contact pair with decomposable ϕ always makes a constant angle with each of the two Reeb vector fields.

4 Minimality

We now turn to our main result on minimality of ϕ -invariant submanifolds of normal metric contact pairs with orthogonal characteristic foliations.

Theorem 3 *Any ϕ -invariant submanifold of dimension ≥ 2 of a normal metric contact pair with decomposable ϕ is minimal.*

Proof Consider a connected ϕ -invariant submanifold N of a normal metric contact pair with decomposable ϕ . When the dimension of N is ≥ 2 , N satisfies one of the cases (1), (3) or (4) enumerated in Theorem 2. Take then the question case by case and use some partial results from [7] to conclude. Assume that N has even dimension (Case 1). Then it is tangent to both the Reeb vector fields and one can readily show that N is also J -invariant. Moreover, the normality of the metric contact pair implies the integrability of J . By Bande and Hadjar [7], when J is integrable a J -invariant submanifold is minimal if and only if it is tangent to the Reeb distribution. So this applies to N and then N is minimal.

Another result from [7] states that, on a normal metric contact pair with decomposable ϕ , a ϕ -invariant submanifold tangent to one Reeb vector field and orthogonal to the other one (Case 3) is minimal.

For the very remaining possible case (Case 4), i.e., when N is leaning, we use the following. On a normal metric contact pair with decomposable ϕ , a ϕ -invariant leaning submanifold is minimal if and only if the angle between one Reeb vector field, say Z_1 , and its tangential part Z_1^T is constant along the line curves of Z_1^T (see [7]). Now by Lemma 3, for our case this angle is constant on the whole N , and then, N is minimal. \square

Observe that by Proposition 3, a connected one-dimensional submanifold is ϕ -invariant if and only if it is contained in one leaf of the vertical foliation \mathcal{V} . This foliation is totally geodesic and the geodesics are integral curves of nonzero vertical vector fields $c_1 Z_1 + c_2 Z_2$ with c_i constant functions, since we have $\nabla_{Z_i} Z_j = 0$ for $i, j = 1, 2$ (see [4]). Hence, for the Case 2 of Theorem 2 we have

Theorem 4 *A connected one-dimensional ϕ -invariant submanifold of a metric contact pair is minimal if and only if it is tangent to a vector field of the form $c_1 Z_1 + c_2 Z_2$ with c_i real numbers.*

This means that minimal one-dimensional ϕ -invariant submanifolds are exactly vertical geodesics.

5 Induced structures on ϕ -invariant submanifolds

In general, an even-dimensional ϕ -invariant submanifold of a metric contact pair does not inherit necessarily a contact pair structure [7]. Anyway we can observe that it still carries some interesting structure.

Proposition 4 *On a metric contact pair a ϕ -invariant submanifold N tangent to both Reeb vector fields carries a metric f -structure with two complemented frames. A normal metric contact pair on the ambient manifolds induces a \mathcal{K} -structure on N .*

Here the notion of f -structure is meant in the sense of Yano [21] with $f = \phi$ restricted to N , and the complemented frames are the restrictions of the Reeb vector fields to N . For

the definition of \mathcal{K} -structure, see [10]. The proof of this proposition is a straightforward computation and will be omitted.

For the odd-dimensional case, when ϕ is decomposable, a ϕ -invariant submanifold tangent to one Reeb vector field and orthogonal to the other one inherits a contact metric structure, and it is Sasakian when the metric contact pair is normal [11]. The following statement concerns the remaining class of ϕ -invariant submanifolds, i.e., those which are leaning.

Theorem 5 *Let $(M, \alpha_1, \alpha_2, \phi, g)$ be a metric contact pair of type (h, k) with decomposable ϕ , and N a ϕ -invariant submanifold of M of dimension $2n + 1 \geq 3$ nowhere tangent and nowhere orthogonal to the Reeb vector fields Z_1 and Z_2 (then $h \geq n$ and $k \geq n$). Set $\zeta = \frac{1}{\|Z_1^T\|} Z_1^T$ and $\omega = g(\zeta, \cdot)$ along N , where Z_1^T is the tangential part of Z_1 . Then*

1. ω induces a contact form on N with Reeb vector field ζ ,
2. (ϕ, ζ, ω, g) induces an almost contact metric structure on N , which is not contact metric.

If the metric contact pair is normal, the induced almost contact structure on N is normal.

For an almost contact manifold (N, ω, ζ, ϕ) , we use the terminology of [9]. A metric is said to be *associated* when $g(X, \zeta) = \omega(X)$ and $g(X, \phi Y) = d\omega(X, Y)$ for all X, Y , while *compatible* means just that $g(\phi X, \phi Y) = g(X, Y) - \omega(X)\omega(Y)$ for all X, Y . By the conclusion (2) in our Theorem 5, the induced metric is compatible but it is not associated. Actually at each point of N , we have $g(X, \phi Y) \neq d\omega(X, Y)$ for some X, Y tangent to N at the same point.

Proof As we know $\zeta = (\cos \theta_1)Z_1 + (\sin \theta_1)Z_2$ for some constant function θ_1 with $0 < \theta_1 < \pi/2$, this gives along N

$$\omega = (\cos \theta_1)\alpha_1 + (\sin \theta_1)\alpha_2.$$

The 1-form ω induces a contact form on N when for all X tangent to N , $\omega(X) = 0$ and $i_X d\omega = 0$ imply that $X = 0$. Take any X tangent to N such that $\omega(X) = 0$; then, X is orthogonal to ζ so that X is horizontal. Put $X = X_1 + X_2$ with $X_i \in \mathcal{H} \cap T\mathcal{F}_j$, for $j \neq i$. If in addition $i_X d\omega = 0$, we have

$$\begin{aligned} 0 &= -d\omega(X, \phi X) = -(\cos \theta_1)d\alpha_1(X, \phi X) - (\sin \theta_1)d\alpha_2(X, \phi X) \\ &= -(\cos \theta_1)d\alpha_1(X_1, \phi X_1) - (\sin \theta_1)d\alpha_2(X_2, \phi X_2) \\ &= -(\cos \theta_1)(d\alpha_1 + d\alpha_2)(X_1, \phi X_1) - (\sin \theta_1)(d\alpha_1 + d\alpha_2)(X_2, \phi X_2) \\ &= (\cos \theta_1)\|X_1\|^2 + (\sin \theta_1)\|X_2\|^2. \end{aligned}$$

Now since $\cos \theta_1 > 0$ and $\sin \theta_1 > 0$, we get $X_1 = X_2 = 0$ and then $X = 0$ as desired. Hence, ω induces a contact form on N .

It is clear that $\omega(\zeta) = 1$. We also have $i_\zeta d\omega = 0$ because ζ is vertical and then lying in the kernels of $d\alpha_1$ and $d\alpha_2$. Hence, ζ is the Reeb vector field of the contact form on N .

To prove that $\phi^2 = -I + \omega \otimes \zeta$, first remark that $\phi\zeta = 0$ and $\omega(\zeta) = 1$. Next any X tangent to N and orthogonal to ζ is horizontal and then satisfies $\phi^2 X = -X = -X + \omega(X)\zeta$. Hence, (ϕ, ζ, ω) induces an almost contact structure on N .

To get the compatibility of the metric with the almost contact structure, we need to prove that $g(\phi X, \phi Y) = g(X, Y) - \omega(X)\omega(Y)$ for every X and Y tangent to N . Such a vector X decomposes as $X = \omega(X)\zeta + X_H$ for some horizontal vector X_H tangent to N , and the

same for Y . Replacing in $g(\phi X, \phi Y) = g(X, Y) - \alpha_1(X)\alpha_2(Y) - \alpha_1(Y)\alpha_2(X)$, we obtain

$$\begin{aligned} g(\phi X, \phi Y) &= g(X, Y) - \alpha_1(\omega(X)\zeta)\alpha_2(\omega(Y)\zeta) - \alpha_1(\omega(Y)\zeta)\alpha_2(\omega(X)\zeta) \\ &= g(X, Y) - \omega(X)\omega(Y) (\alpha_1^2(\zeta) + \alpha_2^2(\zeta)) \\ &= g(X, Y) - \omega(X)\omega(Y) \end{aligned}$$

Hence, (ϕ, ζ, ω, g) induces an almost contact metric structure on N .

Now we prove that the induced metric is not associated. Since N is of dimension ≥ 3 , at every point of N there exists a nonzero vector X tangent to N and orthogonal to ζ . By Theorem 1, $X = X_1 + X_2$ with X_i nonzero horizontal and tangent to \mathcal{F}_j , for $j \neq i$. Choosing $Y = \phi X$ let us compare $g(X, \phi Y)$ with $d\omega(X, Y)$. On the one hand $-g(X, \phi Y) = g(X, X) = \|X_1\|^2 + \|X_2\|^2$, and on the other hand as before we have

$$\begin{aligned} -d\omega(X, Y) &= -d\omega(X, \phi X) \\ &= (\cos \theta_1)\|X_1\|^2 + (\sin \theta_1)\|X_2\|^2. \end{aligned}$$

The difference being $(1 - \cos \theta_1)\|X_1\|^2 + (1 - \sin \theta_1)\|X_2\|^2 > 0$, we get $g(X, \phi Y) \neq d\omega(X, Y)$. Hence, the metric is not associated for the almost contact structure of N .

When the contact pair structure $(\alpha_1, \alpha_2, \phi)$ is normal, for every X and Y tangent to N we have

$$[\phi, \phi](X, Y) + 2d\alpha_1(X, Y)Z_1 + 2d\alpha_2(X, Y)Z_2 = 0$$

and for its orthogonal projection on N using the ϕ -invariance of N we get

$$\begin{aligned} 0 &= [\phi, \phi](X, Y) + 2(\cos \theta_1)d\alpha_1(X, Y)\zeta + 2\sin(\theta_1)d\alpha_2(X, Y)\zeta \\ &= [\phi, \phi](X, Y) + 2d\omega(X, Y)\zeta, \end{aligned}$$

which means that the almost contact structure on N is normal. This completes the proof. □

We can summarize our discussion above concerning the induced structures on ϕ -invariant manifolds as follows.

Theorem 6 *Let $(M, \alpha_1, \alpha_2, \phi, g)$ be a metric contact pair with decomposable ϕ and Reeb vector fields Z_1 and Z_2 , and N a connected ϕ -invariant submanifold of M of dimension ≥ 2 .*

1. *When N has even dimension, it inherits a metric f -structure with two complemented frames.*
2. *Suppose that N has odd dimension. Along N let ζ be the normalized vector field of a nonzero vector field among Z_1^T and Z_2^T (the tangential parts of Z_1 and Z_2), and set $\omega = g(\zeta, \cdot)$. Then*

(a) *ω induces a contact form on N with Reeb vector field ζ , and (ϕ, ζ, ω, g) induces an almost contact metric structure on N .*

(b) *the induced almost contact metric on N is contact metric if and only if N is tangent to one Reeb vector field and orthogonal to the other one.*

In all cases, if the metric contact pair on M is normal, the induced structure on N is normal.

6 Characteristic leaves of $d\alpha_1$ and $d\alpha_2$

Consider a metric contact pair $(M, \alpha_1, \alpha_2, \phi, g)$ of type (h, k) with decomposable ϕ and Reeb vector fields Z_1 and Z_2 . In [4], it has been shown that the two-dimensional vertical

foliation \mathcal{V} tangent to the Reeb distribution is totally geodesic. Actually, this is even true in general for any compatible metric g with respect to a contact pair structure $(\alpha_1, \alpha_2, \phi)$ without decomposability condition on ϕ . It is also known that the characteristic foliations \mathcal{F}_1 and \mathcal{F}_2 of the 1-forms α_1 and α_2 , respectively, are orthogonal, and their leaves are minimal [6].

Now recall the existence of two other remarkable foliations in the manifold M , which are the characteristic foliations \mathcal{G}_1 and \mathcal{G}_2 of the 2-forms $d\alpha_1$ and $d\alpha_2$, respectively. Their corresponding subbundles are

$$T\mathcal{G}_i = \{X : d\alpha_i(X, Y) = 0 \ \forall Y\}, \quad i = 1, 2.$$

Each leaf of \mathcal{G}_1 (respectively, \mathcal{G}_2) inherits a metric contact pair of type $(0, k)$ (respectively, $(h, 0)$), and is foliated by leaves of \mathcal{V} and also by leaves of \mathcal{F}_1 (respectively, \mathcal{F}_2) [4]. Moreover, the leaves of \mathcal{G}_1 and \mathcal{G}_2 are ϕ -invariant and tangent to both Z_1 and Z_2 , and we have the following minimality theorem.

Theorem 7 *On a metric contact pair $(M, \alpha_1, \alpha_2, \phi, g)$ with decomposable ϕ , the leaves of the characteristic foliations \mathcal{G}_1 and \mathcal{G}_2 of the 2-forms $d\alpha_1$ and $d\alpha_2$ are minimal.*

By normality condition on the metric contact pair of the ambient manifold M , it has been shown in [7] that the leaves of \mathcal{G}_1 and \mathcal{G}_2 are minimal. In our theorem, the normality condition is not needed.

Proof To prove the minimality of the leaves of \mathcal{G}_1 in the Riemannian manifold (M, g) , we use the minimality criterion of Rummel [18]. If the type numbers of the contact pair are (h, k) , the dimension of this foliation is $2k + 2$. The volume element of the metric g can be written as [6]

$$dV = \frac{(-1)^{h+k}}{2^{h+k} h! k!} \alpha_1 \wedge (d\alpha_1)^h \wedge \alpha_2 \wedge (d\alpha_2)^k,$$

so that the characteristic $(2k + 2)$ -form of the foliation \mathcal{G}_1 is, up to a constant, $\omega = \alpha_1 \wedge \alpha_2 \wedge (d\alpha_2)^k$. Since the Reeb vector field Z_1 is tangent to the foliation \mathcal{G}_1 and

$$i_{Z_1} d\omega = i_{Z_1} \left(d\alpha_1 \wedge \alpha_2 \wedge (d\alpha_2)^k \right) = 0,$$

the characteristic $(2k + 2)$ -form of the foliation \mathcal{G}_1 is closed on the subbundle $T\mathcal{G}_1$, giving that the leaves of \mathcal{G}_1 are minimal. The same argument applies to \mathcal{G}_2 , completing the proof. □

Example 5 In Example 1 the characteristic subbundle $T\mathcal{G}_1$ of $d\alpha_1$ (respectively, $T\mathcal{G}_2$ of $d\alpha_2$) is spanned by Y_1, Y_2, Y_4 and Y_6 (respectively, Y_1, Y_2, Y_3 and Y_5). Observe that the leaves of \mathcal{G}_2 are totally geodesic while those of \mathcal{G}_1 are minimal but not totally geodesic.

7 Metric contact pairs of type $(h, 0)$

On a metric contact pair with decomposable endomorphism ϕ the leaves of the two characteristic foliations are ϕ -invariant submanifolds. They are minimal and a priori they are not totally geodesic [6]. However, when the contact pair is of type $(h, 0)$, we can state the following.

Theorem 8 *Consider a metric contact pair $(M, \alpha_1, \alpha_2, \phi, g)$ of type $(h, 0)$ and Reeb vector fields Z_1 and Z_2 . If Z_2 is Killing, then the metric contact pair M is locally the product of a contact metric manifold with \mathbb{R} .*

We observed that \mathcal{P} -manifolds of Vaisman are metric contact pairs of type $(h, 0)$ where Reeb vector fields are Killing. Theorem 8 is a generalization to metric contact pairs of a result stated for \mathcal{P} -manifolds by Vaisman [19].

Proof The two characteristic foliations are orthogonal and complementary. The leaves of \mathcal{F}_1 are the integral curves of Z_2 which are geodesics [4]. We have just to prove that the leaves of \mathcal{F}_2 are totally geodesic. Any leaf F of \mathcal{F}_2 is a submanifold of codimension one, and the vector field Z_2 restricted to F is the normal to the leaf. When Z_2 is Killing, any geodesic γ of (M, g) starting from a point p of F and tangent to F at p satisfies

$$\dot{\gamma} g(\dot{\gamma}, Z_2) = g(\nabla_{\dot{\gamma}} \dot{\gamma}, Z_2) + g(\dot{\gamma}, \nabla_{\dot{\gamma}} Z_2) = 0.$$

Therefore, $g(\dot{\gamma}, Z_2) = 0$ because $\dot{\gamma}$ and Z_2 are orthogonal at p . Hence, the geodesic γ remains in the leaf F . \square

Here we used the simple fact that on a Riemannian manifold an immersed hypersurface normal to a Killing vector field is totally geodesic.

We know that the normality of a metric contact pair implies that the Reeb vector fields are Killing [2]. The following corollary is an immediate consequence of our previous theorem, and it has been already stated in equivalent terms of \mathcal{PK} -manifolds (nowadays called non-Kähler Vaisman manifolds) by Vaisman.

Corollary 1 (Vaisman [19]) *A normal metric contact pair of type $(h, 0)$ is locally the product of a Sasakian manifold with \mathbb{R} .*

8 Comments

As we said in Remark 1, a normal metric contact pair is a special metric f -structure with two complemented frames. A manifold carrying such a structure has even dimension and the two corresponding frames are nothing but the Reeb vector fields Z_1 and Z_2 of the contact pair. There are many papers studying submanifolds of f -manifolds. Among them, we can cite [1, 12, 14, 16, 17].

The ϕ -invariant submanifolds, of metric contact pairs, for which the results of these papers may apply are all Cauchy–Riemann submanifolds tangent to the Reeb distribution spanned by Z_1 and Z_2 , and then of even dimensions. In our work, the main results concern only ϕ -invariant submanifolds of odd dimensions with their tangent bundles intersecting the Reeb distribution along a line bundle.

Acknowledgements The second author was supported by a Visiting Professor fellowship at Université de Haute Alsace, Mulhouse, in June 2014 and PRIN 2010/11 “Varietà reali e complesse: geometria, topologia e analisi armonica”—Italy; GNSAGA–INdAM, Italy

References

1. Abatangelo, L.M., Dragomir, S.: Principal toroidal bundles over Cauchy–Riemann products. *Int. J. Math. Math. Sci.* **13**, 299–310 (1990)
2. Bande, G., Blair, D.E., Hadjar, A.: On the curvature of metric contact pairs. *Mediterr. J. Math.* **10**, 989–1009 (2013)
3. Bande, G., Hadjar, A.: Contact pairs. *Tôhoku Math. J.* **57**, 247–260 (2005)
4. Bande, G., Hadjar, A.: Contact pair structures and associated metrics. In: *Differential Geometry—Proceedings of the 8th International Colloquium*, pp. 266–275. World Science Publication, Hackensack (2009)

5. Bande, G., Hadjar, A.: On normal contact pairs. *Int. J. Math.* **21**, 737–754 (2010)
6. Bande, G., Hadjar, A.: On the characteristic foliations of metric contact pairs. In: Loubeau, E., Montaldo, S (eds.) *Harmonic Maps and Differential Geometry*. Contemporary Mathematics, vol. 542, pp. 255–259. American Mathematical Society, Providence (2011)
7. Bande, G., Hadjar, A.: Minimality of invariant submanifolds in metric contact pair geometry. *Ann. Mat. Pura Appl.* **194**, 1107–1122 (2015)
8. Bande, G., Kotschick, D.: Contact pairs and locally conformally symplectic structures. In: Loubeau, E., Montaldo, S (eds.) *Harmonic Maps and Differential Geometry*. Contemporary Mathematics, vol. 542, pp. 85–98. American Mathematical Society, Providence (2011)
9. Blair, D.E.: *Riemannian Geometry of Contact and Symplectic Manifolds*. Progress in Mathematics, vol. 203, 2nd edn. Birkhäuser, Boston (2010)
10. Blair, D.E., Ludden, G.D., Yano, K.: Differential geometric structures on principal toroidal bundles. *Trans. Am. Math. Soc.* **181**, 175–184 (1973)
11. Blair, D.E., Ludden, G.D., Yano, K.: Geometry of complex manifolds similar to the Calabi–Eckmann manifolds. *J. Differ. Geom.* **9**, 263–274 (1974)
12. Dragomir, S., Grimaldi, R.: Cauchy–Riemann submanifolds of manifolds carrying f -structures with complemented frames. *Soochow J. Math.* **16**, 193–209 (1990)
13. Dragomir, S., Ornea, L.: *Locally Conformal Kähler Geometry*. Progress in Mathematics, vol. 155. Birkhäuser, Boston (1998)
14. Goldberg, S.I.: Invariant submanifolds of codimension 2 of framed manifolds. *An. Sti. Univ. “Al. I. Cuza” Iasi* **17**, 157–165 (1971)
15. Gordon, C.S., Wilson, E.N.: The spectrum of the Laplacian on Riemannian Heisenberg manifolds. *Mich. Math. J.* **33**, 253–271 (1986)
16. Mihai, I.: CR-submanifolds of a framed f -manifold. *Stud. Cerc. Mat.* **35**, 127–136 (1983)
17. Ornea, L.: Generic CR-submanifolds in S -manifolds. *Stud. Cerc. Mat.* **36**, 435–443 (1984)
18. Rummeler, H.: Quelques notions simples en géométrie riemannienne et leurs applications aux feuilletages compacts. *Comment. Math. Helv.* **54**, 224–239 (1979)
19. Vaisman, I.: Locally conformal Kähler manifolds with parallel Lee form. *Rend. Mat.* **12**, 263–284 (1979)
20. Vaisman, I.: Generalized Hopf manifolds. *Geom. Dedic.* **13**, 231–255 (1982)
21. Yano, K.: On a structure defined by a tensor field f of type $(1, 1)$ satisfying $f^3 + f = 0$. *Tensor* **1**, 99–109 (1963)