

Inventory of black hole binaries

Jerome A. Orosz¹

*Sterrekundig Instituut, Universiteit Utrecht,
Princetonplein 5, NL-3584 CC Utrecht, Nederland*

Abstract. A small group of X-ray binaries currently provides the best evidence for the existence of stellar-mass black holes. These objects are interacting binary systems where the X-rays arise from accretion of material onto a compact object (*i.e.*, an object with a radius of less than a few hundred km). In some favourable cases, optical studies of the companion star lead to dynamical mass estimates for both components. In 17 cases, the mass of the compact object in an X-ray binary has been shown to exceed the maximum mass of a stable neutron star (about $3 M_{\odot}$), which leads to the conclusion that these objects are black holes. In this contribution I will review the basic properties of these black hole binaries.

1. Definition and source selection

Black holes represent an extreme manifestation of Einstein's theory of general relativity. As an observational astronomer, I will not consider any of the detailed theory of black holes but will instead consider only the practical question of how to find them. The usual route is by the process of elimination. In the current universe, black holes must form via the gravitational collapse of a dying star. Three outcomes are possible, depending on the mass of the degenerate core: a white dwarf, a neutron star, or a black hole. The mass of a white dwarf cannot exceed the well-known Chandrasekhar limit, and a neutron star has a somewhat analogous upper mass limit, generally thought to be on the order of $3 M_{\odot}$ (Rhoades & Ruffini 1974; Kalogera & Baym 1996). Once the mass of the degenerate core exceeds $\sim 3 M_{\odot}$, no known force can halt the gravitational collapse, and a black hole must be formed. Given this, we have a relatively straightforward observational definition of a black hole: *A black hole is a compact object with a mass greater than three solar masses*, where 'compact' in this context means an object with a radius smaller than about 100 km (*i.e.*, much less than the radius of normal stars).

Since black holes are dark, the only way one could hope to observe them is through their gravitational influence on surrounding matter. Early attempts to search catalogs of spectroscopic binaries to look for single-lined binaries with massive and undetected companions were not successful (Zel'dovich & Guseynov 1966; Trimble & Thorne 1969). A far more efficient approach for source selection has its roots in the 1960s when Zel'dovich and others realized that if a black

¹Present address: Department of Astronomy, San Diego State University, 5500 Campanile Drive, San Diego, CA 92182-1221, USA

hole accreted material (either from a nearby companion star or the interstellar medium), then it might shine brightly in X-rays and γ rays. Today we know of a few hundred bright X-ray sources which must be powered by accretion onto a compact object, either a neutron star or a black hole (*i.e.*, rapid variability in many cases indicates a size scale of a few hundred km or less). All of the black hole candidates discussed below were selected on the basis of their X-ray activity, although two of the sources, V404 Cyg and V4641 Sgr, were previously known (optically) variable stars that were essentially ignored until they were associated with bright X-ray sources. Thus, one should keep in mind that owing to special circumstances needed to observe them (*i.e.*, interacting binaries), the list of presently known stellar-mass black holes is probably a very biased sample and may not be representative of the general population of black holes.

2. Mass limits

By my definition above, to establish the presence of a black hole in a given system one must show that there is a compact object with a mass greater than $3 M_{\odot}$. Establishing a lower limit to the mass is relatively easy since in a single-lined spectroscopic binary, the observed radial velocity curve of the visible star can be used to set a firm lower limit on the mass of the unseen companion:

$$f(M) \equiv \frac{PK_2^3}{2\pi G} = \frac{M_{\text{BH}}^3 \sin^3 i}{(M_{\text{BH}} + M_2)^2}, \quad (1)$$

where P is the binary orbital period, K_2 is the semiamplitude of the companion's radial velocity curve, G is the gravitation constant, i is the inclination of the orbit to the line of sight ($i = 90^\circ$ for an orbit seen edge-on), and where M_{BH} and M_2 are the masses of the invisible object and the companion star, respectively. One can easily see that $f(M)$ (generally called the 'optical mass function' or 'mass function') is always *less than* M_{BH} since $M_2 > 0$ and $\sin i \leq 1$ always. Thus a firm lower limit on the mass of the compact object can be obtained from two relatively simple measurements.

In real life, of course, things are always a bit more complicated. For starters, in many cases one cannot even see spectral lines from the companion star in many of the persistently X-ray bright sources. Needless to say this precludes any attempts at dynamical mass estimates. In other cases, the companion star may not be uniformly bright over its surface owing to heating by the X-ray source. If so, then the 'centre of light' will not be the same as the 'centre of mass' and the observed radial velocity curve will be different than the true one. Since the value of the mass function depends on K_2^3 , small changes in K_2 can lead to relatively large changes in $f(M)$. Fortunately, many of the potential difficulties related to X-ray heating are largely absent in the subset of the X-ray binaries known as 'X-ray transients'. As the name suggests, the X-ray luminosity in these sources is highly variable (dynamic ranges of 10^6 to 10^7 are not uncommon). Dynamical observations of these systems are generally reliable, provided one *observes the system after the X-ray activity has completely ceased* (see Phillips, Shahbaz & Podsiadlowski 1999; Shahbaz *et al.* 1999a; and Shahbaz *et al.* 2000 for discussions regarding X-ray heating in GRO J1655–40).

Table 1 lists the mass functions for 17 systems that are widely believed to contain black holes. Fourteen of these systems are transient X-ray sources

Table 1. Basic system parameters for the fourteen transient and three persistent black hole binaries.

source	P (d)	$f(M)$ (M_{\odot})	$v_{\text{rot}} \sin i$ (km s^{-1})	i (deg)	q (M_{BH}/M_2)	M_{BH} (M_{\odot})	refs.
GRO J0422+32	0.2121600(2)	1.19 ± 0.02	90^{+22}_{-27}	44 ± 2	3.2 - 13.2	3.66 - 4.97	1-3
A 0620-00	0.3230160(5)	2.72 ± 0.06	83 ± 5	40.8 ± 3.0	13.3 - 18.3	8.70 - 12.86	4-6
GRS 1009-45	0.285206(2)	3.17 ± 0.12	...	677	6.3 - 8.07	3.64 - 4.747	7,8
XTE J 1118+480	0.169930(4)	6.1 ± 0.3	114 ± 4	81 ± 2	22.7 - 28.8	6.48 - 7.19	9-11
GS 1124-683	0.432606(3)	3.01 ± 0.15	106 ± 13	54 ± 2	4.8 - 8.8	6.47 - 8.18	12-15
4U 1543-47	1.116407(3)	0.25 ± 0.01	46 ± 2	20.7 ± 1.5	3.1 - 4.0	8.45 - 10.39	16
XTE J 1550-564	1.5435(5)	6.86 ± 0.71	90 ± 107	72 ± 5	> 12	8.36 - 10.76	17
GRO J 1655-40	2.6219(2)	2.73 ± 0.09	99 ± 3	70.2 ± 1.2	2.4 - 2.7	6.03 - 6.57	18-20
H 1705-250	0.521(1)	4.86 ± 0.13	< 79	> 60	> 18.9	5.64 - 8.30	21-23
SAX J 1819.3-2525	2.81730(1)	3.13 ± 0.13	98.9 ± 1.5	75 ± 2	2.22 - 2.39	6.82 - 7.42	24,25
XTE J 1859+226	0.382(3)	7.4 ± 1.1	7.6 - 12.07	26
GRS 1915+105	34(2)	9.5 ± 3.0	...	70 ± 27	...	10.0 - 18.07	27-29
GS 2000+25	0.3440915(9)	5.01 ± 0.12	86 ± 8	64.0 ± 1.3	18.9 - 28.9	7.15 - 7.78	30-32
GS 2023+338	6.4714(1)	6.08 ± 0.06	38.8 ± 1.1	56 ± 4	16.1 - 18.9	10.06 - 13.38	33,34
LMC X-3	1.70479(4)	2.29 ± 0.32	130 ± 20	67 ± 3	1.1 - 2.0	5.94 - 9.17	35-37
LMC X-1	4.2288(6)	0.14 ± 0.05	...	> 63?	0.3 - 0.77	4.0 - 10.07	38
Cyg X-1	5.59983(2)	0.244 ± 0.005	94 ± 5	35 ± 5	0.50 - 0.57	6.85 - 13.25	39-42

References: 1: Webb *et al.* 2000; 2: Harlaftis *et al.* 1999; 3: Gelino & Harrison 2002; 4: Leibowitz, Hemar & Orlo 1998; 5: Marsh, Robinson & Wood 1994; 6: Gelino, Harrison & Orosz 2001; 7: Filippenko *et al.* 1999; 8: Gelino 2002; 9: Wagner *et al.* 2001; 10: Orosz 2001; 11: McClintock *et al.* 2001; 12: Orosz *et al.* 1996; 13: Casares *et al.* 1997; 14: Shahbaz, Naylor & Charles 1997; 15: Gelino, Harrison & McNamara 2001; 16: Orosz *et al.* 2002b; 17: Orosz *et al.* 2002a; 18: Greene, Bailyn & Orosz 2001; 19: Shahbaz *et al.* 1999a; 20: Israeli *et al.* 1999; 21: Remillard *et al.* 1996; 22: Filippenko *et al.* 1997; 23: Harlaftis *et al.* 1997; 24: Orosz *et al.* 2001; 25: Orosz *et al.* 2002c; 26: Filippenko & Chornock 2001; 27: Greiner, Cuby & McCaughrean 2001; 28: Mirabel & Rodriguez 1994; 29: Fender *et al.* 1999; 30: Chevalier & Ilovaisky 1993; 31: Harlaftis, Horne & Filippenko 1996; 32: Gelino 2001; 33: Casares & Charles 1994; 34: Shahbaz *et al.* 1994; 35: van der Klis *et al.* 1985; 36: Cowley *et al.* 1983; 37: Kuiper, van Paradijs & van der Klis 1988; 38: Hutchings *et al.* 1987; 39: Brocksopp *et al.* 1999; 40: Ninkov, Walker & Yang 1987; 41: Gies & Bolton 1986; 42: Herrero *et al.* 1995.

with relatively low-mass companions ($M_2 \lesssim 1 M_{\odot}$ generally), and the remaining three systems have massive O/B companions whose bolometric luminosity is comparable to or exceeds the X-ray luminosity of the accreting object. Six of the systems have mass functions which are above $\sim 5 M_{\odot}$, and six more have mass functions more than $\sim 3 M_{\odot}$. Based on the mass functions alone, we can safely conclude these systems contain black holes since the compact objects in them are too massive to be neutron stars. Although the other five systems have relatively small mass functions, we can be reasonably sure these systems also contain black holes since the currently best estimates for the compact object masses (see the next section) are all above $\sim 3.5 M_{\odot}$. In addition to the sources listed in Table 1, there are a few dozen sources which are believed to contain a black hole on the basis of their X-ray spectra and temporal properties (*e.g.*, Tanaka & Lewin 1995; Liu, van Paradijs & van den Heuvel 2001).

3. Towards actual masses

If possible, we want to do better than simply establishing a lower limit to the mass of a black hole. If we think of the companion star as a test particle orbiting the black hole, then clearly in order to measure the actual mass of the black hole after its mass function has been measured, one must be able to convert *projected* radial velocities into *actual* radial velocities (*i.e.*, the inclination i is needed) and to find out how far the star is from the binary centre-of-mass (*i.e.*, one needs to know the mass ratio *or* the mass of the companion star).

The procedure to derive astrophysically interesting parameters of a binary system is conceptually simple. One can make a model of a close binary system

that is specified by several parameters such as the orbital period, the separation between the stars, the mass ratio, the relative radii of the stars, the temperatures of the stars, the inclination to the line of sight, etc. Then, given this model, one can compute various *observable* properties of this binary such as the radial velocity curves, the light curves in various bandpasses, eclipse duration(s) if any, the rotational velocities of the stars, etc. The goal then is to find a set of input model parameters that can best match the set of observed properties of the binary of interest. Given the optimal input parameters, one can then compute astrophysically interesting quantities such as the masses of the components.

Models of close binary stars are relatively easy to construct, and have been used for over thirty years (*e.g.*, Wilson & Devinney 1971; Lyutyi, Syunyaev, Cherepashchuk 1973; Hutchings 1974; Avni & Bahcall 1975). The difficult part has been related to the *inverse* problem of parameter optimisation since the parameter space to search is usually vast and often parameters can be tightly correlated. Common ways to tackle the inverse problem include 'differential corrections' and related techniques (*e.g.*, Wilson & Devinney 1971) and systematic 'grid searches' (*e.g.*, Kuiper, van Paradijs & van der Klis 1988). More recently, I have been using optimisation techniques based on genetic algorithms (Orosz *et al.* 2002a). Genetic algorithms are very robust, and can allow for accurate estimation of uncertainties on the fitted and derived parameters.

In the case of the black hole binaries, the main observables usually are a radial velocity curve for the companion star (this basically sets the scale of the binary), a light curve for the companion in one or more bandpasses (these mainly constrain the inclination), and in most cases a measurement of the mean projected rotational velocity of the companion star (this sets the mass ratio). In nearly cases (with the possible exception of SAX J1819.3–2525), the X-ray source is not eclipsed, and one can set an upper limit to the inclination. Given these observations, it is not an unduly difficult problem (in principle!) to find a good binary model for most of the black hole systems listed in Table 1.

However, as before, in real life things are more complicated. In many cases, the observed light curve is not entirely due to the companion star, owing to the presence of extra sources of light (usually from an accretion disk). In the optical, the mean light curve can change from one year to the next (*e.g.*, Orosz *et al.* 1996; Leibowitz, Hemar & Orio 1998; Pavlenko *et al.* 2001). Presumably these long-term changes are related to the accretion disk. From spectroscopy, one can estimate the light attributable to the accretion disk (*e.g.*, Marsh, Robinson & Wood 1994) and correct for it in the models. Alternatively, it is generally *assumed* that the contamination from the accretion disk is small in the infrared *J*, *H*, and *K* bands (*e.g.*, Shahbaz *et al.* 1994; Gelino, Harrison & McNamara 2001). There have been a few attempts to directly measure the disk contamination in the infrared with spectroscopy, but the results so far are not terribly constraining (Shahbaz *et al.* 1996; Shahbaz, Bandyopadhyay & Charles 1999b).

I have summarised in Table 1 the best (in my opinion) measurements of the rotational velocities and inclinations for the 17 black hole binaries. Relatively uncertain values are denoted by '?'. In some cases there is reasonably good agreement between inclination measurements made by different groups (*e.g.*, GS 1124–683), while in other cases there have been a wide range of measurements (*e.g.*, Cyg X-1, GRO J0422+32). I have computed $\sim 1\sigma$ ranges for the black




Figure 1. Black hole masses (*top*, Table 1 and cited references) and selected neutron star masses (*bottom*, Thorsett & Chakrabarty 1999). Ranges are 1σ in most cases. Arrows in the top indicate measured mass functions.

hole mass and mass ratio using a simple Monte Carlo code that makes use of numerical Roche lobe integrations to map a rotational velocity into a mass ratio. In some cases, rotational velocities are not available and instead I have guessed the mass of the companion in order to arrive at a mass range for the black hole. Figure 1 shows graphically the black hole mass ranges and, for comparison, masses for neutron stars. All things considered, many of the masses are known reasonably well, but one should always keep in mind potential systematic errors such as those related to unstable light curves, *etc.*

References

- Avni, Y., Bahcall, J.N. 1975, *ApJ* 197, 675
 Brocksopp, C., Tarasov, A.E., Lyutyi, V.M., Roche, P. 1999, *A&A* 343, 86
 Casares, J., Charles, P. A. 1994, *MNRAS (Letters)* 271, L5
 Casares, J., Martin, E.L., Charles, P.A., *et al.* 1997, *New Astron.* 1, 299
 Chevalier, C., Ilovaisky, S.A. 1993, *A&A* 269, 301
 Cowley, A.P., Crampton, D., Hutchings, J.B., *et al.* 1983, *ApJ* 272, 118
 Fender, R.P., Garrington, S.T., McKay, D.J., *et al.* 1999, *MNRAS* 304, 865

- Filippenko, A.V., Matheson, T., Leonard, D.C., *et al.* 1997, PASP 109, 461
- Filippenko, A.V., Leonard, D.C., Matheson, T., *et al.* 1999, PASP 111, 969
- Filippenko, A.V., Chornock, R. 2001, IAU Circular No. 7644
- Gelino, D.M., Harrison, T.E., McNamara, B.J. 2001, AJ 122, 971
- Gelino, D.M., Harrison, T.E., Orosz, J.A. 2001, AJ 122, 2668
- Gelino, D.M. 2001, PhD thesis, New Mexico State University, USA
- Gelino, D.M. 2002, BAAS 200, #08.11
- Gelino, D.M., Harrison, T.E. 2002, ApJ in press
- Gies, D.R., Bolton, C.T. 1986, ApJ 304, 371
- Greene, J., Bailyn, C.D., Orosz, J.A. 2001, ApJ 554, 1290
- Greiner, J., Cuby, J.G., McCaughrean, M.J. 2001, Nature 414, 522
- Harlaftis, E.T., Horne, K., Filippenko, A.V. 1996, PASP 108, 762
- Harlaftis, E., Steeghs, D., Horne, K., Filippenko, A.V. 1997, AJ 114, 1170
- Harlaftis, E., Collier, S., Horne, K., Filippenko, A.V. 1999, A&A 341, 491
- Herrero, A., Kudritzki, R.-P., Gabler, R., *et al.* 1995, A&A 297, 556
- Hutchings, J.B. 1974, ApJ (Letters) 193, L61
- Hutchings, J.B., Crampton, D., Cowley, A.P., *et al.* 1987, AJ 94, 340
- Israelian, G., Rebolo, R., Basri, G., Casares, J., Martin, E.L. 1999, Nature 401, 142
- Kalogera, V., Baym, G. 1996, ApJ (Letters) 470, L61
- Kuiper, L., van Paradijs, J., van der Klis, M. 1988, A&A 203, 79
- Leibowitz, E.M., Hemar, S., Orio, M. 1998, MNRAS 300, 463
- Liu, Q.Z., van Paradijs, J., van den Heuvel, E.P.J. 2001, A&A 368, 1021
- Lyutyi, V.M., Syunyaev, R.A., Cherepashchuk, A.M. 1973, Astron. Zh. 50, 3 (= Soviet Astron. 17, 1)
- Marsh, T.R., Robinson, E.L., Wood, J.H. 1994, MNRAS 266, 137
- McClintock, J.E., Garcia, M.R., Caldwell, M., *et al.* 2001, ApJ (Letters) 551, L147
- Mirabel, I.F., Rodriguez, L.F. 1994, Nature 371, 46
- Ninkov, Z., Walker, G.A.H., Yang, S. 1987, ApJ 321, 425
- Orosz, J.A. 2001, ATel #67
- Orosz, J.A., Bailyn, C.D., McClintock, J.E., Remillard, R.A. 1996, ApJ 468, 380
- Orosz, J.A., Kuulkers, E., van der Klis, M., McClintock, J.E., Garcia, M.R., Callanan, P.J., Bailyn, C.D., Jain, R.K., Remillard, R.A. 2001, ApJ 555, 489
- Orosz, J.A., Groot, P.J., van der Klis, M., McClintock, J.E., Garcia, M.R., Zhao, P., Jain, R.K., Bailyn, C.D., Remillard, R.A. 2002a, ApJ 568, 845
- Orosz, J.A., *et al.* 2002b, in preparation
- Orosz, J.A., *et al.* 2002c, in preparation
- Pavlenko, E.P., Kuznetsova, Y.G., Shugarov, S.Yu., Petrov, V.S. 2001, Ap&SS Suppl. 276, 65
- Phillips, S.N., Shahbaz, T., Podsiadlowski, Ph. 1999, MNRAS 304, 839
- Remillard, R.A., Orosz, J.A., McClintock, J.E., Bailyn, C.D. 1996, ApJ 459, 226
- Rhoades, C.E., Jr., Ruffini, R. 1974, Phys. Rev. Letters 32, 324
- Shahbaz, T., Ringwald, F.A., Bunn, J.C., *et al.* 1994, MNRAS 271, L10
- Shahbaz, T., Bandyopadhyay, R., Charles, P.A., Naylor, T. 1996, MNRAS 282, 977
- Shahbaz, T., Naylor, T., Charles, P.A. 1997, MNRAS 285, 607
- Shahbaz, T., van der Hooft, F., Casares, J., Charles, P.A., van Paradijs, J. 1999a, MNRAS 306, 89

- Shahbaz, T., Bandyopadhyay, R.M., Charles, P.A. 1999b, *A&A* 346, 82
- Shahbaz, T., Groot, P., Phillips, S.N., Casares, J., Charles, P.A., van Paradijs, J. 2000, *MNRAS* 314, 747
- Tanaka, Y., Lewin, W.H.G. 1995, in: W.H.G. Lewin, J. van Paradijs, E.P.J. van den Heuvel (eds.), *X-ray Binaries* (Cambridge: CUP), p. 126
- Thorsett, S.E., Chakrabarty, D. 1999, *ApJ* 512, 288
- Trimble, V.L., Thorne, K.S. 1969, *ApJ* 156, 1013
- van der Klis, M., Clausen, M., Jensen, K., Tjemkes, S., van Paradijs, J. 1985, *A&A* 151, 322
- Wagner, R.M., Foltz, C.B., Shahbaz, T., Casares, J., Charles, P.A., Starrfield, S.G., Hewett, P. 2001, *ApJ* 556, 42
- Webb, N.A., Naylor, T., Ioannou, Z., Charles, P.A., Shahbaz, T. 2000, *MNRAS* 317, 528
- Wilson, R.E., Devinney, E.J. 1971, *ApJ* 166, 605
- Zel'dovich, Ya.B., Guseynov, O.H. 1966, *ApJ* 144, 840

Discussion

CHEREPASHCHUK: Recently Antokhina, Cherepashchuk & Shalba proposed a new method of determination of parameters of X-ray binaries with black holes based on the variability of absorption line profiles of the optical star with the phase of orbital period. Such variability is due to complicated temperature distribution over the surface of tidally deformed star. What is your feeling about the possibility to realize this method, using new generation telescopes like VLT, Keck, *etc.*?

OROSZ: The method is in principle a good one, but in practice one needs high resolution, high signal-to-noise optical spectra. We have a few VLT-UVES spectra of the brightest system, V4641 Sgr, but ideally we would want spectra over the whole orbit. Apart from V4641 Sgr, it is not feasible to do this for any other X-ray transient.

KOENIGSBERGER: Are the ages of all these systems (after the SN explosion) actually large enough for them to have circularized and synchronized ?

OROSZ: The X-ray transients with low mass, Roche lobe filling companions can be quite old, easily 10^9 yr, so there has been ample time for circularization and synchronization. As for Cyg X-1, it's orbit is circular, and we would expect it to be synchronous as well, since in general that that timescale is shorter.

MOFFAT: I know this is an old question ...: Is the low observed number of BH + massive companion (O-type) binaries in the Galaxy compatible with the assumption that all stars of $M_i = 40 M_\odot$ become BHs, given the complexities of accretion and the possibility of smothered or sleeping systems? What about Cyg X-3?

OROSZ: Norbert Langer shared in his presentation that it is very difficult to make relatively massive black holes in close binaries. (Recall that we need close binaries to have an X-ray binary that can be selected.) However, given the small numbers of sources and the uncertainties in the models, we are not prepared to make firm comparisons between theory and observations. Cyg X-3 is certainly a black hole candidate. However, I am not aware of a reliable velocity curve for the companion. That system is difficult to observe owing to the large extinction and the strong wind from the companion.