

BMJ Open Investigating locally relevant risk factors for *Campylobacter* infection in Australia: protocol for a case-control study and genomic analysis

Liana Varrone,¹ Russell J Stafford,² Kim Lilly,³ Linda Selvey,¹ Kathryn Glass,⁴ Laura Ford,⁴ Dieter Bulach,^{5,6} Martyn D Kirk,⁴ On behalf of The CampySource Project Team

To cite: Varrone L, Stafford RJ, Lilly K, *et al.* Investigating locally relevant risk factors for *Campylobacter* infection in Australia: protocol for a case-control study and genomic analysis. *BMJ Open* 2018;**8**:e026630. doi:10.1136/bmjopen-2018-026630

► Prepublication history and additional material for this paper are available online. To view these files, please visit the journal online (<http://dx.doi.org/10.1136/bmjopen-2018-026630>).

Received 17 September 2018
Revised 3 October 2018
Accepted 3 October 2018



© Author(s) (or their employer(s)) 2018. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

For numbered affiliations see end of article.

Correspondence to

Dr Martyn D Kirk;
Martyn.Kirk@anu.edu.au

ABSTRACT

Introduction The CampySource project aims to identify risk factors for human *Campylobacter* infection in Australia. We will investigate locally relevant risk factors and those significant in international studies in a case-control study. Case isolates and contemporaneous isolates from food and animal sources will be sequenced to conduct source attribution modelling, and findings will be combined with the case-control study in a source-assigned analysis.

Methods and analysis The case-control study will include 1200 participants (600 cases and 600 controls) across three regions in Australia. Cases will be recruited from campylobacteriosis notifications to health departments. Only those with a pure and viable *Campylobacter* isolate will be eligible for selection to allow for whole genome sequencing of isolates. Controls will be recruited from notified cases of influenza, frequency matched by sex, age group and geographical area of residence. All participants will be interviewed by trained telephone interviewers using a piloted questionnaire. We will collect *Campylobacter* isolates from retail meats and companion animals (specifically dogs), and all food, animal and human isolates will undergo whole genome sequencing. We will use sequence data to estimate the proportion of human infections that can be attributed to animal and food reservoirs (source attribution modelling), and to identify spatial clusters and temporal trends. Source-assigned analysis of the case-control study data will also be conducted where cases are grouped according to attributed sources.

Ethics and dissemination Human and animal ethics have been approved. Genomic data will be published in online archives accompanied by basic metadata. We anticipate several publications to come from this study.

INTRODUCTION

Campylobacter infection is the most commonly notified cause of foodborne gastroenteritis in Australia,¹⁻³ as well as a leading cause of bacterial gastroenteritis worldwide.⁴ At the introduction of Australia's National Notifiable Diseases Surveillance System in 1991 the incidence rate of notified campylobacteriosis

Strengths and limitations of this study

- Case-control study is well powered to identify locally relevant risk factors.
- Linking genomic data to the case-control study strengthens the analysis by enabling source attribution and source-assigned analyses to be conducted.
- Case-control questionnaire questions are being validated in a separate study, demonstrating the reliability of participant recall.
- Potential reporting bias due to inaccurate recall of study participants is a potential weakness of the study.
- Case-control study lacks efficiency for risk factors with high levels of exposure in the study population.

cases was 79.1/100 000 population,⁵ and despite notification rates plateauing in recent years, incidence had risen to 139.7/100 000 population in Australia in 2015,⁵ with an estimated 10 cases for every notified case within the community.⁶ Similarly, the incidence rate of campylobacteriosis in New Zealand in 2014 was 150.3/100 000 population,⁷ with an estimated 10–30 cases in the community for every notified case.⁸ *Campylobacter* notification rates in Australia and New Zealand are still among the highest in the world across high-income countries. Most countries in the European Union consistently report annual campylobacteriosis notification rates below 100/100 000 population.²

Two species of *Campylobacter*—*Campylobacter jejuni* and *C. coli*—contribute to approximately 95% of human campylobacteriosis.⁹ These *Campylobacter* spp are commonly detected in sewage and surface water,¹⁰ reside in the gastrointestinal tract of birds and animals¹¹ and are frequently found in raw meat, particularly poultry, and raw milk.^{12 13} Campylobacteriosis is mostly foodborne, with an

estimated 77% of cases transmitted via food consumption in Australia.^{14,15} Direct and indirect zoonotic transmission can occur via animal contact (direct) or faecally contaminated water or environments (indirect). Person-to-person transmission is considered rare.¹⁶ The majority of cases are thought to be sporadic, with outbreaks less commonly detected.¹⁷ Most outbreaks are linked to the consumption of poultry, raw milk or contaminated water.^{17,18}

Targeted control of foodborne bacterial pathogens generally depends on identification of sources and routes of transmission. Since *Campylobacter* are ubiquitous in the environment and most cases are sporadic, identifying sources is difficult. Source attribution methods require isolation of strains from reservoirs to compare *Campylobacter* strain diversity in foods and animals with that in human infections. Beef, sheep and pig meat have a lower prevalence of *Campylobacter* contamination than chicken meat (<5%–14%),^{19–21} but a higher prevalence is found in animal offal such as liver,²² thus making offal a valuable source of host-associated strains of *Campylobacter* in low-prevalence meats.

STUDY RATIONALE

In the USA, evidence from case–control studies has led to policy change, including changes to chicken slaughtering techniques. The incidence of human *Campylobacter* infection has declined in the USA since this policy was introduced in 1997.²³ More recently, evidence from source attribution analyses in New Zealand has led to the development of poultry production policies and practices aimed at reducing the risk of *Campylobacter* transmission via poultry food products.²⁴ New Zealand has seen a 74% reduction in the number of campylobacteriosis cases attributed to poultry in the region, as well as a 54% reduction in cases overall.²⁵

Source attribution modelling enables us to determine which foods and animals are the most likely sources of infection with each *Campylobacter* strain type, and the proportion of cases attributed to each source. This can be done with simple proportional similarity index (PSI) calculations, or by using more complex models.²⁴ Source attribution also allows for human campylobacteriosis cases to be grouped by potential source, increasing the specificity of risk factor analyses. These source-assigned analyses combine the epidemiological information gained through the traditional case–control study with source attribution modelling to provide greater explanatory power to investigate locally relevant risk factors.

OBJECTIVES

This study aims to:

1. Identify dietary, environmental and behavioural risk factors for *Campylobacter* infection in Australia.
2. Strengthen the epidemiological evidence for previously identified risk factors in Australia.

3. Identify strain-specific risk factors for infection using whole genome sequencing (WGS) data from case isolates.

HYPOTHESES

We will test several hypotheses regarding specific risk factors for *Campylobacter* infection in Australia. The hypotheses are based on exposures that have previously been identified as risk factors for *Campylobacter* infection in Australia as well as internationally.

We hypothesise that:

1. Persons who consume undercooked meats, particularly chicken, are at increased risk of infection.
2. Persons who consume offal are at increased risk of infection.
3. Persons who own companion animals (especially puppies) are at increased risk of infection.
4. Poor food hygiene and handling practices in the home increase the risk of infection.
5. Most human infections will be attributed to consumption of chicken meat.
6. There will be a high level of genetic diversity among *Campylobacter* strains.

STUDY DESIGN

We will conduct a case–control study including genomic testing over a 2-year period in three sentinel sites: the state of Queensland (QLD), the Australian Capital Territory (ACT) and the Hunter New England (HNE) region of New South Wales (figure 1). Sporadic cases of culture-positive *Campylobacter* infection will be identified either through state notifiable disease registers, from local pathology service databases or local notification databases. An isolate from each case will be paired with epidemiological data from the case interview. One control will be recruited for each case who participates in the study, with trained interviewers conducting telephone interviews with both cases and



Figure 1 Map of Australian states and territories including the Hunter New England region of New South Wales.

Table 1 Sample size estimates for an unmatched case–control study

Risk factor	Prevalence of exposure among controls (%)	Prevalence of exposure among cases (%)	OR	Number of required study subjects
Beef	78	85	1.6	960
Pork	52	60	1.4	1130
Lamb	42	50	1.4	1120
Chicken	80	87	1.6	1040
Offal	2.0	5.0	2.6	1154
Puppies	2.1	5.4	2.7	1040

80% power and $\alpha=0.05$.

controls. Participants will be interviewed using a questionnaire that has been specifically designed to collect information on known potential risk factors. This questionnaire will include a selection of questions being validated in a separate study (LV: Validation of questions designed for investigation of gastroenteritis). For cases, the questions will cover the 7 days prior to the onset of illness, while controls will be questioned on the 7 days prior to interview. Meanwhile, *Campylobacter* isolates will also be collected from food and animal samples. All human and non-human isolates will undergo WGS for comparison in source attribution modelling. Data for this study will be collected from 1 March 2017 to 1 March 2019.

Patient and public involvement

To develop the study, we engaged state and territory health departments, food safety agencies and industry to establish research questions and methods. The process involved a dedicated workshop, followed by teleconferences and an iterative process of drafting study documentation. We also established a reference panel, which includes representatives from senior levels of government and industry bodies. No patients or other members of the public were involved in the development of this study.

STUDY POPULATION

The three sentinel sites cover a population of approximately 6.1 million people. Based on notification and diagnostic pathology data, we expect approximately 8650 *Campylobacter* cases to be notified across these sites during the study period.

DEFINITION AND SELECTION OF CASES

Case definition

We define a case as a person from any of the three participating sites with a history of acute diarrhoea and a culture-positive stool result for *Campylobacter*.

SAMPLE SIZE

We used risk factor prevalence data from a previous national *Campylobacter* case–control study in 2001/2002 to estimate sample size for this study.²⁶ For example, the prevalence of chicken consumption among controls in 2001/2002 was 80%. A sample size of approximately 1040 subjects (520 cases; 520 controls) would enable the study to detect an association between chicken consumption and illness with an OR of 1.6, at 80% power and $\alpha=0.05$, as reported in the previous study. Sample size estimates for other potential risk factors are listed in table 1.

From these calculations, we estimate that a study of 1200 subjects (600 cases; 600 controls) will adequately detect significant associations of these magnitudes for potential risk factors of interest. QLD and HNE sites will each enrol at least 250 cases into the study, while ACT will enrol at least 100 cases. Based on the previous Australian case–control study,²⁶ we expect approximately 80% of selected notified cases to be eligible and participate in the study (table 2).

In QLD, we will obtain cases from one private pathology provider reporting approximately 40% of the state's *Campylobacter* notifications. We estimate that this provider will notify 2800 cases during the study period with an estimated 45% of these being culture positive (1260 notified cases). In ACT, approximately 600 *Campylobacter* notifications are expected during the study period; 130 are expected from the participating pathology laboratory. In

Table 2 Sampling method for cases in each site

State	Expected number of notified cases during study period	Estimated cases from participating pathology laboratory	Culture-positive cases	Sequential sampling of notified cases	Total number of cases	Expected number to be recruited (~80% participation rate)
QLD	7000	2800	1260	Select every fourth case	315	250
ACT	600	130	130	Include all notified cases	130	100
NSW (Hunter New England)	1050	313	313	Include all notified cases	313	250
Total	8650	3243	1703		758	600

ACT, Australian Capital Territory; NSW, New South Wales; QLD, Queensland.

Table 3 Eligibility criteria for cases and controls

Criteria	Cases	Controls
Had diarrhoea (≥ 3 loose bowel movements in 24 hours)	Include	Exclude
Known date of illness onset	Include	NA
Household members positive for <i>Campylobacter</i> in 4 weeks prior to onset of illness	Exclude	Exclude (4 weeks prior to interview date)
Household members experiencing diarrhoea in 4 weeks prior to onset of illness	Exclude	Exclude (4 weeks prior to interview date)
Travelled outside of Australia in 2 weeks prior to onset of illness	Exclude	Exclude (2 weeks prior to interview date)
Travelled interstate for the entire 2 weeks prior to onset of illness	Exclude	Exclude (2 weeks prior to interview date)
Cannot speak English	Exclude	Exclude
Not able to answer questions for some other reason (eg, intellectually disabled)	Exclude	Exclude
Not contactable after six telephone attempts	Exclude	Exclude
Live outside the catchment areas	Exclude	Exclude
Do not have a telephone number available for their primary residence, or a mobile phone	Exclude	Exclude
An enteric pathogen other than <i>Campylobacter</i> was isolated/detected in their stool (excluding <i>Blastocystis hominis</i> and <i>Dientamoeba fragilis</i>)	Exclude	NA

NA, not applicable.

HNE, approximately 1050 *Campylobacter* notifications are expected during the study period; 313 of these notifications will be from the participating pathology laboratory.

Enrolment of cases

We will enrol all cases who meet the eligibility criteria (table 3). Each site will check for new notifications of culture-positive *Campylobacter* infection daily, with only culture-positive *Campylobacter* cases eligible for this study. If a case refuses to participate in the study, we will select a subsequent case for inclusion. Enrolment of cases will require consent from the patient, or in the event of a child aged less than 18 years, consent from either one of the parents or the child's guardian. We will interview cases as soon as possible by telephone, preferably within 2 weeks of notification from the laboratory. It will be at the parent's or guardian's discretion as to whether a child aged between 15 and 17 years is interviewed directly. The parent or guardian will be interviewed for cases aged less than 15 years.

DEFINITION AND SELECTION OF CONTROLS

We will recruit controls from notified cases of influenza, frequency matched by sex, age group and geographical area of residence by statistical area level 4 (SA4). These controls will be selected with a delay of at least 6 months from their influenza infection to ensure that controls have returned to eating their customary diet.

Each participating site (QLD, ACT or HNE) will establish a database of controls (previous influenza cases). All cases of influenza notified to the health department in each site between 1 January and 31 December 2017 will be entered into this control database. The age bands are 0–4

years, 5–14 years, 15–34 years, 35–54 years, 55–74 years and ≥ 75 years. An appropriate control will be randomly selected from the database within 30 days of interview of the notified case.

Case and control recruitment

Interviewers trained in computer-assisted telephone interviewing will conduct telephone interviews. A maximum of six attempts will be made to contact any one case or control, with no more than three attempts in any one day. Three calls will be attempted between 09:00 and 15:59, and three attempts between 16:00 and 20:00. A text message will be sent to the potential participant after three failed call attempts, indicating that Public Health is trying to contact them. This protocol will be continued until the person is enrolled or excluded.

QUESTIONNAIRES

We will use specific case and control questionnaires for all participants (see online supplementary appendix 1). Cases will be asked additional questions about the clinical course of their illness and treatment. Interviewers will ask identical questions regarding exposures such as foods consumed, dining locations, water sources, domestic food handling techniques and exposure to animals of cases and controls. Questions on foods consumed, dining locations, water consumed and animal and pet exposures will be asked based on a 7-day history. Questions on international travel will be asked based on a 2-week history. Antibiotic and antacid consumption, immunosuppressive treatment and household history of diarrhoea will be based on a 4-week history. Questions on food handling and general kitchen practices will be based on usual

practices rather than recent history. Demographic information will be collected from cases and controls. Contact information required to conduct interviews will be stored in a password-protected Excel document with only those needing to contact individuals given access. Piloted questionnaires were modified to remove repetitions, improve clarity and to ensure that interviews could be conducted within 20 min.

DATA HANDLING AND RISK FACTOR ANALYSIS

We will undertake descriptive reporting of campylobacteriosis incidence by person, place and time. We will also describe the severity of symptoms, treatment and burden of illness.

Risk factor analysis will involve the examination of 2×2 contingency tables with X^2 or exact tests to determine the presence of univariable associations between variables and disease. To measure the strength of an association, we will estimate ORs and calculate 95% CIs in a univariable analysis, followed by multivariable logistic regression modelling to adjust for potential confounders. Risk factors selected for inclusion in the regression model will include age, season and geographic area, variables with a significant univariable association with disease, and variables with a p value ≤ 0.25 that are biologically plausible and of interest to the research team.

LABORATORY ANALYSES

Human samples

As outlined in table 2, it is expected that 250 human isolates from HNE, 250 from QLD and 100 from ACT will be sequenced, with an additional 100 isolates being sourced from Victoria. The initial isolation and confirmation of *Campylobacter* infection will be performed locally in each state/territory. Only samples with a pure and viable culture will undergo WGS.

We will also collect an additional 20–30 human isolates from four Australian jurisdictions not participating in this case–control study to undergo WGS. This will be done over a 2-month period that overlaps with the case–control study sample collection, and is planned to help inform the generalisability of the case–control study.

Animal and food samples

We will collect samples from chicken meat (covering the two production methods of continually housed and free

range/housed), beef, lamb, pork and pet dogs. Given low prevalence of *Campylobacter* in meats other than chicken, samples will be collected from offal (preferably liver) from bovine, ovine and porcine sources to ensure sufficient positive samples are obtained for the study. Given the rising importance of chicken liver pate as a source of outbreaks in Australia,²⁷ chicken offal will also be sampled. Sample sizes by source are based on data from two states to ensure 50 positive samples per food source, and 30 samples in companion animals (table 4). We will also contact veterinary clinics and teaching hospitals to ensure sufficient *Campylobacter*-positive samples from dogs. Water samples have been omitted from the genomic aspect of this study due to logistical constraints in sampling untreated water sources across the large geographical area involved in this study, and the complexity of designing an appropriate sampling frame. As there is a lack of evidence implicating municipal drinking water as sources of *Campylobacter* infection in Australia^{1 26} we excluded water sampling from this study.

The initial isolation and confirmation of *Campylobacter* will be performed locally at laboratories in each state/territory, with isolates forwarded to the Microbiological Diagnostic Unit Public Health Laboratory for WGS, except QLD isolates which will be sequenced at Queensland Health. To detect seasonal and temporal variation in *Campylobacter* genetic types, 1041 food and animal samples (estimated to produce 330 *Campylobacter* isolates) will be collected over a period of 1 year in QLD, and 2 years in New South Wales. To assess latitudinal variation in chicken meat samples across eastern Australia, 105 chicken samples (70 chicken meat and 35 chicken offal) will be collected over a 6-month period in Victoria. Food samples will be collected monthly from retail premises, using protocols from surveys undertaken in 2014 by partner organisations, with a pilot of 30 isolates in QLD.

SEQUENCING AND SEQUENCE DATA PROCESSING

Campylobacter isolates selected for sequencing will be re-purified on solid medium and a single colony selected for preparation of genomic DNA. A sequencing library will be prepared from the genomic DNA for sequencing on the Illumina sequencing platform (MiSeq or NextSeq). A sample of the selected colony will be regrown and cryopreserved (resuspended in liquid medium supplemented with 10% glycerol and stored at -80°C). In some cases,

Table 4 Sampling to ensure 50 isolates per food source and 30 isolates from companion animals

	Foods						Animals	
	Chicken			Beef	Lamb	Pork	Dogs	Total
	Continually housed	Free range	Offal	Offal	Offal	Offal		
Assumed prevalence	0.7	0.7	0.7	0.14	0.6	0.22	0.2	
Samples required	72	72	72	286	100	272	150	1041
Positive isolates	50	50	50	40	60	60	30	330

Campylobacter enrichment cultures will be cryopreserved to enable future investigation of the genetic diversity of *Campylobacters* present. The short-read, paired-end data set produced by the Illumina Instrument from the genomic DNA of each isolate will be processed to produce a draft genome sequence for the isolate using a de novo assembler such as MEGAHIT.²⁸ The draft genome sequence will be annotated using Prokka.²⁹ We will use the draft genome sequence to perform the initial subspecies classification by deriving a multilocus sequence type (MLST) using the ‘*Campylobacter jejuni/coli*’ typing scheme (pubmlst.org). Again, using the draft genome sequence, further typing, for example, virulence factors (<http://www.mgc.ac.cn/VFs/>) or antimicrobial resistance genotype (<https://cge.cbs.dtu.dk/services/ResFinder/>), will be performed using Abricate (<https://github.com/tseemann/abricate>). We will perform comparative genomics to examine the genetic relationships between selected subgroups of isolates in more detail using Nullarbor (<https://github.com/tseemann/nullarbor>).

SOURCE ATTRIBUTION MODELLING

We will analyse the epidemiological data within designated MLST groups or other typing groups derived from the genomic sequence data. Source attribution modelling and source-assigned analyses will be conducted.

Source attribution models combine typing data from isolates from food, animals and humans to estimate the proportion of human infections that can be attributed to animal and food reservoirs.^{30–31} Once inferred MLSTs have been ascertained, the PSI²⁵ will be used to assess similarities by source. We will then undertake source attribution analyses by adapting the asymmetric island model which has previously been applied to MLST data^{25,32} using Markov chain Monte Carlo methods³³ implemented using the free software WinBUGS.³⁴ These methods will first be applied to MLST data extracted from whole genome sequences (the aforementioned ‘inferred MLSTs’), and then compared with structured phylogenetic modelling approaches^{35–36} that provide scope to infer interhost transmission.

We will then group cases according to putative source based on these source attribution methods.³⁷ For example, all isolates attributed to chicken will be grouped together, regardless of differing strains. These cases attributed to chicken will then be compared with all controls in a risk factor analysis to produce a source-assigned analysis.

SPATIAL CLUSTERS AND TEMPORAL TRENDS

We will use newly designated WGS-based MLSTs to assess heterogeneity in isolates from food sources and companion animals in QLD and New South Wales, and in isolates from chicken meat and humans across QLD, New South Wales, Victoria and ACT. A 2-year sampling framework in New South Wales, 1 year of sampling in QLD and previous survey work in these states will allow

us to assess the extent of seasonal and temporal trends. Postcode-level data associated with human illnesses will be used to detect space-time clusters using a scan statistic implemented in the free software SaTScan, at the SAI.³⁸ We will use a retrospective space-time permutation model to detect high-risk clusters by comparing the observed number of illnesses with the expected number in that geographic zone and time period.³⁹

Author affiliations

¹Faculty of Medicine, University of Queensland, Herston, Queensland, Australia

²Communicable Diseases Branch, Queensland Health, Brisbane, Queensland, Australia

³Hunter New England Population Health, Newcastle, New South Wales, Australia

⁴National Centre for Epidemiology and Population Health, The Australian National University, Canberra, Australian Capital Territory, Australia

⁵College of Health and Medicine, Melbourne Bioinformatics, Melbourne, Victoria, Australia

⁶Microbiological Diagnostic Unit Public Health Laboratory, The Peter Doherty Institute, Melbourne, Victoria, Australia

Collaborators

Study linkages and collaborations: The CampySource Project Team comprises three working groups and a reference panel. The working groups focus on: food and animal sampling, epidemiology and modelling, and genomics. The reference panel includes expert representatives from government and industry. The study is supported by the following partner organisations: the Australian National University, Massey University, University of Melbourne, Queensland Health, Queensland Health Forensic and Scientific Services, New South Wales Health, Hunter New England Health, Victorian Department of Health and Human Services, Food Standards Australia New Zealand, Commonwealth Department of Health and AgriFutures Australia—Chicken Meat Program. CampySource is also supported by collaboration with the following organisations: ACT Health, Sullivan Nicolaides Pathology, University of Queensland, Primary Industries and Regions South Australia, Department of Health and Human Services Tasmania, Meat and Livestock Australia, and New Zealand Ministry for Primary Industries. The CampySource Project Team consists of: Nigel P French, Massey University, New Zealand; Mary Valcanis, The University of Melbourne; Dieter Bulach, The University of Melbourne; Emily Fearnley, The Australian National University; Russell Stafford, Queensland Health; John Bates, Queensland Health; Trudy Graham, Queensland Health; Keira Glasgow, Health Protection NSW; Kirsty Hope, Health Protection NSW; Arie H Havelaar, The University of Florida, USA; Joy Gregory, Department of Health and Human Services, Victoria; James Flint, Hunter New England Health; Simon Firestone, The University of Melbourne; James Conlan, Food Standards Australia New Zealand; James J Smith, Queensland Health; Sally Symes, Department of Health and Human Services, Victoria; Barbara Butow, Food Standards Australia New Zealand; Liana Varrone, The University of Queensland; Linda Selvey, The University of Queensland; Deborah Denehy, ACT Health; Radomir Krsteski, ACT Health; Natasha Waters, ACT Health; Kim Lilly, Hunter New England Health; Julie Collins, Hunter New England Health; Tony Merritt, Hunter New England Health; Joanne Barfield, Hunter New England Health; Ben Howden, The University of Melbourne; Kylie Hewson, AgriFutures Australia—Chicken Meat Program; Laura Ford, The Australian National University; Liz Walker, The Australian National University; Cameron Moffatt, The Australian National University; Martyn Kirk, The Australian National University; and Kathryn Glass, The Australian National University.

Contributors MDK conceived the original idea for this study. All authors contributed to the study design and analysis plan. LV and RJS wrote the first draft with contributions from all authors. LF was heavily involved in determining timing and logistics in and between all sites. KL assisted in questionnaire design and flow. DB developed the bioinformatics analysis protocol. LV, RJS, LS, MDK and KG were involved in multiple revisions. The final version of the manuscript was approved by all authors.

Funding This CampySource project was supported by an NHMRC partnership grant 1116294 and contributions from Queensland Health, Food Standards Australia New Zealand, AgriFutures Australia—Chicken Meat Program, Commonwealth Department of Health, and New South Wales Department of Primary Industries. MDK is supported by a National Health and Medical Research Council Fellowship (GNT1145997). While undertaking studies, LV is supported through an Australian Government Research Training Program (RTP) Scholarship.

Competing interests None declared.

Patient consent Not required.

Ethics approval The Australian National University's Human Research Ethics Committee (ethics ID: 2016/426). The Hunter New England Human Research Ethics Committee (ethics ID: 17/08/16/4.03). The University of Melbourne's Animal Ethics Committee (ethics ID: 1714156).

Provenance and peer review Not commissioned; peer reviewed for ethical and funding approval prior to submission.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

REFERENCES

- OzFoodNet Working Group. Monitoring the incidence and causes of diseases potentially transmitted by food in Australia: annual report of the OzFoodNet network, 2011. *Commun Dis Intell Q Rep* 2015;39:E236–64.
- Kaakoush NO, Castaño-Rodríguez N, Mitchell HM, *et al*. Global epidemiology of campylobacter infection. *Clin Microbiol Rev* 2015;28:687–720.
- Ford L, Kirk M, Glass K, *et al*. Sequelae of foodborne illness caused by 5 pathogens, Australia, circa 2010. *Emerg Infect Dis* 2014;20:1865–71.
- Fernandes AM, Balasegaram S, Willis C, *et al*. Partial failure of milk pasteurization as a risk for the transmission of campylobacter from cattle to humans. *Clin Infect Dis* 2015;61:903–9.
- Commonwealth of Australia. National notifiable diseases surveillance system. 2018.
- Hall G, Yohannes K, Raupach J, *et al*. Estimating community incidence of Salmonella, Campylobacter, and Shiga toxin-producing Escherichia coli infections, Australia. *Emerg Infect Dis* 2008;14:1601–9.
- The Institute of Environmental Science and Research Ltd. *Notifiable diseases in New Zealand: annual report 2014*. Porirua, New Zealand: The Institute of Environmental Science and Research Ltd, 2015.
- Cressey P, Lake R. *Estimated incidence of foodborne illness in New Zealand*. New Zealand: Application of overseas models and multipliers, 2011.
- Lastovica AJ, Allos BM. Clinical Significance of Campylobacter and Related Species Other Than Campylobacter jejuni and Campylobacter coli. In: Nachamkin I, Szymanski CM, Blaser MJ, eds. *Campylobacter*. Third Edition: American Society of Microbiology, 2008.
- Fricke CR, Park RWA. A two-year study of the distribution of 'thermophilic' campylobacters in human, environmental and food samples from the Reading area with particular reference to toxin production and heat-stable serotype. *J Appl Bacteriol* 1989;67:477–90.
- Blaser MJ, Taylor DN, Feldman RA. Epidemiology of campylobacter jejuni infections. *Epidemiol Rev* 1983;5:157–76.
- Stern NJ, Hernandez MP, Blankenship L, *et al*. Prevalence and distribution of campylobacter jejuni and campylobacter coli in retail meats. *J Food Prot* 1985;48:595–9.
- Wood RC, MacDonald KL, Osterholm MT. Campylobacter enteritis outbreaks associated with drinking raw milk during youth activities. A 10-year review of outbreaks in the United States. *JAMA* 1992;268:3228–30.
- Vally H, Glass K, Ford L, *et al*. Proportion of illness acquired by foodborne transmission for nine enteric pathogens in Australia: an expert elicitation. *Foodborne Pathog Dis* 2014;11:727–33.
- Kirk M, Ford L, Glass K, *et al*. Foodborne illness, Australia, circa 2000 and circa 2010. *Emerg Infect Dis* 2014;20:1857–64.
- Heymann DL, ed. *Control of communicable diseases manual*. 19th edn. Maryland: United Book Press, Inc, 2008.
- Pebody RG, Ryan MJ, Wall PG. Outbreaks of campylobacter infection: rare events for a common pathogen. *Commun Dis Rep CDR Rev* 1997;7:R33–7.
- Tauxe RV, Hargrett-Bean N, Patton CM, *et al*. Campylobacter isolates in the United States, 1982–1986. *MMWR CDC Surveill Summ* 1988;37:1–13.
- Sammarco ML, Ripabelli G, Fanelli I, *et al*. Prevalence and biomolecular characterization of Campylobacter spp. isolated from retail meat. *J Food Prot* 2010;73:720–8.
- Whyte P, McGill K, Cowley D, *et al*. Occurrence of Campylobacter in retail foods in Ireland. *Int J Food Microbiol* 2004;95:111–8.
- Wong TL, Hollis L, Cornelius A, *et al*. Prevalence, numbers, and subtypes of Campylobacter jejuni and Campylobacter coli in uncooked retail meat samples. *J Food Prot* 2007;70:566–73.
- Noormohamed A, Fakhr MK. A higher prevalence rate of campylobacter in retail beef livers compared to other beef and pork meat cuts. *Int J Environ Res Public Health* 2013;10:2058–68.
- Samuel M, Reilly K, Shallow S, *et al*. *Burden of Campylobacter infection in the United States and declining trend in California, FoodNet 1996–1998*. Atlanta, GA: International Conference on Abstract Emerging Infectious Diseases, 2000.
- French NP, Marshall JC. *The molecular epidemiology and public health laboratory*. Massey University: Source Attribution Studies for Campylobacteriosis in New Zealand, 2014.
- Mullner P, Spencer SE, Wilson DJ, *et al*. Assigning the source of human campylobacteriosis in New Zealand: a comparative genetic and epidemiological approach. *Infect Genet Evol* 2009;9:1311–9.
- Stafford RJ, Schluter P, Kirk M, *et al*. A multi-centre prospective case-control study of campylobacter infection in persons aged 5 years and older in Australia. *Epidemiol Infect* 2007;135:978–88.
- Merritt T, Combs B, Pingault N. Campylobacter outbreaks associated with poultry liver dishes. *Commun Dis Intell Q Rep* 2011;35:299–300.
- Li D, Liu CM, Luo R, *et al*. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics* 2015;31:1674–6.
- Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;30:2068–9.
- Glass K, Fearnley E, Hocking H, *et al*. Bayesian source attribution of salmonellosis in South Australia. *Risk Anal* 2016;36:561–70.
- Hald T, Vose D, Wegener HC, *et al*. A Bayesian approach to quantify the contribution of animal-food sources to human salmonellosis. *Risk Anal* 2004;24:255–69.
- Sears A, Baker MG, Wilson N, *et al*. Marked campylobacteriosis decline after interventions aimed at poultry, New Zealand. *Emerg Infect Dis* 2011;17:1007–15.
- Mullner P, Jones G, Noble A, *et al*. Source attribution of food-borne zoonoses in New Zealand: a modified Hald model. *Risk Anal* 2009;29:970–84.
- Lunn DJ, Thomas A, Best N, *et al*. WinBUGS - A Bayesian modelling framework: Concepts, structure, and extensibility. *Stat Comput* 2000;10:325–37.
- Dearlove BL, Cody AJ, Pascoe B, *et al*. Rapid host switching in generalist Campylobacter strains erodes the signal for tracing human infections. *Isme J* 2016;10:721–9.
- Mather AE, Vaughan TG, French NP. Molecular approaches to understanding transmission and source attribution in nontyphoidal salmonella and their application in Africa. *Clin Infect Dis* 2015;61 Suppl 4(suppl_4):S259–S265.
- Mughini Gras L, Smid JH, Wagenaar JA, *et al*. Risk factors for campylobacteriosis of chicken, ruminant, and environmental origin: a combined case-control and source attribution analysis. *PLoS One* 2012;7:e42599.
- Kulldorff M, Hefner R, Hartman J, *et al*. A space-time permutation scan statistic for disease outbreak detection. *PLoS Med* 2005;2:e59.
- Touray K, Adetifa IM, Jallow A, *et al*. Spatial analysis of tuberculosis in an urban west African setting: is there evidence of clustering? *Trop Med Int Health* 2010;15:664–72.