

 Open access • Journal Article • DOI:10.1039/C7FD00014F

Investigation of the reaction of ozone with isoprene, methacrolein and methyl vinyl ketone using the HELIOS chamber. — [Source link](#)

[Yangang Ren](#), [Benoit Grosselin](#), [Véronique Daële](#), [Abdelwahid Mellouki](#)

Institutions: [Centre national de la recherche scientifique](#)

Published on: 22 Aug 2017 - [Faraday Discussions](#) (The Royal Society of Chemistry)

Topics: [Methacrolein](#), [Isoprene](#), [Methyl vinyl ketone](#) and [Ozonolysis](#)

Related papers:

- [Atmospheric fates of Criegee intermediates in the ozonolysis of isoprene](#)
- [Estimates of global terrestrial isoprene emissions using MEGAN \(Model of Emissions of Gases and Aerosols from Nature\)](#)
- [A review of Secondary Organic Aerosol \(SOA\) formation from isoprene](#)
- [Global data set of biogenic VOC emissions calculated by the MEGAN model over the last 30 years](#)
- [Isoprene photooxidation via the hydroperoxyl pathway](#)

Share this paper:    

View more about this paper here: <https://typeset.io/papers/investigation-of-the-reaction-of-ozone-with-isoprene-1v8ks9ueh1>



HAL
open science

Investigation of the reaction of ozone with isoprene, methacrolein and methyl vinyl ketone using the HELIOS chamber

Yangang Ren, Benoit Grosselin, Véronique Daële, Abdelwahid Mellouki

► To cite this version:

Yangang Ren, Benoit Grosselin, Véronique Daële, Abdelwahid Mellouki. Investigation of the reaction of ozone with isoprene, methacrolein and methyl vinyl ketone using the HELIOS chamber. *Faraday Discussions*, Royal Society of Chemistry, 2017, 200, pp.289-311. 10.1039/c7fd00014f . insu-01588945

HAL Id: insu-01588945

<https://hal-insu.archives-ouvertes.fr/insu-01588945>

Submitted on 24 May 2018

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Investigation of the reaction of ozone with isoprene, methacrolein and methyl vinyl ketone using the HELIOS Chamber

Yangang Ren, Benoit Grosselin, Véronique Daële and Abdelwahid Mellouki

Institut de Combustion, Aérodynamique, Réactivité et Environnement (ICARE), CNRS (UPR 3021), 1C Avenue de la Recherche Scientifique, 45071 Orléans Cedex 2, France

Abstract:

The rate constants for the ozonolysis of isoprene (ISO), methacrolein (MACR) and methyl vinyl ketone (MVK) have been measured using the newly built large volume atmospheric simulation chamber at CNRS-Orleans (France), HELIOS (cHambre de simuLation atmosphérique à Irradiation naturelle d'Orléans). The OH radical yields from the ozonolysis of isoprene, MACR and MVK have been also determined as well as the gas phase stable products and their yields. The secondary organic aerosol yield for the ozonolysis of isoprene has been tentatively measured in presence and absence of OH radicals scavenger. The measurements have been performed under different experimental conditions with and without adding cyclohexane (cHX) as OH radical scavenger. All experiments have been conducted at 760 torr of purified dry air ($RH \approx 2.5\%$) and ambient temperature ($T = 281-295$ K). The data obtained are discussed and compared with those from the literature. The use of the HELIOS facility and its associated analytical equipment enables to derive kinetic parameters as well as mechanistic information in near realistic atmospheric conditions.

Keywords: simulation chamber HELIOS, gas-phase, kinetic, ozonolysis, isoprene, methacrolein, methyl vinyl ketone.

Corresponding author: A. Mellouki (mellouki@cnrs-orleans.fr)

1 Introduction

Atmospheric simulation chambers are among the most advanced tools for investigating the atmospheric processes to derive physico-chemical parameters which are required for air quality and climate models. Recently, the ICARE-CNRS at Orléans (France) has set up a new large outdoor simulation chamber, HELIOS (cHambRE de simuLation atmosphérique à Irradiation naturelle d'OrléanS). The new facility has been used to study the ozonolysis of isoprene (ISO), one of the most important volatile organic compounds in the atmosphere, and its major oxidation products, methacrolein (MACR) and methyl vinyl ketone (MVK). Isoprene is the most abundant emitted non-methane hydrocarbon (NMHC) into the atmosphere; it originates mainly from biogenic sources, primarily from terrestrial vegetation. Human activities may affect the lifecycle of these biogenic species and hence change their source capacity. Isoprene is sufficiently active to affect oxidant levels in the lower troposphere and boundary. It is removed from the atmosphere mainly through reaction with OH radicals during daytime and NO₃ radicals during nighttime. However, the reaction with ozone occurs throughout the day and night and hence could have a substantial contribution to the overall removal of isoprene from the atmosphere¹. The ozonolysis of isoprene has been the subject of large number of studies, in which the reaction rate constant value has been reported at room temperature, using both absolute²⁻¹² and relative¹³⁻¹⁵ methods. However, only a limited number of investigations have dealt with the temperature dependence near atmospheric conditions. The few mechanistic studies have indicated that the ozonolysis of isoprene (CH₂C(CH₃)CHCH₂) leads the formation of methacrolein (CH₂C(CH₃)CCHO), methyl vinyl ketone (CH₂CHC(O)CH₃), and formaldehyde (HCHO) among the carbonyl products in addition to a series of intermediates

including Criegee intermediates (CIs) which are presently subject to high interest to the atmospheric scientific community ^{10, 17-20}. Indeed, the CIs can react with a number of trace species in the atmosphere to form hydroperoxides, organic acids as well as aerosols ^{3,4}.

Using the new and well equipped HELIOS facility, we have initiated studies to investigate the chemistry of isoprene and its main oxidation products under conditions close to atmospheric ones. In this first work, we report the rate constants for the reactions of O₃ with isoprene, methacrolein and methyl vinyl ketone as well as the yields of the main products formed. The OH radicals yield from these reactions has been also determined as well as the secondary organic aerosol yield from the ozonolysis of isoprene. The data obtained are discussed and compared to the ones from previous studies. While several studies have been carried out earlier to investigate the reaction of ozone with isoprene, only a limited number have been performed under realistic atmospheric conditions and most of them have been conducted in flow tube system/small chamber using high reactant concentration. (e.g. high initial reactants concentrations) ⁵. The reaction of ozone with methacrolein and methyl vinyl ketone have been investigated only in a few studies ⁵. The present work provides new insight to the atmospheric importance of these two reactions. The chemistry of the Criegee intermediates and the subsequent reactions products are not discussed in the present paper, it is subject of an ongoing work in our laboratory.

2-Experimental:

Experiments were carried out using the newly built large simulation chamber at CNRS-Orleans, HELIOS (cHambRE de simuLation atmosphérique à Irradiation naturelle d'OrléanS): The

facility consists of 90 m³ hemispherical outdoor simulation chamber (47°50'18.39N; 1°56'40.03E) made of FEP Teflon film. Two fans installed in the chamber ensure a rapid mixing of reactants (within 90 seconds). Purified air is supplied by a pure air generation system (AADCO Instruments, Inc., 737 series). Pressure (P), relative humidity (RH) and temperature (T) were continuously measured by a three-axis Ultrasonic Anemometer (Delta Ohm, HD 2003) installed in the center of the chamber. In addition, six thermocouples (PT-100), spatially and equally placed in the chamber, were used to measure continuously the temperature distribution, they were found to be within ± 1 K. The chamber is protected from “severe” weather conditions such as rain and strong wind by a mobile protective housing which is also used to keep the chamber in full dark conditions in order to conduct ozonolysis experiments such as those reported in the present work. The chamber can be fully exposed to sunlight when needed within 30 s by automatically moving the protective housing.

Organic compounds were monitored by in situ Fourier transform infrared spectrometry (Bruker Vertex70 spectrometer) coupled to a White-type multipass cell (320.6 m optical path length). Infrared spectra were recorded every 3 minutes by co-adding 250 interferograms with a resolution of 0.4 cm⁻¹. Quantitative analysis of infrared spectra was performed either by subtraction or integration of the peak area using calibrated spectra. The gas phase mixtures were also analyzed using a gas chromatography coupled to a mass spectrometer (GC-MS, PekinElmer Clarus 600 C). Gas samples were collected from the chamber onto Air Toxics trap and analyzed through a thermal desorber (TurboMatrix™ 150 ATD), with split mode, followed by a thermal desorption at 300 °C (5 min) delivering the sample to a 60-m column (GasPro diameter 0.320mm). The temperature of the GC oven was programmed as follows 25°C min⁻¹

from 180°C to 250°C and held for 25 min. Ozone concentrations were measured continuously by a chemiluminescence analyzer (HORIBA, APNA 360). The organics were also monitored by PTR-ToF-MS (Proton Transfer Reaction - Time of Flight-Mass Spectrometer, IONICON 8000). PTR-ToF-MS spectra were analyzed by the PTR-ToF Data Analyzer ⁶. HCHO was monitored continuously by an Aerolaser A4021moniror using Hantzsch reaction (Aerolaser GMBH). The detection limits for the main species of interest (isoprene, MACR, MVK, HCHO, and cyclohexanone) were typically \approx 1-2 ppb by FTIR analysis, \approx 0.2-0.5 ppb by GCMS, and \approx 0.1-0.2 ppb by PTR-ToF-MS. The precisions were \leq 7%. The measurements of HCHO by the A4021moniror had a precision and detection limit of 2% and \approx 100 ppt, respectively. Ozone concentrations measurement the chemiluminescence analyzer (HORIBA, APOA 370) had a detection limit of 1 ppb.

Isoprene (ISO), methacrolein (MACR), methyl vinyl ketone (MVK) and cyclohexane (cHX) were introduced into the chamber by placing known volumes in a bubbler and flushed by a stream of purified air. Their concentrations were derived by considering the volume of the liquid introduced, the pressure and the temperature using the ideal gas law. O₃ was generated either through a Trailigaz® ozone generator or by using a Pen-Ray® Mercury Lamp radiation through a flow of O₂ prior to be introduced into the chamber. Gaseous reactants (i.e., SF₆) were injected into the chamber using a calibrated gas cylinder equipped with capacitance manometers. In order to compensate sampling flows and leaks, a slight flow of purified air (15-25 L/min) was added continuously during all experiments maintaining a slight overpressure in

the chamber, avoiding any contamination from outside air. The dilution rate in the chamber was determined by monitoring the decay of introduced amount of SF₆ by FT-IR and was found to be typically $k_{\text{SF}_6} = (4.6 \pm 0.1) \times 10^{-6} \text{ s}^{-1}$.

Between each experiment, the chamber was cleaned by flushing pure air (800 L/min) for at least 12 hours. Background concentrations in the chamber were systematically checked and found to be below the detection limits of the available analytical instruments (e.g., [NO_x] < 1.3 × 10¹⁰, [O₃] < 1.3 × 10¹⁰ and [VOC] < 1.3 × 10⁸ molecule cm⁻³).

Chemicals. The chemicals used in this work and their stated purities were: Isoprene (Aldrich, 99%), cHX (Aldrich, 99.5%), MACR (Aldrich, 95%), MVK (Aldrich, 98%), SF₆ (Mitry-Mory 99.95%) and O₂ (Alphagaz, 99.9999%).

3- Results and discussion

3-1 Kinetic measurements:

It is well established that the ozonolysis of unsaturated organic compounds constitutes a potential non-photolytic source of OH radicals under atmospheric conditions⁷⁻⁹. Hence, in order to take that into account during our measurements, we have conducted the experiments using three different strategies, (S1) [ISO] in excess over [O₃] in the absence of cHX; (S2): [O₃] in excess over [ISO/MACR/MVK] in the presence of cHX (used as OH scavenger) and (S3): [O₃] in excess over [ISO/MACR/MVK] in the absence of cHX. Typically, initial O₃ and ISO/MACR/MVK concentrations for the (S2) and (S3) strategies were in the range 110-1000 ppb and 9-90 ppb, respectively while initial O₃ and isoprene concentrations for (S1) were 13-

35 ppb and 100-350 ppb, respectively. When added, cHX concentrations were in the range 1.5-17 ppm. Under (S2) and (S3) conditions, ISO/MACR/MVK were introduced into the chamber firstly to derive their losses in the absence of ozone which represent basically the wall loss and dilution. Under (S1) conditions, O₃ was introduced first into the chamber and its loss measured in the absence of organic reactants. Rate constants for the gas-phase reaction of O₃ with Organic (ISO/MACR/MVK) were determined by monitoring the enhanced decay rates of the O₃ or ORG (organic reactant) depending on the initial concentrations conditions. When the organic reactant is in excess, the decay of the ozone concentration can be expressed as $[O_3]_t = [O_3]_0 \times \exp(-k't)$ where $k' = k \times [ORG]_0 + k'_0$, where k (in cm³ molecule⁻¹ s⁻¹) is the rate coefficient of the ozone reaction with organic, k'_0 (in s⁻¹) is the pseudo-first order decay rate of ozone in the absence of the organic reactant and $[ORG]_0$ is the initial concentration of organic. Similarly, when O₃ was in excess, the decay of the organic is expressed as $[ORG]_t = [ORG]_0 \times \exp(-k't)$ where $k' = k \times [O_3]_0 + k'_0$, where k'_0 is the pseudo-first order decay rate of organic in the absence of O₃ and $[O_3]_0$ is the initial concentration of ozone. In our experimental conditions, $k'_{0_ISO} = (4.9 \pm 0.7) \times 10^{-6}$, $k'_{0_MACR} = (6.0 \pm 1.0) \times 10^{-6}$, $k'_{0_MVK} = (5.7 \pm 1.0) \times 10^{-6}$ and $k'_{0_O_3} = (5.0 \pm 0.4) \times 10^{-6}$ s⁻¹. In excess of O₃ when cHX was not added to scavenge OH radicals, the loss of the organic due to the reaction with OH radicals was taken into account to correct the measured $k_{O_3 + ORG}$.

Figures 1(a-d) displays examples of the pseudo-first order rate constants versus the initial concentrations of the species in excess (O₃ or organic) obtained. The slopes of these plots were used to derive the reactions rate constants. The initial experimental conditions together with the measured rate constants for the reaction of O₃ with three organics

(ISO/MACR/MVK) are listed in Tables 1-3. The runs were performed under ambient temperatures, 280 to 295 K, which were the outdoor temperature during the experiments period. Consequently, the results have been assembled by averaging the values from different runs at the same temperature ($\pm 3\text{K}$).

In the experiments where isoprene concentrations were in excess over that of O_3 , a rate constant value of $k_{\text{O}_3+\text{isoprene}} = (8.6 \pm 0.5) \times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at $285 \pm 1 \text{ K}$ was obtained. Under conditions where O_3 was in excess over the organic in the presence of cyclohexane as OH scavenger, several runs were performed at $T = 294 \pm 2$ and $285 \pm 2 \text{ K}$ and a single run at $278 \pm 1 \text{ K}$ for the reaction of O_3 with isoprene. The obtained values are: $k_{\text{O}_3+\text{isoprene}} = (11.3 \pm 1.7)$ and $(9.3 \pm 0.7) \times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ at 294 ± 2 and $285 \pm 2 \text{ K}$, respectively, in excellent agreement with the IUPAC panel recommendations using the Arrhenius expression $k = 1.03 \times 10^{-14} \exp(-1995/T)$ in the range 240-360 K, $k_{\text{O}_3+\text{isoprene}} = 11.9$ and $9.4 \times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. The rate constant value obtained in the single run at $278 \pm 1 \text{ K}$, $(6.7 \pm 0.9) \times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, is $\approx 15 \%$ lower than the IUPAC recommendation $7.9 \times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. In the absence of scavenger, the reaction rate constants values obtained at 284 ± 1 , 281 ± 1 , and $288 \pm 1 \text{ K}$ (a single run), respectively, $k_{\text{O}_3+\text{isoprene}} = (10.8 \pm 1.1)$, (9.7 ± 0.7) , and $(11.9 \pm 1.8) \times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$, have been found to be systematically $\approx 15 \%$ higher than those from the IUPAC recommendations 9.16×10^{-18} , 8.5×10^{-18} , and $10.9 \times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$. The reason for the observed differences is attributed to the contribution of the OH reaction to the consumption of isoprene when O_3 was in excess.

Regarding the O_3 reactions with MACR and MVK, experiments were conducted only in excess of O_3 in the presence and absence of OH scavenger. In the presence of scavenger, the rate

constant values obtained at $T=285\pm 1$ K for the reaction of O_3 with MACR is $k_{O_3+MACR} = (7.1 \pm 0.6) \times 10^{-19} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ which is slightly lower than the IUPAC recommended value ($k = 8.8 \times 10^{-19}$) using the expression $k = 1.4 \times 10^{-15} \exp(-2100/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ over the temperature range 240-330 K. The single run carried out at $T=287\pm 1$ K leads to slightly lower value, $k=7.9 \times 10^{-19}$, compared to the recommendation, 9.3×10^{-19} . The runs performed in absence of scavenger led to higher values: $k_{282} = (12 \pm 1) \times 10^{-19}$ and $k_{289} = (15 \pm 2) \times 10^{-19}$ compared to those in the presence of scavenger and also to IUPAC recommendations which are $k_{282} = 8.2 \times 10^{-19}$ and $k_{289} = 8.8 \times 10^{-19} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.

The O_3+MVK rate constant measured at $T=289\pm 3$ K in the presence of scavenger was found to be $k_{O_3+MVK} = (4.5 \pm 0.1) \times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ in excellent agreement with the recommended value, $k_{O_3+MVK} = 4.4 \times 10^{-18}$, using the Arrhenius expression $k = 8.5 \times 10^{-16} \exp(-1520/T) \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$ over the temperature range 240-330 K. The experiment performed in absence of the scavenger at 287 ± 2 K led to $k = (5.1 \pm 0.1) \times 10^{-18}$ which is $\approx 20\%$ higher than the IUPAC preferred one, $k = 4.3 \times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$.

3-2 Products measurements

3-2-1 OH formation yields

Cyclohexane was used to scavenge OH radicals formed during the ozonolysis of ISO, MARC and MVK. The yield of cyclohexanone produced from the reaction of OH with cyclohexane enabled to derive the OH yields during the ozonolysis of the investigated organics. Cyclohexanone was monitored by GC-MS ($m/z=98$) and PTR-ToF-MS ($m/z=81.0463$ and 99.0465). The OH yields were obtained from the equation:

$$Y_{OH} = \frac{[\text{cyclohexanone}]/\Delta[\text{organic}]}{[\text{cyclohexanone}]/\Delta[\text{cHX}]} = \frac{[\text{cyclohexanone}]/\Delta[\text{organic}]}{0.5}$$

in which cyclohexanone formation yield of $(50 \pm 7) \%$ from OH+cyclohexane reported by Atkinson et al. ¹⁰ was used.

Figure 2 shows the formation of cyclohexanone versus the consumed organics during the course of the experiments and Table 4 summarizes the experimental conditions and the obtained yield values. The results obtained are $Y_{OH} = 24.0 \pm 2.0$; 14.3 ± 3.5 and 13.4 ± 4.1 for the reactions of O_3 with isoprene, MACR and MVK, respectively. Y_{OH} from the reaction of O_3 +isoprene is in excellent agreement with recent measurement by Malkin et al. ¹¹ and Nguyen et al. ² who reported 26 ± 2 and $28 \pm 5 \%$, respectively. It is also excellent agreement with the IUPAC recommended value using the set of the literature data reported before 2005, $Y_{OH} = 25 \%$ ^{5, 12-14}. The obtained OH formation yields for O_3 +MACR and O_3 +MVK have been found to be similar, $Y_{OH} = 14\%$, in agreement with the only existing values from Aschmann et al. ¹⁵ and Paulson et al. ¹².

3-2-2 Gas phase stable products formation yields

Identified oxidation products and corresponding formation yields obtained with different analytical tools are listed in Table 5. Figures 3a-c display the typical IR spectra in the wavenumber region $750\text{-}4000 \text{ cm}^{-1}$ obtained during the experiments carried out. Isoprene, MACR, MVK and SF_6 have been monitored at 893.8 , 2730 , 998 and 948 cm^{-1} , respectively. Ozone was measured also by FTIR (at 1042 cm^{-1}) in addition to the measurement through the Horiba APOA monitor. Panels A show the spectra of organics/ O_3 / SF_6 / air mixtures at the start

of the experiments (after typically ≈ 5 min of mixing the reactants). Panels B show the spectra after few hours of reactions while panels C display the spectra of the reactions products after subtraction of the initial organic reactants/ O_3 and SF_6 . Comparison of panels C with reference spectra of formaldehyde (HCHO), carbon monoxide (CO), formic acid (HCOOH) and methylglyoxal ($CH_3C(O)C(O)H$) in the remaining panels indicates the formation of these products. In a number of runs, the PTR-ToF-MS (ISO at m/z 69.064, MACR/MVK at m/z 71.0442, cyclohexanone at m/z 81.0463 and 99.0465) and GC-MS (ISO at m/z 67, MACR/MVK at m/z 70, cyclohexanone at m/z 98) were also used to monitor the reactants and products.

Figure 4 displays examples of the temporal profiles of the reactants and products from O_3 +ISO, O_3 +MACR and O_3 +MVK obtained by FTIR, PTR-ToF-MS and HCHO-monitor. As shown, the experiments last typically for more than 20 hours each. Table 5 summarizes the experimental conditions and the obtained results along with the literature values. It has to be noted that the experiments presented in this work have been carried out at lower initial reactant concentrations compared to those reported in the previous studies. On the other hand, the experiments were performed in the temperature range 281-295 K while the literature data were mostly conducted in the range 293-298 K as shown in Table 5.

For O_3 +ISO, we have determined the formation yields for MACR, MVK, HCHO, CO and HCOOH. The measured MACR and MVK concentrations were corrected for reaction with O_3 . The yields of MACR and MVK obtained under different experimental conditions (isoprene in excess or O_3 in excess with or without OH scavenger) are in

general agreement with the literature values ^{2, 16-22} ranging from ≈ 30 to ≈ 40 % and ≈ 11 to ≈ 19 %, respectively, excluding the data from Paulson et al. ¹⁸ in which higher values have been reported in absence of OH scavenger. Formaldehyde yield obtained in the present work was found to be between 44 % and 90 % depending on the experimental conditions. $Y_{\text{HCHO}} = 44 \pm 8$ % in the runs with isoprene in excess over O_3 (in absence of cyclohexane), when O_3 was in excess $Y_{\text{HCHO}} = 69 \pm 10$ % in the presence of cyclohexane and 90 ± 2 % in its absence. The observed difference might be an indication of the occurrence of additional sources/sinks to formaldehyde under such conditions. However, as shown in Table 5, the literature data ^{2, 16, 17, 20, 21} report Y_{HCHO} ranging from 55 to 90 %. HCHO may be produced through different mechanisms involving Criegee intermediates as well as the chemistry of the OH radicals scavengers, cyclohexane or methyl-cyclohexane used in different studies. Ongoing work in our laboratory is devoted to the understanding of the specific formation of formaldehyde through the investigated reactions. The CO formation yield measured in absence of OH scavenger and in excess of isoprene, $Y_{\text{CO}} = 26 \pm 6$ %, is in excellent agreement with the earlier work by Sauer et al. ²¹ under the same conditions who reported: 26 ± 1 %. The experiments conducted under excess of O_3 in the absence and presence of OH scavenger led to higher values: $Y_{\text{CO}} = 38 \pm 2$ and 54 ± 4 %, respectively. Formic acid yields, Y_{HCOOH} , under different experimental conditions were similar, $Y_{\text{HCOOH}} = 4 \pm 1$ %, which is in excellent agreement with the measurements by Nguyen et al. ², Sauer et al. ²¹ and Neeb et al. ²³, $Y_{\text{HCOOH}} = 5 \pm 1$ %. Other products have been observed from the ozonolysis of isoprene but not mentioned here such as H_2O_2 and hydroxymethyl hydroperoxide (HMHP). This is a part of an ongoing work in our laboratory

associated to the fate of the Criegee intermediates from a series of alkenes and dienes under atmospheric conditions using HELIOS chamber.

As for O_3 +MACR and O_3 +MVK reactions, the only studies reported so far are those from Grosjean et al.²⁴ who conducted the experiments in excess of the organics in the presence of cyclohexane as the OH scavenger. They have reported yields for formaldehyde ($Y_{HCHO} = 12 \pm 3$ and 5 %) and methylglyoxal ($Y_{MGLYOX} = 58 \pm 6$ and 87 ± 5 %) for O_3 +MACR and O_3 +MVK, respectively. While a good agreement is observed between the present work and that from Grosjean et al.²⁴ on the yields of methylglyoxal, a very large discrepancy exists in the formaldehyde yields as shown in Table 5.

As mentioned above, the ozonolysis of isoprene has been subject to numerous studies under different conditions (RH, presence or absence of OH scavengers). The general reaction scheme is similar to the ozonolysis of alkenes. The initial step involves the 1,3-dipolar addition of O_3 to C=C bond (cyclo-addition), which gives rise to the production of a 1,2,3-trioxolane (primary ozonide, POZ). The POZ is a short lived species that undergoes cyclo-reversion to form carbonyl oxides or Criegee intermediates (CIs) and carbonyls (aldehydes and ketones). Two reaction pathways exist for CIs, part of CIs have sufficient internal energy and are subjected to prompt unimolecular reaction to form a hydroperoxide intermediates through H migration which subsequently decomposes or isomerizes to give OH radical, carbonyls, CO_2 and other products, some of which are potential SOA precursors. The other part of CIs would go through collisional stabilization (SCI). SCI may undergo ring closure to form dioxirane that subsequently decomposes to HO_2 radical and other products via “hot acid intermediate”. Thermally stabilized SCI may

also undergo bimolecular reactions with H₂O, HCHO and other species in the atmosphere. As isoprene is a conjugated diene, four possible product sets are formed due to two classes of cycloreversion pathway: methacrolein (MACR) and CH₂OO, formaldehyde (HCHO) and MACR oxide, methyl vinyl ketone (MVK) and CH₂OO, HCHO and MVK oxide following the formation of the two types of primary ozonides. MACR, MVK and HCHO are the dominant primary carbonyl products. The chemically activated MACR oxide and MVK oxide subsequently undergo decomposition and isomerization to form a number of products. The measured higher yield of MACR compared to MVK indicates that the O₃ reaction with isoprene occurs predominately through the attack on the CH₂=C(CH₃)- group.

The mechanisms of the ozonolysis of MACR and MVK have been subject to only one investigation each and from the same group, Grosjean et al.²⁴. O₃+MACR leads to two channels, CH₃COCHO (methylglyoxal) + [CH₂COO]* and HCHO + [CH₃COOCHO]*, while O₃+MVK leads to CH₃COCHO (methylglyoxal) + [CH₂COO]* and HCHO + [CH₃COCHOO]*. In the present work, the experiments were performed in excess of O₃ over MACR and MVK. CO, HCHO, HCOOH and methylglyoxal have been observed from both reactions. The presence or absence of scavenger did not affect significantly the measured yields of CO and HCHO but HCOOH and methylglyoxal were below detection limit our instrumentation in the experiments carried out in the presence of cyclohexane as OH scavenger. Formaldehyde formation yields were significantly higher than that reported from Grosjean et al.²⁴. who conducted the experiments in excess of organics and added cyclohexane. Methylglyoxal yields obtained in the present work in absence of cyclohexane but using O₃ in excess are in agreement with those reported by Grosjean et al.²⁴ conducted in the presence of cyclohexane under the

organics excess conditions. A large discrepancy is observed between the present measurements of the HCHO yields and those reported by Grosjean et al.²⁴. Would this be due to some difficulties in analyzing HCHO in one of the two sets of experiments? Grosjean et al.²⁴ have used HPLC analysis while we have used both in-situ FTIR and the sensitive and specific HCHO-AL4021 monitor based on Hantzsch reaction. Ongoing experiments in our laboratory are dedicated to check this possibility.

3-2-3 Secondary organic formation

A limited number of runs were carried out to investigate the occurrence of secondary organic formation (SOA) during the ozonolysis of isoprene. The main aim of these runs was to check the capabilities of our new built chamber to study SOA formation. Experiments were conducted under the same experimental conditions as those used in the kinetic and products studies (excess of isoprene or O₃ and presence or absence of OH scavenger) in absence of added aerosol seeds. Particle size distributions from 10 to 490 nm were measured with a scanning mobility particle sizer (SMPS; Model 3934, TSI Inc.). Total particle number concentrations were monitored with condensation particle counter (CPC, TSI, Inc., 3022A) along with a differential mobility analyzer (TSI, Inc., 3081). Table 6 summarizes the experimental conditions and the aerosols yields obtained and Figure 5 displays examples of temporal profiles of SOA formation distribution under various experimental conditions (different initial concentrations of isoprene and O₃, with/without OH radical scavengers). The SOA yield (Y_{SOA}) was defined as the ratio of maximum SOA produced (ΔM_0 , $\mu\text{g}/\text{m}^3$) to the mass concentration consumed ($\Delta[\text{isoprene}]$, $\mu\text{g}/\text{m}^3$), $Y_{SOA} = \Delta M_0 / (\Delta[\text{isoprene}])$ as Kleindienst et al.²⁵. A density of $1 \text{ g}/\text{cm}^3$ was applied to convert the integrated SOA volume to mass concentration. The

chamber wall loss of SOA was taken into account by applying a first order loss rate obtained from the decay of the particle volume concentration after reaching its maximum value for each individual experiment.

Figures 5 and 6 show, respectively, the prompt formation of SOA with the initiation of the reaction and their growth as function of the consumed isoprene under the three experimental conditions used in this study. The SOA yields obtained depend on the experimental conditions as shown in Table 6. In excess of isoprene and absence of cyclohexane, the SOA yield was $Y_{\text{SOA}} \approx 3.5 \pm 2.5 \%$, higher than that obtained during the experiments conducted in excess of ozone in both, without and with cyclohexane added, $Y_{\text{SOA}} \approx 1.0 \pm 0.2 \%$ and $< 1\%$, respectively. These data have to be considered as preliminary and more experiments need to be carried out under wider experimental conditions in order to characterize more precisely the SOA yields. However, it has to be noted that the earlier studies conducted on the aerosol formation from the ozonolysis of isoprene have reported formation yields of $\approx 1 \%$ (Kleindienst et al.)²⁵ or less (e.g. Jang et al. 2002, Czoschke et al. 2003)^{26, 27} depending on the experimental conditions.

4 Conclusions and future work

A series of experiments were carried out using the new built simulation chamber, HELIOS, to check its capacities in investigating complex gas phase processes. The characteristics of HELIOS enable us to conduct studies under ambient temperatures, typically around $\approx 10 \text{ }^\circ\text{C}$ ($\approx 283 \text{ K}$) from late autumn to early spring in the Orleans (France) area. The analytical equipment at the facility makes investigations under wide range concentrations of the reactants and products, from ppb to ppm levels, possible. In the current paper, we describe a first set of

data obtained on the ozonolysis of isoprene, one of the most important VOCs in the atmosphere, and its main oxidation products, methacrolein and methyl vinyl ketone. Reactions rate constants for the reactions of ozone with the above organic species have been measured in the ambient temperature range 281-295 K under different experimental conditions such as initial reactants concentrations, excess of O₃ over the organics and vis-versa in presence or absence of OH scavenger. The obtained values for $k_{O_3+organic}$ have been found in good agreement with the recommended values calculated using the Arrhenius expressions from IUPAC panel.

A section of the present work is dedicated to the OH radical and stable products formation from the investigated reactions. OH radical formation yield is reported for the three reactions, $Y_{OH} = 24 \pm 2\%$ from O₃+ISO, $14.3 \pm 3.5\%$ from O₃+MACR and $13.4 \pm 4.1\%$ from O₃+MVK. The OH formation yield has been subject to numerous studies^{2, 10, 12, 13, 18, 20, 22, 28-30}. Our measured value is in excellent agreement with the preferred value from IUPAC panel, $Y_{OH}=25\%$. The OH formation yield from O₃+MACR and O₃+MVK have been measured earlier in a very limited number of studies. The value obtained here for O₃+MACR reaction is in agreement with that from Aschmann et al.¹⁵, 20^{+10}_{-13} and that for O₃+MVK is in excellent agreement with the one measured by Aschmann et al.¹⁵, $18 \pm 8\%$ and by Paulson et al.¹² $16 \pm 5\%$. Yields for series stable oxidation products are reported and compared to the literature data. We report here formation yields for MACR, MVK, HCHO, HCOOH and CO. The general trends of the obtained values are in line with the recommended IUPAC values for O₃ reaction with isoprene. The product yields obtained

for the reactions of O₃ with MACR and MVK are also compared to those from the single available study on these reactions from Grosjean et al.²⁴.

In addition to the gas phase product, a limited number of runs were performed to estimate the SOA formation yields from ozonolysis of isoprene which showed that the yields values depend on the initial experimental conditions. However, the experiments conducted in presence of OH scavenger ($\leq 1\%$) were lower than the ones in its absence (1-3.5%) which indicates a potential contribution of OH chemistry in the SOA production observed.


Work is ongoing in our laboratory to investigate the complete chemistry of the studied reactions by using wider set of analytical equipment for the analysis of unstable species as well as peroxides and hydroperoxydes. To this aim, new instrumentation such as API-ToF-CIMS and UHPLC-MS have been recently connected to the HELIOS chamber for the analysis of the missing organic fraction as well as characterizing the aerosol composition. The chamber is large enough to enable the collection of sufficient aerosol mass for chemical analysis. In addition, the CNRS-Orleans CIMS dedicated to the OH and HO₂ measurements will be used to conduct the ongoing work.

Acknowledgements

This work is supported by Labex Voltaire (ANR-10-LABX-100-01), ANR (13-BS06-0002-01, COGNAC), ARD PIVOTS program (supported by the Centre-Val de Loire regional council), and the European Union's Horizon 2020 research and innovation programme under grant agreement No 730997 "Eurochamp2020". RY is grateful to the China Scholarship Council for the financial support.

1. R. Atkinson and J. Arey, *Chemical reviews*, 2003, **103**, 4605-4638.
2. T. B. Nguyen, G. S. Tyndall, J. D. Crouse, A. P. Teng, K. H. Bates, R. H. Schwantes, M. M. Coggon, L. Zhang, P. Feiner, D. O. Milller, K. M. Skog, J. C. Rivera-Rios, M. Dorris, K. F. Olson, A. Koss, R. J. Wild, S. S. Brown, A. H. Goldstein, J. A. de Gouw, W. H. Brune, F. N. Keutsch, J. H. Seinfeld and P. O. Wennberg, *Physical Chemistry Chemical Physics*, 2016, **18**, 10241-10254.
3. M. Sipila, T. Jokinen, T. Berndt, S. Richters, R. Makkonen, N. M. Donahue, R. L. Mauldin, III, T. Kurten, P. Paasonen, N. Sarnela, M. Ehn, H. Junninen, M. P. Rissanen, J. Thornton, F. Stratmann, H. Herrmann, D. R. Worsnop, M. Kulmala, V. M. Kerminen and T. Petaja, *Atmospheric Chemistry and Physics*, 2014, **14**, 12143-12153.
4. M. J. Newland, A. R. Rickard, L. Vereecken, A. Muñoz, M. Ródenas and W. J. Bloss, *Atmos. Chem. Phys. Discuss.*, 2015, **15**, 8839-8881.
5. R. Atkinson, D. L. Baulch, R. A. Cox, J. N. Crowley, R. F. Hampson, R. G. Hynes, M. E. Jenkin, M. J. Rossi, J. Troe and I. Subcommittee, *Atmos. Chem. Phys.*, 2006, **6**, 3625-4055.
6. M. Müller, T. Mikoviny, W. Jud, B. D'Anna and A. Wisthaler, *Chemometrics and Intelligent Laboratory Systems*, 2013, **127**, 158-165.
7. C. Schäfer, O. Horie, J. N. Crowley and G. K. Moortgat, *Geophysical Research Letters*, 1997, **24**, 1611-1614.
8. A. A. Chew and R. Atkinson, *Journal of Geophysical Research: Atmospheres*, 1996, **101**, 28649-28653.
9. M. S. Alam, A. R. Rickard, M. Camredon, K. P. Wyche, T. Carr, K. E. Hornsby, P. S. Monks and W. J. Bloss, *The Journal of Physical Chemistry A*, 2013, **117**, 12468-12483.
10. R. Atkinson and S. M. Aschmann, *Environmental Science & Technology*, 1993, **27**, 1357-1363.
11. T. L. Malkin, A. Goddard, D. E. Heard and P. W. Seakins, *Atmos. Chem. Phys.*, 2010, **10**, 1441-1459.
12. S. E. Paulson, M. Chung, A. D. Sen and G. Orzechowska, *Journal of Geophysical Research*, 1998, **103**, 25533.
13. P. Neeb and G. K. Moortgat, *The Journal of Physical Chemistry A*, 1999, **103**, 9003-9012.
14. J. H. Kroll, T. F. Hanisco, N. M. Donahue, K. L. Demerjian and J. G. Anderson, *Geophysical Research Letters*, 2001, **28**, 3863-3866.
15. S. M. Aschmann, J. Arey and R. Atkinson, *Atmospheric Environment*, 1996, **30**, 2939-2943.
16. R. M. Kamens, M. W. Gery, H. E. Jeffries, M. Jackson and E. I. Cole, *International Journal of Chemical Kinetics*, 1982, **14**, 955-975.
17. H. Niki, P. D. Maker, C. M. Savage and L. P. Breitenbach, *Environmental Science & Technology*, 1983, **17**, 312A-322A.
18. S. E. Paulson, R. C. Flagan and J. H. Seinfeld, *International Journal of Chemical Kinetics*, 1992, **24**, 103-125.
19. S. M. Aschmann and R. Atkinson, *Environmental Science & Technology*, 1994, **28**, 1539-1542.
20. R. Gutbrod, E. Kraka, R. N. Schindler and D. Cremer, *Journal of the American Chemical Society*, 1997, **119**, 7330-7342.
21. F. Sauer, C. Schäfer, P. Neeb, O. Horie and G. K. Moortgat, *Atmospheric Environment*, 1999, **33**, 229-241.
22. R. Iannone, R. Koppmann and J. Rudolph, *Atmospheric Environment*, 2010, **44**, 4135-4141.
23. P. Neeb, F. Sauer, O. Horie and G. K. Moortgat, *Atmospheric Environment*, 1997, **31**, 1417-1423.

24. D. Grosjean, E. L. Williams and E. Grosjean, *Environmental Science & Technology*, 1993, **27**, 830-840.
25. T. E. Kleindienst, M. Lewandowski, J. H. Offenberg, M. Jaoui and E. O. Edney, *Geophysical Research Letters*, 2007, **34**, L01805.
26. M. Jang, N. M. Czoschke, S. Lee and R. M. Kamens, *Science*, 2002, **298**, 814-817.
27. N. M. Czoschke, M. Jang and R. M. Kamens, *Atmospheric Environment*, 2003, **37**, 4287-4299.
28. N. M. Donahue, J. H. Kroll, J. G. Anderson and K. L. Demerjian, *Geophysical Research Letters*, 1998, **25**, 59-62.
29. A. R. Rickard, D. Johnson, C. D. McGill and G. Marston, *Journal of Physical Chemistry A*, 1999, **103**, 7656-7664.
30. A. G. Lewin, D. Johnson, D. W. Price and G. Marston, *Physical Chemistry Chemical Physics*, 2001, **3**, 1253-1261.



Figures 1(a-d) Plot of decay rate ($k' - k'_0$) as a function of $[O_3]_0$ or $[Isoprene]_0$, (a) [isoprene] in excess over $[O_3]$ in the absence of cHX; (b) $[O_3]$ in excess over [isoprene] in the presence of cHX; (c) $[O_3]$ in excess over [MACR] in the presence of cHX; (d) $[O_3]$ in excess over [MVK] in the presence of cHX.




Figure 2: Plot of cyclohexanone concentration with respect to consumed isoprene, MACR and MVK (TOF, GC: data obtained using PTR-ToF-MS or GC-MS, respectively)




Figure 3a – O_3 + isoprene: FTIR spectra acquired after 5 minutes of reaction (A) and 2 hours (B), panel C = B-A (to identify the products), panel D = is the HCHO reference spectrum. Panel E shows the residual spectrum after subtraction of features attributable to formaldehyde. Reference spectra are shown for MACR (F), CO (G), and HCOOH (H).




Figure 3b – O_3 + MACR: FTIR spectra acquired after 5 minutes of reaction (A) and 5 hours (B), panel C = B-A (to identify the products). Panel D is the HCHO reference spectrum. Panel E shows the residual spectrum after subtraction of features attributable to formaldehyde. Reference spectra are shown for methylglyoxal (F), CO (G), and HCOOH (H).




Figure 3c – $O_3 + MVK$: FTIR spectra acquired after 5 minutes of reaction (A) and 5 hours (B), panel C = B-A (to identify the products). Panel D is the HCHO reference spectrum. Panel E shows the residual spectrum after subtraction of features attributable to formaldehyde. Reference spectra are shown for methylglyoxal (F), CO (G), and HCOOH (H).




Figure 4 –Temporal profiles of reactants (isoprene/MACR/MVK, O_3) and observed products in the reactions of O_3 -isoprene (a), O_3 -MACR (b), O_3 -MVK (c). Isoprene, O_3 , CO, HCOOH and MGLYOX were monitored by FTIR, MACR and MVK were measured by PTR-ToF-MS, and HCHO was monitored by Aerolaser-4021.




Figure 5 – Examples of temporal profiles of SOA formation (number concentration) under various experimental conditions including different initial concentrations of isoprene and O_3 , with/without OH radical scavengers.




Figure 6 – SOA growth as a function of consumed isoprene concentration under different experimental conditions.

Table 1: Reaction of O₃ with isoprene: Initial experimental conditions and results from the kinetic studies

Experimental conditions	T (K)	[O ₃] ₀ (molecule cm ⁻³)	[Isoprene] ₀ (molecule cm ⁻³)	[Cyclohexane] ₀ (molecule cm ⁻³)	$k'-k'_0$ ($\pm 1\delta$) ($\times 10^{-5}$ s ⁻¹)
Isoprene in excess, without OH scavenger	286±1	3.4×10 ¹²	24.7×10 ¹¹	0	1.9±0.1
	285±1	5.6×10 ¹²	40.8×10 ¹¹	0	3.2±0.2
	285±1	5.4×10 ¹²	58.7×10 ¹¹	0	4.9±0.6
	285±1	9.0×10 ¹²	85.6×10 ¹¹	0	7.1±0.3
	283±1	6.9×10 ¹²	88.8×10 ¹¹	0	7.5±0.7
	285±1	Average: $k=(8.6\pm 0.5)\times 10^{-18}$ cm ³ molecule ⁻¹ s ⁻¹			
	294±1	4.1×10 ¹²	3.9×10 ¹¹	8.1×10 ¹³	4.7±2.2
	295±1	7.7×10 ¹²	6.6×10 ¹¹	15.9×10 ¹³	9.8±2.7/9.5±0.1 ^a
	295±1	16.5×10 ¹²	14.4×10 ¹¹	18.4×10 ¹³	19.1±2.9/19.5±0.2 ^a
	291±1	16.9×10 ¹²	14.6×10 ¹¹	29.5×10 ¹³	17.9±2.4
	294±1	23.8×10 ¹²	21.1×10 ¹¹	45.1×10 ¹³	28.1±2.9
	294±2	Average: $k=(11.3\pm 1.7)\times 10^{-18}$ cm ³ molecule ⁻¹ s ⁻¹			
O ₃ in excess, with OH scavenger	286±1	4.0×10 ¹²	2.6×10 ¹¹	7.0×10 ¹³	3.7±2.0
	282±1	5.7×10 ¹²	3.9×10 ¹¹	9.4×10 ¹³	4.5±1.4/4.4±0.1 ^a
	284±1	11.3×10 ¹²	5.6×10 ¹¹	12.8×10 ¹³	9.5±2.7/8.9±0.2 ^a
	286±1	12.6×10 ¹²	11.2×10 ¹¹	15.0×10 ¹³	11.7±1.7/10.5±0.3 ^a
	284±1	16.2×10 ¹²	8.4×10 ¹¹	22.4×10 ¹³	15.2±2.4/14.8±0.5 ^a
	283±1	28.0×10 ¹²	23.7×10 ¹¹	51.8×10 ¹³	25.2±2.0/24.5±0.5 ^a
285±2	Average: $k=(9.3\pm 0.7)\times 10^{-18}$ cm ³ molecule ⁻¹ s ⁻¹				
	278±1	15.3×10 ¹²	13.8×10 ¹¹	24.7×10 ¹³	10.4±1.2/10.1±0.2 ^a
	k=(6.7±1.0)×10 ⁻¹⁸ cm ³ molecule ⁻¹ s ⁻¹				
	283±1	3.3×10 ¹²	4.3×10 ¹¹	0	3.2±0.6
	283±1	4.0×10 ¹²	3.4×10 ¹¹	0	5.0±0.7
	285±1	11.9×10 ¹²	10.7×10 ¹¹	0	13.1±1.1
	283±1	12.8×10 ¹²	5.8×10 ¹¹	0	14.5±2.0/13.5±0.3 ^a
285±1	15.8×10 ¹²	16.3×10 ¹¹	0	17.1±2.0/17.0±0.4 ^a	
284±1	Average: $k=(10.8\pm 1.1)\times 10^{-18}$ cm ³ molecule ⁻¹ s ⁻¹				
O ₃ in excess, without OH scavenger	280±1	10.7×10 ¹²	5.3×10 ¹¹	0	10.0±2.7/9.8±0.2 ^a
	282±1	12.6×10 ¹²	12.3×10 ¹¹	0	12.3±1.5
	279±1	14.2×10 ¹²	6.9×10 ¹¹	0	13.9±3.1
	281±1	20.3×10 ¹²	9.9×10 ¹¹	0	20.0±2.6/19.2±0.4 ^a
	281±1	25.3×10 ¹²	23.0×10 ¹¹	0	24.1±2.1
	281±1	Average: $k=(9.7\pm 0.7)\times 10^{-18}$ cm ³ molecule ⁻¹ s ⁻¹			
288±1	21.3×10 ¹²	18.8×10 ¹¹	0	25.4±2.4/25.4±0.5 ^a	
k=(11.9±1.8)×10 ⁻¹⁸ cm ³ molecule ⁻¹ s ⁻¹					

^a value from PTR-ToF-MS

Table 2: Reactions of O₃ with Methacrolein (MACR): Initial experimental conditions and results from the kinetic studies

	T (K)	[ozone] ₀ (molecule cm ⁻³)	[MACR] ₀ (molecule cm ⁻³)	[cyclohexane] ₀ (molecule cm ⁻³)	$k'-k'_{loss}$ ($\pm I\delta$) ($\times 10^{-6}$ s ⁻¹)
O ₃ in excess with OH scavenger	285±1	24.9×10 ¹²	9.8×10 ¹¹	18.9×10 ¹³	16.7±1
	284±1	15.8×10 ¹²	7.7×10 ¹¹	18.4×10 ¹³	10.4±0.7
	285±1	13.1×10 ¹²	5.3×10 ¹¹	17.3×10 ¹³	9.1±0.6
	285±1	20.5×10 ¹²	32.3×10 ¹¹	29.8×10 ¹³	14.2±0.5
	285±1	6.6×10 ¹²	2.8×10 ¹¹	27.1×10 ¹³	4.7±0.8
	285±1	8.8×10 ¹²	2.4×10 ¹¹	29.0×10 ¹³	6.6±0.4
	285±1	Average: $k=(7.1\pm 0.6)\times 10^{-19}$ cm ³ molecule ⁻¹ s ⁻¹			
	287±1	18.1×10 ¹²	10.4×10 ¹¹	28.5×10 ¹³	14.2±0.8
		k=(7.9±1.2)×10 ⁻¹⁹ cm ³ molecule ⁻¹ s ⁻¹			
O ₃ in excess without OH scavenger	280±1	7.1×10 ¹²	5.9×10 ¹¹	0	8.0±0.4
	280±1	5.1×10 ¹²	2.6×10 ¹¹	0	5.5±0.3
	284±1	8.2×10 ¹²	7.6×10 ¹¹	0	10.4±0.6
	282±1	3.8×10 ¹²	2.4×10 ¹¹	0	4.7±0.4
	283±1	24.7×10 ¹²	10.1×10 ¹¹	0	30.5±1.8
	282±2	Average: $k=(1.2\pm 0.1)\times 10^{-18}$ cm ³ molecule ⁻¹ s ⁻¹			
	290±1	15.4×10 ¹²	6.2×10 ¹¹	0	24.2±0.8
	289±1	17.4×10 ¹²	11.8×10 ¹¹	0	26.5±1.2
	289±1	11.2×10 ¹²	5.1×10 ¹¹	0	17.5±0.6
	289±1	14.1×10 ¹²	5.9×10 ¹¹	0	22.2±0.8
288±1	20.0×10 ¹²	7.1×10 ¹¹	0	30.7±1.5	
289±1	Average: $k=(1.5\pm 0.2)\times 10^{-18}$ cm ³ molecule ⁻¹ s ⁻¹				

Table 3: Reactions of O₃ with Methyl vinyl ketone (MVK): Initial experimental conditions and results from the kinetic studies

	T (K)	[ozone] ₀ (molecule cm ⁻³)	[MVK] ₀ (molecule cm ⁻³)	[cyclohexane] ₀ (molecule cm ⁻³)	$k'-k'_{loss} (\pm I\delta)$ ($\times 10^{-5} \text{ s}^{-1}$)
	286±1	2.9×10 ¹²	2.3×10 ¹¹	3.9×10 ¹³	1.1±0.1
O ₃ in excess	288±1	5.1×10 ¹²	4.7×10 ¹¹	4.1×10 ¹³	2.1±0.1
with OH	288±1	6.9×10 ¹²	5.5×10 ¹¹	5.9×10 ¹³	2.8±0.2
scavenger	290±1	3.8×10 ¹²	2.8×10 ¹¹	3.9×10 ¹³	1.5±0.1
	292±1	5.4×10 ¹²	4.9×10 ¹¹	5.8×10 ¹³	2.3±0.2
	289±3	Average: $k=(4.5\pm 0.1)\times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$			
	286±1	8.5×10 ¹²	7.6×10 ¹¹	0	4.2±0.2
O ₃ in excess	287±1	8.0×10 ¹²	7.0×10 ¹¹	0	4.0±0.5
without OH	289±1	6.4×10 ¹²	6.3×10 ¹¹	0	3.1±0.4
scavenger	289±1	4.2×10 ¹²	3.0×10 ¹¹	0	2.0±0.1
	287±2	Average: $k=(5.1\pm 0.1)\times 10^{-18} \text{ cm}^3 \text{ molecule}^{-1} \text{ s}^{-1}$			

Table 4: The OH yields from the ozonolysis of isoprene, MACR and MVK: experimental conditions and results

[cHX]/ [organic] ₀	[cyclohexanone]/ Δ[organic]	Y _{OH} (%)	Method	Reference
Isoprene+O₃				
145	(12.49±0.21)×10 ⁻²	25.0±0.4	cHX as Scavenger	
130	(13.07±0.24)×10 ⁻²	26.1±0.5	PTR-ToF-MS	
240	(14.02±0.16)×10 ⁻²	28.0±0.3		
245	(13.60±0.29)×10 ⁻²	27.2±0.7		
245	(11.12±0.24)×10 ⁻²	22.2±0.5		
270	(11.99±0.20)×10 ⁻²	24.0±0.4		
220	(12.91±0.16)×10 ⁻²	25.8±0.3		
250	(12.53±0.21)×10 ⁻²	25.1±0.4		
200	(11.64±0.65)×10 ⁻²	23.3±1.3	cHX as Scavenger, GC-MS	
230	(11.31±1.21)×10 ⁻²	22.6±2.5		
180	(10.85±0.48)×10 ⁻²	21.7±0.9		
Average		24.0±2.0 *	cHX as Scavenger	This work
		26±2	cHX as Scavenger, TMB tracer, FAGE	11
		28±5	LIF and FAGE	2
		25	Recommendation	IUPAC
MACR+O₃				
192	(8.56±0.66)×10 ⁻²	17.1±1.8	cHX as Scavenger	
240	(7.99±0.66)×10 ⁻²	16.0±1.9	PTR-ToF-MS	
330	(5.01±0.37)×10 ⁻²	10.0±1.0		
240	(7.84±0.58)×10 ⁻²	15.6±1.7		
270	(6.33±0.51)×10 ⁻²	12.7±1.4		
Average		14.3±3.5	cHX as Scavenger	This work
		20 ⁺¹⁰ ₋₁₃	cHX as Scavenger GC-MS/GC-FID/GC-FTIR	15
MVK+O₃				
82	(5.47±0.66)×10 ⁻²	11.0±1.9	cHX as Scavenger,	
86	(9.39±1.08)×10 ⁻²	18.8±2.2	PTR-ToF-MS	
107	(3.8±0.70)×10 ⁻²	7.6±1.6		
140	(9.82±1.30)×10 ⁻²	19.6±2.9		
119	(5.03±1.18)×10 ⁻²	10.1±1.1		
Average		13.4±4.1	cHX as Scavenger	This work
		16±8	cHX as Scavenger GC-MS/GC-FID/GC-FTIR	15
		16±5	tracers, GC-FID	12

Errors quoted are standard deviation (SD) obtained in the regression analysis combined with estimated overall uncertainties in the PTR-ToF-MS and GC-MS response factors for isoprene, MACR, MVK and cyclohexanone. * Average values determined by PTR-ToF-MS and GC-MS.

Table 5: The product yields of the ozonolysis of isoprene, MACR and MVK under different experimental conditions.

Exp.	[organic]	[O ₃]	Y _{MACR} (%)	Y _{MVK} (%)	Y _{CO} (%)	Y _{HCOOH} (%)	Y _{HCHO} (%)	Y _{MGLYOX} (%)	T(K)	Reference
Isoprene+O ₃										
No Scavenger	2.5-8.9	0.3-0.9	29±6	10±1	26±6	3±1	45±9	-	283-286	This work*
No Scavenger	0.4-2.3	3.3-25.3	36±7	13±3	38±2	4±1	69±10	-	281-288	
w/ cHX	0.4-1.5	4.1-23.8	32±5	11±1	54±4	4±1	90±2	-	286-295	
w/ cHX	2.4	14.4	42±6	18±6	-	5	81±6	-	295	2
w/ CO	400	24	33.4±4.2	15.2±0.3	-	-	-	-	295	22
No Scavenger	127	55.2	34±1	14±1	26±1	5±1	68±3	-	295±2	21
No Scavenger	120	55			-	4	-	-		23
w/ CO	504-576	200-230	30	20	-	-	55	-	298	20
No Scavenger	504-576	200-230	28	21	-	-	54	-	298	20
w/ cHX	45-48	5	38.7±3	15.9±1.3	-	-	-	-	296±2	19
No Scavenger	45-48	5	33.9±2.6	19.1±1.5					296±2	19
w/methyl-cHX	240-272	5760	37	17	-	-	-	-	298±8	18
No Scavenger	312-408	5760	67±9	26±6					298±8	18
No Scavenger	55	77	>33	>13	-	-	85	-		17
No Scavenger	22.3	16.9	41	18	-	-	90±5	-	295	16
			39-44	16-17	-	-	90	-		IUPAC
MACR+O ₃										
No Scavenger	0.4-0.9	7.1-24.9			55±4	3±1	57±8	59±9	281-290	This work*
w/ cHX	0.3-0.8	8.7-24.9			59±4		66±4	-	285-288	
w/ cHX	21	1.4-2.1					12±3	58±6	293	24
MVK+O ₃										
No Scavenger	0.7-0.8	6.4-8.5			28±4	4±1	38±6	71±6	286-289	This work*
w/ cHX	0.5-0.8	5.4-8.7			30±8	-	44±5	-	282-292	
w/ cHX	20	2.1			-	-	5	87±5	293	24

The units of [organic] and [O₃] in $\times 10^{12}$ molecule cm^{-3} , MGLYOX=methylglyoxal. * Average values determined from different experiments. Errors quoted are 1 standard deviation (SD) of different experiments.

Table 6. Initial experimental conditions and results of secondary organic aerosol (SOA) mass concentration (ΔM_0) and SOA yield (Y_{SOA})^a.

Experimental conditions	[ISO] ₀ (ppb)	[O ₃] ₀ (ppb)	cHX (ppm)	[ISO] _{consumed} (ppb)	ΔM_0 (max) ($\mu\text{g}/\text{m}^3$)	Y_{SOA} (%)	T (K)
	354.6	27.4	-	29.0±1.6	2.9±0.15	3.3±0.3	283±1
ISO in excess w/o cHX	158.2	33.7	-	23.5±2.5	0.8±0.1	1.1±0.1	281±1
	220.6	42.5	-	35.1±3.6	6.4±0.5	6.1±1.0	281±1
O ₃ in excess w/o cHX	75.1	849.0	-	64.7±0.9	2.0±0.1	1.0±0.05	288±1
	64.6	627.5	-	54.6±1.1	1.9±0.2	1.1±0.2	285±1
	45.0	504.1	6.6	22.6±2.8	0.06±0.01	0.09±0.02	286±1
O ₃ in excess w/ cHX	58.6	670.9	7.5	31.4±4.5	0.25±0.01	0.26±0.04	295±1
	90.1	965.4	16.9	78.3±1.8	2.1±0.12	0.89±0.09	294±1
	58.4	672.5	11.8	32.6±2.9	0.52±0.05	0.53±0.07	291±1

^a Stated uncertainties were from scatter in particle volume measurements; ^b Assuming a density of 1.0 g/cm³; ^c SOA yields were obtained from the maximum aerosol volume;