



Invisible decays of ultra-high energy neutrinos

Luis Dorame¹, Omar G. Miranda^{2*} and J. W. F. Valle¹

¹ Astroparticle and High Energy Physics Group, Instituto de Física Corpuscular, Consejo Superior de Investigaciones Científicas/Universitat de València, Valencia, Spain

² Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN México, DF, Mexico

Edited by:

Stefano Morisi, University of Würzburg, Germany

Reviewed by:

Alexander Merle, University of Southampton, UK

Alessandro Mirizzi, University of Hamburg, Germany

*Correspondence:

Omar G. Miranda, Departamento de Física, Centro de Investigación y de Estudios Avanzados del IPN, Av. IPN 2508, Apdo. Postal 14-740, México, DF, Mexico
e-mail: omr@fis.cinvestav.mx

Gamma-ray bursts (GRBs) are expected to provide a source of ultra high energy cosmic rays, accompanied with potentially detectable neutrinos at neutrino telescopes. Recently, IceCube has set an upper bound on this neutrino flux well below theoretical expectation. We investigate whether this mismatch between expectation and observation can be due to neutrino decay. We demonstrate the phenomenological consistency and theoretical plausibility of the neutrino decay hypothesis. A potential implication is the observability of majoron-emitting neutrinoless double beta decay.

Keywords: neutrino interactions, cosmic rays

The source of ultra high energy cosmic rays remains a mystery. In gamma-ray burst (GRB) models such as the fireball model cosmic-ray acceleration should be accompanied by neutrinos produced in the decay of charged pions created in interactions between the high-energy protons and γ -rays [39]. Recently the Ice-Cube collaboration reported an upper limit on the flux of energetic neutrinos associated with GRBs almost four times below this prediction [3].

Various possible explanations have been considered to explain the non-observation of this ultra high energy neutrino flux. For example, a complete detailed numerical analysis of the fireball neutrino model predicts a neutrino flux that is one order of magnitude lower than the analytical computations [24]. On the other hand, another recent computation [22] of the neutrino flux in the fireball model gives a mild reduction in the neutrino flux if a relation between the bulk Lorentz factor, Γ , and the burst energy is assumed. Finally, based on the specific case of GRB 130427A, it has been argued that the low neutrino flux can be explained with relatively large values for the bulk Lorentz factor ($\Gamma \geq 500$) and for the dissipation radius ($R_d \geq 10^{14}$ cm); it was shown in the same reference that both the internal shock and the baryon photosphere models satisfied these conditions.

Here we focus on a different approach to explain the neutrino flux deficit. Instead of studying the astrophysical mechanism of the source objects, we look for a high energy physics explanation. Some mechanisms involving new physics in order to explain a possible deficit in the observed neutrino flux have already been suggested. For instance, the possibility of an oscillation involving a quasi Dirac neutrino [38] has been considered in Reference [13]; a specific model for this case has been studied in [25] and the possibility of a resonance effect has also been discussed [29]. Another mechanism recently discussed has been the case of a spin precession into a sterile neutrino as a result of a non-zero neutrino magnetic moment [35] and the strong magnetic fields expected to be present in a GRB [8].

Here we speculate on the plausibility of the neutrino decay hypothesis as a possible explanation for the mismatch between observation and expectation. The most attractive possibility involves invisible decays, which have been considered theoretically since the eighties [17, 18, 21, 34, 37], and recently revisited for the case of GRB neutrino fluxes [31]. These decays arise in models with spontaneous violation of ungauged lepton number [12], though typically suppressed [34]. A natural scenario to test neutrino stability are astrophysical objects [7, 9, 10, 27]. In particular, limits on Majoron couplings from solar and supernova neutrinos have been obtained in Reference [26]. For non-astrophysical constraints, for example from $0\nu\beta\beta$ searches, see [5, 16, 28, 36]. Moreover, as already mentioned, recent results from the Pierre Auger Observatory (PAO) [4], ANTARES [11] and IceCube [2] have placed strong constraints on the neutrino flux coming from distant ultra high energy (UHE) neutrino sources.

Here we explore the phenomenological plausibility and theoretical consistency of the decay hypothesis within a class of low-scale $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ seesaw schemes with spontaneous family-dependent lepton number violation. We show that the required neutrino decay lifetime range hinted by the non-observation of UHE muon neutrinos is theoretically achievable for the majoron-emitting neutrino decays and, moreover, consistent with all existing phenomenological constraints.

The decay rate $\nu_i \rightarrow \nu_j + J$ in the rest frame of ν_i is

$$\Gamma(\nu_i \rightarrow \nu_j + J) = \frac{g_{ij}^2}{16\pi} \frac{(m_i + m_j)^2}{m_i^3} (m_i^2 - m_j^2), \quad (1)$$

where ν_i and ν_j are active neutrinos and J is a massless or very light majoron associated to the spontaneous violation of ungauged lepton number. Taking $m_j = 0$, we can estimate the decay length (in meters) for a relativistic neutrino as given by

$$L = c\tau = \frac{cE_i}{m_i\Gamma} \simeq 1 \times 10^9 \left(g_{ij}^{-2}\right) \left(\frac{E_i}{100\text{TeV}}\right) \left(\frac{1\text{eV}}{m_i}\right)^2 \text{ m}. \quad (2)$$

For typical AGN distances we obtain the required values of g_{ij} for a given neutrino mass m_i which would cause decay before reaching the detector. In **Figure 1** we took AGN distances from 3.6 Mpc (the distance to Centaurus A) up to 100 Mpc. In the bottom panel of the same **Figure 1** we have plotted the corresponding result for GRBs, at typical distances of $10\text{--}10^3$ Mpc. The vertical lines correspond to the relevant region for m_2 and m_3 for the case $m_1 = 0$. We have explicitly verified that, for the GRB case, this approximation is in agreement with more detailed estimates of neutrino lifetime ranges [6].

As we will discuss below, putting into a theory context, such couplings are fairly large to achieve theoretically. In order to have an estimate of the neutrino flux reduction resulting from neutrino decay we note that, since coherence is lost, the final flux of a given neutrino flavor will be

$$\phi_{\nu_\alpha}(E) = \sum_{i\beta} \phi_{\nu_\beta}^{\text{source}}(E) |U_{\beta i}|^2 |U_{\alpha i}|^2 e^{-L/\tau_i(E)}, \quad (3)$$

where ϕ 's are neutrino fluxes at production and detection, L the travel distance, τ_i the neutrino lifetime in the laboratory frame and $U_{\alpha i}$ the elements of the lepton mixing matrix [33]. Typical neutrino energies lie in the range of 10^5 TeV and 10^3 TeV for AGNs and GRBs, respectively. Note that, in the limit that $L \gg \tau_i$ where only the stable state survives

Equation (3) becomes



$$\phi_{\nu_\alpha}(E) = \sum_{i(\text{stable})\beta} \phi_{\nu_\beta}^{\text{source}}(E) |U_{\beta i}|^2 |U_{\alpha i}|^2. \quad (4)$$

Here we take a normal hierarchy neutrino mass spectrum, the disappearance of all states except the lightest (in this case ν_1) is allowed. The final flux of ν_e , ν_μ and ν_τ can be computed from Equation (4) and will depend on the three mixing angles and the Dirac CP phase δ . In particular, we can calculate the suppression of the muon neutrino flux, ϕ_{ν_μ} , using the ratio

$$R_{\nu_e:\nu_\mu} \equiv \frac{\phi_{\nu_e}}{\phi_{\nu_\mu}} = \left(\frac{\cos \theta_{12} \cos \theta_{13}}{|-\sin \theta_{12} \cos \theta_{23} - \sin \theta_{13} \sin \theta_{23} \cos \theta_{12} e^{i\delta}|} \right)^2, \quad (5)$$

where θ_{12} , θ_{23} and θ_{13} are the neutrino mixing angles determined in neutrino oscillation experiments. The left panel of **Figure 2** shows the expected values for this ratio when the neutrino mixing angles lie within the 1σ bands from their current global best fit values [14, 15, 20]. One can also see, in the right panel of the same **Figure 2** that by allowing these parameters to vary up to their three sigma ranges, $R_{\nu_e:\nu_\mu}$ can be as large as 25 or as low as 2. Very similar results are found for global fit of Reference [20], as shown in **Table 1**.

It is important to notice that, in this picture, the neutrino decay will lead to a decrease in the muon neutrino flux while



the electron neutrino flux will increase. Alternatively in the presence of light sterile states one can envisage a scenario where the muon neutrino decays to the sterile state. Here we do not consider this case. Recent data from Icecube reports the observation of two neutrinos with energies around 10^{15} eV, probably electron neutrinos [1]. Moreover, there have been recent announcements of more neutrino events detected in IceCube [40].

We now turn to the issue of theoretical consistency of the decay hypothesis. In most $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$ seesaw models with spontaneous lepton number violation when one diagonalizes the neutrino mass matrix one also diagonalizes, to first approximation, the coupling of the resulting Nambu-Goldstone boson to the mass eigenstate neutrinos [34]. The exact form of the light-neutrino majoron couplings can be determined explicitly by perturbative diagonalization of the seesaw mass matrix, or by using a more general approach using only the symmetry properties. The result is [34]

$$g_{ij} = -\frac{m_i}{v_1} \delta_{ij} + \left[\frac{m_i}{v_1} \left(V_1^\dagger D^* M^{*-1} M^{-1} D^T V_1 \right)_{ij} \right]_S + \dots \quad (6)$$

where the subscript S denotes symmetrization, D and M are the Dirac and Majorana mass terms in, say, the type-I seesaw scheme and V_1 is the light neutrino diagonalization matrix. One sees that the majoron couples proportionally to the light neutrino mass, hence the coupling matrix is diagonal to first approximation. The off-diagonal part of the g_{ij} is inversely proportional to three powers of lepton number violation scale $v_1 \equiv \langle \sigma \rangle$, since $M \propto v_1$. This is tiny, the only hope being to use a seesaw scheme that allows for a very low lepton number violation scale, such as the inverse seesaw [30]. The particle content is the same as that of the Standard Model (SM) except for the addition of a pair of two component gauge singlet leptons, $\bar{\nu}_i^c$ and S_i , within each of the three generations, labeled by i . The isodoublet neutrinos ν_i and the fermion singlets S_i have the same lepton number, opposite with respect to that of the three singlets $\bar{\nu}_i^c$ associated to the “right-handed” neutrinos. In the ν, ν^c, S basis the 9×9 neutral lepton mass matrix M_ν has the form:

$$M_\nu = \begin{bmatrix} 0 & m_D^T & 0 \\ m_D & 0 & M^T \\ 0 & M & \mu \end{bmatrix}, \quad (7)$$

Table 1 | Allowed values of $R_{\nu_e:\nu_\mu}$ according to different neutrino oscillation fits.

Global fit	1st octant	2nd octant	At 3σ
[15]	2–7	3–14	2–25
[20]	2–7	3–12	2–23
[14]	2–6	–	2–22

The second column shows the allowed region for the case of a θ_{23} in the first octant, while the third column shows the second octant case. Finally, the fourth column displays the allowed region if we consider the 3σ confidence level for θ_{23} .

where $m_D \propto \langle \Phi \rangle$ is the standard Dirac term coming from the SM Higgs vev and M is a bare mass term. The term $\mu \propto \langle \sigma \rangle$, the vacuum expectation value of σ responsible for spontaneous low-scale lepton number violation as proposed in [21]. This gives rise to a majoron J ,

$$J = \sqrt{2} \text{Im}\sigma. \quad (8)$$

As a result of diagonalization one obtains an effective light neutrino mass matrix. Note that lepton number symmetry is recovered as $\mu \rightarrow 0$, making the three light neutrinos strictly massless. The majoron couplings of the light mass eigenstate neutrinos are determined again as a sum of two pieces as in Equation (6). Detailed calculation shows that its off-diagonal part behaves as $P' \sim \mu^2 D^2 M^{-4}$. Even if the M can be significantly lower than that of the standard high-scale type-I seesaw it is clear that this is way too small in order to produce neutrino decay within the relevant astrophysical scales.

The only way out is to induce a mismatch between the neutrino mass basis and the coupling basis. This can be achieved by making lepton number a family-dependent symmetry [17, 18]. The model is by no means unique, here we give an example based on $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \otimes U(1)_H$ assigned as shown in **Table 2**.

The $SU(3)_c \otimes SU(2)_L \otimes U(1)_Y \otimes U(1)_H$ invariant Lagrangian would be

$$\mathcal{L}_\nu = m_{Dij} \bar{L}_i \nu_{Rj} h + M_{ij} \bar{\nu}_{Ri}^c S_j + M_{\sigma ij} \bar{\nu}_{Ri}^c S_j \sigma + \mu_{ij} \bar{S}_i S_j + \mu_{\sigma ij} \bar{S}_i S_j \sigma^* + h.c., \quad (9)$$

where the relevant sub-matrices are

$$m_D = \begin{bmatrix} m_a & m_b & 0 \\ m_c & m_d & 0 \\ 0 & 0 & m_e \end{bmatrix}, \quad M = \begin{bmatrix} 0 & 0 & M_1 \\ 0 & 0 & M_2 \\ 0 & 0 & 0 \end{bmatrix}, \quad (10)$$


$$M_\sigma = \begin{bmatrix} M_3 & M_4 & 0 \\ M_5 & M_6 & 0 \\ 0 & 0 & M_7 \end{bmatrix} \quad (11)$$

$$\mu = \begin{bmatrix} M_8 & M_9 & 0 \\ M_{10} & M_{11} & 0 \\ 0 & 0 & 0 \end{bmatrix}, \quad \mu_\sigma = \begin{bmatrix} 0 & 0 & M_{12} \\ 0 & 0 & M_{13} \\ M_{14} & M_{15} & 0 \end{bmatrix} \quad (12)$$

One can check explicitly that the first term in Equation (6) is already non-diagonal and, for sufficiently low values of the $U(1)_H$


Table 2 | Model field representation content and transformation properties.

	\bar{L}_e	\bar{L}_μ	\bar{L}_τ	ν_{Re}	$\nu_{R\mu}$	$\nu_{R\tau}$	h	S_1	S_2	S_3	σ
$SU(2)$	2	2	2	1	1	1	2	1	1	1	1
$U(1)_H$	-2	-2	-4	2	2	4	0	0	0	-2	-2



breaking scale can induce a decay sufficiently fast as to suppress the flux of ϕ_{ν_μ} to account for its non-observation by IceCube.

As an additional interesting feature of this scheme, we propose an indirect test of our neutrino decay hypothesis through the single majoron-emitting $J0\nu\beta\beta$ decay mode [19]



The decay rate for single Majoron emission is given by [32]

$$\Gamma^{0\nu} = |\langle g_{ee} \rangle|^2 |\mathcal{M}_J|^2 G_J(Q, Z), \quad (14)$$

where $|\langle g_{ee} \rangle|$ is an averaged coupling constant, $G_J(Q, Z)$ accounts for the phase space factor and the nuclear matrix element (NME) \mathcal{M}_J depends on the mechanism and the relevant nucleus. For single Majoron emission one can use the same NMEs from the standard $0\nu\beta\beta$ decay [23].

We can see from Equation (14) that the decay width for single Majoron emission in neutrinoless double beta decay depends on the coupling constant g_{ee} and it is therefore an indirect relation with the expression in Equation (1) through the coupling g_{12} .

Indeed if the majoron exists and its coupling to the electron neutrino is not expected to significantly differ from the one required to explain the muon neutrino deficit in IceCube through the neutrino decay hypothesis, there will be a correlation between $R_{\nu_e:\nu_\mu}$ and $\langle g_{ee} \rangle$. This correlation is depicted in **Figure 3**. We plot in the left panel the correspondence between g_{ee} and g_{12} when we fix the neutrino mixing angles at their best fit values and we consider a Dirac mass entry at $m_D \sim 10$ GeV, $M \sim 1$ TeV $M_\sigma \sim 1$ TeV and $\mu \sim 1$ keV where the \sim sign takes into account order one differences among the various flavor components of each block. In the right panel of the same figure, we take one of the points shown in the left panel and vary its CP phase from 0 to 2π in order to obtain an estimate for the ratio $R_{\nu_e:\nu_\mu}$ relevant at IceCube. The dotted (blue) curve corresponds to the case when we consider the second octant of the atmospheric mixing angle, particularly its central value $\sin^2 \theta_{23} = 0.61$ (we have also chosen the central values of the other mixing angles, $\sin^2 \theta_{12} = 0.320$ and $\sin^2 \theta_{13} = 0.0246$); from this case it is possible to see that, for example, for a coupling constant $\langle g_{ee} \rangle = 4.33 \times 10^{-6}$ a reduction by a factor five in the muon flux can be

obtained for an appropriate value of the CP phase, $\delta = 2.5$, while the suppression could be as high as a factor 10. The solid (red) line corresponds to a point in the first octant ($\sin^2 \theta_{23} = 0.427$). Although in this particular case the values of the ratio, $R_{\nu_e:\nu_\mu}$ are lower than for the second octant, one can still achieve an important suppression; in particular, we can see that a reduction by a factor five is again possible (for the values $\langle g_{ee} \rangle = 7.34 \times 10^{-6}$ and $\delta = 1.6$) This is an interesting observation, considering that currently the “preferred” octant is not yet uniquely determined by the neutrino oscillation fits [14, 15, 20]. Moreover, we can see that even with the central values of the neutrino mixing angles one can obtain a suppression factor of five or higher that could be sufficient to explain the limits reported by the IceCube collaboration [3].

In conclusion one sees that the decay hypothesis invoked to account for the IceCube results may be tested in the upcoming searches for the $J0\nu\beta\beta$ decay.

Work supported by MINECO grants FPA2011-22975 and MULTIDARK Consolider CSD2009-00064, by Prometeo/2009/091 (Gen. Valenciana), and by EPLANET. L. D. is supported by JAE Predoctoral fellowship. O. G. M. was supported by CONACyT grant 132197.

REFERENCES

1. Aartsen MG, Abbasi R, Abdou Y, Ackermann M, Adams J, Aguilar JA, et al. [IceCube Collaboration], First observation of PeV-energy neutrinos with IceCube. *Phys Rev Lett.* (2013) **111**:021103 [arXiv:1304.5356 [astro-ph.HE]]. doi: 10.1103/PhysRevLett.111.021103
2. Abbasi R, Abu-Zayyad T, Allen M, Amann J, Archbold G, Belov K, et al. An upper limit on the electron-neutrino flux from the HiRes detector. (2008). [arXiv:0803.0554]. doi: 10.1086/590335
3. Abbasi, R., Abdou, Y., Abu-Zayyad, T., Ackermann, M., Adams, J., Aguilar, J. A., et al. An absence of neutrinos associated with cosmic-ray acceleration in γ -ray bursts. *Nature* (2012) **484**:351–3. doi: 10.1038/nature11068
4. Abreu P, Aglietta M, Ahn EJ, Albuquerque IFM, Allard D, Allekotte I, et al. The pierre auger observatory III: other astrophysical observations. (2011). [arXiv:1107.4805].
5. Argyriades J, Arnold R, Augier C, Baker J, Barabash AS, Basharina-Freshville A, et al. Measurement of the two neutrino double beta decay half-life of Zr-96 with the NEMO-3 detector. *Nucl Phys.* (2010) **A847**:168–79. doi: 10.1016/j.nuclphysa.2010.07.009
6. Baerwald P, Bustamante M, Winter W. Neutrino decays over cosmological distances and the implications for neutrino telescopes. *JCAP* (2012) **1210**:020. doi: 10.1088/1475-7516/2012/10/020

7. Bahcall JN, Petcov S, Toshev S, Valle J. Tests of neutrino stability. *Phys Lett.* (1986) **B181**:369–74. doi: 10.1016/0370-2693(86)90065-1
8. Barranco J, Miranda O, Moura C, Parada A. A reduction in the UHE neutrino flux due to neutrino spin precession. *Phys Lett.* (2012) **B718**:26–9. doi: 10.1016/j.physletb.2012.10.024
9. Beacom JF, Bell NF. Do solar neutrinos decay? *Phys Rev.* (2002) **D65**:113009. doi: 10.1103/PhysRevD.65.113009
10. Beacom JF, Bell NF, Hooper D, Pakvasa S, Weiler TJ. Decay of high-energy astrophysical neutrinos. *Phys Rev Lett.* (2003) **90**:181301. doi: 10.1103/PhysRevLett.90.181301
11. Biagi S. Upper limit on the diffuse flux of cosmic ν_μ with the antare neutrino telescope. *Nucl Phys Proc Suppl.* (2011) **212–213**:109–114. doi: 10.1016/j.nuclphysbps.2011.03.015
12. Chikashige Y, Mohapatra RN, Peccei R. Are there real goldstone bosons associated with broken lepton number? *Phys Lett.* (1981) **B98**:265. doi: 10.1016/0370-2693(81)90011-3
13. Esmaili A, Farzan Y. Implications of the pseudo-dirac scenario for ultra high energy neutrinos from GRBs. *JCAP* (2012) **1212**:014. doi: 10.1088/1475-7516/2012/12/014
14. Fogli G, Lisi E, Marrone A, Montanino D, Palazzo AM, Rotunno AM. Global analysis of neutrino masses, mixings and phases: entering the era of leptonic CP violation searches. *Phys Rev.* (2012) **D86**:013012. doi: 10.1103/PhysRevD.86.013012
15. Forero D, Tortola M, Valle J. Global status of neutrino oscillation parameters after Neutrino-2012. *Phys Rev.* (2012) **D86**:073012. doi: 10.1103/PhysRevD.86.073012
16. Gando A, Gando Y, Hanakago H, Ikeda H, Inoue K, Kato R, et al. Limits on majoron-emitting double-beta decays of Xe-136 in the KamLAND-Zen experiment. *Phys Rev.* (2012) **C86**:021601. doi: 10.1103/PhysRevC.86.021601
17. Gelmini G, Schramm DN, Valle J. Majorons: a simultaneous solution to the large and small scale dark matter problems. *Phys Lett.* (1984) **B146**:311. doi: 10.1016/0370-2693(84)91703-9
18. Gelmini G, Valle J. Fast invisible neutrino decays *Phys Lett.* (1984) **B142**:181. doi: 10.1016/0370-2693(84)91258-9
19. Georgi HM, Glashow SL, Nussinov S. Unconventional model of neutrino masses. *Nucl Phys.* (1981) **B193**:297. doi: 10.1016/0550-3213(81)90336-9
20. Gonzalez-Garcia M, Maltoni M, Salvado J, Schwetz T. Global fit to three neutrino mixing: critical look at present precision. *JHEP* (2012) **1212**:123. doi: 10.1007/JHEP12(2012)123
21. Gonzalez-Garcia M, Valle J. Fast decaying neutrinos and observable flavor violation in a new class of majoron models. *Phys Lett.* (1989) **B216**:360. doi: 10.1016/0370-2693(89)91131-3
22. He H-N, Liu R-Y, Wang X-Y, Nagataki S, Murase K, Dai Z-G. Icecube non-detection of GRBs: constraints on the fireball properties. *Astrophys J.* (2012) **752**:29. doi: 10.1088/0004-637X/752/1/29
23. Hirsch M, Klapdor-Kleingrothaus H, Kovalenko S, Pas H. On the observability of majoron emitting double beta decays. *Phys Lett.* (1996) **B372**:8–14. doi: 10.1016/0370-2693(96)00038-X
24. Hummer S, Baerwald P, Winter W. Neutrino emission from gamma-ray burst fireballs, revised. *Phys Rev Lett.* (2012) **108**:231101. doi: 10.1103/PhysRevLett.108.231101
25. Joshipura AS, Mohanty S, Pakvasa S. Pseudo-dirac neutrinos via mirror-world and depletion of UHE neutrinos. (2013). [arXiv:1307.5712].
26. Kachelriess M, Tomas R, Valle J. Supernova bounds on Majoron emitting decays of light neutrinos. *Phys Rev.* (2000) **D62**:023004. doi: 10.1103/PhysRevD.62.023004
27. Keranen P, Maalampi J, Peltoniemi J. Testing neutrino instability with active galactic nuclei. *Phys Lett.* (1999) **B461**:230–4. doi: 10.1016/S0370-2693(99)00862-X
28. Lessa A, Peres O. Revising limits on neutrino-majoron couplings. *Phys Rev.* (2007) **D75**:094001. doi: 10.1103/PhysRevD.75.094001
29. Miranda O, Moura C, Parada A. Sterile neutrinos, dark matter, and resonant effects in ultra high energy regimes. (2013). [arXiv:1308.1408].
30. Mohapatra R, Valle J. Neutrino mass and baryon number nonconservation in superstring models. *Phys Rev.* (1986) **D34**:1642. doi: 10.1103/PhysRevD.34.1642
31. Pakvasa S, Joshipura A, Mohanty S. Explanation for the low flux of high energy astrophysical muon-neutrinos. *Phys. Rev. Lett.* (2013) **110**:171802. doi: 10.1103/PhysRevLett.110.171802
32. Rodejohann W. Neutrino-less double beta decay and particle physics. *Int J Mod Phys.* (2011) **E20**:1833–1930. doi: 10.1142/S0218301311020186
33. Schechter J, Valle J. Neutrino masses in SU(2) x U(1) theories. *Phys Rev.* (1980) **D22**:2227. doi: 10.1103/PhysRevD.22.2227
34. Schechter J, Valle J. Neutrino decay and spontaneous violation of lepton number. *Phys Rev.* (1982) **D25**:774. doi: 10.1103/PhysRevD.25.774
35. Schechter J, Valle JWF. Majorana neutrinos and magnetic fields. *Phys Rev.* (1981) **D24**:1883.
36. Tomas R, Pas H, Valle J. Generalized bounds on Majoron-neutrino couplings. *Phys Rev.* (2001) **D64**:095005. doi: 10.1103/PhysRevD.64.095005
37. Valle J. Fast neutrino decay in horizontal majoron models. *Phys Lett.* (1983) **B131**:87. doi: 10.1016/0370-2693(83)91097-3
38. Valle J. Neutrinoless double beta decay with quasi dirac neutrinos. *Phys Rev.* (1983) **D27**:1672–4. doi: 10.1103/PhysRevD.27.1672
39. Waxman E, Bahcall JN. High-energy neutrinos from cosmological gamma-ray burst fireballs. *Phys Rev Lett.* (1997) **78**:2292–5. doi: 10.1103/PhysRevLett.78.2292
40. Whitehorn N, Kopfer C, Neilson NK. *Talk Given at the IceCube Particle Astrophysics Symposium (IPA)*. Madison, WI. (2013).

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Received: 03 October 2013; accepted: 14 November 2013; published online: 03 December 2013.

Citation: Dorame L, Miranda OG and Valle JWF (2013) Invisible decays of ultra-high energy neutrinos. *Front. Physics* 1:25. doi: 10.3389/fphy.2013.00025

This article was submitted to High-Energy and Astroparticle Physics, a section of the journal *Frontiers in Physics*.

Copyright © 2013 Dorame, Miranda and Valle. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.