

Update on Salt Stress

Ion Homeostasis in NaCl Stress Environments¹

Xiaomu Niu, Ray A. Bressan, Paul M. Hasegawa*, and José M. Pardo

Center for Plant Environmental Stress Physiology, Purdue University, 1165 Horticulture Building, West Lafayette, Indiana 47907–1165 (X.N., R.A.B., P.M.H.); and Instituto de Recursos Naturales y Agrobiología, Consejo Superior de Investigaciones Científicas, Apdo 1052, Sevilla-41080, Spain (J.M.P.)

Homeostasis can be defined as the tendency of a cell or an organism to maintain internal steady state, even in response to any environmental perturbation or stimulus tending to disturb normality, because of the coordinate responses of its constituent components. Typically, ions constantly flux in and out of cells in a controlled fashion with net flux adjusted to accommodate cellular requirements, thus creating an ionic homeostasis. When plant cells are exposed to salinity, mediated by high NaCl concentrations, kinetic steady states of ion transport for Na⁺ and Cl⁻ and other ions, such as K⁺ and Ca²⁺, are disturbed (Binzel et al., 1988). High apoplastic levels of Na⁺ and Cl⁻ alter aqueous and ionic thermodynamic equilibria, resulting in hyperosmotic stress, ionic imbalance, and toxicity. Thus, it is vital for the plant to re-establish cellular ion homeostasis for metabolic functioning and growth, that is, to adapt to the saline environment.

Comparisons of what have been interpreted to be adaptive responses among various species lead to the conclusion that some salt-tolerant plants have evolved specialized complex mechanisms that allow adaptation to saline stress conditions. In fact, these unique mechanisms, such as salt glands, exist in few plant species and cannot be presumed to be ubiquitously functional for salt adaptation of all plants. However, intrinsically cellular-based mechanisms appear to be common to all genotypes and are a requisite for salt tolerance. Of paramount importance are those mechanisms that function to regulate ion homeostasis while mediating osmotic adjustment through the accumulation and intracellular compartmentation of ions that are predominant in the external environment. In this update we will focus principally on Na⁺ homeostasis in sodic environments; however, we also include discussions of H⁺, K⁺, Ca²⁺, and Cl⁻ because of the interrelationship of these ions with Na⁺ homeostasis. Ion transport processes across the plasma membrane and the tonoplast will be empha-

sized because these are presumed to be most essential for the control of intracellular Na⁺ uptake and vacuolar compartmentation.

ION TRANSPORT ACROSS PLANT CELL MEMBRANES

Since ions are hydrated in solution and do not readily traverse the hydrophobic lipid bilayer of membranes, flux across the plasma membrane and tonoplast occurs via transport proteins. Ion flux across membranes is dependent on the thermodynamic gradient ($\Delta\mu$). $\Delta\mu$ consists of two components, the electrical gradient or membrane potential and the chemical gradient (Nobel, 1991). Transport of ions down the $\Delta\mu$ is passive, whereas transport against the gradient is active. The transport proteins that mediate ion flux can be generally categorized as pumps, carriers, and channels (Sussman and Harper, 1989). Pumps directly utilize metabolic energy for vectorial transport, whereas carriers couple uphill transport of one solute to the downhill movement of another, either in the same (symporter) or opposite (antiporter) direction. Channels mediate passive transport, i.e. movement down a free energy gradient, although movement may be electrophoretic flux resulting from an energy-dependent process. In plants many of these transport proteins have been identified based on physiological evidence and biochemical characterization, and in some instances, the genes encoding these proteins have been cloned (summarized in Table I).

Currently, it is presumed that energy-dependent flux of most ions in plants is mediated by the $\Delta\mu_{\text{H}^+}$ that generates the pH gradient (ΔpH) and is principally responsible for the $\Delta\Psi$ (Sze, 1985). The proton motive force, which results from the conversion of chemical energy ($\Delta\mu_{\text{H}^+}$) to mV, is $\Delta p = \Delta\mu_{\text{H}^+}/F = \Delta\Psi - 59\Delta\text{pH}$ (at 25°C), where F is the Faraday constant (Sze, 1985). Thus, carrier-mediated ion transport in plants is coupled to the downhill flux of H⁺. Ca²⁺-ATPases in the plasma membrane and endomembranes are notable exceptions to this generalization about $\Delta\mu_{\text{H}^+}$ -mediated ion transport, as we discuss below. H⁺-ATPases in the plasma membrane and tonoplast and the tonoplast H⁺-pyrophosphatase are primary pumps that couple the free energy of hydrolysis of ATP or PPI, respectively, to vectorial H⁺ transport and generation of $\Delta\mu_{\text{H}^+}$.

Abbreviations: $\Delta\mu$, electrochemical potential; $\Delta\mu_{\text{H}^+}$, H⁺ electrochemical gradient; $\Delta\Psi$, membrane potential.

¹ This research was supported in part by U.S. Department of Agriculture/National Research Initiative Competitive Grants Program grant No. 92-37100-7738, and by Comision Interministerial de Ciencia y Tecnología grant No. BIO94-0622. This is journal article No. 14,705 from the Purdue Agricultural Experiment Station.

* Corresponding author; e-mail hasegawa@vm.cc.purdue.edu; fax 1-317-494-0391.

Table 1. Plant genes encoding transport proteins that are presumed to function in establishment and maintenance of ion homeostasis for salt adaptation

In instances where yeast genes are listed, the transport process mediated is assumed to be so fundamental that a functional higher plant homolog must exist. Only one isogene is listed per species.

Category	Name	Location	Gene	Characteristics	Species	Reference	
Pumps	H ⁺ -ATPase	Plasma membrane	<i>AHA3</i>	12 isogenes (?)	<i>Arabidopsis thaliana</i>	Pardo and Serrano (1989)	
			<i>PMA2</i>	4 isogenes	<i>Nicotiana glauca</i>	Boutry et al. (1989)	
			<i>LHA1</i>	2 isogenes	<i>Lycopersicon esculentum</i>	Ewing et al. (1990)	
	Tonoplast			<i>OSA1</i>		<i>Oryza sativa</i>	Wada et al. (1992)
				<i>Unnamed</i>	Subunit A (69 kD)	<i>Daucus carota</i>	Zimniak et al. (1988)
				<i>CVA69.24</i>	Subunit A (69 kD)	<i>Gossypium hirsutum</i>	Wilkins (1993)
				<i>At57</i>	Subunit B (57 kD)	<i>A. thaliana</i>	Manolson et al. (1988)
				<i>HTB1</i>	Subunit B, 2 isogenes	<i>Hordeum vulgare</i>	Berkelman et al. (1993)
				<i>Unnamed</i>	Subunit B (60 kD)	<i>G. hirsutum</i>	Wan and Wilkins (1994)
				<i>VATP-P1</i>	Subunit c (16 kD)	<i>Avena sativa</i>	Lai et al. (1991)
H ⁺ -PPase	Tonoplast	<i>AVP</i>	3 isogenes	<i>A. thaliana</i>	Sarafian et al. (1992)		
Ca ²⁺ -ATPase	ER	<i>LCA1</i>		<i>L. esculentum</i>	Wimmers et al. (1992)		
Na ⁺ -ATPase	Plasma membrane	<i>ENA1</i>	2 isogenes	<i>Saccharomyces cerevisiae</i>	Haro et al. (1991)		
Carriers	Na ⁺ /H ⁺ antiporter	Plasma membrane	<i>SOD2</i>		<i>Schizosaccharomyces pombe</i>	Jia et al. (1992)	
			<i>Unnamed</i>		<i>Zygosaccharomyces rouxii</i>	Watanabe et al. (1991) ^a	
	K ⁺ -H ⁺ symporter	Plasma membrane	<i>HKT1</i>		<i>Triticum aestivum</i>	Schachtman and Schroeder (1994)	
Channels	K ⁺ inward	Plasma membrane	<i>KAT1</i>		<i>A. thaliana</i>	Anderson et al. (1992)	
			<i>AKT1</i>		<i>A. thaliana</i>	Sentenac et al. (1992)	
			<i>KST1</i>		<i>Solanum tuberosum</i>	Müller-Röber et al. (1995)	

^a GenBank accession No. D43629.

(Maathuis and Sanders, 1992). The H⁺ pump in the plasma membrane is a P-type ATPase (Michelet and Boutry, 1995) that establishes a ΔpH of about 1.5 to 2 units (pH 5.5 to 5.0 in the apoplast) and is principally responsible for the inside negative membrane potential of -120 to -200 mV across this membrane under physiologically steady-state conditions (Sze, 1985). From the Nernst equation, $\Delta\psi$ values in this range would establish free-energy gradients that would result in the concentration of monovalent cations (K⁺ or Na⁺) 100- to 1000-fold or greater in the cytosol (Nobel, 1991). Conversely, these free-energy gradients would make intracellular uptake of anions thermodynamically untenable. The two H⁺ pumps present in the tonoplast, vacuolar-type H⁺-ATPase (Sze, 1985) and H⁺-pyrophosphatase (Rea and Sanders, 1987), establish a $\Delta\mu_{\text{H}^+}$ across this membrane. This $\Delta\mu_{\text{H}^+}$ is predominantly the result of ΔpH (about 2 units between the cytosol and inside the vacuole), since a $\Delta\psi$ of 0 to +20 mV (inside positive) normally occurs across the tonoplast.

Under typical physiological conditions, homeostatic concentrations of ions in the cytosol are 100 to 200 mM K⁺, 1 to 10 mM Na⁺ and Cl⁻, and 100 to 200 nM Ca²⁺ (Binzel et al., 1988; Bush, 1995). The $\Delta\mu_{\text{K}^+}$ across the plasma membrane is near equilibrium, and K⁺ homeostasis is achieved by the gating of inward- and outward-rectifying K⁺ channels and by the influx activity of a high-affinity K⁺-H⁺ symporter (Schachtman and Schroeder, 1994; Schroeder et al., 1994). Uptake of Na⁺ and Ca²⁺ occurs passively across the plasma membrane, and efflux is presumably due to the

activities of a Na⁺/H⁺ antiporter and a Ca²⁺-ATPase, respectively. Cl⁻ uptake is assumed to be coupled to a H⁺ symporter because of the large inside negative $\Delta\psi$ (Poole, 1988). Extrusion of Cl⁻ is by electrophoretic flux.

IMPACT OF NaCl STRESS ON ION HOMEOSTASIS AND ADAPTIVE RESPONSES

When plants are exposed to NaCl, ions reduce the apoplastic water potential and accumulate excessively in the cytosol (Binzel et al., 1988). Plant cells adjust to the water relations imbalance through osmotic adjustment by synthesizing compatible organic solutes and accumulating ions from the external environment. Osmotic adjustment must be achieved without undue concentration, in the cytosol, of ions from the external environment, and the ion activities of those that do accumulate must be attenuated by osmoprotectants. Presumably, genotypes that are most adapted to salt tightly regulate ion uptake across the plasma membrane at a rate that is compatible with the capacity for vacuolar compartmentation (Binzel et al., 1988). Thus, transport processes at the plasma membrane and tonoplast that regulate ion influx and efflux, particularly those involved in the control of Na⁺ uptake and vacuolar compartmentation, are of crucial importance to salinity adaptation.

Sodium Influx, Association with Potassium Uptake

The Na⁺ electrochemical gradient dictates that Na⁺ influx across the plasma membrane is passive and efflux is

active at physiological $\Delta\Psi$ values. When the NaCl concentration increases in the surrounding environment, the high extracellular level of Na^+ (relative to the cytosol) and inside-negative $\Delta\Psi$ establish a steep thermodynamic gradient for Na^+ influx. Vacuolar compartmentation of Na^+ , which reduces cytosolic levels of this ion, further facilitates the energetically downhill influx across the plasma membrane.

The mechanism of Na^+ influx across the plasma membrane is unknown. Na^+ acts as a competitor of K^+ uptake (Watad et al., 1991; Schroeder et al., 1994), suggesting that the uptake mechanisms for both cations are similar. Plant roots utilize two systems for K^+ acquisition. System 1 has a high affinity for K^+ (K_m of 10–30 μM) to allow uptake at low K^+ concentrations; it is not inhibited by Na^+ (Rains and Epstein, 1967). System 2 mediates uptake at higher external K^+ concentrations (mM) and has a less pronounced K^+/Na^+ selectivity. Na^+ influx into plants likely occurs via the low-affinity rather than the high-affinity K^+ uptake system (Rains and Epstein, 1967). Presumably, system 1 is an active transporter (perhaps a K^+-H^+ symporter; Schachtman and Schroeder, 1994) because of thermodynamic constraints to passive influx at very low external K^+ concentrations, whereas inward-rectifying K^+ channels may mediate K^+ uptake by system 2 at external concentrations above 0.3 mM (Maathuis and Sanders, 1993; Schroeder et al., 1994). However, available data indicate that plasma membrane inward-rectifying K^+ channels have extremely high K^+/Na^+ selectivity, whereas the K^+-H^+ symporter is less selective (Anderson et al., 1992; Schachtman and Schroeder, 1994).

Schachtman et al. (1991) have suggested that Na^+ uptake may occur through outward-rectifying cation channels. Plasma membrane depolarization and exposure to high external NaCl increases the open probability of outward-rectifying cation channels in wheat root and tobacco cells, thereby allowing Na^+ influx to occur down its steep electrochemical gradient. Any regulatory process that decreases the open probability of these outward-rectifying cation channels would reduce both the entry of Na^+ into the cell and the leakage of K^+ out of the cell, thus hypothetically representing a mechanism of adaptation to NaCl stress.

The high-affinity K^+ uptake system in barley roots can be induced by low (μM) and inhibited by elevated (mM) extracellular K^+ concentrations (Fernando et al., 1992, and refs. therein). Whether the functions of root K^+ uptake systems 1 and 2 are also regulated by Na^+ stress, thereby altering Na^+ influx into the root, remains to be determined. Suggested by analogy with yeast (discussed below), the prediction is that high external Na^+ would trigger, either by excessive Na^+ influx or reduced K^+ uptake, the activation of system 1 and the inhibition of system 2. Consistent with this hypothesis, tobacco cells adapted to NaCl exhibited enhanced capacity for K^+ uptake relative to wild-type cells, presumably due to a higher degree of coupling between H^+ efflux and K^+ influx across the plasma membrane (Watad et al., 1991). Further, no inhibition of K^+ uptake by NaCl was observed in salt-adapted cells, indi-

cating that these cells had a higher K^+/Na^+ selectivity at the plasma membrane (Watad et al., 1991). Therefore, increased K^+/Na^+ selectivity of the K^+ uptake system might represent a significant adaptation to high concentrations of NaCl. At the whole plant level, it is generally accepted that increased K^+/Na^+ selectivity during uptake and reduced Na^+ translocation from the root to the shoot contribute to the overall salt tolerance of glycophytes.

Sodium Efflux and Vacuolar Compartmentation

Active efflux of Na^+ from the cytosol across the plasma membrane and the tonoplast is essential for the regulation of net intracellular uptake and vacuolar compartmentation. Physiological and biochemical data support the likelihood that Na^+/H^+ antiporters in the plasma membrane and tonoplast mediate these fluxes (DuPont, 1992, and refs. therein). Mechanistically, the Na^+/H^+ antiporters in the plasma membrane and tonoplast are coupled to the ΔpH s generated by the H^+ pumps located in these membranes.

Induction of plasma membrane Na^+/H^+ antiport activities by NaCl occurred in the halotolerant alga *Dunaliella salina* (Katz et al., 1992) and the halophyte *Atriplex nummularia* (Hassidim et al., 1990). The H^+ -translocating activities of the plasma membrane ATPase increased in response to NaCl treatment in tobacco cells and roots of the halophyte *A. nummularia* (Braun et al., 1986; Watad et al., 1991). This up-regulation may be mediated, at least in part, by increased gene expression (Niu et al., 1993a, 1993b; Perez-Prat et al., 1994). The levels of the plasma membrane pump mRNA were substantially higher in the elongation and differentiation zones after NaCl treatment (Niu et al., 1993a). The gene expression and physiological responsiveness of the plasma membrane H^+ -ATPase to NaCl is positively correlated with salt tolerance, since halotolerant cells and plants exhibited higher transcript levels and/or pump activities than intolerant counterparts (Braun et al., 1986; Niu et al., 1993a, 1993b; Perez-Prat et al., 1994).

Plasma membrane H^+ -ATPase mRNA accumulation is induced only during NaCl adaptation, since lower and similar mRNA levels were detected in samples obtained from cells that were either growing in the absence of salt or had been adapted to salt for several hundred generations (Niu et al., 1993b; Perez-Prat et al., 1994). These results indicate that salt-induced plasma membrane H^+ -ATPase gene expression, which is assumed to be a basis for at least part of higher H^+ transport activity, occurs during stress adaptation but not after the new adaptive state has been achieved. Presumably, after adaptation (when cells are growing actively in salt), the pump re-establishes a H^+ transport steady state that is the same as for cells growing in a nonsaline environment. Thus, during the period of stress adaptation, active plasma membrane transport mechanisms (e.g. Na^+/H^+ antiporter) are induced or activated, requiring higher pump activity, whereas after adaptation and when ion gradients have been established across the membrane, altered membrane permeability (e.g. reduced passive transport restricting Na^+ influx) may be the primary basis for ion homeostasis (Binzel et al., 1988; Watad et al., 1991).

Vacuolar compartmentation of Na^+ and Cl^- is an essential mechanism for salt tolerance, since it results in lower cytosolic ion levels and facilitates osmotic adjustment required for cell expansion and maintenance of turgor. Tobacco cells adapted to and growing in 428 mM NaCl accumulated 780 mM Na^+ and 624 mM Cl^- in the vacuole, whereas cytoplasmic levels of both ions were below 100 mM (Binzel et al., 1988). Na^+/H^+ antiport activities of tonoplast vesicles from beet cells and barley roots (DuPont, 1992) were induced by NaCl. Antibodies that reacted to a 170-kD polypeptide and that accumulated in tonoplast vesicles after salt treatment inhibited Na^+/H^+ activity of these membrane vesicles. Similarly, H^+ transport by the vacuolar H^+ -ATPase increased during adaptation to NaCl in several plants (DuPont, 1992, and refs. therein). In barley roots, increased H^+ transport activity did not peak until 3 d after the onset of NaCl treatment, and this induction was inhibited by protein synthesis inhibitors, indicating that de novo synthesis was required. Accordingly, 70-kD subunit (subunit B) mRNA of the vacuolar H^+ -ATPase accumulated in response to NaCl treatment (Narasimhan et al., 1991). These data imply that salt-induced H^+ transport activity of the tonoplast ATPase is mediated through increased gene expression. The induction of tonoplast H^+ -ATPase activity by NaCl apparently occurs only during stress adaptation but not after adaptation, since both vesicle H^+ pumping activities and 70-kD subunit mRNA levels were similar in wild-type and salt-adapted cells (Narasimhan et al., 1991). The tonoplast H^+ -ATPase of salt-adapted cells exhibited altered biochemical characteristics consistent with a higher specific H^+ pumping activity. Enhanced activity of the tonoplast pyrophosphatase in NaCl-stressed carrot cells has also been reported (Colombo and Cerana, 1993).

Cl^-

Little is known about intracellular uptake and vacuolar compartmentation of Cl^- (Binzel et al., 1988). The extreme inside negative membrane potential across the plasma membrane that occurs when ion homeostasis is established and maintained is a substantial thermodynamic barrier to Cl^- influx, even at relatively high external concentrations. However, active uptake could be mediated by a Cl^-/H^+ symporter (Poole, 1988). If Na^+ influx depolarizes the $\Delta\Psi$ across the plasma membrane, the Cl^- can be taken up passively through an anion channel (Skerrett and Tyerman, 1994). Vacuolar compartmentation of Cl^- is an essential adaptation for NaCl tolerance. Cl^- movement from the cytosol to the vacuole may be achieved through channels with transport driven by the electrophoretic flux generated by H^+ pumps across the tonoplast, i.e. under saline conditions, Cl^- may be the counterion to H^+ , since the $\Delta\mu_{\text{H}^+}$ across this membrane is primarily due to ΔpH . Slow-vacuolar-type channels can account for all ion conductances at elevated cytosolic ($>1 \mu\text{M}$) Ca^{2+} levels (Plant et al., 1994). An alternative possibility is a H^+/anion antiporter that would actively transport Cl^- across the tonoplast (Rea and Sanders, 1987).

Ca^{2+}

The role of Ca^{2+} in NaCl stress adaptation is complex and not well defined, although this cation and its homeostasis are thought to be essential. Externally supplied Ca^{2+} ameliorates NaCl stress through an unknown function that preserves K^+/Na^+ selectivity (Zhong and Läuchli, 1994). Moreover, Ca^{2+} inhibits inward-rectifying K^+ channels, that may reduce the Na^+ influx mediated by the low-affinity component of K^+ uptake (Schroeder et al., 1994). Cytosolic Ca^{2+} levels are usually maintained at 100 to 200 nM by active transport, and small increases in concentration typically initiate very specific signal transduction cascades (Bush, 1995). NaCl causes a rapid increase in cytosolic Ca^{2+} that probably acts as a general stress signal (Lynch et al., 1989). Although it is not clear if this increase is an effector of salt tolerance, increases in cytosolic Ca^{2+} that mediate salt adaptation must be transitory, as is the case for all signals. Current information indicates that re-establishment of cytosolic Ca^{2+} homeostasis is a requisite for adaptation.

The rise in cytosolic Ca^{2+} may occur via influx from the apoplast through stretch-activated Ca^{2+} channels similar to those in guard cells that respond to tension produced by osmotic stress (Cosgrove and Hedrich, 1991). Additionally, inositol (1,4,5)-triphosphate-regulated channels in the tonoplast and other endomembranes may release Ca^{2+} from internal pools (Alexandre et al., 1990), perhaps in response to G-protein activation and inositol (1,4,5)-triphosphate formation. Several transport mechanisms facilitate Ca^{2+} efflux from the cytosol, including Ca^{2+} -ATPases in the plasma membrane and endomembranes (Bush, 1995) and a tonoplast $\text{Ca}^{2+}/\text{H}^+$ antiporter. Plant Ca^{2+} -ATPase cDNAs have been isolated that encode putative ER pumps. Transcripts detected by these cDNAs accumulated in response to NaCl treatment, and Ca^{2+} -ATPase mRNA levels remained elevated after adaptation (Perez-Prat et al., 1992). These results suggest that the increase in cytosolic Ca^{2+} that follows exposure to NaCl may be lowered by increased activity of the Ca^{2+} -ATPase.

THE FUNGAL MODEL

Ion transport in plants and fungi shares many common features, and it has been suggested by Haro et al. (1993) that this organism can be a model for the characterization of salt-tolerant mechanisms in plant cells because of similarities with higher plants and the available powerful molecular genetic tools. Genetic and molecular studies of yeast have shown that restricted Na^+ uptake, rapid Na^+ efflux, and efficient Na^+ compartmentation into the vacuole are important salt-tolerant determinants similar to those in plant cells. As has been established for plant roots, fungi exhibit a dual K^+ uptake system, with high or low affinity of K^+ , that adjusts in response to environmental stimuli. In NaCl-stressed *Saccharomyces cerevisiae* cells, the K^+ uptake system changes to a state with increased affinity for K^+ , whereas the affinity for Na^+ is relatively unaltered, thereby effectively reducing Na^+ influx (Ramos et al., 1985). The high-affinity K^+ uptake system depends on

TRK1, a gene encoding a membrane protein presumed to be the K^+ transporter, although direct biochemical evidence for such a function is not yet available (Gaber et al., 1988). A *trk1* mutant, deficient in the high-affinity K^+ transport mode, is sensitive to Na^+ and Li^+ (Haro et al., 1993), consistent with the presumption that enhanced K^+/Na^+ selectivity is required for NaCl tolerance.

Two different Na^+ efflux systems have been characterized in yeast. The *SOD2* gene of *Schizosaccharomyces pombe* encodes a plasma membrane Na^+/H^+ antiporter (Jia et al., 1992). In *S. cerevisiae*, Na^+ efflux is mediated by a novel plasma membrane P-type Na^+ -ATPase, the ENA protein (Haro et al., 1991). Apparently, *SOD2* and ENA proteins are the only Na^+ efflux mechanisms present in *S. pombe* and *S. cerevisiae*, respectively, because null mutations in the corresponding genes completely abolished Na^+ efflux and rendered cells highly sensitive to NaCl. Gene amplification of *SOD2* enhanced Na^+ efflux and increased tolerance to NaCl, indicating that Na^+ efflux capacity delineates the upper limit for Na^+ tolerance in yeast (Jia et al., 1992). Theoretically, overexpression of the ENA ATPase might increase Na^+ tolerance even further because the efflux of Na^+ would be less dependent on the transmembrane Na^+ and H^+ gradients.

In *S. cerevisiae*, the tonoplast has a functional role in salt and osmotic stress tolerances. Some *vpt* and all *ssv* mutants, with altered tonoplast function or morphology, were sensitive to high concentrations of salts and polyols, indicating that correct vacuolar function is required for osmoregulation (Banta et al., 1988; Latterich and Watson, 1991). Besides this osmoregulatory role, ion compartmentation in the large vacuolar space may also prevent the buildup of toxic levels of ions in the cytoplasm. A *vatC/vma3* mutant defective in vacuolar H^+ -ATPase function, and therefore having disabled energization of the tonoplast, shows a reduced capacity for Na^+ accumulation. Furthermore, it was sensitive to low levels of Na^+ and Li^+ that do not represent a significant osmotic stress (Haro et al., 1993).

The genetic analysis of NaCl tolerance in the yeast *S. cerevisiae* has produced important insights into the regulatory mechanisms controlling adaptation to a saline environment. A screening for genes that enhance tolerance to NaCl led to the isolation of *HAL1* (Gaxiola et al., 1992). The overaccumulation of the HAL1 protein resulted in increased intracellular K^+ levels in salinized media. Preliminary data suggested that plants might also contain a *HAL1* homolog whose expression is induced by NaCl and ABA (Gaxiola et al., 1992). Additionally, the isolation of yeast mutants unable to adapt to mild NaCl stress led to the identification of the Ca^{2+} /calmodulin-dependent protein phosphatase, calcineurin, as a key component of the signaling pathway controlling adaptation to NaCl stress (Mendoza et al., 1994). A calcineurin-deficient cell was unable to fully induce *ENA1* gene expression and convert the K^+ uptake system to the high-affinity state. Hence, calcineurin seems to be a common intermediate in signaling pathways leading to the control of both Na^+ efflux and influx that regulate net uptake of this ion across the plasma membrane.

CONCLUSIONS AND PERSPECTIVES

Figure 1 illustrates a typical scenario of the consequences of NaCl stress to plant cells and the transport processes that are involved in the establishment of ion homeostasis. NaCl stress environments impose water deficit and ion imbalance on plants, and both must be alleviated for survival and growth. Maximizing utilization of the predominant ions in the stress environment for osmotic adjustment and controlling cytosolic levels of these ions (i.e. ion activities) to minimize metabolic toxicity are principal salt-adaptation mechanisms. During the initial period of NaCl stress adaptation, a primary necessity is to evacuate Na^+ and Cl^- from the cytosol by efflux to the apoplast and the vacuole. Activation of the H^+ pumps in the plasma membrane and tonoplast generates $\Delta\mu_{H^+}$ required for active transport of Na^+ across these membranes and compartmentation of Cl^- in the vacuole. Concurrent with H^+ pump activation must be adaptations that more effectively regulate passive transport of Na^+ across the plasma membrane and, perhaps, passive efflux of Na^+ and Cl^- from the vacuole. The control of passive transport processes is required to maintain Na^+ and Cl^- gradients that result from active transport and electrophoretic flux mediated by the re-establishment of the plasma membrane $\Delta\Psi$. These new gradients can be maintained by an ion flux steady state that does not require the high level of active transport necessary for their establishment. Consequently, the energy costs are not substantially greater than before adaptation (Schnapp et al., 1991), presumably due in large part to the down-regulation of the H^+ pumps from the levels induced by salt exposure (Niu et al., 1993a). Therefore, membrane transport proteins and their regulators are critical salt-tolerant determinants.

The immediate research challenge is to identify and characterize the genes controlling ion homeostasis in saline environments. Plant genes encoding putative K^+ transport systems in the plasma membrane (Table I) (Schroeder et al., 1994) can now be used for molecular genetic and biochemical experimentation to determine if these mediate Na^+ uptake. Perhaps molecular modifications can be made to these proteins that would increase K^+/Na^+ selectivity. Current evidence indicates that Na^+ efflux across the plasma membrane in plants occurs via a Na^+/H^+ antiporter. It is not known whether a Na^+ pump analogous to yeast ENA exists in higher plants, but a Na^+ -activated plasma membrane P-type ATPase has been described in the marine alga *Heterosigma akashiwo* (Wada et al., 1989), and a primary Na^+ pump has been implicated in the halophilic alga *D. salina* (Katz et al., 1991). Theoretically, overexpression of a plasma membrane Na^+ efflux transport protein from a heterologous system could establish the significance of active Na^+ efflux across the plasma membrane in salt adaptation. The identification of a 170-kD polypeptide as a probable constituent of the tonoplast Na^+/H^+ antiporter (Barkla and Blumwald, 1991) should lead eventually to the isolation of a gene critical to vacuolar compartmentation of Na^+ .

A key factor in the genetic capacity for salt adaptation must involve an appropriately responsive and coordinated





Figure 1. Ionic status of plant cells and proposed transport processes that control cytosolic Na^+ levels in low NaCl (A) and high NaCl during (B) and after (C) adaptation. Simplified diagrams illustrated plasma membrane and tonoplast pumps, carriers, and channels presumed to be involved in the regulation of Na^+ , Cl^- , K^+ , Ca^{2+} , and H^+ transport required to establish and maintain ion homeostasis in a saline environment. Tonoplast $\Delta\Psi$ is relative to the cytosol. Illustrated are comparisons of relative Na^+ and K^+ pools before NaCl stress, immediately after stress imposition, and after stress adaptation.

signal transduction system. As we have indicated, signaling molecules in cascades that directly regulate proteins involved in ion flux or regulate the expression of the genes that encode transport proteins are principal components of salt adaptation and probably limit the efficiency of salt adaptation in many important crop plants. Physiological evidence implicates Ca^{2+} as a secondary messenger in the signal transduction of salt-stress perception to the mechanisms that control ion homeostasis in plants. In yeast, the

Ca^{2+} /calmodulin-dependent phosphatase calcineurin is a key intermediate in the regulation of Na^+ influx and efflux systems (Mendoza et al., 1994) that are responsible for salt adaptation. Perhaps a functional calcineurin homolog facilitates the effect of Ca^{2+} on transport processes that regulate K^+/Na^+ selectivity in plants. There is still little known about the genes involved in such signal transduction pathways. However, overcoming the limitations placed on ion homeostasis adaptability by a poorly respon-

sive or inadequately coordinated signal transduction system remains a major objective in understanding and achieving salt-stress tolerance in plants.

An important consideration for future research will be the possible impact of manipulating the K^+/Na^+ selectivity of transporters on K^+ nutrition. Higher K^+/Na^+ selectivity may reduce the ability to transport K^+ rapidly and thereby affect K^+ uptake to the extent that growth rates are compromised. An interesting speculation is that there is a relationship between K^+/Na^+ selectivity, K^+ uptake capacity, and the well-noted slow growth rates of highly Na^+ -tolerant halophytic species. In other words, have highly Na^+ -tolerant species sacrificed K^+ uptake capacity for reduced Na^+ uptake, thereby obtaining Na^+ tolerance but losing rapid growth capability? Only further research will answer these important questions.

Received May 19, 1995; accepted August 11, 1995.

Copyright Clearance Center: 0032-0889/95/109/0735/08.

LITERATURE CITED

- Alexandre L, Lassalles JP, Kado RT (1990) Opening of Ca^{2+} channels in isolated red beet root vacuole membrane by inositol-1,4,5-triphosphate. *Nature* **343**: 567–570
- Anderson JA, Huprikar SS, Kochian LV, Lucas WJ, Gaber RF (1992) Functional expression of a probable *Arabidopsis thaliana* potassium channel in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci USA* **89**: 3736–3740
- Banta LM, Robinson JS, Klionsky DJ, Emr SD (1988) Organelle assembly in yeast: characterization of yeast mutants defective in vacuolar biogenesis and protein sorting. *J Cell Biol* **107**: 1369–1383
- Barkla BJ, Blumwald E (1991) Identification of a 170-kDa protein associated with the vacuolar Na^+/H^+ antiport of *Beta vulgaris*. *Proc Natl Acad Sci USA* **88**: 11177–11181
- Berkelman T, Houtchens KA, DuPont FM (1994) Two cDNA clones encoding isoforms of the B subunit of the vacuole ATPase from barley roots. *Plant Physiol* **104**: 287–288
- Binzel ML, Hess FD, Bressan RA, Hasegawa PM (1988) Intracellular compartmentation of ions in salt adapted tobacco cells. *Plant Physiol* **86**: 607–614
- Boutry M, Baudin M, Goffeau A (1989) Molecular cloning of a family of plant genes encoding a protein homologous to plasma membrane H^+ -translocating ATPases. *Biochem Biophys Res Commun* **162**: 567–574
- Braun Y, Hassidim M, Lerner HR, Reinhold L (1986) Studies on H^+ -translocating ATPase in plants of varying resistance to salinity. I. Salinity during growth modulates the proton pump in the halophyte *Atriplex nummularia*. *Plant Physiol* **81**: 1050–1056
- Bush DS (1995) Calcium regulation in plant cells and its role in signaling. *Annu Rev Plant Physiol Plant Mol Biol* **46**: 95–122
- Colombo R, Cerana R (1993) Enhanced activity of tonoplast pyrophosphatase in NaCl-grown cells of *Daucus carota*. *J Plant Physiol* **142**: 226–229
- Cosgrove DJ, Hedrich R (1991) Stretch-activated chloride, potassium, and calcium channels coexisting in plasma membranes of guard cells of *Vicia faba* L. *Planta* **186**: 143–153
- DuPont FM (1992) Salt-induced changes in ion transport: Regulation of primary pumps and secondary transporters. In DT Cooke, DT Clarkson eds, *Transport and Receptor Proteins of Plant Membranes*. Plenum Press, New York, pp 91–100
- Ewing NN, Wimmers LE, Meyer DJ, Chetelat RT, Bennett AB (1990) Molecular cloning of tomato plasma membrane H^+ -ATPase. *Plant Physiol* **94**: 1874–1881
- Fernando M, Mehroke J, Glass ADM (1992) *De novo* synthesis of plasma membrane and tonoplast polypeptides of barley roots during short-term K^+ deprivation. In search of the high-affinity K^+ transport system. *Plant Physiol* **100**: 1269–1276
- Gaber RF, Styles CA, Fink GR (1988) *TRK1* encodes a plasma membrane protein required for high-affinity potassium transport in *Saccharomyces cerevisiae*. *Mol Cell Biol* **8**: 2848–2859
- Gaxiola R, de Larrinoa IF, Villalba JM, Serrano R (1992) A novel and conserved salt-induced protein is an important determinant of salt tolerance in yeast. *EMBO J* **11**: 3157–3164
- Haro R, Garcíadeblas B, Rodríguez-Navarro A (1991) A novel P-type ATPase from yeast involved in sodium transport. *FEBS Lett* **291**: 189–191
- Haro R, Bañuelos MA, Quintero FJ, Rubio F, Rodríguez-Navarro A (1993) Genetic basis of sodium exclusion and sodium tolerance in yeast. A model for plants. *Physiol Plant* **89**: 868–874
- Hassidim M, Braun Y, Lerner HR, Reinhold L (1990) Na^+/H^+ and K^+/H^+ antiport in root membrane vesicles isolated from the halophyte *Atriplex* and the glycophyte cotton. *Plant Physiol* **94**: 1795–1801
- Jia ZP, McCullough N, Martel R, Hemmingsen S, Young PG (1992) Gene amplification at a locus encoding a putative Na^+/H^+ antiporter confers sodium and lithium tolerance in fission yeast. *EMBO J* **11**: 1631–1640
- Katz A, Bentall M, Dengani H, Avron M (1991) *In vivo* pH regulation by a Na^+/H^+ antiporter in the halotolerant alga *Dunaliella salina*. *Plant Physiol* **96**: 110–115
- Katz A, Pick U, Avron M (1992) Modulation of Na^+/H^+ antiporter activity by extreme pH and salt in the halotolerant alga *Dunaliella salina*. *Plant Physiol* **100**: 1224–1229
- Lai S, Watson JC, Hansen JN, Sze H (1992) Molecular cloning and sequencing of cDNAs encoding the proteolipid subunit of the vacuolar H^+ -ATPase from a higher plant. *J Biol Chem* **266**: 16078–16084
- Latterich M, Watson MD (1991) Isolation and characterization of osmosensitive vacuolar mutants of *Saccharomyces cerevisiae*. *Mol Microbiol* **5**: 2417–2426
- Lynch J, Polito VS, Läuchli A (1989) Salinity stress increases cytoplasmic Ca activity in maize root protoplasts. *Plant Physiol* **90**: 1271–1274
- Maathuis FJM, Sanders D (1992) Plant membrane transport. *Curr Opin Cell Biol* **4**: 661–669
- Maathuis FJM, Sanders D (1993) Energization of potassium uptake in *Arabidopsis thaliana*. *Planta* **191**: 302–307
- Manolson MF, Ouellette BF, Filion M, Poole RJ (1988) cDNA sequence and homologies of the “57-kDa” nucleotide-binding subunit of the vacuolar ATPase from *Arabidopsis*. *J Biol Chem* **263**: 17987–17994
- Mendoza I, Rubio F, Rodríguez-Navarro A, Pardo JM (1994) The protein phosphatase calcineurin is essential for NaCl tolerance of *Saccharomyces cerevisiae*. *J Biol Chem* **269**: 8792–8796
- Michelet B, Boutry M (1995) The plasma membrane H^+ -ATPase. A highly regulated enzyme with multiple physiological functions. *Plant Physiol* **108**: 1–6
- Müller-Röber B, Ellenberg J, Provart N, Willmitzer L, Busch H, Becker D, Dietrich P, Hoth S, Hedrich R (1995) Cloning and electrophysiological analysis of KST1, an inward rectifying K^+ channel expressed in potato guard cells. *EMBO J* **14**: 2409–2416
- Narasimhan M, Binzel ML, Perez-Prat E, Chen Z, Nelson DE, Singh NK, Bressan RA, Hasegawa PM (1991) NaCl regulation of tonoplast ATPase 70-kilodalton subunit mRNA in tobacco cells. *Plant Physiol* **97**: 562–568
- Niu X, Narasimhan ML, Salzman RA, Bressan RA, Hasegawa PM (1993a) NaCl regulation of plasma membrane H^+ -ATPase gene expression in a glycophyte and a halophyte. *Plant Physiol* **103**: 713–718
- Niu X, Zhu J-K, Narasimhan ML, Bressan RA, Hasegawa PM (1993b) Plasma membrane H^+ -ATPase gene expression is regulated by NaCl in cells of the halophyte *Atriplex nummularia* L. *Planta* **190**: 433–438
- Nobel PS (1991) *Physiochemical and Environmental Plant Physiology*. Academic Press, New York
- Pardo JM, Serrano R (1989) Structure of a plasma membrane H^+ -ATPase gene from the plant *Arabidopsis thaliana*. *J Biol Chem* **264**: 8557–8562

- Perez-Prat E, Narasimhan ML, Binzel ML, Botella MA, Chen Z, Valpuesta V, Bressan RA, Hasegawa PM (1992) Induction of a putative Ca^{2+} -ATPase mRNA in NaCl-adapted cells. *Plant Physiol* **100**: 1471-1478
- Perez-Prat E, Narasimhan ML, Niu X, Botella MA, Bressan RA, Valpuesta V, Hasegawa PM, Binzel ML (1994) Growth cycle stage-dependent NaCl induction of plasma membrane H^{+} -ATPase mRNA accumulation in de-adapted tobacco cells. *Plant Cell Environ* **17**: 327-333
- Plant PJ, Gelli A, Blumwald E (1994) Vacuolar chloride regulation of an anion-selective tonoplast channel. *J Membr Biol* **140**: 1-12
- Poole RJ (1988) Plasma membrane and tonoplast. In DA Baker, JL Hall, eds, *Solute Transport in Plant Cells and Tissues*. John Wiley & Sons, New York, pp 83-105
- Rains DW, Epstein E (1967) Sodium absorption by barley roots. Its mediation by mechanism 2 of alkali cation transport. *Plant Physiol* **42**: 319-323
- Ramos J, Contreras P, Rodriguez-Navarro A (1985) A potassium transport mutant of *Saccharomyces cerevisiae*. *Arch Microbiol* **143**: 88-93
- Rea PA, Sanders D (1987) Tonoplast energization: two pumps, one membrane. *Physiol Plant* **81**: 131-141
- Sarafian V, Kim Y, Poole RJ, Rea PA (1992) Molecular cloning and sequence of cDNA encoding the pyrophosphate-energized vacuolar membrane proton pump of *Arabidopsis thaliana*. *Proc Natl Acad Sci USA* **89**: 1775-1779
- Schachtman DP, Schroeder JI (1994) Structure and transport mechanism of a high-affinity potassium uptake transporter from higher plants. *Nature* **370**: 655-658
- Schachtman DP, Tyerman SD, Terry BR (1991) The $\text{K}^{+}/\text{Na}^{+}$ selectivity of a cation channel in the plasma membrane of root cells does not differ in salt-tolerant and salt-sensitive wheat species. *Plant Physiol* **97**: 598-605
- Schnapp SR, Curtis WR, Bressan RA, Hasegawa PM (1991) Growth yields and maintenance coefficients of unadapted and NaCl-adapted tobacco cells grown in semicontinuous culture. *Plant Physiol* **96**: 1289-1293
- Schroeder JI, Ward JM, Gassmann W (1994) Perspectives on the physiology and structure of inward-rectifying K^{+} channels in higher plants: biophysical implications for K^{+} uptake. *Annu Rev Biophys Biomol Struct* **23**: 441-471
- Sentenac H, Bonneaud N, Minet M, Lacroute F, Salmon J-M, Gaymard F, Grignon C (1992) Cloning and expression in yeast of a plant potassium ion transport system. *Science* **256**: 663-665
- Skerrett M, Tyerman SD (1994) A channel that allows inwardly directed fluxes of anions in protoplasts derived from wheat roots. *Planta* **192**: 295-305
- Sussman MR, Harper JF (1989) Molecular biology of the plasma membrane of higher plants. *Plant Cell* **1**: 953-960
- Sze H (1985) H^{+} -translocating ATPase. *Advances using membrane vesicles*. *Annu Rev Plant Physiol* **36**: 175-208
- Wada M, Satoh S, Kasamo K, Fujii T (1989) Presence of a Na^{+} -activated ATPase in the plasma membrane of the marine Raphidophycean *Heterosigma akashiwo*. *Plant Cell Physiol* **30**: 923-928
- Wada M, Takano M, Kasamo K (1992) Nucleotide sequence of a complementary DNA encoding plasma membrane H^{+} -ATPase from rice (*Oryza sativa* L.). *Plant Physiol* **99**: 794-795
- Wan CY, Wilkins TA (1994) Isolation of multiple cDNAs encoding the vacuolar H^{+} -ATPase subunit B from developing cotton (*Gossypium hirsutum* L.) ovules. *Plant Physiol* **106**: 393-394
- Watad AA, Reuveni M, Bressan RA, Hasegawa PM (1991) Enhanced net K^{+} uptake capacity of NaCl-adapted cells. *Plant Physiol* **95**: 1265-1269
- Wilkins TA (1993) Vacuolar H^{+} -ATPase 69-kilodalton catalytic subunit cDNA from developing cotton (*Gossypium hirsutum*) ovules. *Plant Physiol* **102**: 679-680
- Wimmers LE, Ewing NN, Bennett AB (1992) Higher plant Ca^{2+} -ATPase: primary structure and regulation of mRNA abundance by salt. *Proc Natl Acad Sci USA* **89**: 9205-9209
- Zhong H, Läuchli A (1994) Spatial distribution of solutes, K, Na, Ca and their deposition rates in the growth zone of primary cotton roots: effects of NaCl and CaCl_2 . *Planta* **194**: 34-41
- Zimniak L, Dittrich P, Gogarten JP, Kibak H, Taiz L (1988) The cDNA sequence of the 69-kDa subunit of the carrot vacuolar H^{+} -ATPase. Homology to the beta-chain of FOF1-ATPases. *J Biol Chem* **263**: 9102-9112