



iPTF Discovery of the Rapid “Turn-on” of a Luminous Quasar

S. Gezari^{1,2}, T. Hung¹, S. B. Cenko^{3,2}, N. Blagorodnova⁴, Lin Yan^{5,6}, S. R. Kulkarni⁴, K. Mooley⁷, A. K. H. Kong⁸, T. M. Cantwell⁹, P. C. Yu¹⁰, Y. Cao¹¹, C. Fremling¹², J. D. Neill⁴, C.-C. Ngeow¹⁰, P. E. Nugent^{13,14}, and P. Wozniak¹⁵

¹Department of Astronomy, University of Maryland, Stadium Drive, College Park, MD 20742-2421, USA; suvi@astro.umd.edu

²Joint Space-Science Institute, University of Maryland, College Park, MD 20742, USA

³NASA Goddard Space Flight Center, Mail Code 661, Greenbelt, MD 20771, USA

⁴Department of Astronomy, California Institute of Technology, Pasadena, CA 91125, USA

⁵Caltech Optical Observatories, Cahill Center for Astronomy and Astrophysics, California Institute of Technology, Pasadena, CA 91125, USA

⁶Infrared Processing and Analysis Center, California Institute of Technology, Pasadena, CA 91125, USA

⁷Astrophysics, Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH, UK

⁸Institute of Astronomy, National Tsing Hua University, Hsinchu 30013, Taiwan

⁹Jodrell Bank Centre for Astrophysics, Alan Turing Building, Oxford Road, Manchester M13 9PL, UK

¹⁰Graduate Institute of Astronomy, National Central University, Taoyuan City 32001, Taiwan

¹¹eScience Institute and Astronomy Department, University of Washington, Seattle, WA 98195, USA

¹²Department of Astronomy, The Oskar Klein Center, Stockholm University, AlbaNova, SE-10691, Stockholm, Sweden

¹³Department of Astronomy, University of California, Berkeley, CA 94720-3411, USA

¹⁴Lawrence Berkeley National Laboratory, 1 Cyclotron Road, MS 50B-4206, Berkeley, CA 94720, USA

¹⁵Los Alamos National Laboratory, MS D436, Los Alamos, NM, 87545, USA

Received 2016 November 4; revised 2016 December 13; accepted 2016 December 13; published 2017 January 24

Abstract

We present a radio-quiet quasar at $z = 0.237$ discovered “turning on” by the intermediate Palomar Transient Factory (iPTF). The transient, iPTF 16bco, was detected by iPTF in the nucleus of a galaxy with an archival Sloan Digital Sky Survey spectrum with weak narrow-line emission characteristic of a low-ionization nuclear emission-line region (LINER). Our follow-up spectra show the dramatic appearance of broad Balmer lines and a power-law continuum characteristic of a luminous ($L_{\text{bol}} \approx 10^{45} \text{ erg s}^{-1}$) type 1 quasar 12 yr later. Our photometric monitoring with PTF from 2009–2012 and serendipitous X-ray observations from the *XMM-Newton* Slew Survey in 2011 and 2015 constrain the change of state to have occurred less than 500 days before the iPTF detection. An enhanced broad $\text{H}\alpha/[\text{O III}] \lambda 5007$ line ratio in the type 1 state relative to other changing-look quasars also is suggestive of the most rapid change of state yet observed in a quasar. We argue that the >10 increase in Eddington ratio inferred from the brightening in UV and X-ray continuum flux is more likely due to an intrinsic change in the accretion rate of a preexisting accretion disk than an external mechanism such as variable obscuration, microlensing, or the tidal disruption of a star. However, further monitoring will be helpful in better constraining the mechanism driving this change of state. The rapid “turn-on” of the quasar is much shorter than the viscous infall timescale of an accretion disk and requires a disk instability that can develop around a $\sim 10^8 M_{\odot}$ black hole on timescales less than 1 yr.

Key words: accretion, accretion disks – black hole physics – galaxies: active – surveys

1. Introduction

Variability is a ubiquitous property of quasars on timescales of hours to years and is likely attributed to processes in the accretion disk fueling the central supermassive black hole (SMBH; Pereyra et al. 2006; Kelly et al. 2009). More dramatic changes in accretion activity are also expected to occur on much longer timescales. Hydrodynamic simulations of quasar fueling and feedback reveal a “duty cycle” of accretion activity on timescales of a megayear (Novak et al. 2011) and may explain the lack of clear observational evidence for causal connections between active galactic nucleus (AGN) activity and star formation in galaxy studies (Hickox et al. 2014). Observations of ionization nebulae on the outskirts of galaxies (or lack thereof) have indicated the shutoff (or turn-on) of quasar engines on timescales of tens of thousands of years (e.g., Schawinski et al. 2010, 2015), and our own Galactic center shows signs of X-ray reflection from enhanced nuclear activity on a timescale of only hundreds of years ago (Ponti et al. 2010). However, as we explore the behavior of AGNs more systematically in the time domain with optical imaging and spectroscopic surveys, we are finding evidence for significant accretion state changes on even shorter timescales.

In the AGN unification model, type 1 (spectra with narrow and broad lines) and type 2 (spectra with narrow lines only) classifications are explained as a viewing angle effect due to nuclear obscuration of the broad-line region and accretion disk continuum (Antonucci 1993). A challenge to this paradigm is the rare class of “changing-look” AGNs, which change their spectral class with the appearance and/or disappearance of broad Balmer lines, accompanied by large-amplitude changes in the AGN continuum. Most of the changing-look AGN cases reported to date have been spectroscopically known Seyferts that have shown a dramatic appearance or disappearance of their broad Balmer lines in follow-up spectra, resulting in a “change of state” between a type 1 and a type 1.8–2 spectrum. A type 1.8 or 1.9 Seyfert classification depends on the presence of weak broad $\text{H}\beta$ or broad $\text{H}\alpha$, respectively (Osterbrock 1981).

Notable examples of changing-look AGNs include the appearance of broad, double-peaked $\text{H}\alpha$ and $\text{H}\beta$ lines in LINER galaxy NGC 1097 (Storchi-Bergmann et al. 1993) and the complete disappearance of the broad $\text{H}\beta$ line in Seyfert 1 galaxy Mrk 590 (Denney et al. 2014). More recently the ASAS-SN optical time domain survey discovered an outburst from NGC 2617 at $z = 0.0142$, which was accompanied by a

transition from a Seyfert 1.8 to a Seyfert 1 optical spectrum on a timescale of ~ 10 yr (Shappee et al. 2014). And recent follow-up observations of the changing-look AGN Mrk 1018, which had changed from a type 1.9 to type 1 Seyfert in 1984, revealed a change back to a type 1.9 Seyfert 30 yr later (McElroy et al. 2016). The fact that the variable UV/optical continuum in Mrk 1018 followed the $L \sim T^4$ relation expected for thermal emission from a disk, and that there was no evidence for neutral hydrogen absorption in its X-ray spectrum, favored intrinsic changes in the accretion flow instead of an obscuration event (Husemann et al. 2016), which has been inferred to be the cause for rapid drops in the UV continuum (Guo et al. 2016) and X-ray flux (Risaliti et al. 2009; Marchese et al. 2012) in some AGNs.

The first case of a changing-look AGN with the luminosity of a quasar (which we define as $L_{\text{bol}} > 10^{44}$ erg s $^{-1}$) was SDSS J0159+0033 at $z = 0.312$ (LaMassa et al. 2015), which was discovered to change from a type 1 to type 1.9 spectrum between its Sloan Digital Sky Survey (SDSS) DR1 spectrum in 2001 and its SDSS-III Baryon Oscillation Spectroscopic Survey (BOSS) spectrum in 2010. Since then, there was a systematic search by Ruan et al. (2016) of SDSS quasars with multiple epochs of spectra for which the spectroscopic pipeline classification changed between a “QSO” and a “GALAXY,” or vice versa, which recovered SDSS J0159+0033 and revealed two more cases of changing-look quasars (at $z = 0.198$ and $z = 0.243$) with a dimming in their continuum and disappearance of the broad H β line on a timescale of 5–7 yr in the rest frame. An archival search of SDSS quasars with multiple epochs of spectra and large amplitudes of variability ($\Delta g > 1$ mag) by MacLeod et al. (2016) also yielded SDSS J0159+0033 and nine more changing-look quasars at $z = 0.2$ – 0.6 , five of which show the *appearance* of broad H β between the SDSS and BOSS spectral epochs. New epochs of spectra from the Time-Domain Spectroscopic Survey (TDSS; Morganson et al. 2015) have also revealed one new case of a transition from a type 1 to type 1.9 quasar at $z = 0.246$ (Runnoe et al. 2016).

Here we report the rapid (< 1 yr) emergence of a type 1 radio-quiet quasar from a galaxy at $z = 0.237$ with weak narrow-line emission in its pre-event spectrum characteristic of a LINER nucleus that does not require an AGN to power the line emission. Throughout this paper, we refer to quasars as radio-quiet or radio-loud AGNs above a bolometric luminosity of 10^{44} erg s $^{-1}$. We also adopt a cosmology where $H_0 = 70$ km s $^{-1}$, $\Omega_M = 0.3$, and $\Omega_\Lambda = 0.7$, yielding a luminosity distance for iPTF 16bco of $d_L = 1186$ Mpc.

2. Observations

2.1. SDSS Archival Imaging and Spectroscopy

The source SDSS J155440.25+362952.0 was imaged by SDSS on UT 2003 April 29 (all days hereafter are in the UT system) and morphologically classified as a galaxy with $r = 18.18 \pm 0.01$ mag. It was targeted in the SDSS spectroscopic legacy survey as a *ugri*-selected quasar, selected for lying more than 4σ from the stellar locus (Richards et al. 2002) at high Galactic latitude (QSO_CAP). It turns out that while the colors measured in the SDSS survey are outside the stellar locus, with $u - g = +0.44 \pm 0.04$ mag, $g - r = +0.92 \pm 0.01$ mag, and $r - i = +0.48 \pm 0.01$ mag, given the known redshift of the galaxy, they are inconsistent

with the quasar color–redshift relation measured for the SDSS spectroscopic sample (Schneider et al. 2007).

The SDSS legacy spectrum, obtained on 2004 June 16, was determined by the spectroscopic pipeline to have a $z = 0.2368$ and a spectroscopic classification of a “GALAXY,” with a stellar velocity dispersion of $\sigma_* = 176 \pm 14$ km s $^{-1}$. The galaxy classification was due to the presence of strong galaxy absorption features (Ca H & K, G band, Mg I, Na D), with only weak [O III] and [N II] emission lines detected. According to the $M_{\text{BH}} - \sigma_*$ scaling relation, this velocity dispersion corresponds to a central black hole mass of $(1_{-0.7}^{+2}) \times 10^8 M_\odot$ (McConnell & Ma 2013).

We find no evidence for significant flux variations between the SDSS image in 2003 and legacy spectrum in 2004. The synthetic *r*-band magnitude of the 2004 SDSS spectrum (measured by projecting the best-fit spectral template onto the *r*-band filter) is $\text{spectroSynFlux}_r = 18.82 \pm 0.02$ mag. The corresponding fiber magnitude for the SDSS imaging in 2003 (measured with an aperture equal to the 3'' diameter spectroscopic fiber) is $r = 18.95$ mag. Since the SDSS spectrum in 2004 is dominated by host-galaxy starlight in the wavelength range of the *r* band ($\lambda_{\text{eff}} = 6231$ Å), we conclude that the SDSS image in 2003 is also dominated by host-galaxy starlight.

2.2. GALEX Archival Imaging

The source was observed by the GALEX All-Sky Imaging Survey on 2004 May 15 with a 6'' (4 pixel) radius aperture magnitude corrected for the total energy enclosed (Morrissey et al. 2007) of $\text{NUV} = 21.61 \pm 0.35$ mag, a 5σ point-source upper limit of $\text{FUV} > 20.6$ mag for $t_{\text{exp}} = 112$ s, and a background of 2.6×10^{-3} counts s $^{-1}$ pixel $^{-1}$. Note that the color measured by GALEX and SDSS of $\text{NUV} - r = +3.4 \pm 0.35$ mag is entirely consistent with normal galaxies with a similar luminosity on the blue sequence (Wyder et al. 2007).

2.3. iPTF Detection

iPTF 16bco was discovered as a transient detection by the Palomar 48-in telescope (P48) on 2016 June 1 during an (intermediate) Palomar Transient Factory (iPTF) *g+r*-band experiment by the real-time difference-imaging pipeline run at LBNL (Cao et al. 2016), with $g = 19.4$ mag and $r = 19.6$ mag. The source was flagged as a “nuclear” transient given the measured offset of 0''.44 from its host galaxy, within our centroiding accuracy of 0''.8. The data were re-reduced with the PTFIDE pipeline run at IPAC (Masci et al. 2016).

2.4. Follow-up Spectroscopy

On the next day after the discovery (2016 June 2), the transient iPTF 16bco was followed up with the robotic low-resolution ($R = \lambda/\Delta\lambda \sim 100$) Integral Field Unit (IFU) spectrograph, part of the Spectral Energy Distribution Machine (SEDM) instrument on the Palomar 60-in telescope (P60). The spectrum was strikingly different from its archival SDSS spectrum, and three more epochs of spectroscopy were obtained with DEIMOS (Faber et al. 2003) on the Keck-II telescope on 2016 June 4 and the DeVeny spectrograph on the Discovery Channel Telescope (DCT) on 2016 June 13 and 2016 July 9.




Figure 1. Series of follow-up spectra of iPTF 16bco. Spectra are normalized and have been offset vertically for clarity. Dotted lines show the wavelengths of broad Balmer lines at $z = 0.2368$. The uncorrected telluric A-band absorption feature at ~ 7600 Å in the DCT spectra is also marked.

The SEDM IFU obtained 2700 s exposures on 2016 June 2 and July 24, and data reduction was performed by the SEDM pipeline.¹⁶ Sky subtraction was performed using “A–B” extraction: two exposures of the same length are taken offset by a few arcseconds, so that the object of interest lies on an empty sky region on the next exposure, and then subtracted from each other to remove the sky lines. The spectrum is extracted in each exposure separately, and then the fluxes are summed to get the final spectrum.

The Keck spectrum was obtained with a $0''.8$ slit, and the exposure time was 240 s. Data were reduced with usual procedures from the DEEP2 pipeline in IDL (Cooper et al. 2012; Newman et al. 2013) and in PyRAF. The flux calibration and telluric correction were done with the flux standard star BD 262606. The spectrum covers wavelengths ranging from 4550 to 9550 Å with a spectral resolution of ~ 4 Å. The DCT spectra were obtained with a $1''.5$ slit and an exposure time of 600 s and 1200 s, respectively, with a spectral resolution of ~ 9 Å. Data were reduced in standard IRAF routines, and flux calibration was performed with the flux standard star BD +40d4032.

Figure 1 shows the series of follow-up spectra, which demonstrate strong, broad, Balmer emission lines characteristic of a type 1 quasar at $z = 0.2368$.

2.5. Palomar 60-inch Imaging

We also monitored the source in three filters (g , r , i) with the SEDM on the P60 telescope. The data were host-subtracted using FPipe (Fremling et al. 2016). One epoch of the P60 data points on MJD 57,576 is from the GRBCam. We do not plot the i -band GRBCam data from this night, due to the large difference in the shape of its filter transmission curve in this band. The light curve of the transient iPTF 16bco is presented in Figure 2, and the photometry is given in Table 1. We adjust the g -band P48 photometry by +0.25 mag in order to match the P60 photometry. This offset is attributed to the difference in filter curves in the g band and the strong blue continuum. Note that since its discovery by iPTF on 2016 June 1 in the g and r bands, iPTF 16bco has retained a blue color, $g - r \approx -0.1$




Figure 2. Light curve of iPTF 16bco. Optical g -, r -, and i -band difference-imaging photometry is from the Palomar 48-in (solid-circles) and 60-in telescopes (filled squares: SEDM, open squares: GBMCam), while ultraviolet aperture photometry in the $uvw2$ band is from the *Swift* telescope (stars), with a negligible contribution from host-galaxy light. The dashed red line shows the mean r -band upper limit measured during PTF observations between 2009 and 2012. The epochs for which spectra were taken are marked with an S.

mag, and demonstrated a rise in brightness of ~ 0.5 mag over a timescale of ~ 1 month.

2.6. PTF Historical Light Curve

The source was observed in the PTF survey in 2009–2012, and no variability was detected in this time frame, with a median transient point-source upper limit over 151 epochs of $r < 20.9$ mag, with the last nondetection on 2012 May 28. When adding in the host-galaxy flux measured by SDSS, this corresponds to a total magnitude of $r < 18.10$ mag, or a $\Delta r < 0.08$ mag during this time period. The P48 observations constrain the onset of the nuclear transient to be after the last nondetection on 2012 May 28, 4 yr before the iPTF discovery.

2.7. Swift Observations

The source was observed with our *Swift* Key Project program for UV follow-up of iPTF nuclear transients (PI: Gezari) on 2016 June 21 and July 5. We extracted the source from a $5''.0$ region with a background region of $20''.0$ radius using the task `uvotsource` in HEASoft, which includes a correction for the enclosed energy in the aperture. The source was observed in the $uvw2$ filter with ~ 1 ks exposures and detected with 19.30 ± 0.05 mag and 19.11 ± 0.05 mag in the AB system, respectively. The corresponding UV–optical color of iPTF 16bco (with a negligible contribution of UV flux from the host) is $NUV - r \sim -0.5$ mag, notably bluer than the near-UV $-r$ colors of low-redshift quasars measured by *GALEX* and SDSS ($NUV - r = 0.0$ – 0.5 mag; Bianchi et al. 2005; Agüeros et al. 2005).

Simultaneous *Swift* XRT observations were processed with the UK *Swift* Data Science Centre¹⁷ pipeline that takes into account dead columns and vignetting to extract counts from the source in the energy range of 0.3–10 keV. The X-ray count rate on June 21 and July 5 is 0.025 ± 0.0057 and 0.013 ± 0.0059 counts s^{-1} , respectively. We further obtained a 1.7 ks *Swift* XRT exposure on 2016 October 21, and the source was

¹⁶ <http://www.astro.caltech.edu/sedm>

¹⁷ http://www.swift.ac.uk/user_objects/

Table 1
iPTF Photometry

Telescope+Camera	Filter	MJD	Magnitude	Error
P48+CFH12k	<i>g</i>	57,540.336	19.66	0.07
P48+CFH12k	<i>g</i>	57,540.371	19.77	0.07
P48+CFH12k	<i>g</i>	57,547.328	19.58	0.11
P48+CFH12k	<i>g</i>	57,547.359	19.55	0.07
P48+CFH12k	<i>g</i>	57,551.375	19.67	0.06
P48+CFH12k	<i>g</i>	57,555.297	19.83	0.12
P48+CFH12k	<i>g</i>	57,555.324	19.69	0.12
P48+CFH12k	<i>r</i>	57,540.305	19.63	0.10
P48+CFH12k	<i>r</i>	57,547.293	19.72	0.13
P48+CFH12k	<i>r</i>	57,555.270	19.63	0.12
P60+SEDM	<i>g</i>	57,546.410	19.68	0.03
P60+SEDM	<i>g</i>	57,547.211	19.76	0.03
P60+SEDM	<i>g</i>	57,553.465	19.92	0.05
P60+SEDM	<i>g</i>	57,558.398	19.86	0.08
P60+SEDM	<i>g</i>	57,564.219	19.88	0.32
P60+SEDM	<i>g</i>	57,579.262	19.43	0.02
P60+SEDM	<i>g</i>	57,582.199	19.33	0.02
P60+SEDM	<i>g</i>	57,584.281	19.26	0.03
P60+SEDM	<i>g</i>	57,586.215	19.24	0.03
P60+SEDM	<i>g</i>	57,588.258	19.25	0.04
P60+SEDM	<i>g</i>	57,590.227	19.23	0.07
P60+SEDM	<i>g</i>	57,592.266	19.20	0.02
P60+SEDM	<i>g</i>	57,593.266	19.14	0.02
P60+SEDM	<i>g</i>	57,594.258	19.09	0.02
P60+SEDM	<i>g</i>	57,595.258	19.13	0.02
P60+SEDM	<i>g</i>	57,596.281	19.12	0.02
P60+SEDM	<i>g</i>	57,599.246	19.10	0.02
P60+SEDM	<i>g</i>	57,601.211	19.06	0.02
P60+SEDM	<i>g</i>	57,607.199	19.10	0.01
P60+SEDM	<i>g</i>	57,611.289	19.08	0.02
P60+SEDM	<i>g</i>	57,613.254	19.14	0.03
P60+SEDM	<i>g</i>	57,615.242	19.04	0.04
P60+SEDM	<i>g</i>	57,618.230	19.09	0.04
P60+SEDM	<i>g</i>	57,620.223	19.03	0.04
P60+SEDM	<i>g</i>	57,622.266	19.01	0.04
P60+SEDM	<i>g</i>	57,624.199	19.08	0.02
P60+SEDM	<i>g</i>	57,626.188	19.11	0.02
P60+SEDM	<i>r</i>	57,546.406	19.59	0.03
P60+SEDM	<i>r</i>	57,547.207	19.71	0.04
P60+SEDM	<i>r</i>	57,551.395	19.88	0.03
P60+SEDM	<i>r</i>	57,553.461	19.74	0.05
P60+SEDM	<i>r</i>	57,558.391	19.64	0.07
P60+SEDM	<i>r</i>	57,564.215	19.72	0.02
P60+SEDM	<i>r</i>	57,579.258	19.59	0.04
P60+SEDM	<i>r</i>	57,582.191	19.43	0.02
P60+SEDM	<i>r</i>	57,584.277	19.32	0.04
P60+SEDM	<i>r</i>	57,586.211	19.28	0.03
P60+SEDM	<i>r</i>	57,588.254	19.27	0.03
P60+SEDM	<i>r</i>	57,592.262	19.26	0.02
P60+SEDM	<i>r</i>	57,593.258	19.26	0.02
P60+SEDM	<i>r</i>	57,594.254	19.25	0.02
P60+SEDM	<i>r</i>	57,595.254	19.21	0.02
P60+SEDM	<i>r</i>	57,596.277	19.19	0.02
P60+SEDM	<i>r</i>	57,599.242	19.17	0.02
P60+SEDM	<i>r</i>	57,601.207	19.16	0.02
P60+SEDM	<i>r</i>	57,607.191	19.17	0.02
P60+SEDM	<i>r</i>	57,609.301	19.20	0.02
P60+SEDM	<i>r</i>	57,611.285	19.19	0.02
P60+SEDM	<i>r</i>	57,613.250	19.18	0.03
P60+SEDM	<i>r</i>	57,615.238	19.19	0.05
P60+SEDM	<i>r</i>	57,618.223	19.08	0.06
P60+SEDM	<i>r</i>	57,620.219	19.12	0.03
P60+SEDM	<i>r</i>	57,622.262	19.17	0.05
P60+SEDM	<i>r</i>	57,624.191	19.15	0.02

Table 1
(Continued)

Telescope+Camera	Filter	MJD	Magnitude	Error
P60+SEDM	<i>r</i>	57,626.184	19.17	0.02
P60+SEDM	<i>i</i>	57,546.406	19.60	0.05
P60+SEDM	<i>i</i>	57,547.211	19.73	0.05
P60+SEDM	<i>i</i>	57,551.398	19.82	0.09
P60+SEDM	<i>i</i>	57,553.465	19.84	0.10
P60+SEDM	<i>i</i>	57,558.395	19.69	0.06
P60+SEDM	<i>i</i>	57,564.219	19.75	0.07
P60+SEDM	<i>i</i>	57,579.258	19.53	0.04
P60+SEDM	<i>i</i>	57,582.195	19.39	0.03
P60+SEDM	<i>i</i>	57,584.281	19.35	0.04
P60+SEDM	<i>i</i>	57,586.215	19.31	0.03
P60+SEDM	<i>i</i>	57,588.254	19.30	0.04
P60+SEDM	<i>i</i>	57,592.262	19.19	0.03
P60+SEDM	<i>i</i>	57,593.262	19.19	0.03
P60+SEDM	<i>i</i>	57,594.254	19.20	0.03
P60+SEDM	<i>i</i>	57,595.254	19.19	0.03
P60+SEDM	<i>i</i>	57,596.277	19.17	0.03
P60+SEDM	<i>i</i>	57,599.242	19.19	0.02
P60+SEDM	<i>i</i>	57,601.211	19.18	0.03
P60+SEDM	<i>i</i>	57,607.195	19.19	0.02
P60+SEDM	<i>i</i>	57,611.285	19.22	0.04
P60+SEDM	<i>i</i>	57,613.254	19.18	0.03
P60+SEDM	<i>i</i>	57,615.238	19.20	0.06
P60+SEDM	<i>i</i>	57,618.227	19.12	0.04
P60+SEDM	<i>i</i>	57,620.223	19.13	0.04
P60+SEDM	<i>i</i>	57,622.266	19.11	0.04
P60+SEDM	<i>i</i>	57,624.195	19.17	0.02
P60+SEDM	<i>i</i>	57,626.188	19.16	0.02
P60+GRBCam	<i>g</i>	57,576.348	19.50	0.03
P60+GRBCam	<i>r</i>	57,576.348	19.58	0.03

detected with 0.027 ± 0.004 counts s^{-1} . This confirms the lack of significant X-ray variability between the *Swift* observations. Furthermore, the combined spectrum of all the data can be modeled with an absorbed power law with a spectral index of $\Gamma = 2.1 \pm 0.5$ and N_H fixed at the Galactic value of 1.63×10^{20} cm^{-2} (Dickey & Lockman 1990). The average unabsorbed 0.2–10 keV flux of the source is 9×10^{-13} erg $s^{-1} cm^{-2}$, corresponding to a luminosity of 1.5×10^{44} erg s^{-1} by assuming an absorbed power law with $N_H = 1.63 \times 10^{20}$ cm^{-2} and $\Gamma = 2.1$ at $z = 0.2368$.

2.8. Archival X-Ray Observations

The *ROSAT* upper limit in the 0.1–2.4 keV band from the All-Sky Survey in 1990–1991 (Voges et al. 1999) is 0.1 counts s^{-1} , which corresponds to an unabsorbed flux of $\sim 7 \times 10^{-13}$ erg $s^{-1} cm^{-2}$. There are even more constraining 2σ upper limits from the XMM Slew Survey¹⁸ of <0.601 and <0.817 counts s^{-1} in the 0.2–12.0 keV band on 2011 February 27 and 2015 February 08, corresponding to $<2.7 \times 10^{-13}$ and $<3.2 \times 10^{-13}$ erg $s^{-1} cm^{-2}$, respectively. The latest XMM Slew Survey upper limit implies a factor of >3 increase in flux in the *Swift* XRT detection on a timescale of <1.1 yr in the quasar rest frame.

¹⁸ <http://xmm.esac.esa.int/UpperLimitsServer/>




Figure 3. $H\beta$ (left) and $H\alpha$ (right) regions of iPTF 16bc0 during its pre-event spectrum from SDSS. The red line shows our galaxy template fit, and the blue line shows the emission-line component fit. The residual from the galaxy template fit is also shown, along with the emission-line component fit. The individual Gaussian components of the $[N\ II]+H\alpha$ complex are plotted with dashed lines. The emission lines are all fitted with a narrow Gaussian with $\sigma = 420\text{ km s}^{-1}$, with no evidence for a broad $H\alpha$ or $H\beta$ line.

2.9. Radio Observations

We observed iPTF 16bc0 with the AMI-LA at 15.5 GHz on 2016 October 16.63. The source is not detected, with the 3σ upper limit of $68\ \mu\text{Jy}$. We can also convert this to a 1.4 GHz upper limit of $370\ \mu\text{Jy}$ using a spectral index of -0.7 . This is consistent with the nondetection in the VLA FIRST survey (Becker et al. 1995) from 1999, which gives an independent 1.4 GHz upper limit of $500\ \mu\text{Jy}$.

3. Analysis

3.1. Host-galaxy Classification

The archival SDSS spectrum from 2004 was fitted by the automated spectroscopic pipeline (Bolton et al. 2012) with a combination of stellar, galaxy, and quasar templates plus emission lines. After visual inspection of the pipeline fit, we found a poor fit to the $H\alpha+[N\ II]$ and $[O\ III]$ emission line complexes and refitted the spectrum with host-galaxy template and emission-line gas components using `ppxf` (Cappellari & Emsellem 2004; Cappellari 2016), which uses the MILES stellar template library (Vazdekis et al. 2010). The emission-line fits are shown in Figure 3 and are fitted with a narrow Gaussian with a $\sigma = 420\text{ km s}^{-1}$, with no evidence for broad $H\alpha$ or $H\beta$ components. The narrow-line ratios of $\log([O\ III]\lambda 5007/H\beta) = 0.74 \pm 0.13$ and $\log([N\ II]\lambda 6583/H\alpha) = 0.47 \pm 0.07$, together with $L([O\ III]\lambda 5007) = (1.0 \pm 0.1) \times 10^{41}\text{ erg s}^{-1}$, classify the SDSS spectrum as a type 2 AGN in the LINER region (Kewley et al. 2006) in the diagnostic narrow-line diagrams (Baldwin et al. 1981; Veilleux & Osterbrock 1987; Kauffmann et al. 2003) shown in Figure 4. Note that the archival *WISE* colors from the all-sky survey in 2010 (Cutri et al. 2012) of $W1 - W2 = 0.48 \pm 0.04\text{ mag}$ and $W2 - W3 = 1.6 \pm 0.2\text{ mag}$, where $W1$, $W2$, and $W3$ are 3.4, 4.6, and $12\ \mu\text{m}$, respectively, place the host in the region of Seyfert and star-forming galaxies (Yan et al. 2013). However, in the WHAN diagram (shown in Figure 4) for emission-line galaxies (Cid Fernandes et al. 2010, 2011), the weak equivalent width of $H\alpha$ ($W_{H\alpha} = 1.6 \pm 0.3\ \text{\AA}$) classifies the galaxy as a “retired galaxy” powered by hot, low-mass, evolved (post-asymptotic giant branch) stars and *not* an AGN.




Figure 4. Diagnostic narrow-line ratio diagrams for iPTF 16bc0 during its pre-event spectrum from SDSS. Top: BPT diagram, with the lines demarcating star-forming galaxies from AGNs. The solid line is the theoretical curve from Kewley et al. (2001), and the dashed line is the empirical curve based on the SDSS spectroscopic sample from Kauffmann et al. (2003). The line demarcating Seyferts from LINER galaxies from Cid Fernandes et al. (2010) is plotted with a dotted line. Bottom: diagnostic WHAN diagram defined by Cid Fernandes et al. (2010), with the regions demarcating star-forming galaxies, Seyferts, weak AGNs (wAGNs), and “fake AGNs” powered by stars from Cid Fernandes et al. (2011).




Figure 5. Left: dramatic change in spectrum between the archival SDSS legacy spectrum obtained on 2004 June 16 and the follow-up spectrum obtained by Keck 2 +DEIMOS on 2016 June 4. Right: difference spectrum corrected for Galactic extinction. Tick marks show the broad Balmer lines ($H\alpha$, $H\beta$, and $H\delta$), as well as the broad Fe II complexes, and He I $\lambda 5877$. The dashed blue line shows a power-law fit to the continuum, $f_\lambda \propto \lambda^{\alpha_\lambda}$, with $\alpha_\lambda = -1.45$.

3.2. Change of State

The transient iPTF 16bco shows two remarkable changes: a factor of 10 increase in UV flux, and a transformation from a LINER galaxy to a luminous type 1 quasar. Figure 5 shows the dramatic change of state between the SDSS spectrum in 2004 and the follow-up Keck spectrum in 2016.

3.2.1. Continuum Variability

According to the empirical correlation between hard X-ray emission and [O III] luminosity for AGNs (Heckman et al. 2005; Ueda et al. 2015), from the [O III] $\lambda 5007$ luminosity measured in the pre-event SDSS spectrum in 2004, one would expect $L(2-10)\text{ keV} \sim 10^{43} \text{ erg s}^{-1}$. This could be even lower if the narrow-line emission in iPTF 16bco is powered by stars, and it infers a pre-event X-ray luminosity at least an order of magnitude below the X-ray luminosity of $1.5 \times 10^{44} \text{ erg s}^{-1}$ measured by *Swift* XRT during the type 1 state of iPTF 16bco in 2016. The change in near-UV flux between the *GALEX* AIS measurement in 2004 and the *Swift* UVOT measurement in 2016 also indicates a brightening by $\Delta m = -2.4 \pm 0.4$ mag. Allowing for some fraction of the near-UV flux in the *GALEX* measurement to be from star formation in the host galaxy, this yields a lower limit of a factor of ~ 10 increase in flux.

When adding back in the host-galaxy r -band magnitude measured by SDSS (see Section 2.1), the amplitude of optical variability since 2003 is only 0.3 mag. However, since the SDSS photometry in 2003 is dominated by host-galaxy starlight, the 0.3 mag of variability is only a *lower* limit to the true amplitude of the continuum increase in the optical.

Thus, we conclude that the continuum in iPTF 16bco had an increase in X-ray/UV/optical flux by a factor of >10 between 2004 and 2016. It is not entirely surprising that such a large increase in the photoionizing continuum was also accompanied by dramatic spectral changes.

3.2.2. Spectral Variability

The spectra of iPTF 16bco demonstrate a strong blue continuum and broad Balmer-line, Fe II, and He I features characteristic of a type 1 quasar. Figure 5 also shows the difference between the Keck spectrum in 2016 and the archival

SDSS spectrum from 2004. After correcting for Galactic extinction of $E(B-V) = 0.021$ mag from the Schlegel et al. (1998) dust extinction map and using the extinction curve of Cardelli et al. (1989), the continuum in the difference spectrum (after masking the broad emission lines) is reasonably fitted with a single power law of $f_\lambda \propto \lambda^{\alpha_\lambda}$, where $\alpha_\lambda = -1.45$. This is shallower than the standard theoretical thin accretion disk spectrum with $f_\nu \propto \nu^{\alpha_\nu}$, where $\alpha_\nu = 1/3$, and $\alpha_\lambda = -(\alpha_\nu + 2) = -2.33$, which is well fitted to difference spectra (Wilhite et al. 2005; MacLeod et al. 2016) and difference spectral energy distributions (Hung et al. 2016) of quasars. However, it is close to the power law observed in *averaged optical* quasar spectra ($\alpha_\lambda = -1.56$) blueward of $H\beta$ (Vanden Berk et al. 2001; Wilhite et al. 2005). A power-law index similar to the average quasar spectrum in the difference spectrum is another indication that the low-state spectrum has very little contribution from a nonstellar continuum.

The [O III] $\lambda 5007$ luminosity in 2016 is $(1.6 \pm 0.3) \times 10^{41} \text{ erg s}^{-1}$, consistent within the errors of the [O III] luminosity measured in SDSS. The broad $H\alpha$ flux, in contrast, dramatically appears, with $L(H\alpha, \text{ broad}) = (8.22 \pm 0.09) \times 10^{42} \text{ erg s}^{-1}$ with an FWHM of $4048 \pm 36 \text{ km s}^{-1}$. This difference in behavior between the broad and narrow lines can be explained by the fact the [O III] line luminosity traces the average AGN continuum over a much longer timescale than the flaring event detected by iPTF. While the emissivity decay time (dominated by recombination charge transfer) for [O III] $\lambda 5007$ is only a few years, light-travel time effects prolong the response time to hundreds of years (the light-crossing time of the narrow-line region; Eracleous et al. 1995). Thus, the narrow-line region has not yet had the chance to respond to the continuum flux changes happening on the timescale of $\lesssim 1$ yr.

The monochromatic optical luminosity in the 2016 Keck spectrum, after subtracting the starlight using the archival SDSS spectrum, is $\lambda L_\lambda(5100 \text{ \AA}) = \lambda f_\lambda 4\pi d_L^2 (1+z) = 1.6 \times 10^{-16} \text{ erg s}^{-1} \text{ \AA}^{-1} 4\pi d_L^2 (1+z) = 1.7 \times 10^{44} \text{ erg s}^{-1}$. This is in excellent agreement with the nearly linear correlation between broad $H\alpha$ luminosity and optical continuum luminosity in AGNs (Greene & Ho 2005), which for this broad $H\alpha$ luminosity one would expect to be $\lambda L_\lambda(5100 \text{ \AA}) \sim 1.5 \times 10^{44} \text{ erg s}^{-1}$.

The FWHM of the broad $H\beta$ line, $4770 \pm 200 \text{ km s}^{-1}$, and the monochromatic luminosity at 5100 \AA can be used to

estimate the central black hole mass to be $2_{-1.5}^{+4.0} \times 10^8 M_\odot$ (Vestergaard & Peterson 2006), in good agreement with M_{BH} inferred from the host-galaxy stellar velocity dispersion. We adopt a bolometric correction factor for the monochromatic optical luminosity of 8.1 (Runnoe et al. 2012) to get $L_{\text{bol}} = 1.4 \times 10^{45} \text{ erg s}^{-1}$. We then derive an Eddington ratio of $\lambda_{\text{Edd}} = L_{\text{bol}}/L_{\text{Edd}} = 0.05$ during the type 1 quasar state.

3.3. Nature of the Variability

3.3.1. Variable Obscuration or Microlensing

The Eddington ratio inferred for iPTF 16fnl from its nearly constant narrow [O III] $\lambda 5007$ line luminosity is in disagreement with the Eddington ratio observed in its type 1 quasar state ($\lambda_{\text{Edd}} \sim 0.05$). Therefore, we favor a change in accretion rate (intrinsic change of state) in the quasar as opposed to variable obscuration (extrinsic change of state) of a nonvariable quasar.

Timescale arguments also disfavor a variable line-of-sight extinction due to the disappearance of an intervening absorber to explain the appearance of the broad H α line in iPTF 16bco. In the case of an obscuring cloud, the distance between the nucleus and the cloud must be larger than the radius of the broad-line region it is obscuring. Using the radius–luminosity relation measured for AGNs from reverberation mapping studies of broad lines (Bentz et al. 2013), the luminosity of iPTF 16bco would have H β broad-line emission with a characteristic radius of $R_{\text{BLR}} \sim 45$ days. Following the argument of LaMassa et al. (2015, their Equation (4)), this translates to a crossing time on a circular, Keplerian orbit, $t_{\text{cross}} = \Delta\phi/\omega_K$, where $\Delta\phi = \arcsin(r_{\text{src}}/r_{\text{orb}})$ is the angular length of the arc and $\omega_K = \frac{\sqrt{GM}}{2\pi} r^{-3/2}$ is the Keplerian frequency, which for $r_{\text{orb}} > R_{\text{BLR}}$ yields $t_{\text{cross}} > 15$ yr, which is much longer than the timescale over which the continuum appeared in iPTF 16bco.

Finally, we note that lensing of a background broad-line quasar by a star in an intervening galaxy (Lawrence et al. 2016) is also ruled out, since the redshift of the quasar is the same as the galaxy in its dim state.

3.3.2. Tidal Disruption Event (TDE)

One mechanism to rapidly increase the mass accretion rate onto an SMBH is to get a new supply of gas from a star that wanders close enough to the SMBH to be torn apart by tidal forces. In a TDE, roughly half of the disrupted stellar debris remains bound to the black hole, falls back onto the SMBH, circularizes through shocks, and is accreted (Rees 1988). The characteristic timescale of a TDE is the orbital period of the most tightly bound debris, which is given by $\Delta t = 0.35 M_7^{1/2} m_*^{-1} r_*^{3/2}$ yr, where $M_7 = M_{\text{BH}}/10^7 M_\odot$, $m_* = M_*/M_\odot$, and $r_* = R_*/R_\odot$ (Lodato & Rossi 2011). The peak mass accretion rate is given by $\dot{M}_{\text{acc}} = (1/3)(M_*/\Delta t)$ and can exceed the Eddington rate ($\dot{M}_{\text{Edd}} = 0.2 M_7 (\eta/0.1)^{-1} M_\odot \text{ yr}^{-1}$, where η is the radiative efficiency) for black holes $< 10^7 M_\odot$. Following the peak, a TDE has a characteristic $t^{-5/3}$ power-law decay determined by the fallback rate of the stellar debris to pericenter (Rees 1988; Evans & Kochanek 1989; Phinney 1989).

However, we point out several issues with interpreting the flaring state of iPTF 16bco with a TDE: (1) For an $M_{\text{BH}} \gtrsim 10^8 M_\odot$, the tidal disruption radius of a solar-type star

($R_T = R_*(M_{\text{BH}}/M_*)^{1/3} = 3.23 \times 10^{13} M_8^{1/3} \text{ cm}$) is smaller than the Schwarzschild radius ($R_S = 2GM_{\text{BH}}/c^2 = 2.95 \times 10^{13} M_8 \text{ cm}$), and the star crosses the event horizon before being disrupted. (2) The X-ray power-law ($\Gamma = 2.1$) continuum of iPTF 16bco is unlike the extremely soft, thermal X-ray spectra observed in TDEs (Komossa 2002; Miller et al. 2015). (3) The light curve of iPTF 16bco shows a complex shape uncharacteristic of a TDE, with a month-long plateau, followed by a 2-week rise to another plateau. (4) The broad Balmer lines in iPTF 16bco are narrower and stronger than have been observed in TDEs, and iPTF 16bco does not have strong, broad He II $\lambda 4686$, which is characteristic of TDE spectra (Gezari et al. 2012; Arcavi et al. 2014; Holoien et al. 2014, 2016a, 2016b). (5) Finally, the broad-line emission and X-ray continuum in iPTF 16bco are consistent with radio-quiet quasars in the ‘‘Eigenvector 1’’ parameter space: the FWHM velocity width of the broad H β line versus the ratio of the equivalent widths of the Fe II $\lambda 4570$ complex to broad H β strength ($R_{\text{Fe II}}$) (Sulentic et al. 2000) favors a change in \dot{M}_{acc} of a preexisting accretion disk, instead of a newly formed debris disk from a TDE. Continued photometric monitoring can determine whether the light curve of iPTF 16bco eventually evolves into a power-law decline expected for a TDE.

3.4. Accretion Disk Instabilities

We now investigate the scenario that iPTF 16bco was the result of a change of state in a preexisting quasar accretion disk. Interestingly, however, iPTF 16bco puts stringent limits on the timescale over which such accretion rate changes must occur. In the rest frame, iPTF 16bco demonstrates a dramatic change in continuum flux over a timescale of $\Delta t < 4/(1+z) = 3.23$ yr, or in < 1.1 yr based on the archival X-ray upper limits. The timescale by which an accretion disk can change its accretion rate should be determined by the viscous radial inflow timescale, t_{infl} . Furthermore, this timescale is expected to be longer for a quasar ‘‘turning on’’ instead of ‘‘turning off,’’ since it scales as $\lambda_{\text{Edd}}^{-2}$ (LaMassa et al. 2015). The t_{infl} corresponding to the Eddington ratio and black hole mass estimated for iPTF 16bco in its dim state, and assuming a radiation-pressure-dominated inner region of a Shakura–Sunyaev disk, is,

$$t_{\text{infl}} = 1300 \text{ yr} \left[\frac{\alpha}{0.1} \right]^{-1} \left[\frac{\lambda_{\text{Edd}}}{0.005} \right]^{-2} \left[\frac{\eta}{0.1} \right]^2 \left[\frac{r}{10 r_g} \right]^{7/2} \left[\frac{M_8}{2.0} \right],$$

a much longer timescale than the observed rapid change in continuum flux in iPTF 16bco.

One possibility could be an accretion disk eruption as the result of thermal-viscous instabilities in a partial ionization zone, analogous to the outbursts observed in cataclysmic variables (CVs) and X-ray novae (Siemiginowska et al. 1996). However, while such instabilities can produce amplitudes of a factor of 10^4 , the expected durations scale with the central mass, and so while CVs and X-ray novae show large-amplitude outbursts on the timescale of weeks to months, for an accretion disk around a $10^8 M_\odot$ black hole, this corresponds to timescales of $\sim 10^5$ yr. However, state changes on \sim minute timescales have been observed in some X-ray binaries (Fender 2001; Fender & Belloni 2004), which would imply similar changes in an SMBH on the timescale of only ~ 10 yr.

In the accretion disk instability model, the more narrow the unstable zone, the more rapid the timescale of variability. The size of the unstable zone depends on the accretion rate: the

lower the accretion rate, the closer in the unstable region is to the inner edge of the disk. However, the smaller the unstable zone, the smaller the expected amplitude of variability. For models that demonstrate a factor of ~ 10 variability ($\alpha = 0.1$, $\dot{M} = 3 \times 10^{-5} M_{\odot} \text{ yr}^{-1}$) in Siemiginowska et al. (1996), they have a “turn-on” timescale of $\sim 10^3$ yr. This is still 3 orders of magnitude longer than we require for the “turn-on” timescale of iPTF 16bco.

Interestingly, the thermal timescale itself, $t_{\text{th}} \sim 1/\alpha\Omega_K$, is much shorter, adopting Equation (8) from (Siemiginowska et al. 1996):

$$t_{\text{th}} \sim 2.7(\alpha/0.1)^{-1} M_8^{-0.5} (r/10^{16} \text{ cm})^{1.5} \text{ yr.}$$

A disk with local thermal fluctuations, potentially driven by the magnetorotational instability, could be consistent with the rapid timescale of the continuum variability in iPTF 16bco (Dexter & Agol 2011). While an inhomogeneous disk has been demonstrated to fit composite difference spectra (Ruan et al. 2014) and color variability of quasars (Schmidt et al. 2012), Hung et al. (2016) find a simple disk model adequate to fit difference-flux UV/optical spectral energy distributions of individual quasars. Moreover, Kokubo (2015) argues that the tight interband correlations observed in SDSS quasar light curves are inconsistent with the inhomogeneous disk model. However, whether or not these thermal fluctuations can be coherent enough to produce a large-amplitude outburst is still to be determined.

An instability that arises in a radiation-pressure-dominated disk (Lightman & Eardley 1974) has been used to model recurrent flares in accreting Galactic X-ray binaries (Belloni et al. 1997) and has recently been applied by Saxton et al. (2015) and Grupe et al. (2015) to explain the large-amplitude soft X-ray flares in Seyfert galaxies NGC 3599 and IC 3599, respectively. In this scenario, when the internal radiation pressure of the disk become greater than the gas pressure, a heating wave propagates through the disk. This results in an enhanced local viscosity, scale height, and accretion rate, which rapidly drains the disk. The instability recurs when the inner disk fills back in. The rise time and recurrence timescale can be as short as a year and hundreds of years for a $10^8 M_{\odot}$ black hole, respectively.

Another mechanism for driving large-amplitude variability in a quasar accretion disk could be related to the presence of a binary SMBH. Hydrodynamical simulations show that in a circumbinary disk, streams penetrate the disk cavity to feed the primary and secondary black hole at a periodic rate, and that at close to equal mass ratios, the perturbed circumbinary disk has an enhanced accretion rate that can be quite bursty on a timescale of ~ 5 times the orbital period of the SMBH binary (Farris et al. 2014). Such a scenario could in principle be testable, due to the intrinsically periodic nature of the outbursts. Furthermore, one could look for periodic changes in the broad-line profiles, if they originate in the circumbinary disk. However, the circumbinary disk outbursts in Farris et al. (2014) have a sawtooth pattern, with a rise time much shorter than the decay time. Already the light curve of iPTF 16bco is not in good agreement with this model, given its rebrightening by ~ 0.5 mag during its high state (see Figure 2).

The continued intrinsic variability during the high state of iPTF 16bco, as also seen in the study of changing-look quasars by MacLeod et al. (2016), may also provide insight as to the

nature of what caused the “changing look” of the quasar. The optical variability amplitudes of these sources in their type 1 quasar states of ~ 0.5 – 1.0 mag are on the high-amplitude tail of what is typically observed on these timescales for quasars of a similar luminosity range (MacLeod et al. 2012), although we note that in both of our studies the changing-look quasars were selected by their optical variability. However, intrinsic variability is also observed in the changing-look quasar presented in LaMassa et al. (2015), which was selected based on its spectral changes. In fact, the rapid rise and power-law decline of its light curve were interpreted as a signature of a TDE (Merloni et al. 2015). However, the light curves of iPTF 16bco and the other changing-look quasars with archival photometry have large fluctuations that are inconsistent with the smooth, power-law decline observed in the optical light curves of known TDEs (e.g., Gezari et al. 2008; van Velzen et al. 2011; Gezari et al. 2012; Arcavi et al. 2014; Holoien et al. 2014, 2016a, 2016b). The erratic intraburst behavior in the two-state (hot and cold) α instability models of Siemiginowska et al. (1996) could be promising. A potential testable prediction is that in these models, the accretion disks spend the majority of their time in the low state. The variability during the enhanced accretion state in iPTF 16bco could also be a signature of clumpy accretion in an advection-dominated accretion flow (ADAF), for which cold clumps form in the accretion flow owing to instabilities in the radiation-dominated regions of the disk (Wang et al. 2012)

3.5. Disk–Jet Connection

The dramatic change in accretion rate from $\lambda_{\text{Edd}} \lesssim 0.005$ to $\lambda_{\text{Edd}} \sim 0.05$ inferred for iPTF 16bco could be accompanied by a structural change in the accretion flow if the quasar accretion disk is transitioning from a radiatively inefficient to radiatively efficient mode. Note that such changes in X-ray binaries are often accompanied by changes in jet activity. The implied high-state radio-to-optical flux density ratio for iPTF 16bco of $R = \log(S_{1.4 \text{ GHz}}/S_{\text{opt}}) < 0.8$ and radio luminosity $L_R < 3 \times 10^{22} \text{ W Hz}^{-1}$ are typical of radio-quiet AGNs (e.g., Padovani et al. 2011). While these values are consistent with the LINER classification from the optical spectrum, it is surprising that if the accretion event in iPTF 16bco were triggered by a disk instability, there would be no evidence for a jet or outflow during its high state in the radio. This is in contrast with X-ray binaries and CVs, which generally show flaring at radio, optical, and X-ray wavelengths alongside strong Balmer emission lines. Similarly, the fundamental plane of black hole activity (e.g., Plotkin et al. 2012; Saikia et al. 2015) predicts L_R significantly greater than $\sim 10^{22} \text{ W Hz}^{-1}$ when $L_X = 1.5 \times 10^{44} \text{ erg s}^{-1}$.

3.6. Comparison with Other Changing-look Quasars

iPTF 16bco is one of only a dozen other changing-look quasars (here defined as $M_i < -22$ mag, $L([\text{O III}]) > 10^{41} \text{ erg s}^{-1}$), roughly half of which have been caught in the act of “turning on” by demonstrating the sudden appearance of broad lines. Figure 6 shows the redshift and [O III] luminosity of all the changing-look quasars in the literature that pass our [O III] luminosity cut (thus, we exclude SDSS J0126-0839 and SDSS J2336+0017 from Ruan et al. 2016), color-coded by whether they show appearing broad lines or disappearing broad lines. We also do not include three changing-look quasars from the




Figure 6. Comparison of iPTF 16bco with other changing-look quasars that have disappearing (red circles) and appearing (blue circles) broad-line emission. Left: $[\text{O III}] \lambda 5007$ luminosity during the high state of the quasar vs. redshift. Right: broad $\text{H}\alpha$ luminosity during the high state of the quasar vs. the $[\text{O III}] \lambda 5007$ luminosity during the high state. Contours show the distribution of 20%, 50%, and 90% of the quasars’ $L(\text{H}\alpha)$ vs. $L([\text{O III}])$ ratio for the DR7 SDSS quasar sample from Shen et al. (2011). iPTF 16bco is an outlier of this distribution, with a high broad $\text{H}\alpha$ luminosity relative to its $[\text{O III}] \lambda 5007$ luminosity.

MacLeod et al. (2016) sample that do not have good coverage of the broad $\text{H}\alpha$ line in its high state (appearing SDSS J214613 at $z = 0.62$, disappearing SDSS J022562 at $z = 0.63$, and both appearing and disappearing SDSS J022556 at $z = 0.50$). Given that all the other appearing changing-look quasars are from MacLeod et al. (2016), there appears to be a bias toward finding appearing broad lines in higher-redshift galaxies. This is likely due to the fact that the BOSS spectra extend to longer wavelengths than the SDSS spectra, and thus finding an appearing broad $\text{H}\alpha$ line in the BOSS spectrum is more likely at higher redshift than finding a disappearing broad $\text{H}\alpha$ line in the SDSS spectrum.

Figure 6 also shows the $[\text{O III}] \lambda 5007$ versus broad $\text{H}\alpha$ luminosity in all the changing-look quasars in their type 1 state, in comparison to the full SDSS quasar sample. We determine the $[\text{O III}] \lambda 5007$ luminosity from the line flux measured in the SDSS DR7 *SpecLine* table and the DR12 interactive spectrum line measurement table. We use the broad $\text{H}\alpha$ fluxes from the Shen et al. (2011) catalog for broad-line quasars, or from the literature when available. All luminosities are calculated for our adopted cosmology. Note that iPTF 16bco is on the edge of the normal quasar distribution, while the other changing-look quasars reported in the literature appear to lie squarely in the distribution of normal quasars in this parameter space.

The enhanced broad $\text{H}\alpha$ luminosity observed in iPTF 16bco relative to $[\text{O III}]$ in comparison to normal quasars, as well as the previously discovered changing-look quasars, is likely a signature of its rapid transition to a type 1 state. As discussed in Section 3.2.2, given the extended size of the narrow-line region, the $[\text{O III}]$ line will lag in its response to a continuum flare in comparison to the broad emission lines due to light-travel time effects. Interestingly, in the MacLeod et al. (2016) sample, the rise time in the continuum flux of the quasars was serendipitously measured by photometric monitoring to be

~ 1000 days. Given the significantly weaker $[\text{O III}]$ line emission relative to broad $\text{H}\alpha$ in iPTF 16bco compared to these objects, this would imply an even shorter “turn-on” timescale, in agreement with the inferred “turn-on” timescale for the continuum in iPTF 16bco of $\lesssim 1$ yr.

One interesting aspect of changing-look quasars is the lack of strong changes in Mg II line emission, despite the dramatic changes in the Balmer lines. This was explained by MacLeod et al. (2016) as being due to the relatively weak responsivity of the Mg II line to continuum flux changes, as has been measured in rest-frame UV reverberation mapping studies (Cackett et al. 2015). Given that Mg II is a low-ionization line, its weak responsivity can be explained as a consequence of the stratification of the broad-line region (Korista & Goad 2004). Unfortunately, we do not have short-enough wavelength coverage in the archival or follow-up spectra to determine the presence and/or response of the Mg II line during the change of state in iPTF 16bco.

4. Summary

We present the rapid “turn-on” of a luminous broad-line quasar at $z = 0.237$ discovered from its nuclear optical variability in the iPTF survey (iPTF 16bco) and identified as a newly emerged quasar from comparison of follow-up spectroscopy with an archival SDSS spectrum from over a decade earlier, which shows LINER narrow-line emission potentially powered by stars. Pre-event optical, UV, and X-ray imaging indicates that the quasar continuum increased by a factor of > 10 on a timescale of $\lesssim 1$ yr in the quasar rest frame. The broadband properties of iPTF 16bco in its high state are best explained by an intrinsic change of state to a radio-quiet type 1 quasar, rather than variable obscuration or a TDE. However, continued monitoring will help further constrain the nature of its rapid brightening. The dramatic appearance of broad Balmer lines during the high state of iPTF 16bco, with

no significant change in the [O III] λ 5007 line, is explained as a delayed response of gas in the narrow-line region to the flare in photoionizing continuum due to light-travel time effects. The enhanced broad H α /narrow [O III] λ 5007 ratio in iPTF 16bco relative to normal quasars and previously reported changing-look quasars is further evidence that iPTF 16bco may have demonstrated the most rapid change of state yet observed in a quasar. iPTF 16bco pushes the limits of accretion disk theory and may represent a new class of state changes in quasars that will be discovered more routinely in regular monitoring of millions of quasars with the next generation of optical time domain surveys (Zwicky Transient Facility and the Large Synoptic Survey Telescope), together with follow-up spectroscopy triggered by flaring events.

We thank the anonymous referee for their helpful comments that improved the manuscript. S.G. is supported in part by NSF CAREER grant 1454816 and NASA Swift Cycle 12 grant NNX16AN85G. S.G. thanks Mike Koss for help with the X-ray data archives. These results made use of the Discovery Channel Telescope at Lowell Observatory. Lowell is a private, non-profit institution dedicated to astrophysical research and public appreciation of astronomy and operates the DCT in partnership with Boston University, the University of Maryland, the University of Toledo, Northern Arizona University, and Yale University. The W. M. Keck Observatory is operated as a scientific partnership among the California Institute of Technology, the University of California, and NASA; the Observatory was made possible by the generous financial support of the W. M. Keck Foundation. This research used resources of the National Energy Research Scientific Computing Center, a DOE Office of Science User Facility supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. K.M.'s research is supported by the Oxford Centre for Astrophysical Surveys, which is funded through generous support from the Hintze Family Charitable Foundation. The AMI telescope gratefully acknowledges support from the European Research Council under grant ERC-2012- StG-307215 LODESTONE, the UK Science and Technology Facilities Council (STFC), and the University of Cambridge. We thank the AMI staff for scheduling the observations.

References

- Agüeros, M. A., Ivezić, Ž., Covey, K. R., et al. 2005, *AJ*, **130**, 1022
 Antonucci, R. 1993, *ARA&A*, **31**, 473
 Arcavi, I., Gal-Yam, A., Sullivan, M., et al. 2014, *ApJ*, **793**, 38
 Baldwin, J. A., Phillips, M. M., & Terlevich, R. 1981, *PASP*, **93**, 5
 Becker, R. H., White, R. L., & Helfand, D. J. 1995, *ApJ*, **450**, 559
 Belloni, T., Méndez, M., King, A. R., van der Klis, M., & van Paradijs, J. 1997, *ApJL*, **479**, L145
 Bentz, M. C., Denney, K. D., Grier, C. J., et al. 2013, *ApJ*, **767**, 149
 Bianchi, L., Seibert, M., Zheng, W., et al. 2005, *ApJL*, **619**, L27
 Bolton, A. S., Schlegel, D. J., Aubourg, E., et al. 2012, *AJ*, **144**, 144
 Cackett, E. M., Gültekin, K., Bentz, M. C., et al. 2015, *ApJ*, **810**, 86
 Cao, Y., Nugent, P. E., & Kasliwal, M. M. 2016, *PASP*, **128**, 114502
 Cappellari, M. 2016, arXiv:1607.08538
 Cappellari, M., & Emsellem, E. 2004, *PASP*, **116**, 138
 Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, **345**, 245
 Cid Fernandes, R., Stasińska, G., Mateus, A., & Vale Asari, N. 2011, *MNRAS*, **413**, 1687
 Cid Fernandes, R., Stasińska, G., Schlickmann, M. S., et al. 2010, *MNRAS*, **403**, 1036
 Cooper, M. C., Griffith, R. L., Newman, J. A., et al. 2012, *MNRAS*, **419**, 3018
 Cutri, R. M., et al. 2012, VizieR Online Data Catalog, 2311
 Denney, K. D., De Rosa, G., Croxall, K., et al. 2014, *ApJ*, **796**, 134
 Dexter, J., & Agol, E. 2011, *ApJL*, **727**, L24
 Dickey, J. M., & Lockman, F. J. 1990, *ARA&A*, **28**, 215
 Eracleous, M., Livio, M., & Binette, L. 1995, *ApJL*, **445**, L1
 Evans, C. R., & Kochanek, C. S. 1989, *ApJL*, **346**, L13
 Faber, S. M., Phillips, A. C., Kibrick, R. I., et al. 2003, *Proc. SPIE*, **4841**, 1657
 Farris, B. D., Duffell, P., MacFadyen, A. I., & Haiman, Z. 2014, *ApJ*, **783**, 134
 Fender, R., & Belloni, T. 2004, *ARA&A*, **42**, 317
 Fender, R. P. 2001, *MNRAS*, **322**, 31
 Fremling, C., Sollerman, J., Taddia, F., et al. 2016, *A&A*, **593**, A68
 Gezari, S., Basa, S., Martin, D. C., et al. 2008, *ApJ*, **676**, 944
 Gezari, S., Chornock, R., Rest, A., et al. 2012, *Natur*, **485**, 217
 Greene, J. E., & Ho, L. C. 2005, *ApJ*, **630**, 122
 Grupe, D., Komossa, S., & Saxton, R. 2015, *ApJL*, **803**, L28
 Guo, H., Malkan, M. A., Gu, M., et al. 2016, *ApJ*, **826**, 186
 Heckman, T. M., Ptak, A., Hornschemeier, A., & Kauffmann, G. 2005, *ApJ*, **634**, 161
 Hickox, R. C., Mullaney, J. R., Alexander, D. M., et al. 2014, *ApJ*, **782**, 9
 Holoien, T. W.-S., Prieto, J. L., Bersier, D., et al. 2014, *MNRAS*, **445**, 3263
 Holoien, T. W.-S., Kochanek, C. S., Prieto, J. L., et al. 2016a, *MNRAS*, **463**, 3813
 Holoien, T. W.-S., Kochanek, C. S., Prieto, J. L., et al. 2016b, *MNRAS*, **455**, 2918
 Hung, T., Gezari, S., Jones, D. O., et al. 2016, arXiv:1609.06307
 Husemann, B., Urrutia, T., Tremblay, G. R., et al. 2016, *A&A*, **593**, L9
 Kauffmann, G., Heckman, T. M., Tremonti, C., et al. 2003, *MNRAS*, **346**, 1055
 Kelly, B. C., Bechtold, J., & Siemiginowska, A. 2009, *ApJ*, **698**, 895
 Kewley, L. J., Dopita, M. A., Sutherland, R. S., Heisler, C. A., & Trevena, J. 2001, *ApJ*, **556**, 121
 Kewley, L. J., Groves, B., Kauffmann, G., & Heckman, T. 2006, *MNRAS*, **372**, 961
 Kokubo, M. 2015, *MNRAS*, **449**, 94
 Komossa, S. 2002, in *Lighthouses of the Universe: The Most Luminous Celestial Objects and Their Use for Cosmology*, ed. M. Gilfanov, R. Sunyaev, & E. Churazov (Berlin: Springer), 436
 Korista, K. T., & Goad, M. R. 2004, *ApJ*, **606**, 749
 LaMassa, S. M., Cales, S., Moran, E. C., et al. 2015, *ApJ*, **800**, 144
 Lawrence, A., Bruce, A. G., MacLeod, C., et al. 2016, arXiv:1605.07842
 Lightman, A. P., & Eardley, D. M. 1974, *ApJL*, **187**, L1
 Lodato, G., & Rossi, E. M. 2011, *MNRAS*, **410**, 359
 MacLeod, C. L., Ivezić, Ž., Sesar, B., et al. 2012, *ApJ*, **753**, 106
 MacLeod, C. L., Ross, N. P., Lawrence, A., et al. 2016, *MNRAS*, **457**, 389
 Marchese, E., Braitto, V., Della Ceca, R., Caccianiga, A., & Severgnini, P. 2012, *MNRAS*, **421**, 1803
 Masci, F., Laher, R., Rebbapragada, U., et al. 2016, arXiv:1608.01733
 McConnell, N. J., & Ma, C.-P. 2013, *ApJ*, **764**, 184
 McElroy, R. E., Husemann, B., Davis, T. A., et al. 2016, *A&A*, **593**, L8
 Merloni, A., Dwelly, T., Salvato, M., et al. 2015, *MNRAS*, **452**, 69
 Miller, J. M., Kaastra, J. S., Miller, M. C., et al. 2015, *Natur*, **526**, 542
 Morganson, E., Green, P. J., Anderson, S. F., et al. 2015, *ApJ*, **806**, 244
 Morrissey, P., Conrow, T., Barlow, T. A., et al. 2007, *ApJS*, **173**, 682
 Newman, J. A., Cooper, M. C., Davis, M., et al. 2013, *ApJS*, **208**, 5
 Novak, G. S., Ostriker, J. P., & Ciotti, L. 2011, *ApJ*, **737**, 26
 Osterbrock, D. E. 1981, *ApJ*, **249**, 462
 Padovani, P., Miller, N., Kellermann, K. I., et al. 2011, *ApJ*, **740**, 20
 Pereyra, N. A., Vanden Berk, D. E., Turnshek, D. A., et al. 2006, *ApJ*, **642**, 87
 Phinney, E. S. 1989, in *IAU Symp. 136, The Center of the Galaxy*, ed. M. Morris (Dordrecht: Kluwer), 543
 Plotkin, R. M., Markoff, S., Kelly, B. C., Kording, E., & Anderson, S. F. 2012, *MNRAS*, **419**, 267
 Ponti, G., Terrier, R., Goldwurm, A., Belanger, G., & Trap, G. 2010, *ApJ*, **714**, 732
 Rees, M. J. 1988, *Natur*, **333**, 523
 Richards, G. T., Fan, X., Newberg, H. J., et al. 2002, *AJ*, **123**, 2945
 Risaliti, G., Miniutti, G., Elvis, M., et al. 2009, *ApJ*, **696**, 160
 Ruan, J. J., Anderson, S. F., Cales, S. L., et al. 2016, *ApJ*, **826**, 188
 Ruan, J. J., Anderson, S. F., Dexter, J., & Agol, E. 2014, *ApJ*, **783**, 105
 Runnoe, J. C., Brotherton, M. S., & Shang, Z. 2012, *MNRAS*, **422**, 478
 Runnoe, J. C., Cales, S., Ruan, J. J., et al. 2016, *MNRAS*, **455**, 1691
 Saikia, P., Kording, E., & Falcke, H. 2015, *MNRAS*, **450**, 2317
 Saxton, R. D., Motta, S. E., Komossa, S., & Read, A. M. 2015, *MNRAS*, **454**, 2798
 Schawinski, K., Evans, D. A., Virani, S., et al. 2010, *ApJL*, **724**, L30
 Schawinski, K., Koss, M., Berney, S., & Sartori, L. F. 2015, *MNRAS*, **451**, 2517

- Schlegel, D. J., Finkbeiner, D. P., & Davis, M. 1998, *ApJ*, 500, 525
- Schmidt, K. B., Rix, H.-W., Shields, J. C., et al. 2012, *ApJ*, 744, 147
- Schneider, D. P., Hall, P. B., Richards, G. T., et al. 2007, *AJ*, 134, 102
- Shappee, B. J., Prieto, J. L., Grupe, D., et al. 2014, *ApJ*, 788, 48
- Shen, Y., Richards, G. T., Strauss, M. A., et al. 2011, *ApJS*, 194, 45
- Siemiginowska, A., Czerny, B., & Kostyunin, V. 1996, *ApJ*, 458, 491
- Storchi-Bergmann, T., Baldwin, J. A., & Wilson, A. S. 1993, *ApJL*, 410, L11
- Sulentic, J. W., Zwitter, T., Marziani, P., & Dultzin-Hacyan, D. 2000, *ApJL*, 536, L5
- Ueda, Y., Hashimoto, Y., Ichikawa, K., et al. 2015, *ApJ*, 815, 1
- van Velzen, S., Farrar, G. R., Gezari, S., et al. 2011, *ApJ*, 741, 73
- Vanden Berk, D. E., et al. 2001, *AJ*, 122, 549
- Vazdekis, A., Sánchez-Blázquez, P., Falcón-Barroso, J., et al. 2010, *MNRAS*, 404, 1639
- Veilleux, S., & Osterbrock, D. E. 1987, *ApJS*, 63, 295
- Vestergaard, M., & Peterson, B. M. 2006, *ApJ*, 641, 689
- Voges, W., Aschenbach, B., Boller, T., et al. 1999, *A&A*, 349, 389
- Wang, J.-M., Cheng, C., & Li, Y.-R. 2012, *ApJ*, 748, 147
- Wilhite, B. C., Vanden Berk, D. E., Kron, R. G., et al. 2005, *ApJ*, 633, 638
- Wyder, T. K., Martin, D. C., Schiminovich, D., et al. 2007, *ApJS*, 173, 293
- Yan, L., Donoso, E., Tsai, C.-W., et al. 2013, *AJ*, 145, 55