

Research Article

Iron Content of Some Popular Cooked Foods Consumed by the Rural School Children in Sri Lanka

D. Ruwani G. Perera ¹, **Dilantha Gunawardana**,¹ **Renuka Jayatissa**,²
and **A. Buddhika G. Silva** ²

¹Department of Botany, University of Sri Jayewardenepura, Nugegoda 10250, Sri Lanka

²Department of Nutrition, Medical Research Institute, Borella 00800, Sri Lanka

Correspondence should be addressed to A. Buddhika G. Silva; buddhika-gayani@mri.gov.lk

Received 10 May 2019; Revised 26 August 2019; Accepted 1 October 2019; Published 3 November 2019

Academic Editor: Luis Patarata

Copyright © 2019 D. Ruwani G. Perera et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Iron is an important micronutrient that can be found in different food sources in varying quantities. Iron deficiency is common in populations consuming cereal-based diets with little or no animal-derived food products. Rice is the staple for Sri Lankans, which may not provide sufficient bioavailable iron. Sri Lankan food composition table does not contain data on iron in home-cooked foods. Aim of the present study was to determine the iron content in commonly consumed home-cooked foods (rice, vegetables, green leaves, pulses, fish, etc.) by children aged 15-16 years. The study was carried out in eight schools at the Divisional Secretariat, Horana. The average iron values of cooked rice, vegetables, green leaves, pulses, and fish were 23.29 ± 14.46 , 19.34 ± 9.90 , 29.56 ± 13.64 , 45.16 ± 22.28 , and 46.72 ± 14.53 mg/kg, respectively. A serving of cooked red raw rice (rathu kekulu) provided the highest Estimated Daily Intake (EDI) of iron. The total EDI per serving for all categories was 6.39 ± 1.06 mg, contributing to the recommended daily allowance (RDA) of 42.62% of girls and 58.12% of boys as defined by the United States Department of Agriculture. Meanwhile, based on the Sri Lankan RDA, dietary iron contribution per serving ranges from 16.39% to 17.76% in girls and 16.39% to 21.31% in boys in the 15-16 years age category. Therefore, these home-cooked foods which were prepared using local recipes have high iron content and can replenish a major proportion of recommended quotidian values for iron intake.

1. Introduction

Micronutrient malnutrition is a widespread problem throughout the world [1, 2]. Prevention of micronutrient malnutrition is a basic requirement in ensuring the rapid and appropriate national development [3-5]. The achievement and maintaining of the desirable level of nutritional quality in the national food supply is an important public health objective [3]. To achieve it, improving the dietary habits in the current society is a major challenge. An analytical assessment of the staple diet is necessary to identify healthy dietary strategies within the population. Iron is an important micronutrient [6-8], and it can cause ill effects with either lower or higher levels than the desired level [4, 9]. Iron is found to be necessary in certain quantities in food and beverages including drinking water. Therefore,

determination of iron content in food is essential for both food safety and nutritional considerations [7].

Sri Lankan staple diet is mainly based on plant-based food, and they are deficient in iron [4]. More than 40% of children, adolescent girls, and women of childbearing age including pregnant women are currently suffering from iron deficiency. In Sri Lanka, anaemia has become a moderate public health problem among preschool, nonpregnant, and pregnant populations as the prevalence are 33%, 39%, and 34%, respectively [4]. Iron deficiency anaemia is the most common type of anaemia throughout the world, and it is the common cause of nutritional anaemia in Sri Lanka as well [9].

The required amount of iron can be taken through iron supplements or by eating foods that are rich in iron [5]. Different food sources (plants or animals) contain iron in

different quantities [5]. Depending on the cooking method, the iron content can be changed [5]. Furthermore, there is limited available knowledge on nutrient composition of the cooked foods in Sri Lanka [3]. Therefore, it is necessary to analyze the nutrient composition of cooked foods which are prepared according to the Sri Lankan domestic/local recipes. The results of this research can be used by the Ministry of Health, Sri Lanka, in policymaking and to increase public health awareness. The aim of this study was to analyze the iron content of selected commonly consumed cooked food among children aged 15-16 years of age.

2. Materials and Methods

2.1. Commonly Consumed Cooked Food by the Target Group.

Data on food consumption were obtained by a 24-h recall method among 36 boys and 36 girls in the Divisional Secretariat, Horana. Recalls were done for 3 consecutive days per person to capture weekdays and weekends for 2-month period. During the field visits, the recipes of commonly consumed cooked food including approximate amounts of ingredients used were recorded. When the same food was prepared using different recipes, the most popular recipe was used for the laboratory studies. Average portion size for each food item was obtained.

2.2. Sample Preparation. The required raw materials and ingredients were purchased from the retail shops and boutiques in the Horana area. All food items analyzed were cooked by using utensils which did not contain iron.

2.3. Determining Iron in Food Samples. Initially, the fresh weight of the sample was recorded. Each sample was dried in an oven for 3 days at 60°C, and the dry weight was recorded. A representative portion from each cooked food sample was crushed to a fine homogeneous powder by using a blender. Each duplicate sample (0.2500 g) was digested using 3 mL of 69%–78% (w/w) nitric acid and 1 mL of hydrogen peroxide (analytical grade) using a microwave digester. The clear digestion was brought up to 25 mL with deionized water, and they were analyzed in triplicates using a flame atomic absorption spectrophotometer (AAS) at a wavelength of 248.32 nm. Calibration standard for iron was prepared using a stock of 1000 µg·mL⁻¹ using the serial dilution technique.

2.4. Quality Control. All glassware used in the assay was cleaned using detergent water, then rinsed with 5% nitric acid, then rinsed several times with deionized water, and was finally oven-dried. All plastic utensils were cleaned with detergent water, 5% nitric acid, and deionized water, respectively. Then, they were allowed to dry under room temperature. Each batch of runs on the atomic absorption spectrophotometry (AAS) was standardized by running standard solutions. The method validation was done by spiking of known concentration of a standard solution with the known weight of the sample and then by estimating the recovery results.

2.5. Data Management and Statistical Analysis. The average iron content and estimated daily intake (EDI) of iron in each food item were recorded as the mean ± standard deviation (SD). Comparison of the average iron content of each food type was performed orderly using one-way ANOVA in SPSS 23 software package. Statistical analysis was designed at $p < 0.05$.

2.6. Estimated Daily Intake Per Serving. The daily intake of an element is a function of the concentration of the selected element in the food sample as well as the amount of food that is consumed daily. The estimated daily intake (EDI) was calculated using the following equation [10, 11]:

$$EDI = C \times M, \quad (1)$$

where C is the mean concentration of element (iron) in each food sample and M refers to the assumed consumption rate [10, 11]. In this research, consumption rate per person was assumed as the weight of one serving of each food (1 tablespoon for vegetable curries, green leaves, pulses, and recommended serving sizes for fish varieties [12]). The obtained values were then compared with the recommended daily allowances (RDA) of children aged 15-16 years. RDA provides the information on average daily intake level that is sufficient to meet the nutrient requirements of nearly all (97-98%) healthy individuals in a particular life stage and a gender group, which is the standard value in assessing the adequacy of daily intake [13]. The values of the RDA for iron for the boys and girls in the 15-16 year age group are presented in Table 1.

2.7. Contribution of Iron. The estimated contribution of various cooked foods to the daily intake of iron was therefore calculated using the following equation [16]:

$$\text{Contribution to RDA per serving} = \frac{\text{EDI per serving}}{\text{RDA}} \times 100\%. \quad (2)$$

3. Results and Discussion

The daily diet of the children comprised three main meals with different accompaniments. The composition of lunch and dinner was similar with rice and curry being the most popular choice. Rice is the staple food when considering main meals rather than wheat flour-based foods. Even though red raw rice (rathu kakulu haal) is popular as the most nutritious variety of rice available in Sri Lanka with a neutral flavor and good cooking qualities, it was not commonly consumed among the selected study population. Instead, white raw rice (sudu kekulu haal) and parboiled small rice grains (samba) were two of the more popular varieties among children.

The vegetables consumed were beetroot (*Beta vulgaris*), pumpkin (*Cucurbita moschata*), yellow cabbage (*Brassera oleracea*), ash plantain (*Plantago lanceolata*), potato (*Solanum tuberosum*), ladies' fingers (*Abelmoschus esculentus*), brinjal (*Solanum melongena*), and carrot (*Daucus*

TABLE 1: RDA for iron in boys and girls in the 15-16 age range.

Gender	RDA USDA (mg/d) [14]	RDA Sri Lanka (mg/d) [15]	
	15-16 year age group	15 years	16 years
Girls	15	36	39
Boys	11	30	39

RDA = recommended daily allowance; USDA = United States Department of Agriculture.

carota), while vegetable hummingbird (*kathurumurunga*) (*Sesbania grandiflora*), sessile joy weed (*mukunuwenna*) (*Alternanthera sessilis*), and centella (*gotukola*) (*Centella asiatica*) were the most commonly consumed green leafy vegetables. A considerable consumption was recorded regarding the nonvegetarian foods such as dried sprats (*Stolephorus commersonii*), salmon (Brand A), sardinella (*salaya*) (*Sardinella gibbosa*), Spanish mackerel (*thora*) (*Scomberomorus commersoni*), sailfish (*thalapath*) (*Istiophorus platypterus*), yellowfin tuna (*kelawallo*) (*Thunnus albacares*), red snapper (*rathu gal malu*) (*Lutjanus argentimaculatus*), and queenfish (*kattawa*) (*Scomberoides commersonianus*) were the most commonly consumed fish varieties. The most popular pulse varieties among the selected population were lentils and chickpeas which were also selected for the analysis.

While rice with a variety of vegetables, animal sources, and coconut sambol were frequent items in their main meals, wheat flour-based products and cereals were also included in the main meals from time to time. Commonly consumed popular foods among the children and their methods of preparation are described in Table 2.

3.1. Vegetables. Pumpkin curry contained a significantly higher amount of iron ($p < 0.05$) than the rest of the vegetable curries (Table 3). The iron content in the carrot curry was also significantly higher, and it is also higher than the iron content in raw carrots which had been reported by similar studies [17–19]. Except for the pumpkin and carrot curries, no significant difference was observed among the average iron contents of the other vegetable curries. Some studies reported that uncooked beetroot contains 3.7 mg/kg of iron [20], while pickled-drained beetroot contains 3 mg/kg [17] and 5 mg/kg [18] of iron. Therefore, based on the present study, beetroot curry which was cooked according to the local recipe contained a higher amount of iron than raw and pickled-drained beetroot. Yellow cabbage curry has higher iron content than raw cabbage which had been reported in Bruggraber et al. [17], McCance and Widdowson [19], Narain and Ilango [21], and Gooneratne and Kumarapperuma [22]. Raw potato has 7.5 mg/kg [20] of iron while boiled potato has 8.5 mg/kg [17] of iron according to previous studies, while the current study reported a higher iron content in cooked potato compared with the above two. Higher iron content was found to be present in the brinjal curry compared with the value reported in Narin and Ilango (3.16 mg/kg) [21]. Therefore, the average iron contents of cooked vegetables were found to be higher than the amount present in raw vegetables.

3.2. Pulses. The most common type of pulse among the study population was lentil curry. It had a significantly higher iron content (72.49 ± 0.14 mg/kg) than the rest (Table 3). A similar study reported the iron content in uncooked lentils to be 62.8 ± 35.0 mg/kg [3], which is similar to cooked lentils. Daraniyagala and Gunawardena [5], Bruggraber et al. [17], McCance et al. [18], McCance and Widdowson [19], and Karunaratne et al. [23] reported much lower values for the iron content in lentils. Therefore, lentil curry can be considered as a good source of iron. No significant difference was found in the iron contents of the two chickpea samples prepared using different preparation methods. Previous studies revealed that the iron content in uncooked chickpea was 85.0 ± 2 mg/kg [3] which is significantly higher than the iron content in cooked chickpea (Table 3). Karunaratne et al. [23] also reported a higher iron content in cooked chickpeas. Similar studies reported the iron content of boiled chickpea as 34.0 ± 0.0 mg/kg [5] and cooked chickpea as 35.0 ± 5 mg/kg [5], which are in agreement with the current study. According to Bruggraber et al. [17], boiled chickpea had a lower iron content than chickpea curry, which is different from the result obtained in the current study. The iron content in cooked chickpea reported in Daraniyagala and Gunawardena [5] was similar to the present study.

3.3. Green Leaves. Centella (*gotukola* sambol) was found to be the best iron source (45.12 ± 8.33 mg/kg) having a significantly higher average iron content ($p < 0.05$) compared with the other two types (Table 3), which are in accordance with the values reported in Gooneratne and Kumarapperuma [22], but not compatible to results obtained by Deraniyagala et al. [3]. Another study reported the mean iron content of selected fresh green leaves was 68.2 ± 47.0 mg/kg and ranged from 35.6 mg/kg to 132 mg/kg [24]. Values reported in Anthony Swamy et al. [25] of uncooked green leafy vegetables ranged from 0.81 to 0.23 mg/kg and the cooked green leafy vegetables had iron levels ranging from 0.041 to 0.43 mg/kg while these results were not compatible with the present study [25].

3.4. Fish. The highest iron content in 1 kg of food sample was found in fried marinated sardinella (*salaya*) (68.33 ± 4.98 mg/kg), which was significantly higher compared with the curries of other fish varieties (Table 3). The iron content obtained for sardinella (*salaya*) in the present study was compatible with the values reported by Wickramasinghe et al. [26]. Sardinella (*salaya*) curry also has a significantly higher iron content than the rest. Dried sprats have a significantly higher iron content than salmon (Brand

TABLE 2: Commonly consumed foods by the study population and their respective methods of preparation.

Food category	Food type	Method of preparation
Vegetables	Beetroot curry	Cooked 60 g of small cut pieces using coconut milk (60 g of coconut kernel), 0.38 g of chili powder, 0.25 teaspoons (tsp) of curry powder (0.5 g), 0.44 g of turmeric powder, onions, curry leaves
	Pumpkin curry	Cooked 120 g of small cut pieces using coconut milk (100 g of coconut kernel), 0.38 g of chili powder, 0.25 tsp of curry powder (0.5 g), 0.44 g of turmeric powder, garlic, a small piece of cinnamon, black pepper, onions, curry leaves
	Yellow cabbage curry	Same as the beetroot curry
	Ash plantain curry	Cooked 120 g of small cut pieces using coconut milk (120 g of coconut kernel), 0.25 tsp of curry powder (0.5 g), 0.25 tsp of chili powder (0.67 g), 0.44 g of turmeric powder, a small piece of cinnamon, fenugreek seeds, onions, curry leaves
	Potato curry	Same as the ash plantain curry
	Ladies' fingers curry	Cooked 120 g of small cut pieces using coconut milk (120 g of coconut kernel), 0.25 tsp of curry powder (0.5 g), 0.25 tsp of chili powder (0.67 g), 0.44 g of turmeric powder, onions, curry leaves
	Brinjal curry	Cooked 180 g of small cut pieces using a small amount of coconut oil, coconut milk (60 g of coconut kernel), 0.25 tsp of chili powder (0.67 g), 0.5 tsp of curry powder (1.0 g), 0.25 tsp of turmeric powder (0.77 g), onions, curry leaves
	Carrot curry	Same as the brinjal curry
	Pulses	Lentil curry
Chickpea curry		Cooked 150 g using coconut milk (200 g of coconut kernel), 0.25 tsp of chili powder (0.67 g), 0.5 tsp of curry powder (1 g), 0.44 g of turmeric powder, onions, curry leaves
Chickpeas boiled		Cooked 200 g using water
Green leaves	Sessile joy weed (mukunuwenna)	Cooked 60 g of cut leaves with 15 g of coconut scrapings along with 0.25 tsp of turmeric powder (0.77 g), small pieces of green chilies, onions, curry leaves
	Centella (gotukola)	Mixed 60 g of cut leaves with 15 g of coconut scrapings along with small pieces of green chilies, onions, lemon juice
	Vegetable hummingbird (kathurumurunga)	Same as the mukunuwenna malluma.

TABLE 2: Continued.

Food category	Food type	Method of preparation
Fish	Dried sprats curry	Cooked 40 g using coconut milk (120 g of coconut scrapings), 1 tsp of curry powder (2 g), 2 tsp of chili powder (5.4 g), 0.25 tsp of turmeric powder (0.77 g), curry leaves, onions, green chilies
	Salmon (brand A) curry	Cooked 50 g using coconut milk (150 g of coconut scrapings), 1.25 tsp of curry powder (2.5 g), 2 tsp of chili powder (5.4 g), 0.5 tsp of turmeric powder (1.55 g), curry leaves, onions, green chilies
	Sardinella (salaya) fried marinated	Mixed well with 0.25 tsp of turmeric powder (0.77 g) and then deep-fried in coconut oil
	Sardinella (salaya) curry	Cooked 40 g using coconut milk (120 g of coconut scrapings), 1 tsp of curry powder (2 g), 2 tsp of chili powder (5.4 g), 0.25 tsp of turmeric powder (0.77 g), curry leaves, onions, green chilies
	Spanish mackerel (thora) curry	Cooked 80 g using coconut milk (80 g of coconut scrapings), 1 tsp of curry powder (2 g), 1 tsp of chili powder (2.7 g), 0.25 tsp of turmeric powder (0.77 g), curry leaves, onions, green chilies
	Sail fish (thalapath) curry	Cooked 100 g using coconut milk (80 g of coconut scrapings), 1.5 tsp of chili powder (4.05 g), 1 tsp of curry powder (2 g), 0.25 tsp of turmeric powder (0.77 g), curry leaves, onions, green chilies
	Yellowfin tuna (kelawallo) curry	As for the thalapath curry
	Red snapper (rathu gal malu) curry	Cooked 40 g using coconut milk (120 g of coconut scrapings), 2 tsp of chili powder (5.4 g), 1 tsp of curry powder (2 g), 0.25 tsp of turmeric powder (0.77 g), curry leaves, onions, green chilies
	Queen fish (kattawa) curry	Cooked 20 g using coconut milk (120 g of coconut scrapings), 0.5 tsp of chili powder (1.35 g), 0.25 tsp of turmeric powder (0.77 g), 1/6 tsp of fenugreek seeds, curry leaves, onions, green chilies
Rice	White raw rice (sudu kekulu)	Cooked with water
	Red raw rice (rathu kekulu)	Cooked with water
	Parboiled small rice (samba)	Cooked with water
Other foods	Coconut sambol	Spicy coconut preparation using the scraped coconut kernel
	String hoppers	String hoppers (wheat flour) were taken from two boutiques in the Divisional Secretariat Division, Horana

For all food items, salt was added for taste. Onions, green chilies, and curry leaves were added in equal quantities for all food items as 11.5 g, 4 g, and 2 g, respectively. Weight of 1 tsp of turmeric powder = 3.1 g, weight of 1 tsp of chili powder = 2.7 g, and weight of 1 tsp of curry powder = 2 g.

A) and yellowfin tuna (kelawallo) curries. Karunaratne et al. [23] reported an iron content of 17.9 ± 0.6 mg/kg in dried sprats which was less than the value obtained in the present study.

3.5. Rice. All rice varieties consumed by the studied population were local origin, and the popular varieties were red raw rice (rathu kekulu), white raw rice (sudu kekulu), and parboiled small rice grains (samba). The iron levels of these three varieties ranged from 5.74 to 36.92 mg/kg (Table 3) while data obtained from similar studies reported values ranging from 20 to 35 mg/kg [7]. The iron content obtained for rice samples in the present study was compatible with the values reported by Rajbandari and Subedi [7] and Liang [27]. According to Deraniyagala et al. [3], raw samba and raw rathu kekulu

had lower iron content than cooked samba and rathu kekulu, which is different from the results obtained in the present study. Higher iron contents were found to be present in the cooked rice varieties compared with the values reported in Karunaratne et al. [23]. The level of iron was significantly higher in rathu kekulu (36.92 ± 1.37 mg/kg) compared with samba ($p = 0.014$, 95% confident interval), and there was no significant difference between the average iron content of cooked rathu kekulu and cooked sudu kekulu varieties.

3.6. Other Foods. Wheat flour consumption among the selected population was done in two ways, not leavened (as string hoppers) or leavened (as buns or bread). Based on the present study, the iron content of string hoppers (wheat flour) was 28.26 ± 2.78 mg/kg (Table 3) while 0.34 ± 0.033 mg

TABLE 3: Average iron content of cooked food varieties consumed by the adolescents (15-16 years) in the Divisional Secretariat, Horana.

Type	Iron content (mg/kg)	SD
<i>Vegetables</i>		
Beetroot curry	21.46	0.07
Pumpkin curry	38.79	0.37
Yellow cabbage curry	18.20	0.23
Ash plantain curry	9.83	0.31
Potato curry	15.66	1.26
Ladies' finger curry	7.61	0.24
Brinjal curry	15.36	6.43
Carrot curry	27.80	0.18
<i>Pulses</i>		
Lentil curry	72.49	0.14
Chickpea boiled	36.37	9.00
Chickpea curry	26.64	8.10
<i>Green leaves</i>		
Sessile joy weed (mukunuwenna malluma)	17.14	4.23
Centella (gotukola sambol)	45.12	8.33
Vegetable hummingbird (kathurumurunga malluma)	26.42	5.58
<i>Fish</i>		
Dried sprats curry	57.18	3.32
Salmon (brand A) curry	27.74	1.18
Sardinella (salaya) fried marinated	68.33	4.98
Sardinella (salaya) curry	60.46	3.43
Spanish mackerel (thora) curry	41.42	2.35
Sailfish (thalapath) curry	47.02	6.06
Yellowfin tuna (kelawallo) curry	24.17	5.40
Red snapper (rathu gal malu) curry	43.17	8.13
Queenfish (kattawa) curry	50.93	4.23
<i>Rice</i>		
White raw rice (sudu kekulu) boiled	27.22	2.05
Red raw rice (rathu kekulu) boiled	36.92	1.37
Parboiled small rice (samba) boiled	5.74	4.53
<i>Other foods</i>		
Coconut sambol	30.68	1.45
String hoppers	28.26	2.78

of iron can be obtained per serving (12 g; 1 string hopper). Scraped coconut kernel sambol/coconut sambol was used as an accompaniment to string hoppers as well as rice. The average iron content of 30.68 ± 1.45 mg was found in 1 kg of coconut kernel sambol (Table 3), while it provided 0.77 ± 0.036 mg of iron per serving (1 tbs = 25 g). A similar study reported the iron content of the combination of string hoppers (wheat flour) and coconut sambol to be 26.6 ± 2.8 mg/kg (dry weight) [23]. The iron content of uncooked scraped coconut kernel sambol was reported to be 29.4 ± 1.5 mg/kg [23] which is compatible with the present study.

Based on the USDA guidelines, children in 14–18 years of age in the adolescent age group required the highest amount of RDA [14]. In contrast, based on SL RDA, 15-16 years of children required the highest amount of iron (30 mg and 36 mg for 15-year-old boys and girls, respectively, while 39 mg for 16-year-old boys as well as girls) [15]. Meanwhile, boys and girls in the 15-16 age group were referred as upper secondary students, and they have to prepare for the G. C. E Ordinary Level examination successfully which is conducted by the Sri Lankan government to achieve their goals in

further education. Therefore, the nutritional status of the children in this age group is a necessary factor for themselves and also for the public at large. The selected homemade foods for the analysis were the most popular ones among these children, and the percentage contributions of these foods to their RDAs of iron are summarized in Table 4.

3.7. Vegetable Curries. The highest iron content per serving can be obtained by pumpkin curry (1.16 ± 0.011 mg). The mean estimated iron intake that can be obtained through these eight vegetable curries was 0.52 ± 0.30 mg per serving (Table 4). Based on the USDA guidelines, this contributes to 3.45% and 4.70% of the daily requirement of boys and girls, respectively, in the age group 15-16 years. However, based on the Sri Lankan guidelines, this estimated daily intake of iron per serving contributes to 1.44%, 1.33%, 1.72%, and 1.33% of the daily requirement of 15-year-old girls, 16-year-old girls, 15-year-old boys, and 16-year-old boys, respectively (Table 4). Contribution to the RDA by estimated average intake per serving was found to be higher in boys than in girls.

TABLE 4: Estimated daily intake of iron per serving of each food item and its contribution to RDA.

	Serving size (kg)	EDI per serving (mg)	Contribution to the RDA per serving (%)					
			Based on the USDA guidelines		Based on the SL guidelines			
			Girls 15-16 years	Boys 15-16 years	Girls 15 years	Girls 16 years	Boys 15 years	Boys 16 years
<i>(1) Vegetables</i>								
Beetroot curry	0.025	0.54 ± 0.002	3.58	4.88	1.49	1.38	1.79	1.38
Pumpkin curry	0.030	1.16 ± 0.011	7.76	10.58	3.23	2.98	3.88	2.98
Yellow cabbage curry	0.025	0.46 ± 0.006	3.03	4.14	1.26	1.17	1.52	1.17
Ash plantain curry	0.025	0.25 ± 0.008	1.64	2.23	0.68	0.63	0.82	0.63
Potato curry	0.025	0.39 ± 0.031	2.61	3.56	1.09	1.00	1.31	1.00
Ladies' finger curry	0.025	0.19 ± 0.006	1.27	1.73	0.53	0.49	0.63	0.49
Brinjal curry	0.030	0.46 ± 0.193	3.07	4.19	1.28	1.18	1.54	1.18
Carrot curry	0.025	0.69 ± 0.004	4.63	6.32	1.93	1.78	2.32	1.78
<i>Mean</i>	0.026	0.52 ± 0.30	3.45	4.70	1.44	1.33	1.72	1.33
<i>(2) Pulses</i>								
Lentil curry	0.030	2.17 ± 0.004	14.50	19.77	6.04	5.58	7.25	5.58
Chickpea boiled	0.021	0.76 ± 0.189	5.09	6.94	2.12	1.96	2.55	1.96
Chickpea curry	0.025	0.67 ± 0.202	4.44	6.05	1.85	1.71	2.22	1.71
<i>Mean</i>	0.025	1.20 ± 0.765	8.01	10.92	3.34	3.08	4.00	3.08
<i>(3) Green leaves</i>								
Sessile joy weed (mukunuwenna malluma)	0.010	0.17 ± 0.042	1.14	1.56	0.48	0.44	0.57	0.44
Centella (gotukola sambol)	0.013	0.59 ± 0.108	3.91	5.33	1.63	1.50	1.96	1.50
Vegetable hummingbird (kathurumurunga malluma)	0.010	0.26 ± 0.056	1.76	2.40	0.73	0.68	0.88	0.68
<i>Mean</i>	0.011	0.34 ± 0.203	2.27	3.10	0.95	0.87	1.14	0.87
<i>(4) Fish</i>								
Dried sprats curry	0.015	0.86 ± 0.05	5.72	7.80	2.38	2.20	2.86	2.20
Salmon (brand A) curry	0.030	0.83 ± 0.035	5.55	7.57	2.31	2.13	2.77	2.13
Sardinella (salaya) fried marinated	0.030	2.05 ± 0.15	13.67	18.64	5.69	5.26	6.83	5.26
Sardinella (salaya) curry	0.030	1.81 ± 0.103	12.09	16.49	5.04	4.65	6.05	4.65
Spanish mackerel (thora) curry	0.030	1.24 ± 0.071	8.28	11.30	3.45	3.19	4.14	3.19
Sailfish (thalapath) curry	0.030	1.41 ± 0.182	9.40	12.82	3.92	3.62	4.70	3.62
Yellowfin tuna (kelawallo) curry	0.030	0.73 ± 0.162	4.83	6.59	2.01	1.86	2.42	1.86
Red snapper (rathu gal malu) curry	0.030	1.30 ± 0.244	8.63	11.77	3.60	3.32	4.32	3.32
Queenfish (kattawa) curry	0.030	1.53 ± 0.127	10.19	13.89	4.24	3.92	5.09	3.92
<i>Mean</i>	0.028	1.31 ± 0.45	8.71	11.87	3.63	3.35	4.35	3.35
<i>(5) Rice varieties</i>								
White raw rice (sudu kekulu)	0.130	3.54 ± 0.267	23.59	32.17	9.83	9.07	11.80	9.07
Red raw rice (rathu kekulu)	0.130	4.80 ± 0.178	32.00	43.63	13.33	12.31	16.00	12.31
Parboiled small rice (samba)	0.130	0.75 ± 0.589	4.97	6.78	2.07	1.91	2.49	1.91
<i>Mean</i>	0.130	3.03 ± 1.880	20.19	27.53	8.41	7.76	10.09	7.76
<i>(6) Other foods</i>								
Coconut sambol	0.250	0.77 ± 0.036	5.11	6.97	2.13	1.97	2.56	1.97
String hopper	0.120	0.34 ± 0.033	2.26	3.08	0.94	0.87	1.13	0.87

Values of estimated daily intake per serving are mean ± SD ($n = 2$); EDI = estimated daily intake, RDA = recommended daily allowance, USDA = United States Department of Agriculture, and SL = Sri Lanka; Serving size-vegetables, pulses, green leaves, and coconut sambol: 1 tablespoon (tbs); rice: 1 teacup; fish varieties: based on the food-based dietary guidelines [12]; string hoppers = 1 string hopper.

3.8. Pulses. The highest estimated average intake of iron per serving can be obtained from lentil curry (2.17 ± 0.004 mg). Mean iron content of 1.20 ± 0.77 mg per serving can be obtained from the selected cooked pulses (Table 4). Based on the USDA guidelines, it contributes to 8.01% and 10.92% of the recommended daily requirement of girls and boys, respectively. Based on the SL guidelines, it contributes to 3.34% and 3.08% of the recommended daily requirement of girls in the age group 15 and 16 years and 4.00% and 3.08%

of the recommended daily requirement of boys in the same age group (Table 4).

3.9. Green Leaves. A mean iron content of 0.34 ± 0.203 mg can be obtained per serving from the selected green leaves (Table 4). Based on the USDA guidelines, it contributes to 2.27% and 3.10% of the daily iron requirement of girls and boys, respectively. Based on the SL guidelines, it only

contributes to 0.95%, 0.87%, 1.14%, and 0.87% of the daily requirement of 15-year-old girls, 16-year-old girls, 15-year-old boys, and 16-year-old boys, respectively (Table 4). The total iron content of these green leaf samples was dependent on the preparation method. However, green leafy vegetables are considered as a good source of iron in Sri Lankan diet, particularly among the populations that are largely dependent on a plant-based diet [24].

3.10. Fish. The highest iron content which is 2.05 mg can be obtained by marinated salaya per serving (0.030 kg). A mean iron content of 1.31 ± 0.45 mg per serving can be obtained from the selected fish varieties (Table 4). Based on the USDA guidelines, it contributes to 8.71% and 11.87% of the recommended daily requirement of girls and boys in the 15-16 age category. According to the SL guidelines, it only contributes to 3.63%, 3.35%, 4.35%, and 3.35% of the daily iron requirement of 15-year-old girls, 16-year-old girls, 15-year-old boys, and 16-year-old boys, respectively (Table 4).

3.11. Rice. The serving size of rice was represented by one teacup [12], and the average iron content which can be obtained through these three rice varieties was 3.03 ± 1.88 mg per serving (Table 4). Based on the USDA guidelines, the average estimated daily intake per serving contributes to 20.19% of the recommended daily allowance of girls and 27.53% of the recommended daily allowance of boys. According to the SL guidelines, it contributes to 8.41%, 7.76%, 10.09%, and 7.76% of the recommended daily allowances of 15-year-old girls, 16-year-old girls, 15-year-old boys, and 16-year-old boys, respectively (Table 4).

A considerable amount of iron can be obtained through the selected foods which were cooked according to the domestic/local recipes in Sri Lanka. The amount of iron in these cooked food samples was higher than that of raw foods. To prepare these food items according to the domestic/local recipes, several spices and other condiments must be added to enhance the taste. The chili powder (iron: 174 mg/kg [28] and 1.07 mg/1 tbs [29]), curry powder (iron: 190 mg/kg [28] and 1.86 mg/1 tbs [29]), turmeric powder (iron: 550 mg/kg [28] and 2.82 mg/1 tbs [29]), salt (iron: 0.06 mg/1 tbs [29]), black paper (iron: 96 mg/kg [28], 85.73 mg/kg [21] and 1.85/1 tbs [29]), cinnamon (iron: 0.65 mg/1 tbs [29]), fenugreek seed (iron: 3.72 mg/1 tbs [29]), onion (iron: 4.3 mg/kg [17], 3 mg/kg [18], 3 mg/kg [19], and 958 mg/kg [30]), curry leaves, green chili (iron: 9.79 mg/kg [21]), and garlic (iron: 58 mg/kg [28] and 0.23 mg/1 tbs [29]) are usually added as spices/condiments to prepare the meals in domestic/local recipes. Even though the fractional contribution of spices and sourcing agents was negligible, they provide a significant contribution to the total iron content because of the high iron compaction [21]. In some studies, contribution to dietary iron by spices and condiments was found to be 56%, and it provides the highest contribution to the total iron content in cooked food [21].

The total iron content of each cooked food sample is governed by the preparation method (Table 2) as well as the iron content of spices and other condiments, iron in coconut

milk [5], source of used water (groundwater 0.5–22.5 mg/L [7] and tap water 2.6×10^{-1} mg/kg [21]), utensils (knives and pans [5, 18]), soil particles [3, 18], and processing machinery [18]. Coconut in different forms (scraped kernel, milk, and oil) was commonly used in most preparations and considered as an integral part of Sri Lankan dishes [23]. The iron content of coconut kernel was 29.74 mg/kg [31], while Santos *et al* (2014) [32] reported the coconut milk has iron content ranging from 0.76 mg/kg to 5.52 mg/kg and provided a considerable contribution to the iron content in cooked foods such as curries and green leaves (malluma).

The total average EDI per serving of rice, vegetables, pulses, green leaves, and fish was 6.39 ± 1.06 mg; based on the USDA guidelines, it contributes to 42.62% and 58.12% of the RDA of girls and boys, respectively. According to the SL guidelines, the same iron content contributed to 17.76% and 16.39% of the RDA of girls in the age group 15 and 16 years and 21.31% and 16.39% of the RDA of boys in the same age group. Usually, they consumed more than one serving of them. Therefore, the iron contribution to the RDA of boys and girls aged 15-16 years was fulfilled by these cooked food items. A similar study was carried out by Hettiarachchi *et al.* [33] to assess the daily dietary intake, and they found that the mean iron intakes were 11.5 ± 1.0 mg/day and 11.4 ± 1.0 mg/day among boys and girls, respectively.

Even though the dietary iron of these cooked foods contributed to the daily iron intake, the total amount of iron consumed may not be absorbed into the body. The measured total iron content of this study is referred to as the sum of all the freely available ferric (Fe^{3+}) and ferrous (Fe^{2+}) together with the iron complexes available in the food sample. At the wavelength of 248.32 nm, all the Fe^{2+} , Fe^{3+} (nonheme iron), and iron in complexes (heme iron) in the food sample were measured (using the given procedure for iron analysis in this study, all the Fe^{2+} ions were converted to Fe^{3+} by HNO_3 and H_2O_2) [34].

However, the total iron content may not be absorbed by the body; thus, freely available ferrous ions and heme iron are the more bioavailable forms compared with ferric ions in the nonheme iron group [34]. Meanwhile, the absorption of iron from the meal depends on several factors such as the individual's iron status and requirement [35, 36], nature of the sources of iron [37], total iron content within the meal [37], amount of iron released, and the other associated constituents of the meal [38]. Iron deficiency results when the amount of dietary iron absorbed is insufficient to meet the daily iron requirement of an individual.

4. Conclusions

The meals among the selected study population mainly consisted of rice-based meals with vegetable accompaniments. Lentil curry is the best source of iron among the tested foods. Pumpkin curry has the highest iron content among selected vegetable curries, and centella (gotukola sambol) also contains high iron compaction. Fried marinated sardinella (salaya) provides the highest iron content among selected cooked fish varieties. Red raw rice (rathu kekulu) is also a good source of iron than parboiled small

grain rice (samba) and white raw rice (sudu kekulu) varieties. Scraped coconut kernel sambol is used as a popular accompaniment, and it also has considerable iron compaction. The cooked fish category is the best source of iron than the rest.

These cooked food types which are prepared in local/domestic recipes have higher iron content than raw foods, and they can fulfill the considerable proportion of the daily iron requirement of girls and boys in the 15-16 age category, even by a single serving.

Data Availability

The raw data can be received from the corresponding author upon request.

Conflicts of Interest

The authors declare that they have no conflicts of interest.





Acknowledgments

The study was made possible with the support of the Department of Nutrition, Medical Research Institute, Sri Lanka, and the Department of Botany and Instrument Centre at the University of Sri Jayewardenepura, Sri Lanka.

References

- [1] Food and Agricultural Organization of the United Nations/World Health Organization, *Requirements of Vitamin A, Iron, Folate and Vitamin B12: Report of a Joint of FAO/WHO Expert Consultation*, Food and Agriculture Organization, Rome, Italy, 1988.
- [2] V. G. Shobhana, N. Senthil, K. Kalpana et al., "Comparative studies on the iron and zinc contents estimation using atomic absorption spectrophotometer and grain staining techniques (prussian blue and DTZ) in maize germplasms," *Journal of Plant Nutrition*, vol. 36, no. 2, pp. 329–342, 2013.
- [3] S. A. Deraniyagala, K. T. Mudalige, H. M. Anula, and M. Y. P. R. Jamburuthugoda, "Some micronutrient and macronutrient contents of some Sri Lanka food varieties," *Vidyodaya Journal of Science*, vol. 5, no. 1, pp. 93–102, 1994.
- [4] G. Chathuranga, T. Balasuriya, and R. Perera, "Anaemia among female undergraduates residing in the hostels of University of Sri Jayewardenepura," *Anemia*, vol. 2014, Article ID 526308, 5 pages, 2014.
- [5] S. A. Deraniyagala and P. S. Gunawardena, "Effects of cooking on the mineral and oxalate content of some legumes," *Vidyodaya Journal of Science*, vol. 8, pp. 39–49, 1999.
- [6] M. E. Norhaizan and A. W. Nor Faizadatul Ain, "Determination of phytate, iron, zinc, calcium contents and their molar ratios in commonly consumed raw and prepared food in Malaysia," *Malaysian Journal of Nutrition*, vol. 15, no. 2, 2009.
- [7] A. Rajbhandari and T. Subedi, "Spectrophotometric determination of total iron in rice and maize samples," *Scientific World*, vol. 11, no. 11, pp. 101–104, 2013.
- [8] N. S. Lawal, N. Tajuddeen, and B. B. Garba, "Assessment of some mineral elements in different brands of powdered milk sold in Samaru Zaria, Nigeria," *International Food Research Journal*, vol. 22, no. 6, 2015.
- [9] *Survey Report—Assessment of Anaemia Status in Sri Lanka*, Medical Research Institute of the Health Sciences, Ministry of Health, Nutrition and Welfare, Sri Lanka, 2001.
- [10] T. Pedron, F. R. Segura, F. F. da Silva, A. L. de Souza, H. F. Maltez, and B. L. Batista, "Essential and non-essential elements in Brazilian infant food and other rice-based products frequently consumed by children and celiac population," *Journal of Food Composition and Analysis*, vol. 49, pp. 78–86, 2016.
- [11] E. Pinto, A. Almeida, and I. M. Ferreira, "Essential and non-essential/toxic elements in rice available in the Portuguese and Spanish markets," *Journal of Food Composition and Analysis*, vol. 48, pp. 81–87, 2016.
- [12] Nutrition Division Ministry of Health, *Food Based Dietary Guidelines for Sri Lankans*, Nutrition Division, Ministry of Health, Sri Lanka, 2nd edition, 2011.
- [13] E. Damastuti, W. Y. N. Syahfitri, M. Santoso, and D. D. Lestiani, "Assessment of trace element daily intake based on consumption rate of foodstuffs in Bandung City," *Atom Indonesia*, vol. 38, no. 1, pp. 29–34, 2012.
- [14] United States Department of Health and Human Services, *Iron—Fact Sheet for Health Professionals*, National Institute of Health, U.S., Department of Health and Human Services, Washington, DC, USA, 2019, <https://ods.od.nih.gov/factsheets/Iron-HealthProfessional/#en5>.
- [15] Medical Research Institute, *Recommended Daily Allowances for Sri Lankans*, Department of Nutrition, Medical Research Institute, Colombo, Sri Lanka, 2007.
- [16] W. Y. N. Syahfitri, E. Damastuti, N. Adventini et al., "Essential minerals of rice in West Java Indonesia and its daily intake estimation," *Atom Indonesia*, vol. 44, no. 3, pp. 155–163, 2018.
- [17] S. F. A. Bruggraber, T. P. E. Chapman, C. W. Thane, A. Olson, R. Jugdaohsingh, and J. J. Powell, "A re-analysis of the iron content of plant-based foods in the United Kingdom," *British Journal of Nutrition*, vol. 108, no. 12, pp. 2221–2228, 2012.
- [18] Food Standards Agency, *McCance and Widdowson's the Composition of Foods*, Royal Society of Chemistry, Cambridge, UK, 6th edition, 2002.
- [19] R. McCance and E. Widdowson, *The Chemical Composition of Foods*, His Majesty's Stationery Office, London, UK, 1940.
- [20] A. M. Mayer, "Historical changes in the mineral content of fruits and vegetables," *British Food Journal*, vol. 99, no. 6, pp. 207–211, 1997.
- [21] R. Narain and V. Ilango, "Analysis of iron content of selected vegetarian food items in Dubai, UAE," *International Journal of Science, Environment and Technology*, vol. 4, pp. 2278–3687, 2015.
- [22] J. Gooneratne and S. C. Kumarapperuma, "In-vitro dialysability of iron in green leafy vegetables and seasonal variation of total iron content," *Journal of the National Science Foundation of Sri Lanka*, vol. 35, no. 1, p. 9, 2011.
- [23] A. M. Karunaratne, P. H. Amerasinghe, V. M. Sadagopa Ramanujam, H. H. Sandstead, and P. A. J. Perera, "Zinc, iron and phytic acid levels of some popular foods consumed by rural children in Sri Lanka," *Journal of Food Composition and Analysis*, vol. 21, no. 6, pp. 481–488, 2008.
- [24] M. Kumari, S. Gupta, A. J. Lakshmi, and J. Prakash, "Iron bioavailability in green leafy vegetables cooked in different utensils," *Food Chemistry*, vol. 86, no. 2, pp. 217–222, 2004.
- [25] T. Athoney Swamy, N. L. Kerubo, J. K. Obey, and M. Hillary, "Determination of iron content in selected indigenous green leafy vegetables in Baraton, Kenya," *International Journal of Bioassays*, vol. 5, no. 4, pp. 4500–4504, 2016.
- [26] I. Wickramasinghe, K. K. D. S. Ranaweera, U. G. Chandrika, and A. Bamunaurachchi, "Content and in vitro availability of

- iron in cooked fish commonly consumed Sri Lanka,” *Journal of the National Science Foundation of Sri Lanka*, vol. 36, no. 4, 2008.
- [27] J. Liang, “Iron, zinc and phytic acid in rice from China: wet and dry processing towards improved mineral bioavailability,” Doctoral Thesis, Wageningen University, Wageningen, Netherlands, 2007, https://www.researchgate.net/publication/40105304_Iron_zinc_and_phytic_acid_in_rice_from_China_wet_and_dry_processing_towards_improved_mineral_bioavailability.
- [28] USDA National Nutrient Database, <https://ndb.nal.usda.gov/ndb/>.
- [29] Ferralet, Mission Pharmacal Company, <https://www.ferralet.com/>.
- [30] V. Singh and A. N. Garg, “Availability of essential trace elements in Indian cereals, vegetables and spices using INAA and the contribution of spices to daily dietary intake,” *Food Chemistry*, vol. 94, no. 1, pp. 81–89, 2006.
- [31] A. E. Akpakpa, U. M. Eduok, D. S. Udiog, I. E. Udo, and A. I. Ntukuyoh, “Levels of metals in Kernels and shells of oil palm and coconut fruits,” *International Journal of Modern Chemistry*, vol. 2, no. 1, pp. 20–27, 2012.
- [32] D. C. M. B. Santos, L. S. B. Carvalho, D. C. Lima, D. J. Leão, L. S. G. Teixeira, and M. G. A. Korn, “Determination of micronutrient minerals in coconut milk by ICP OES after ultrasound-assisted extraction procedure,” *Journal of Food Composition and Analysis*, vol. 34, no. 1, pp. 75–80, 2014.
- [33] M. Hettiarachchi, C. Liyanage, R. Wickremasinghe, D. Hilmers, and S. Abrams, “Nutrient intake and growth of adolescents in southern Sri Lanka,” *Ceylon Medical Journal*, vol. 51, no. 3, pp. 89–92, 2006.
- [34] P. Niedzielski, M. Zielinska-Dawidziak, L. Kozak et al., “Determination of iron species in samples of iron-fortified food,” *Food Analytical Methods*, vol. 7, no. 10, pp. 2023–2032, 2014.
- [35] J. D. Cook, “Adaptation in iron metabolism,” *The American Journal of Clinical Nutrition*, vol. 51, no. 2, pp. 301–308, 1990.
- [36] T. H. Bothwell, R. W. Charlton, J. D. Cook, and C. A. Finch, *Iron Absorption and Iron Metabolism in Man*, Blackwell Scientific Publications, London, UK, 1989.
- [37] H. C. Schonfeldt and N. G. Hall, “8th international food data conference,” *Journal of Food Composition and Analysis*, vol. 24, no. 4-5, pp. 469–756, 2011.
- [38] P. E. Adams, “Determining iron content in foods by spectrophotometry,” *Journal of Chemical Education*, vol. 72, no. 7, p. 649, 1995.



Hindawi

Submit your manuscripts at
www.hindawi.com

