

CORRECTION

Correction: Iron deficiency associates with deterioration in several symptoms independently from hemoglobin level among chronic hemodialysis patients

Shuta Motonishi, Kentaro Tanaka, Takashi Ozawa

[Fig 5](#) is incorrect. The authors have provided a corrected version here.



OPEN ACCESS

Citation: Motonishi S, Tanaka K, Ozawa T (2018) Correction: Iron deficiency associates with deterioration in several symptoms independently from hemoglobin level among chronic hemodialysis patients. PLoS ONE 13(9): e0204789. <https://doi.org/10.1371/journal.pone.0204789>

Published: September 24, 2018

Copyright: © 2018 Motonishi et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Fig 5. Relationship between TSAT and total score. The strength of linear association between TSAT and total symptom questionnaire score was analyzed by Spearman's correlation coefficient. TSAT: transferrin saturation.

<https://doi.org/10.1371/journal.pone.0204789.g001>

Reference

1. Motonishi S, Tanaka K, Ozawa T (2018) Iron deficiency associates with deterioration in several symptoms independently from hemoglobin level among chronic hemodialysis patients. *PLoS ONE* 13(8): e0201662. <https://doi.org/10.1371/journal.pone.0201662> PMID: 30071093